

Testing the Potential of Proposed DNA Barcoding Markers in *Nezara viridula* and *Nezara antennata* When Geographic Variation and Closely Related Species Were Considered

Authors: Li, Min, Liu, Qiang, Xi, Li, Liu, Yang, Zhu, Gengping, et al.

Source: Journal of Insect Science, 14(79) : 1-11

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.014.79>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



Testing the potential of proposed DNA barcoding markers in *Nezara viridula* and *Nezara antennata* when geographic variation and closely related species were considered

Min Li^{1,2}, Qiang Liu¹, Li Xi², Yang Liu², Gengping Zhu¹, Yanni Zhao¹, Wenjun Bu^{2a}

¹Tianjin Key Laboratory of Animal and Plant Resistance, Tianjin Normal University, 300387, Tianjin, China

²Institute of Entomology, College of Life Sciences, Nankai University, Tianjin, 300071, China

Abstract

The *COI* gene as the core of the DNA barcoding system for animals has received significant attention. The observed wide overlap between intra- and interspecific sequence variability has led researchers to envisage the primary *COI*-based method. The sequences of 16S rDNA, *COI*, and *Cyt b* genes of *Nezara viridula* (L.) (Hemiptera: Pentatomidae) from 13 countries and the same sequences of *N. antennata* Scott were chosen as molecular markers to analyze the intra- and interspecific relationships between the closely related species in this study. The results support that *Cyt b* gene may be a good candidate alongside *COI*, when the combined factors of geographic variation and closely related species are taken into account.

Keywords: 16S rDNA, *COI*, *Cyt b*, molecular markers

Correspondence: ^a wenjunbu@nankai.edu.cn

Editor: Karl Gordon was editor of this paper.

Received: 27 May 2012 **Accepted:** 10 December 2013 **Published:** 30 May 2014

Copyright: This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

ISSN: 1536-2442 | Vol. 14, Number 79

Cite this paper as:

Li M, Liu Q, Xi L, Liu Y, Zhu G, Zhao Y, Bu W. 2014. Testing the potential of proposed DNA barcoding markers in *Nezara viridula* and *Nezara antennata* when geographic variation and closely related species were considered. *Journal of Insect Science* 14(79). Available online: <http://www.insectscience.org/14.79>

Introduction

DNA barcoding is designed to provide rapid, accurate, and automatable species identifications by using short, standardized gene regions as internal species tags (Hebert and Gregory 2005). It has become a hotspot problem in biological taxonomy and the focus of controversy. The *COI* gene as the core of the global bio-identification system for animals has suffered great disputations (Will and Rubinoff 2004; DeSalle 2005; Hurst and Jiggins 2005; Meier et al. 2006; Memon et al. 2006; Jansen et al. 2009; Sundberg et al. 2010; Yassin et al. 2010). The most important problem is the observed wide overlap between intra- and interspecific sequence variability using *COI* as the molecular marker (Meyer and Paulay 2005; Meier et al. 2006; Memon et al. 2006; Alexander et al. 2009; Jansen et al. 2009). The lack of resolving power of the *COI* sequence has led researchers to envisage the primary *COI*-based method. Hebert and Gregory (2005) concluded that though DNA barcoding does not assure complete taxonomic resolution using a single gene region, in many cases when it fails the results can still be resolved fully with additional genetic or other data. Other genes, such as 16S rDNA (Vences et al. 2004; Steinke et al. 2005; Kappner and Bieler 2006; Aliabadian et al. 2009) and *Cyt b* (Bradley and Baker 2001; Pfunder et al. 2004; DeSalle et al. 2005; Hajibabaei et al. 2007), have been also advocated as standard or as complementary DNA barcoding markers.

When considering complementary barcoding markers, geographic variation and closely related species are factors that both need to be taken into account. Species with a wide geographic distribution often contain a great amount of genetic variability. This variability is not considered by only sampling individuals from a single site, so the distinctness of the

species barcode is easily underestimated (Moritz and Cicero 2004; Will and Rubinoff 2004; Prendini 2005). Conversely, sequence divergences would be overestimated if the closely related species between congeneric taxa were not included (Meyer and Paulay 2005). The DNA barcode database depends on the exhaustiveness of intra-taxon sampling and closely related species selecting. This point stresses a key challenge for the DNA barcoding initiative (Frezal and Leblois 2008).

We approached this issue by studying the southern green stink bug, *Nezara viridula* (L.) (Hemiptera: Pentatomidae) and the oriental green stink bug, *N. antennata* Scott, which are closely related species in morphology (the genus *Nezara* Amyot and Serville only includes three species in China: *N. viridula*, *N. antennata*, and *N. yunnana*). *Nezara viridula* is a polymorphic and cosmopolitan pentatomid pest that causes economic damage to many crop species (Panizzi 2000; Reid 2006; Knight and Gurr 2007; Martin et al. 2007). It is present throughout tropical and subtropical regions of Eurasia, Africa, Australia, and the Americas, located in the latitude between 45°N and 45°S, and is an active invading species (Hoffman et al. 1987; Panizzi 2000). There are a lot of studies on the population differentiation of *N. viridula* from different countries (Kiritani and Yukawa 1963; Bao-ying et al. 2000; Meglič et al. 2001; Sosa-Gómez et al. 2005; Kavar et al. 2006). Based on the current studies, the cryptic species from Botswana has been questioned (Kavar et al. 2006). *Nezara antennata* is distributed mainly in the oriental region and the southeastern edge of the Palaearctic region. It is very closely related to *N. viridula*, which had been considered as the synonym for *N. viridula* (Yang 1962), but studies based on interspecific copulation behavior (Kon et al. 1993), pheromones (Aldrich et al. 1993), and acous-

tical signals (songs) (Kon et al. 1998) between *N. antennata* and *N. viridula* showed that they are two distinct species.

We used the sequences of 16S rDNA, *COI*, and *Cyt b* of *N. viridula* drawn from 13 countries and the same sequences of *N. antennata* to analyze the intra- and interspecific relationships between these closely related species, seeking the proper molecular marker for DNA barcoding from these three genes.

Materials and Methods

Data for *N. viridula* from 11 different countries and regions (Slovenia, France, Greece, Italy, Madeira, Japan, Guadeloupe, Galapagos, California, Brazil, and Botswana) were downloaded from GenBank (www.ncbi.nlm.nih.gov/genbank). The *COI*, 16S rDNA, and *Cyt b* genes were amplified in 29 individual adult *N. viridula* from nine field collections (seven provinces in China: Guangxi, Hubei, Guangdong, Guizhou, Zhejiang, Hunan, Jiangxi; and southern and northern Iran). The same genes of *N. antennata* from individuals from two provinces (Guizhou and Zhejiang, China) were generated by PCR amplification. DNA was extracted from dissected thoracic muscles following the cetyltrimethylammonium bromide protocol (CTAB-based extraction protocol) (Doyle and Doyle 1987; Cullings 1992). The primer sequences are shown in Table 1. Amplification reactions were performed in a 25 μ L volume. Each PCR contained 2.5 μ L of 10 \times PCR buffer, 112.5 μ M MgCl₂, 10 μ M of each dNTP, 2.5 unit of Taq polymerase (Takara, www.takara-bio.com), 10 pM of each primer, 12.5 μ L of distilled water, and 2–3 μ L of DNA template. The PCR thermal regime consisted 5 min initial denaturation at 94°C, followed by 35 cycles of 45 sec denaturation at 94°C, 30 sec annealing at 60°C (16S) or 50°C (*COI*, *Cyt b*),

2 min extension at 72°C, and finalized by 10 min at 72°C. Each PCR product was subsequently purified using the gel extraction kit (Biospin, www.bioer.com.cn) and sequenced on an ABI PRISM 3730 automated sequencer (by Sunbio Company, www.sunbio.com). The voucher specimens were deposited in the Institute of Entomology, College of Life Sciences, Nankai University, Tianjin, China. The details of the taxa and mtDNA sequences informations are shown in Table 2.

Multiple sequences alignments were performed with Clustal W (Thompson et al. 1994). The sequences were compared among them, and only those showing different haplotypes were included in the analysis. Mesquite version 2.74 (Maddison and Maddison, 2010) was used to differentiate haplotypes of the sequences. The distinct sequences obtained in this study were submitted to GenBank, and the accession numbers are provided in Table 2. The intraspecific genetic distances of each of three genes (16S rDNA, *COI* and *Cyt b*) of *N. viridula* and the interspecific distance between *N. viridula* and *N. antennata* were calculated by Taxon DNA 1.0 (Meier et al. 2006). Because the cryptic species from Botswana has been questioned by some recent studies (Kavar et al. 2006), the inter- and intraspecific distances were re-calculated after removing the sequence from Botswana.

To construct trees, we used the neighbour joining (Saitou and Nei 1987) (Kimura 2-parameter model (K2P) for nucleotide, as recommended by Barrett and Hebert (2005)) and maximum parsimony (Swofford and Begle, 1993) methods. The methods were all performed by PAUP 4.0b for Windows (Swofford 2003). Unweighted parsimony analyses of various datasets were performed. Bootstrap values were generated in PAUP from 1000 replicates, each with 10 random-

addition heuristic searches. Sequences of two other pentatomid bugs, *Piezodorus lituratus* (F.) and *Rhaphigaster nebulosa* (Poda), were used as out-groups. As the compared sequence lengths were different from each other, it may have affected the results. We carried out separate analyses for all sequences with equal overlap. All trees were also re-constructed after removing the sequence from Botswana.

Results

The worldwide intraspecific genetic distances of each of the three genes (16S rDNA 448 bp, *COI* 347 bp, and *Cyt b* 460 bp) of *N. viridula* from Europe, Asia, Africa, and the Americas and the interspecies distance between *N. viridula* and *N. antennata* are shown in Table 3, which also contains the results after the sequence from Botswana was removed. The interspecies distances are equal or even shorter than the intraspecies distances based on each data set of three genes. The sequence differences between Botswana and other haplotypes of *N. viridula* are larger than those between *N. viridula* (without Botswana) and *N. antennata*. The overlap still existed in the results of the 16S rDNA data set even when the sequence from Botswana was removed. The ranges of the overlaps were wide (0.01–0.03), with almost 90% of intraspecific distances falling into this interval. The inter- and intraspecies distances of sequences with equal length (347 bp) are shown in Table 4 with similar results of different lengths.

According to the tree reconstruction methods, the sequences of *N. viridula* and *N. antennata* fail to form species-specific clusters (Figure 1). When the sequence from Botswana (NV-11) was removed, the failures still existed in the 16S rDNA data set, but *COI* and *Cyt b* are considered successfully identified, as they clustered with conspecific sequences (Figure




Figure 1. The phylogenetic tree based on three separate markers, *COI*, *Cyt b*, and 16S rDNA. The algorithms of neighbor joining and maximum parsimony (MP) methods were calculated by bootstrap resampling with 1,000 replicates. Bootstrap supports (NJ/MP) are given at the node (Symbols: plus: *COI*; circle: *Cyt b*; rhomb: 16S rDNA). Each data set of *COI*, *Cyt b*, or 16S rDNA (NJ and MP) produced completely resolved trees with similar topologies. High quality figures are available online.




Figure 2. The phylogenetic tree based on separate markers *COI* and *Cyt b*. The sequence of Botswana (NV-11) was excluded. The algorithms of neighbor-joining (NJ) and maximum parsimony (MP) methods were calculated by bootstrap resampling with 1,000 replicates. Bootstrap supports (NJ/MP) are given at the node (Symbols: plus: *COI*; circle: *Cyt b*). Each data set of *COI* or *Cyt b* (NJ and MP) produced completely resolved trees with similar topologies. High quality figures are available online.

2). The trees with a sequence of equal length (347 bp) have similar results with the different lengths data matrix.

Discussion

Both the distance- and tree-based results suggest that the specimen from Botswana may represent a distinct species. We found that the interspecies distances were equal or even shorter than the intraspecies distances based on each data set of three genes in the worldwide populations of *N. viridula* (Tables 3 and 4). The differences between sequences of *N. viridula* (without Botswana) and *N. antennata* were even smaller than those between Botswana and other haplotypes of *N. viridula*. Differences also could be found in the sexual communication system of *N. viridula*, which suggests that a cryptic species might exist (Ryan et al. 1996; Jeraj and Walter 1998). Further sampling in Africa will be necessary, and its reproductive isolation needs to be observed in order to ascertain its taxonomic status.

We found that the results changed when the Botswana sequence was removed (Tables 3 and 4). But, the overlap and failures also existed in 16S rDNA, making it difficult to identify candidate species. The proper marker should show small sequence divergences between intraspecifics and larger distance between interspecifics, and form species-specific clusters. So, 16S rDNA may not be a proper barcoding marker in this group.

In the populations of *N. viridula* (Botswana specimen (Nv-11) removed), *COI* and *Cyt b* are considered successfully identified depending on tree reconstruction techniques, as they clustered with conspecific sequences. The threshold of intraspecific variabilities of *COI* was 5%, and for *Cyt b* it was 3%. Constrained

intraspecific variation is a key finding in the DNA barcode effort. Traditionally, *COI* was considered to have far less variation within species (Hebert and Gregory 2005). In this study we found that the variation of *Cyt b* (0.03) within the species was smaller than *COI* (0.05). So, *Cyt b* may be a good candidate as a DNA barcoding marker along side *COI* in this group.

Hebert (2003) suggested that the thresholds of *COI* for species diagnosis are ordinarily greater than 3%. However, determining the thresholds that distinguish species in other geographical regions and taxonomic groups is important. Thresholds will particularly need to be established for groups with differences in traits, such as dispersal regime or generation length, which are likely to change rates of molecular evolution or the extent of a population subdivision. What is the boundary between a population and a species? Does it exist? To solve this issue, broad-ranging intraspecific sampling should be integrated in the database, and one must consider species boundaries not as a definitive but as a revisable concept. If the geographical structure is ignored, the species delineation will be blurred and distorted.

DNA barcoding using a single gene region does not assure complete taxonomic resolution. We suggest that a number of mitochondrial and nuclear genes may be used as DNA barcoding markers to complement *COI*. *Cyt b* is a good candidate in this group.

Acknowledgments

This project was supported by the Natural Science Foundation of China (No. 31372240, No. J1210005, No. 31201730); Tianjin City High School Science & Technology Fund Planning Project (No. 20110602), Talent introduction program in Tianjin Normal

University (5RL127) and The Doctor Foundation of Tianjin Normal University (No. 52XB1003).

References

- Aldrich JR, Numata H, Borges M, Bin F, Waite GK, Lusby WR. 1993. Artifacts and pheromone blends from *Nezara* spp. and other stink bugs (Heteroptera : Pentatomidae). *Zeitschrift für Naturforschung (Ser: C)* 48: 73–79.
- Alexander LC, Delion M, Hawthorne DJ, Lamp WO, Funk DH. 2009. Mitochondrial lineages and DNA barcoding of closely related species in the mayfly genus *Ephemerella* (Ephemeroptera:Ephemerellidae). *Journal of the North American Benthological Society* 28: 584–595.
- Aliabadian M, Kaboli M, Nijman V, Vences M. 2009. Molecular Identification of Birds: Performance of Distance-Based DNA Barcoding in Three Genes to Delimit Parapatric Species. *PLoS ONE* 4(1): e4119. doi: 10.1371/journal.pone.0004119
- Bao-ying QI, Walter GH, O'Toole AH. 2000. Structural incompatible – an investigation on microstructures of genitalia of *Nezara viridula* (Heteroptera: Pentatomidae) from Australia and Slovenia. *Journal of Inner Mongolia Normal University* 29: 48–54.
- Bradley R, Baker R. 2001. A test of the genetic species concept: cytochrome-b sequences and mammals. *Journal of Mammalogy* 82: 960–973.
- Cullings KW. 1992. Design and testing of a plant-specific PCR primer for ecological and evolutionary studies. *Molecular Ecology* 1: 233–240.
- DeSalle R. 2005. Conservation genetics – Genetics at the brink of extinction. *Heredity* 94: 386–387.
- DeSalle R, Egan MG, Siddall M. 2005. The unholy trinity: taxonomy, species delimitation and DNA barcoding. *Philosophical Transactions Of The Royal Society B-Biological Sciences* 360: 1905–1916.
- Doyle JJ, Doyle JL. 1987. A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochemistry Bulletin* 19: 11–15.
- Frezal L, Leblois R. 2008. Four years of DNA barcoding: Current advances and prospects. *Infection Genetics And Evolution* 8: 727–736.
- Hajibabaei M, Singer GA, Clare EL, Hebert PD. 2007. Design and applicability of DNA arrays and DNA barcodes in biodiversity monitoring. *BMC Biology* 5: 24.
- Hebert PDN, Cywinska A, Ball SL, DeWaard JR. 2003. Biological identifications through DNA barcodes. *Proceedings of the Royal Society of London Series B-Biological Sciences* 270: 313–321.
- Hebert PDN, Gregory TR. 2005. The promise of DNA barcoding for taxonomy. *Systematic Biology* 54: 841–844.
- Hoffman MP, Wilson LT, Zalom FG. 1987. Control of stink bugs in tomatoes. *California Agriculture* 41: 4–6.
- Hua JM, Li M, Dong PZ, Cui Y, Xie Q, Bu WJ. 2008. Comparative and phylogenomic studies on the mitochondrial genomes of

Pentatomomorpha (Insecta: Hemiptera: Heteroptera). *BMC Genomics* 9: 610.

Hurst GDD, Jiggins FM. 2005. Problems with mitochondrial DNA as a marker in population, phylogeographic and phylogenetic studies: the effects of inherited symbionts. *Proceedings of the Royal Society Biological Sciences Series B – Biological Sciences* 272: 1525–1534.

Jansen G, Savolainen R, Vepsäläinen K. 2009. DNA barcoding as a heuristic tool for classifying undescribed Nearctic *Myrmica* ants (Hymenoptera: Formicidae). *Zoologica Scripta* 38: 527–536.

Jeraj M, Walter GH. 1998. Vibrational communication in *Nezara viridula*: response of Slovenian and Australian bugs to one another. *Behavioural Processes* 44: 51–58.

Kappner I, Bieler R. 2006. Phylogeny of venus clams (Bivalvia: Venerinae) as inferred from nuclear and mitochondrial gene sequences. *Molecular Phylogenetics And Evolution* 40: 317–331.

Kavar T, Pavlovic P, Susnik S, Meglic V, Virant-Doberlet M. 2006. Genetic differentiation of geographically separated populations of the southern green stink bug *Nezara viridula* (Hemiptera: Pentatomidae). *Bulletin of Entomological Research* 96: 117–128.

Kiritani K, Yukawa J. 1963. A note on the polymorphism of *N. viridula* and *N. antennata*. *Rostrum* 5: 19–21.

Knight KMM, Gurr GM. 2007. Review of *Nezara viridula* (L.) management strategies and potential for IPM in field crops with emphasis on Australia. *Crop Protection* 26: 1–10.

Kon M, Oe A, Numata H, Hidaka T. 1988. Comparison of the mating behaviour between two sympatric species, *Nezara antennata* and *N. viridula* (Heteroptera: Pentatomidae), with special reference to sound emission. *Journal of Ethology* 6: 91–98.

Kon M, Oe A, Numata H. 1993. Intra- and interspecific copulations in the two congeneric green stink bugs, *Nezara antennata* and *N. viridula* (Heteroptera, Pentatomidae), with references to postcopulatory changes in the spermatheca. *Journal of Ethology* 11: 83–89.

Maddison WP, Maddison DR. 2010 Mesquite: a modular system for evolutionary analysis. Version 2.72. Available online: www.mesquiteproject.org

Martin PAW, Hirose E, Aldrich JR. 2007. Toxicity of chromobacterium subtsugae to southern green stink bug (Heteroptera: Pentatomidae) and corn rootworm (Coleoptera: Chrysomelidae). *Journal of Economic Entomology* 100: 680–684.

Meier R, Shiyang K, Vaidya G, Ng PKL. 2006. DNA barcoding and taxonomy in Diptera: a tale of high intraspecific variability and low identification success. *Systems Biology* 55: 715–728.

Meglić V, Virant-Doberlet M, Šuštar-Vozlič J, Sušnik S, Čokl A, Mikla N, Renou M. 2001. Diversity of the southern green stink bug *Nezara viridula* (L.) (Heteroptera: Pentatomidae). *Journal of Central European Agriculture* 2: 241–250.

Memon N, Meier R, Manan A, Su KFY. 2006. On the use of DNA sequences for determining the species limits of a polymorphic new species in the stink bug genus *Halys*

- (Heteroptera: Pentatomidae) from Pakistan. *Systematic Entomology* 31: 703–710.
- Meyer CP, Paulay G. 2005. DNA barcoding: Error rates based on comprehensive sampling. *PLoS Biol* 3(12): e422. doi: 10.1371/journal.pbio.0030422
- Moritz C, Cicero C. 2004. DNA barcoding: Promise and pitfalls. *PLoS Biol* 2(10): e354. doi: 10.1371/journal.pbio.0020354
- Muraji M, Tachikawa S. 2000. Phylogenetic analysis of water striders (Hemiptera: Gerroidea) based on partial sequences of mitochondrial and nuclear ribosomal RNA genes. *Entomological Science* 3: 615–626.
- Muraji M, Kawasaki K, Shimizu T. 2000. Phylogenetic utility of nucleotide sequences of mitochondrial 16S ribosomal RNA and cytochrome b genes in anthocorid bugs (Heteroptera: Anthocoridae). *Applied Entomology and Zoology* 35: 293–300.
- Panizzi AR. 2000. Suboptimal nutrition and feeding behavior of hemipterans on less preferred plant food sources. *Anais da Sociedade Entomologica do Brasil* 29: 1–12.
- Pfunder M, Holzgang O, Frey J. 2004. Development of microarray-based diagnostics of voles and shrews for use in biodiversity monitoring studies, and evaluation of mitochondrial cytochrome oxidase I vs. cytochrome b as genetic markers. *Molecular Ecology* 13: 1277–1286.
- Prendini L. 2005. Comment on "Identifying spiders through DNA barcodes". *Canadian Journal Of Zoology-Revue Canadienne De Zoologie* 83: 498–504.
- Reid S. 2006. A significant interception of the green vegetable bug, *Nezara viridula* (L.) (Hemiptera: Pentatomidae) in the UK. *Entomologist's Record and Journal of Variation* 118: 123–125.
- Ryan MA, Čokl A, Walter GH. 1996. Differences in vibratory sound communication between a Slovenian and an Australian population of *Nezara viridula* (L.) (Heteroptera: Pentatomidae). *Behavioural Processes* 36: 183–193.
- Sosa-Gómez DR, Silva JJ, Costa F, Binneck E, Marin SRR, Nepomuceno L. 2005. Population structure of the Brazilian southern green stink bug, *Nezara viridula*. *Journal of Insect Science* 5:23. Available online: www.insectscience.org/5.25
- Steinke D, Vences M, Salzburger W, Meyer A. 2005. TaxI: a software tool for DNA barcoding using distance methods. *Philosophical Transactions Of The Royal Society B - Biological Sciences* 360: 1975–1980.
- Sundberg P, Vodoti ET, Strand M. 2010. DNA barcoding should accompany taxonomy - the case of *Cerebratulus* spp (Nemertea). *Molecular Ecology Resources* 10: 274–281.
- Thompson J, Higgins D, Gibson T, et al. 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Research* 22: 4673–4680.
- Vences M, Thomas M, Van der Meijden A, Chiari Y, Vieites DR. 2004. Performance of 16S rRNA in DNA barcoding of amphibians.

Integrative And Comparative Biology 44: 657–657.

Will KW, Rubinoff D. 2004. Myth of the molecule: DNA barcodes for species cannot replace morphology for identification and classification. *Cladistics* 20: 47–55.

Will KW, Rubinoff D. 2004. Myth of the molecule: DNA barcodes for species cannot replace morphology for identification and classification. *Cladistics-The International Journal Of The Willi Hennig Society* 20: 47–55.

Yang WY. 1962. *Economic Entomography of China. Volume II. Hemiptera Pentatomidae*. pp. 126–128. Science Press.

Yassin A, Markow TA, Narechania A, O’Grady PM, DeSalle R. 2010. The genus *Drosophila* as a model for testing tree- and character-based methods of species identification using DNA barcoding. *Molecular Phylogenetics and Evolution* 57: 509–517.

Table 1. Primer used for sequence amplification.

Locus	P primer (5'-3')	Size (bp)	Reference
	R primer (5'-3')		
16S rDNA	CCGGTTTGAAGTCAGATCATGT	560	Muraji and Tachikawa 2000
	CGCCTGTTTAACAAAAACAT		
<i>Cyt b</i>	TAGGATATGTTTACCTTAGGACA	486	Muraji et al. 2000
	TCCTCCTAATTATTAGGAATTG		
<i>COI</i>	GGAACAGGATGAACAGTTTACCTCC	914	Hua et al. 2008
	TCTGAATATCGTCGAGGTATTC		

Table 2. The detailed information for the *Nezara viridula* and *N. antennata* specimens used in this study.
Note: The sequence accession numbers starting with EU and FJ are sequences from from this research.

Species	Composite Haplotypes	Collecting Locality	Haplotype/ Accession numbers		
			16S	COI	Cyt b
<i>Nezara viridula</i>	NV-1	Slovenia, Italy, Madeira, Brazil	NV-16S-1/	NV-COI-1/	NV-Cyt b-1/
			AY839153	AY839161	AY839171
	NV-2	Slovenia	NV-16S-2/	—	—
			AY839154		
	NV-3	France	NV-16S-1/	—	NV-Cyt b-2/
			AY839153		AY839172
	NV-4	Madeira	—	NV-COI-2/	NV-Cyt b-1/
				AY839162	AY839171
	NV-5	Brazil	—	NV-COI-3/	—
				AY839163	
	NV-6	Greece, Guadeloupe, Galapagos, California, Japan, Italy, Brazil	NV-16S-3/	NV-COI-1/	NV-Cyt b-3/
			AY839155	AY839161	AY839174
	NV-7	Japan	NV-16S-4/	—	NV-Cyt b-4/
			AY839156		AY839173
	NV-8	Brazil	NV-16S-1/	NV-COI-4/	NV-Cyt b-1/
			AY839153	AY839164	AY839171
	NV-9	Japan	NV-16S-5/	NV-COI-5/	NV-Cyt b-5/
			AY839157	AY839165	AY839170
	NV-10	Japan	—	NV-COI-6/	—
				AY839166	
	NV-11	Botswana	NV-16S-6/	NV-COI-7/	NV-Cyt b-6/
			AY839158	AY839167	AY839175
	NV-12	China, Guangxi province	16S-C1/	NV-COI-5/	Cyt b-C2/
			EU275199	AY839165	FJ418864
<i>Nezara antennata</i>	NV-13	China, Hubei province	—	COI-C1/	—
				FJ418856	
	NV-14	China, Guangdong province	—	COI-C3/	Cyt b-C1/
				FJ418858	FJ418863
	NV-15	China, Guizhou province	16S-C2/	COI-C2/	Cyt b-C2/
			FJ418869	FJ418857	FJ418864
	NV-16	China, Zhejiang province	16S-C1/	NV-COI-5/	NV-Cyt b-5/
			EU275199	AY839165	AY839170
	NV-17	China, Hunan province	—	COI-C3/	Cyt b-C2/
				FJ418858	FJ418864
	NV-18	China, Guangxi province	—	COI-C4/	NV-Cyt b-5/
				FJ418859	AY839170
<i>Nezara antennata</i>	NV-19	China, Guangdong province	—	NV-COI-5/	Cyt b-C1/
				AY839165	FJ418863
	NV-20	Iran, Northern part	NV-16S-1/	NV-COI-1/	Cyt b-IN/
			AY839153	AY839161	FJ418865
	NV-21	Iran, Northern part	—	—	Cyt b-IS/
					FJ418866
	NV-22	Iran, Southern part	16S-IS/	COI-IS/	—
			FJ418870	FJ418860	
	NA-1	China, Guizhou province	NA-16S/	NA-COI-1/	NA-Cyt b-1/
			FJ418871	FJ418861	FJ418867
<i>Nezara antennata</i>	NA-2	China, Zhejiang province	—	NA-COI-2/	NA-Cyt b-2/
				FJ418862	FJ418868

Table 3. The intra- and interspecifics distance (K2P) of each gene.

Sequence	Complete data set		Botswana removed	
	intraspecific	interspecific	intraspecific	interspecific
16S rDNA	0–0.05	0.01–0.05	0–0.03	0.01–0.05
<i>COI</i>	0–0.12	0.06–0.09	0–0.05	0.06–0.08
<i>Cyt b</i>	0–0.14	0.06–0.14	0–0.03	0.06–0.08

Table 4. The intra- and interspecifics distance (K2P) of 16S rDNA and *Cyt b* with 347 bp.

Sequence	Complete data set		Botswana removed	
	intraspecific	interspecific	intraspecific	interspecific
16S rDNA	0–0.05	0.01–0.05	0–0.04	0.01–0.06
<i>Cyt b</i>	0–0.14	0.06–0.14	0–0.03	0.06–0.08