

Subfamily Coleoscirinae (Acari: Trombidiformes: Cunaxidae), with Description of One New Species from Pakistan

Authors: Bashir, Muhammad Hamid, Afzal, Muhammad, Ashfaq, Muhammad, Ali, Shaukat, Kamran, Muhammad, et al.

Source: Journal of Insect Science, 14(82) : 1-14

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.014.82>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



Subfamily Coleoscirinae (Acari: Trombidiformes: Cunaxidae), with description of one new species from Pakistan

Muhammad Hamid Bashir^{1a*}, Muhammad Afzal^{2b}, Muhammad Ashfaq^{1c}, Shaukat Ali^{3d}, Muhammad Kamran^{2e}, and Sabyan Faris Honey^{1f}

¹Department of Agri. Entomology, University of Agriculture, Faisalabad, Pakistan

²University College of Agriculture, University of Sargodha, Sargodha, Pakistan

³College of Natural Resource and Environment, South China Agricultural University, China

Abstract

The Coleoscirinae (Acari: Trombidiformes: Cunaxidae) from Pakistan are summarized in this paper. Two species of *Scutascirus* Den Heyer (*S. pirus* Chaudhri and Akbar and *S. tactus* Chaudhri and Akbar), ten species of *Coleoscirus* Berlese (*C. bapto*s (Chaudhri and Akbar), *C. carex* (Inayatullah and Shahid), *C. carnus* Muhammad and Chaudhri, *C. comis* Muhammad and Chaudhri, *C. disparis* Muhammad and Chaudhri, *C. irroratus* Muhammad and Chaudhri, *C. mardi* (Inatullah and Shahid), *C. raviensis* Afzal, Ashfaq and Khan, *C. tobaensis* Bashir, Afzal, Ashfaq, and Khan, and *C. trudus* Bashir, Afzal and Akbar), and three species of *Pseudobonzia* Smiley (*P. ashfaqi* Bashir, Afzal and Akbar, *P. numida* Chaudhri and Akbar, and *P. parilus* Chaudhri) have been previously reported. One new species of *Pseudobonzia*, *Pseudobonzia bakeri* sp. n., is herein described and illustrated. A key to the genera of the subfamily and keys to the species in each genus are given to incorporate the new species from Pakistan. Distribution records of all known species in Pakistan are also given.

Keywords: Neoscirulini, Prostigmata

Correspondence: ^a hamid_uaf@yahoo.com, ^b chafzal64@yahoo.com, ^c drashfaqi@yahoo.com, ^d aliscou@scau.edu.cn, ^e Kamran1513@gmail.com, ^f sabyanfarishoney@ymail.com, *Corresponding author

Editor: Takumasa Kondo was editor of this paper.

Received: 26 April 2012 **Accepted:** 12 March 2014 **Published:** 7 July 2014

Copyright: This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

ISSN: 1536-2442 | Vol. 14, Number 82

Cite this paper as:

Bashir MH, Afzal M, Ashfaq M, Ali S, Kamran M, Honey SF. 2014. Subfamily Coleoscirinae (Acari: Trombidiformes: Cunaxidae), with description of one new species from Pakistan. *Journal of Insect Science* 14(82). Available online: <http://www.insectscience.org/14.82>

Introduction

Cunaxidae (Acari: Trombidiformes) are important predatory mites. They have been reported to feed on other small arthropods, including harmful mites and soft bodied insects (Nesbitt 1946; Lord 1949; Schruft 1971; Kuznetsov and Sizova 1978; Youssef et al. 1980; Smiley 1992) and nematodes (Walter and Kaplan 1991). Cunaxidae occupy diverse habitats and have been reported from all parts of the world (Walter and Kaplan 1991; Swift 1996; Walter 1999; Arbabi et al. 2002; Tagore and Putatunda 2003; de-Oliveira and Daemon 2003; Tagore and Putatunda 2003; Sionti 2003; Sergeyenkov 2005, 2006, 2009, 2011; Mejia-Recamier and Palacios-Vargas 2007; Corpuz-Raros 2007; Kaluz 2009; Den Heyer 2011; Skvarla et al. 2011; Skvarla and Dowling 2012).

Coleosirinae was erected by Den Heyer (1979). It encompasses two tribes (Coleosirini and Neosirulini) and six genera (*Neoscirula* Den Heyer, *Pseudobonzia* Smiley, *Coleobonzia* Den Heyer & Castro, *Coleoscirus* Berlese, *Orangesirula* Bu & Li, and *Scutascirus* Den Heyer). Ninety-one species have thus far been described in this subfamily (Den Heyer 2011). *Neoscirula*, which Smiley (1992) transferred to Bonzinae based on characteristics of the hypostomal setae, has recently been placed back into Coleosirinae (Den Heyer 2011).

Berlese (1916) erected the genus *Coleoscirus* and included two species, *C. halacaroides* and *C. corniculatus*, in it. Den Heyer (1978) synonymized *Coleoscirus corniculatus* with *Scirus curtispalpis*. Berlese (1916) designated *Coleoscirus halacaroides* as the type species of genus *Coleoscirus*, in spite of an earlier described species, *Coleoscirus curtispalpis* (Berlese 1888). Ewing (1917) described a new

species *Scirus* (= *Coleoscirus*) *simplex* from the U.S.A.

Thor and Willmann (1941) redescribed and provided drawings of *Coleoscirus simplex*, *C. brevicornis*, *C. curtispalpis*, *C. halacaroides*, and *C. curtispalpis* (as *C. corniculatus*). Baker and Hoffmann (1948) redescribed and provided the drawings of *C. curtispalpis*, *C. brevicornis*, and *C. simplex*. They also described and gave drawings of a new species, *Cunaxa mexicana* (= *Coleoscirus mexicanus* Baker & Hoffmann, 1948) from Mexico and the U.S.A.

Smiley (1975) provided a new genus name, *Pseudocunaxa*, for species related to and including *Coleoscirus simplex*. This new genus name is a synonym of *Coleoscirus* Berlese, 1916 (Den Heyer 1980a). Den Heyer (1979) described two new species, *C. tuberculatus* and *C. magdalenae*, from Africa and gave a key for these two species. Den Heyer (1980a) described three new species, *Coleoscirus coatesi*, *C. buartus*, and *C. breslauensis*, and prepared a comprehensive key. Den Heyer (1980b) included the genus *Coleoscirus* in the subfamily Coleosirinae, in his new tribe Coleosirini, along with the genus *Scutascirus*. He mentioned the genus *Pseudocunaxa* Smiley, 1975 as synonym of genus *Coleoscirus*. Tseng (1980) erected a new genus, *Lapicunaxa*, with *Lapicunaxa horidula* as its type species.

Den Heyer (1979) synonymized *Pseudocunaxa* with *Coleoscirus*. Chaudhri and Akbar (1985) and Inayatullah and Shahid (1993) apparently missed that publication or did not agree with it and described species in the genus *Pseudocunaxa*. In 1992, Smiley gave the new classification system for Cunaxidae family, as he synonymized *Pseudocunaxa* and *Lapicunaxa* with *Coleoscirus* and described

11 species of this genus, including the species of Chaudhri and Akbar (1985). Later, Muhammad and Chaudhri (1992a, b) and Bashir et al. (2006, 2008) contributed to the fauna of this genus from Pakistan.

The genus *Pseudobonzia* was erected by Smiley (1975). He designated *Cunaxa reticulata* Heryford as its type species. Den Heyer (1977) described six new species from the Ethiopian region and placed this genus in the subfamily Coleoscirinae (Den Heyer 1980b). Later, Den Heyer (1980a), Luxton (1982), Liang (1984), Sepasgosarian (1984), Chaudhri and Akbar (1985), Michocka (1987), Smiley (1992), and Bashir et al. (2008) made significant contributions to the fauna of this genus worldwide.

The genus *Scutascirus* was erected by Den Heyer (1976). He designated *Scutascirus polyscutosus* as its type species. Den Heyer (1979, 1980a), Sepasgosarian (1984), Chaudhri and Akbar (1985), Smiley (1992), and Lin et al. (2001) have contributed to the knowledge of Cunaxidae.

Materials and Methods

Sieve collection was used for field collecting. The plant parts, such as leaves, twigs, and inflorescences, were beaten on a sieve held over a white piece of paper. Cunaxids were sorted with the help of a magnifying lens and stored in vials containing 70% alcohol and few drops of glycerin. Materials such as soil and leaf debris that could not be processed in the field were processed in Berlese funnels for at least 24 hours. They were subsequently sorted under a binocular microscope and preserved in 70% ethanol. The specimens were mounted permanently on glass slides using Hoyer's medium and identified using a phase contrast microscope. Illustrations were prepared by

using an ocular grid. The identification of the species was done with the help of existing keys and literature. The setal nomenclature of Kethley (1990) has been adopted. All the measurements (in μm) and ranges are given in the description. The following abbreviations are used in this manuscript:

asl: attenuate solenidion

bsl: blunt ended solenidion

sts: simple tactile setae

T: trichobothrium

Peo: cunaxid peg organ on tarsi I

Nomenclature

This publication and the nomenclature it contains have been registered in Zoobank. The LSID number is:

urn:lsid:zoobank.org:pub:13C6436F-7557-439D-B4A1-A7C8F07D3530

It can be found online by inserting the LSID number after www.zoobank.org/

Results and Discussion

Key to genera of the subfamily Coleoscirinae (Smiley, 1992)

1) Ventral idiosoma with sub triangular plate adjacent to ventrolateral coxal and genital plates-----*Scutascirus* Den Heyer

- Ventral idiosoma without sub triangular plate adjacent to ventrolateral coxal and genital plates-----2

2) Female with sternal and ventrolateral plates. Dorsum with a single shield extending from propodosoma into hysterosomal region-----*Coleoscirus* Berlese

- Female without sternal and ventrolateral plates. Dorsum with a single shield confined to propodosomal region-----*Pseudobonzia* Smiley

Genus *Scutascirus***Key to species of the genus *Scutascirus* known from Pakistan**

1) Gnathosoma with striations at base, palp telofemur with 1 seta -----
----- *tactus* Chaudhri & Akbar

2) Gnathosoma with reticulations at base, palp telofemur with 1 seta and 1 spine-----
----- *pigrus* Chaudhri & Akbar

Scutascirus pigrus Chaudhri & Akbar
Scutascirus pigrus Chaudhri & Akbar, 1985:
231

Known distribution: Pakistan: Faisalabad
Known source: Rawan (*Vigna sinensis*)

Scutascirus tactus Chaudhri & Akbar
Scutascirus tactus Chaudhri & Akbar, 1985:
229

Known distribution: Pakistan: Gujranwala
Known source: Jute (*Corchorus capsularis*)

Genus *Coleoscirus***Key to species of the genus *Coleoscirus* known from Pakistan**

1) Dorsal shield with 4 pairs of setae in hysterosomal region -----2
- Dorsal shield with more than 4 pairs of setae in hysterosomal region-----4

2) Palp tibiotarsus with large spur on inner medial surface; genital shield with 4 pairs of simple setae -----
----- *trudus* Bashir & Afzal
- Palp tibiotarsus with small tubercle on inner medial surface; genital shield with 3 pairs of simple setae-----3

3) Genu IV with 5 setae; tibia I with 7 setae-----
----- *carex* (Inayatullah & Shahid)

- Genu IV with 6 setae; tibia I with 6 setae-----
----- *mardi* (Inayatullah & Shahid)

4) Dorsal shield with 6 pairs of simple setae in hysterosomal region -----5
- Dorsal shield with 5 pairs of simple setae in hysterosomal region -----7

5) Venter with 6 pairs of simple setae between lateral plates and distal part of body in addition to setae of anal and genital region-----
----- *raviensis* Bashir, Afzal, Ashfaq & Khan
- Venter with less than 6 pairs of simple setae between lateral plates and distal part of body in addition to setae of anal and genital region -----6

6) Leg genu IV with 6 setae; venter with 4 pairs of simple setae between lateral plates and genital region-----
----- *tobaensis* Bashir, Afzal, Ashfaq & Khan
- Leg genu IV with 7 setae; Venter with 5 pairs of simple setae between lateral plates and genital region-----
----- *carnus* Muhammad & Chaudhri

7) Coxa II with 3 setae-----8
- Coxa II with 2 setae-----
----- *baptos* (Chaudhri & Akbar)

8) Basifemur III with 4 setae-----9
- Basifemur III with 5 setae-----10

9) Genu I with 8 setae; telofemur I with 4 setae-----
----- *simplex* (Ewing)
- Genu I with 9 setae; telofemora I with 5 setae-----
----- *irroratus* Muhammad & Chaudhri

10) Para anal seta present-----
----- *comis* Muhammad & Chaudhri
- Para anal seta absent-----11

11) Genu I-IV with 8-7-6-6 setae-----
----- *kayfayati* (Inayatullah & Shahid)

- Genu I–IV with 9-8-6-7 setae-----
-----*disparis* Muhammad & Chaudhri

Coleoscirus bapto (Chaudhri & Akbar)
Pseudocunaxa bapto Chaudhri & Akbar,
1985: 223
Known distribution: Pakistan: Charrapani
(Murree)
Known source: Pine (*Pinus* spp.)

Coleoscirus carex (Inayatullah & Shahid)
Pseudocunaxa carex Inayatullah & Shahid,
1993: 318
Known distribution: Pakistan: Peshawar
Known source: Rotten leaves

Coleoscirus carnus Muhammad & Chaudhri
Coleoscirus carnus Muhammad & Chaudhri,
1992: 309
Known distribution: Pakistan: Peshawar
Known source: Pear (*Pyrus communis*)

Coleoscirus comis Muhammad & Chaudhri
Coleoscirus comis Muhammad & Chaudhri,
1992: 99
Known distribution: Pakistan: Faisalabad
Known source: Stored wheat

Coleoscirus disparis Muhammad & Chaudhri
Coleoscirus disparis Muhammad & Chaudhri,
1992: 310
Known distribution: Pakistan: Karachi
Known source: Stored rice (*Oryza sativa*)

Coleoscirus irroratus Muhammad &
Chaudhri
Coleoscirus irroratus Muhammad & Chaud-
hri, 1992: 99
Known distribution: Pakistan: Shahkot, Guj-
ranwala, Faisalabad
Known source: Wheat (*Triticum aestivum*),
sugarcane, debris

Coleoscirus kifayati (Inayatullah & Shahid)

Pseudocunaxa kifayati Inayatullah & Shahid,
1993: 315
Known distribution: Pakistan: Peshawar
Known source: Banana (*Musa paradisiaca*)

Coleoscirus mardi (Inatullah & Shahid)
Pseudocunaxa mardi Inatullah & Shahid,
1993: 316
Known distribution: Pakistan: Mingora (Swat)
Known source: Rice (*Oryza sativa*)

Coleoscirus raviensis Bashir, Afzal, Ashfaq
& Khan
Coleoscirus raviensis Bashir, Afzal, Ashfaq &
Khan, 2008: 453
Known distribution: Pakistan: Lahore,
Kasoor, Sialkot, Bahawalpur
Known source: Plant debris

Coleoscirus simplex Ewing
Scirus simplex Ewing, 1917: 150
Cunaxa simplex (Ewing), Thor & Willmann,
1941: 172; Baker & Hoffmann, 1948: 240;
Muma, 1960: 324; Shiba, 1978: 114
Pseudocunaxa simplex Smiley, 1975: 241;
Chaudhri, 1977:43; 1985:223; Inayatullah &
Shahid, 1993: 315
Coleoscirus simplex (Ewing), den Heyer,
1979c: 524; 1980d: 105; Sepasgosaran, 1984:
143
Known distribution: Pakistan: Faisalabad,
Multan, Khanewal, Layyah, Lodhran, Toba
Tek Singh, Lahore, Okara, Sahiwal, Gujran-
wala,
Known source: Plant debris, cucumber, toma-
to, brinjal, citrus

Coleoscirus tobaensis Bashir, Afzal, Ashfaq
& Khan
Coleoscirus tobaensis Bashir, Afzal, Ashfaq
& Khan, 2008: 455
Known distribution: Pakistan: Faisalabad,
T.T. Singh, Chakwal, Sialkot
Known source: Plant debris

Coleoscirus trusus Bashir, Afzal & Khan
Coleoscirus trusus Bashir, Afzal & Khan,
 2006: 74
 Known distribution: Pakistan: Faisalabad,
 Muzzafarghar, Toba Tek Singh, Kasoor, D.G.
 Khan
 Known source: Leaf debris

Genus *Pseudobonzia*

Key to species of genus *Pseudobonzia* known from Pakistan

- 1) Palp tibiotarsus with thick spine-like seta-----2
 - Palp tibiotarsus without thick spine-like seta-----*bakeri*, **sp. n.**
- 2) Ventral hysterosoma with 5 pairs simple setae between coxae II and distal part of the body excluding setae of anal and genital region-----*parilus* Chaudhri
 - Ventral hysterosoma with more than 5 pairs simple setae between coxae II and distal part of the body excluding setae of anal and genital region-----3
- 3) Venter with 7 pairs simple setae between coxae II and distal part of the body in addition to setae of anal and genital region; coxa IV with 2 setae-----
 -----*numida* Chaudhri & Akbar
 - Venter with 6 pairs simple setae between coxae II and distal part of the body in addition to setae of anal and genital region; coxa IV with 3 setae-----
 -----*ashfaqi* Bashir, Afzal & Akbar

Pseudobonzia ashfaqi Bashir, Afzal & Akbar
Pseudobonzia ashfaqi Bashir, Afzal and Akbar, 2008: 77
 Known distribution: Pakistan: Faisalabad
 Known source: Plant debris

Pseudobonzia numida Chaudhri & Akbar

Pseudobonzia numida Chaudhri & Akbar,
 1985: 220
 Known distribution: Pakistan: Hassanabdal
 Known source: Fungus

Pseudobonzia parilus Chaudhri
Pseudobonzia parilus Chaudhri, 1977: 45
 Known distribution: Pakistan: Sialkot
 Known source: Chili peppers (*Capsicum frutescens*)

Pseudobonzia bakeri, Bashir, Afzal, Ashfaq, Raza & Kamran, **sp. n.**
 (Figures 1–4)

Female

Gnathosoma. Gnathosoma 140 long and 80 wide. Hypostome sub rectangular, cone shaped distally; with 4 pairs hypognathal setae (hg₁–hg₄) (Figure 1A). Palp 5 segmented, measuring 110. Chaetotaxy of palp as follows: trochanter none; basifemur with one simple seta; telofemur with one simple seta; genu with 4 simple setae; tibiotarsus terminating in a claw, with 6 simple setae (Figure 1B).

Chelicerae 135 long, terminating in a claw, with one simple dorsomedial seta, dorsal and ventral sides with papillae (Figure 1C).

Dorsum. Body 370 long (without gnathosoma) and 270 wide. Propodosoma with a weakly sclerotized sub rectangular shield bearing randomly placed different sized papillae. Propodosomal shield with sensillae *vi* and *sce* measuring 100 and 115, respectively, and propodosomal setae *ve* measuring 17, *sci* measuring 11.

Hysterosoma separated from propodosoma by papillae bearing striae. Hysterosoma with setae *c1*, *c2*, *d1*, *e1*, *f1*, *f2*, *h1*, and *h2* measuring 11, 11, 11, 11, 13, 12, 17, and 17, respective-

ly. Hysterosoma with one pair of cupules, *im*, on integument anterior to setae *fl* (Figure 2).

Venter. Venter with papillate striations. Coxae I–II contiguous, connected by small lateral apodemes; coxae III–IV contiguous, broader than coxae I–II. Ventral hysterosoma with 1 pairs propodogastral simple setae and 7 pairs hystergastral setae in addition to setae of anal and genital region. Genital shield with two valves bearing papillae. Each valve with 4 genital setae (*g₁–g₄*) longitudinal aligned and 2 genital suckers. Two pairs of anal setae (*a*) and one pair of paranal setae (*pa*) present. One pair minute pores near anal shield (Figure 3).

Legs. All legs pitted with papillae and blunt ended. Legs I–IV measuring (from trochanter base to the tip of tarsus) 193, 185, 185, and 187 respectively. Chaetotaxy of legs I–IV as follows: Coxae 3-3-3-3 *sts*; trochanters 1-1-2-1 *sts*; basifemora 2-4-4-2 *sts*; telofemora 5-5-4-4 *sts*; genua 8 (3 *asl* + 5 *sts*)-8-6-5 *sts*; tibiae 7 (1 *asl* + 1 *bsl* + 5 *sts*)-6 (1 *bsl* + 5 *sts*)-6 (1 *bsl* + 5 *sts*)-5 (1 *T* + 4 *sts*) and tarsi 25 (3 *asl* + 1 *peo* + 2 *bsl* + 19 *sts*)-25 (1 *asl* + 24 *sts*)-18-16 *sts* (Figure 4).

Male

Unknown.

Type material. Holotype female, collected in Lahore from leaf debris on 28 August 2004 (Hamid) and deposited in the Acarology Research Laboratory, Department of Agri. Entomology, University of Agriculture, Pakistan.

Etymology. The species epithet is in reference to Dr. Edward W. Baker, Research Entomologist (Acarology), Systematic Entomology Laboratory, U.S. Department of Agriculture, Beltsville, Maryland, USA, for his outstanding contribution to the field of Acarology.

Remarks. This new species, *Pseudobonzia bakeri* **sp. n.**, is very similar to *P. summersi* Smiley but can be separated by the following characters: 1) Ventral hysterosoma with 5 pairs hystergastral setae in *P. summersi*, compared to 7 pairs in *P. bakeri* **sp. n.**; 2) Chaetotaxy of legs I–IV in *P. summersi* is: basifemora 4-6-4-2, telofemora 5-5-4-3, genua 8-7-6-6, tibiae 7-6-6-5, and tarsi 18-19-20-18, whereas in *P. bakeri* **sp. n.** the chaetotaxy is: basifemora 2-4-4-2, telofemora 5-5-4-4, genua 8-8-6-5, tibiae 7-6-6-5, and tarsi 25-25-18-16.

Pseudobonzia bakeri **sp. n.** can be separated from *P. clathratus* (Shiba) by the following characters: 1) Palp tibiotarsus with 5 simple setae in *P. clathratus*, whereas with 6 simple setae in *P. bakeri* **sp. n.**; 2) Venter with 6 pairs of hystergastral setae in *P. clathratus*, compared to 7 pairs in *P. bakeri* **sp. n.**; 3) Chaetotaxy of legs I–IV in *P. clathratus* is: basifemora 3-3-2-1, telofemora 6-5-4-3, genua 9-6-6-6, tibiae 8-6-6-5, and tarsi 27-21-18-14, whereas in *P. bakeri* **sp. n.** the chaetotaxy is: basifemora 2-4-4-1, telofemora 4-5-4-4, genua 8-8-5-5, tibiae 7-6-5-5, and tarsi 24-25-18-16.

This new species can also be compared with *P. delfinadobakerae* Smiley, however, they can be separated by the combination of the following features: 1) Palp tibiotarsus with a thick spine-like seta in *P. delfinadobakerae*, whereas it is absent in *P. bakeri* **sp. n.**; 2) Ventral hysterosoma with 6 pairs of hystergastral setae in *P. delfinadobakerae*, compared to 7 pairs in *P. bakeri* **sp. n.**; 3) Chaetotaxy of legs I–IV in *P. delfinadobakerae* is: basifemora 3-3-2-1, telofemora 6-5-4-3, genua 9-7-6-6, tibiae 7-6-6-5, and tarsi 26-21-18-14, whereas in *P. bakeri* **sp. n.** the chaetotaxy is: basifemora 2-4-4-1, telofemora 4-5-4-4, genua 8-8-5-5, tibiae 7-6-5-5, and tarsi 24-25-18-16; 4) Genital shield and coxae

are papillate in *P. bakeri* **sp. n.**, whereas they are reticulated in *P. delfinadobakerae*.

References

Arbabi M, Golmohammadzadeh NK, Askari M. 2002. Plant mite fauna of Sistan-Baluchestan and Hormozgan provinces. *Journal of Entomological society of Iran* 22(1): 87-105.

Baker EW, Hoffmann A. 1948. Acaros de la Familia Cunaxidae. *Anales de la Escuela Nacional de Ciencias Biologicas Mexico* 5(3-4): 229-273.

Bashir MH, Afzal M, Akbar S. 2008. A new species of genus *Pseudobonzia* Smiley (Prostigmata: Acari) from Punjab, Pakistan. *Biologia* 54(1): 77-82.

Bashir MH, Afzal M. 2006. A new species *Coleoscurus trudus* (Acari: Cunaxidae) from Pakistan. *Pakistan Entomologist* 28(2): 73-77.

Bashir, MH, Afzal M, Ashfaq M, Khan BS. 2008. On the identity of two new species of the genus *Coleoscurus* (Acari: Cunaxidae) with annotations on the genus from Pakistan. *Pakistan Journal of Agricultural Sciences* 45(4): 452-458.

Berlese A. 1916. Centuria Seconda di Acari Nuovi. *Redia* 12(1): 127-177

Berlese A. 1888. Acari Austro-Americani quos collegit Aloysius Balzan. *Bollettino della Società Entomologica Italiana* 20: 177-222.

Chaudhri WM, Akbar S. 1985. Studies on Biosystematics and Control of Mites of Field Crops, Vegetables and Fruits Plants in Pakistan. *U.A.F. Technical Bulletin No. 3*. University of Agriculture.

Chaudhri WM. 1977. Description of the mite of the family Cunaxidae (Acarina) from Pakistan. *Pakistan Journal of Agricultural Sciences* 14: 41-52.

Corpuz-Raros LA, 2007. Additional species of Bonziinae and Cunaxoidinae and description of the male of *Coleoscurus horidula* (Tseng) (Coleoscurinae) from the Philippines (Cunaxidae, Acari). *Asia Life Sciences* 16(2): 153-173

Den Heyer J. 1976. *Scutascirus*, a new Cunaxid genus (Prostigmata: Acari) from South Africa. Wetenskaplike Bydraes van die Pu vir Cho Reeks B. *Natuurwetenskappe* 93: 1-10.

Den Heyer J. 1977. Six new species of *Pseudobonzia* Smiley, 1975 (Prostigmata: Acari) from the Ethiopian Region. *Journal of Entomological Society of Southern Africa* 40(2): 171-194.

Den Heyer J. 1978. Coleoscurinae, a new cunaxid subfamily and two New South African species of *Coleoscurus* Berlese, 1916. (Prostigmata: Acari). *Acarologia* 20(4): 522-541.

Den Heyer J. 1980a. Six new species of sub family Coleoscurinae (Cunaxidae: Actinedida: Acarida). *Phytophylactica* 12: 105-128.

Den Heyer J. 1980b. *A classification system for the family Cunaxidae (Actinedida)*. Publication of the University of the North, Series A 23: 1-12.

Den Heyer J. 2011. Some statistics on the taxonomy of the family Cunaxidae (Acari: Prostigmata). In: Moraes GJ de, Proctor H, Editors. *Acarology XIII: Proceedings of the*

International Congress. *Zoosymposia* 6: 1-304.

de-Oliveira AJ, Daemon E. 2003. Qualitative and quantitative assessment of mites (Acari) in domiciliary dust in rural dwellings in the “Zona da Mata” region, Minas Gerais, Brazil. *Revista Brasileira de Zoologica* 20(4): 675-679.

Ewing HE. 1917. New Acarina. Part II, Description of new species and varieties from Iowa, Missouri, Illinois, Indiana and Ohio. *Bulletin of American Museum of Natural History* 37(2): 149-172.

Inayatullah, Shahid M. 1993. Three new predatory mites of genus *Pseudocunaxa* Smiley (Acarina: Cunaxidae) from Pakistan. *Pakistan Journal of Zoology* 25(4): 315-320.

Kaluz S. 2009. Two new Palearctic mite species of the family Cunaxidae (Acari: Prostigmata). *Zootaxa* 2198: 27-40

Kethley J. 1990. Acarina: Prostigmata (Actinedida) In: Dindal DL, Editor. *Soil Biology Guide*. pp. 667–756. John Wiley and Sons.

Kuznetsov NN, Sizova JY. 1978. The predaceous mite fauna of cotton fields in Uzbekistan. *Uzbekskii Biologicheskii Zhurnal* 40: 59-64.

Liang GW. 1984. A new species and a new record of the genus *Pseudobonzia* from China. *Acta Zootaxonomica Sinica* 9: 49-51.

Lin J-Z, Zhang Y-X, Ji J. 2001. Three new species of Cunaxidae from Fujian, China. *Systematic and Applied Acarology* 6: 145-153.

Lord FT. 1949. The influence of spray programs on the fauna of Apple orchards in Nova Scotia. III. Mites and their predators. *Canadian Entomologist* 31: 202-214.

Luxton M. 1982. Some new species of mites from New Zealand peat soils. *New Zealand Journal of Zoology* 9: 325-332.

Mejía-Recamier BE, Palacios-Vargas JG. 2007. Three new species of *Neoscirula* (Prostigmata: Cunaxidae) from a Tropical dry forest in Jalisco, Mexico. *Zootaxa* 1545: 17-31

Muhammad T, Chaudhri WM. 1992a. Two new mite species of genus *Coleoscurus* Berlese (Acarina: Cunaxidae) from Pakistan. *Pakistan Journal of Zoology* 24(4): 309-311.

Muhammad T, Chaudhri WM. 1992b. Genus *Coleoscurus* Berlese (Cunaxidae: Acarina) from Pakistan. *Pakistan Entomologist* 14(1-2): 97-103.


Nesbitt HHJ. 1946. Three new mites from Nova Scotian apple trees. *Canadian Entomologist* 68: 15-22.

Schruff G. 1971. *Haleupalus oliveri* new species, A “Thorn-palped” mite on grape vines (*Vitis* Spp.) Acari: Cunaxidae. *Deutsche Entomologische Zeitschrift* 18(4-5): 379-382.

Sepasgosarian H. 1984. The world genera and species of the family Cunaxidae (Actinedida: Acarida). *Zeitschrift für Angewandte Zoologie* 71: 135-150.

Sergeyenko AL. 2003. A new species of mites of the genus *Cunaxa* (Acarina: Prostigmata: Cunaxidae) from Crimea (Ukraine). *Acarina* 11(2): 225-229.

- Sergeyenko AL. 2005. A new species of the mite genus *Pseudobonzia* Smiley, 1975 (Acarina: Prostigmata: Cunaxidae) from Ukraine. *Acarina* 13(2): 159-163
- Sergeyenko AL. 2006. New species and new record of mites of the genus *Rubroscirus* Den Heyer, 1979 (Acarina: Prostigmata: Cunaxidae) from Ukraine. *Acarina* 14(1): 103-107
- Sergeyenko AL. 2009. New mite species of the genus *Cunaxa* (Acari: Prostigmata: Cunaxidae) from the Crimea, Ukraine. *Zootaxa* 2161: 1-19.
- Sergeyenko AL. 2011. Mites of the genera *Pulaeus* and *Lupaeus* (Acari: Prostigmata: Cunaxidae) of Crimea, Ukraine. *Zootaxa* 3088: 54-68
- Sionti PG, Papadoulis GT. 2003. Cunaxid mites of Greece (Acari: Cunaxidae). *International Journal of Acarology* 29(4): 315-325
- Skvarla MJ, Dowling APG. 2012. Some new Armascirine cunaxids (Acari: Prostigmata: Cunaxidae) from the Eastern United States. *Zootaxa* 3194: 1-34.
- Skvarla MJ, Fisher JR, Dowling APG. 2011. A new species of *Neoscirula* (Acari: Cunaxidae: Coleosirinae) from the Ozark highlands (USA), with a note on biogeography. *Acarologia* 51(3): 283-293.
- Smiley RL. 1975. A generic revision of the mites of the family Cunaxidae (Acarina). *Annals of the Entomological Society of America* 68(2): 227-244
- Smiley RL. 1992. *The predatory mite family Cunaxidae (Acari) of the world with a new classification*. Indira Publishing House.
- Swift SF. 1996. Two new species of *Dactyloscirus* (Acari: Prostigmata: Cunaxidae) in the Hawaiian Islands. *Anales del Instituto Biologia, Universidad Nacional Autonoma de Mexico, Serie Zoologia* 67(2): 225-237.
- Tagore A, Putatunda BN. 2003. Mites associated with some ornamental plants at Hisar, Haryana. *Pest Management and Economic Zoology* 11(1): 37-44.
- Thor S, Willmann C. 1941. Acarina Prostigmata 6-11 (Eupodidae, Penthalodidae, Penthaleidae, Pachygnathidae, Cunaxidae). *Das Tierreich* 71a: 1-186.
- Tseng YH. 1980. Taxonomical study of the mite family Cunaxidae from Taiwan (Acarina: Trombidiformes). *Quarterly Journal of Taiwan Museum* 33(3-4): 253-277.
- Walter DE, Kaplan DT. 1991. Observations on *Coleosirius simplex* (Acarina: Prostigmata), a predatory mite that colonizes greenhouse cultures of rootknot nematode (*Meloidogyne* spp.), and a review of feeding behavior in the Cunaxidae. *Experimental and Applied Acarology* 12(1-2): 47-59.
- Walter DE. 1999. Cryptic inhabitants of a noxious weed: Mites (Arachnida: Acari) on *Lantana camara* L. invading forests in Queensland. *Australian Journal of Entomology* 38: 197-200
- Youssef AA, El-Badry EA, Hykal IH. 1980. Mites inhabiting cotton and associated weeds in Egypt. *Bulletin of the Entomological Society of Egypt* 60: 223-227.






Figure 2. *Pseudobonzia bakeri*, **sp. n.** Dorsal side. High quality figures are available online.




Figure 3. *Pseudobonzia bakeri*, sp. n. Ventral side. High quality figures are available online.




Figure 4. *Pseudobonzia bakeri*, sp. n. Legs I–IV. High quality figures are available online.