

## **A New Exotic Noctuid for the Hawaiian Archipelago: *Feltia subterranea* (Fabricius) (Lepidoptera: Noctuidae: Noctuinae)**

Author: Prestes, Andersonn Silveira

Source: The Journal of the Lepidopterists' Society, 68(3) : 220-221

Published By: The Lepidopterists' Society

URL: <https://doi.org/10.18473/lepi.v68i3.a11>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

A NEW EXOTIC NOCTUID FOR THE HAWAIIAN ARCHIPELAGO: *FELTIA SUBTERRANEA*  
(FABRICIUS) (LEPIDOPTERA: NOCTUIDAE: NOCTUINAE)

**Additional key words:** island, introduced species, new record, Hawaii

The archipelago of Hawaii has a rich endemic biota. As the most isolated group of islands on earth, Hawaii is characterized by a disharmonic and highly endemic fauna and flora. As elsewhere, humans have been heavily modifying its natural habitats since Polynesians first arrived to the islands. Relative to other terrestrial taxa, the Noctuidae of Hawaii represent a rich and interesting fauna. There are 73 described native species in 14 genera (Riotte 1991; Mitchell 1997). On the other hand, the exotic fauna is substantial with 45 species from 35 genera (Riotte 1991). The exotic noctuid fauna can be frequently encountered, both at low or high elevations, and species such as *Mythimna unipuncta* (Haworth) or *Athetis thoracica* (Moore) are often abundant. It is not certain if exotic species are displacing and competing with the natives. But since they are closely related, with similar life histories and ecological affinities, disruption might be expected to some degree. An important conservation factor in the relationship between exotics and natives is the sharing of natural enemies, and there is already evidence of natural enemies introduced to control exotic species preying on natives (Henneman & Memmott 2001). Also, closely related exotic species might hybridize with native species causing serious disruption in the species' integrity, where it might potentially lead to the extinction of the native populations (Hardwick 1965). Exotic species can have a negative economic impact, and may cause serious impacts to crops, quickly becoming pests and widespread. In this context, one more exotic noctuid species has been found in the islands of Hawaii.

Members of the Daniel Rubinoff Laboratory at the University of Hawaii in Honolulu collected 4 specimens of *Feltia subterranea* (Fabricius) from three different islands in the past seven years. There are two samples from Maui, one from Lanai, and one from Hawaii, all of them collected after 2006.

*Feltia subterranea* was formerly placed in *Agrotis*, but Lafontaine's monograph (2004) included an unambiguous morphological character for identification of *Feltia*: a doubly biserrate male antenna with the enlarged apical seta being on a third process. *Feltia subterranea* can be recognized by "the double biserrated male antennae, the small round orbicular

spot connected to the reniform spot by a narrow black bar, and the translucent pearly white hindwing in both sexes" (Lafontaine 2004). The highly polyphagous larvae, commonly known as granulated cutworm, feeds on bean, beet, cabbage, corn, lettuce, peas, potato, tobacco, tomato, and turnip, among others. The larvae are easily recognized by the presence of scattered, raised, conical skin granules (Lafontaine 2004). To my knowledge, the caterpillars have not yet been found on Hawaiian crops, although three of the samples are near disturbed and agricultural lands.

The moth is widespread and common in North America. It is a migratory species, which also occurs in Central and South America, as far south as Peru and Brazil (Lafontaine 2004). As a species with a broad range, including tropical areas, and great mobility, it is possible that Hawaii may become a favorable place to spread and establish. As possible routes of introduction to the islands, the larvae might have travelled in the roots of its hostplants or as adults on a ship (Lafontaine, personal communication).

More information about the species can be found in Lafontaine (2004) or in the website [http://entnemdept.ufl.edu/creatures/veg/granulate\\_cutworm.htm#dist](http://entnemdept.ufl.edu/creatures/veg/granulate_cutworm.htm#dist).

Material examined (at University of Hawaii Insect Museum, UHIM): 1 male, 1 female, USA: HI, East Maui, Makawao Forest Reserve, near banana patch, 762m, 20-21.Jun.2006, W. Haines; 1 male, USA: HI, Hawaii, Waimea, 720m, N20 04 16.2 W155 34 56.8, 31.Out.2008, Rubinoff lab.; 1 male, USA: HI, Lanai, Kanepu'u Reserve, near outplanting, 18-19.Sept.2009, D. Rubinoff, C. King, W. Haines.

#### ACKNOWLEDGMENTS

I thank Daniel Rubinoff for advice and suggestions. Donald Lafontaine for the help on the identification, and Ryan Caesar for suggestions. I thank the Brazilian "Coordenação de Aperfeiçoamento de Pessoal de Nível Superior" (CAPES foundation) for the Doctorate Scholarship. I thank two anonymous reviewers for suggestions.

#### LITERATURE CITED

- LAFONTAINE, J. D. 2004. The Moths of North America: Noctuoidea, Nctuidae (Part). Fascicle 27.1. The Wedge Entomological Research Foundation, 385 pp.
- HARDWICK, D. F. 1965. The corn earworm complex. Mem. Entomol. Soc. Canada 248 pp.

- HENNEMAN, M.L. & MEMMOTT, J. 2001. Infiltration of a Hawaiian community by introduced biological control agents. *Science* 293:1314-1316.
- MITCHELL, A. 1997. The first endemic Hawaiian *Heliothis* (Lepidoptera: Noctuidae): *H. melanoleuca*, a new species from Riparian forest. Occas. pap. Bernice P. Bishop Mus. 48: 78-81.
- RIOTTE, J.C.E. 1991. Reassessment of the Noctuoidea of the Hawaiian Islands. Occas. pap. Bernice P. Bishop Mus. 31: 139-151.

ANDERSONN SILVEIRA PRESTES *University of Hawaii at Manoa, Department of Plant and Environmental Protection Sciences. 3050 Maile Way, Gilmore 310. Honolulu, HI 96822. email: aprestes@hawaii.edu*

*Submitted for publication 28 November 2013; revised and accepted 10 March 2014.*