

## **Introduction and a theoretical basis for using disturbance by forest management activities to sustain aquatic ecosystems**

Authors: Kreutzweiser, David P., Sibley, Paul K., Richardson, John S., and Gordon, Andrew M.

Source: Freshwater Science, 31(1) : 224-231

Published By: Society for Freshwater Science

URL: <https://doi.org/10.1899/11-114.1>

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## BRIDGES

# Introduction and a theoretical basis for using disturbance by forest management activities to sustain aquatic ecosystems

David P. Kreutzweiser<sup>1,4</sup>, Paul K. Sibley<sup>2,5</sup>, John S. Richardson<sup>3,6</sup>, AND Andrew M. Gordon<sup>2,7</sup>

<sup>1</sup>Canadian Forest Service, Natural Resources Canada, Sault Ste Marie, Ontario, Canada P6A 2E5

<sup>2</sup>School of Environmental Sciences, University of Guelph, Guelph, Ontario, Canada N1G 2W1

<sup>3</sup>Department of Forest Sciences, University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z4

**Abstract.** Emulation of natural disturbance (END) is an emerging paradigm for modern, ecosystem-based forest management in North America. On the premise that periodic disturbance is an integral part of natural, determinative processes on forest landscapes, managing forests by emulating natural disturbance is thought to produce landscape patterns that resemble those arising from natural disturbances and that are known to maintain critical processes and habitat for conserving biodiversity. Applying END principles to forest watersheds has implications for the protection of aquatic ecosystems because END can include intentional logging disturbance near water to emulate natural riparian disturbance. Literature shows that logging in watersheds, and especially in riparian areas, can lead to negative abiotic and biotic effects in aquatic ecosystems. However, an integration of the current understanding of land–water linkages in forest watersheds with general disturbance ecology would suggest that periodic watershed and riparian disturbances may be natural renewal processes that are required for long-term sustainability of aquatic ecosystems. Previous syntheses of END in forestry failed to consider the implications for aquatic ecosystems, and most forest-management guidelines default to the protection of water resources by systematic riparian (shoreline) buffers. This paper introduces the concepts of END and provides a theoretical basis for using intentional riparian forest disturbance to sustain aquatic habitat complexity and ecosystem integrity.

**Key words:** natural disturbance emulation, forest watershed, logging impacts, aquatic ecosystem sustainability.

Forest management policies and practices are evolving in North America. Policymakers and practitioners are increasingly adopting whole ecosystem-based management principles as a means to attain sustainable forest management (Kimmins 2004). The assumption is that ecosystem-based management will simultaneously provide the required habitat for most species across the landscape and thereby sustain biodiversity and critical ecosystem processes within their natural range of variation (Hunter 1999, Lindenmayer et al. 2006). An emerging paradigm that underpins ecosystem-based management for forests is

the emulation of natural disturbance, or END (Long 2009). On the premise that periodic disturbance is an integral part of natural, determinative processes on forest landscapes, managing forests to emulate natural disturbance is thought to achieve the goals of ecosystem-based management by providing landscape patterns resembling those that result from natural disturbances and that are known to renew and maintain critical processes and habitat for conserving biodiversity (OMNR 2001). This practice means creating spatial patterns across managed forest landscapes that are similar in size and forest structure to patterns arising from natural disturbance. It is generally accepted that forest ecosystem resilience has evolved in part from a legacy of natural disturbance in the sense that disturbance ultimately fosters the structural and functional complexity of forest

<sup>4</sup> E-mail addresses: dave.kreutzweiser@nrca.gc.ca

<sup>5</sup> psibley@uoguelph.ca

<sup>6</sup> john.richardson@ubc.ca

<sup>7</sup> agordon@uoguelph.ca

landscapes (Gunderson 2000). Under END, management goals seek to sustain forest ecosystem structure and function across a number of spatial scales thereby retaining a capacity for renewal, maintaining desired ecological goods and services, and reducing the probability of shifting to an undesirable state (Drever et al. 2006).

Perera and Buse (2004, p. 4) define END as “an approach in which forest managers develop and apply specific management strategies and practices, at appropriate spatial and temporal scales, with the goal of producing forest ecosystems as structurally and functionally similar as possible to the ecosystems that would result from natural disturbances and that incorporate the spatial, temporal, and random variability intrinsic to natural systems”. Fire is currently the most common disturbance agent being modeled for END in forest management (Hunter 1993), but it is not the only disturbance type that is useful or appropriate in this context. Other disturbances being considered for effectiveness as forest management models include insect defoliation (MacLean 2004), gap phase processes from natural mortality (Seymour et al. 2002), windthrow (Kneeshaw et al. 2011), debris flows (Rood 2006), and a combination of disturbances (Suffling and Perera 2004). Natural disturbance emulation is applicable to any forest system that is disturbance-influenced to varying degrees, ranging from stand-replacing events (like fire) to gap openings from natural mortality. However, END probably will be applied to mimic disturbance-generated features only on landscapes with frequent and moderate disturbances because emulating very large, severe, or infrequent disturbances probably would meet with societal and political resistance regardless of their ecological relevance (Kimmins 2004). Therefore, the application of END through forest management is particularly relevant to disturbance-prone systems, such as the boreal forest, where stand-replacing events can occur at temporal scales of decades to centuries (Hunter 1993).

### **Implications of Natural Disturbance Emulation for Aquatic Ecosystems**

The END approach in forest watersheds has implications for the protection of aquatic ecosystems because water bodies are ubiquitous across most forest regions and are integral parts of forest ecosystems (Richardson and Danehy 2007). Forest water bodies are strongly linked to their surrounding terrestrial catchments (Hynes 1975), so forest management activities that alter the forest structure and other watershed features can influence aquatic ecosystems. Numerous

studies have assessed the impacts on aquatic ecosystems of watershed disturbances from forest management over the past few decades (see reviews by Webster et al. 1992, Prepas et al. 2003, Fortino et al. 2004, Kreutzweiser et al. 2008b, Richardson 2008, Buttle et al. 2009). In general, these studies show that tree removal, ground disturbances, changes in watershed nutrient cycling and export, and road construction can affect receiving waters and their biotic communities and that the magnitude of effects is site- or context-specific. Effects can include reduced canopy cover and shading, increased water temperatures, increased fine-sediment deposition, increased nutrient concentrations, reduced inputs of large wood and fine organic matter, and restricted fish movement, all of which have implications for aquatic ecosystem sustainability. As a result, best management practices for forest operations have been developed and refined to mitigate aquatic impacts of forest management (Table 1) (Vowell and Frydenborg 2004, Schilling 2009, Ice et al. 2010, Richardson et al. 2012).

One of the most broadly applied best management practices in North America for protecting forest water bodies is retention of riparian (shoreline) buffers in which logging operations are prohibited or restricted around water bodies (Phillips et al. 2000, Lee et al. 2004). Forested riparian buffers are usually effective for mitigating many forest management impacts on aquatic systems, particularly those associated with near-water canopy removal or ground disturbance (Barling and Moore 1994, Broadmeadow and Nisbet 2004, Hickey and Doran 2004), but not necessarily those impacts that arise from watershed-level effects (i.e., significant changes in water quality or biotic responses were detected even with intact riparian buffers) (Prepas et al. 2003, Martel et al. 2007, Kreutzweiser et al. 2008a, Lecerf and Richardson 2010).

The systematic application of riparian buffers around water bodies leads to unnatural, linear patterns of older-growth forests across the landscape resulting in ribbons along streams and donuts around lakes (Buttle 2002, Steedman et al. 2004). Protection of riparian forests in managed watersheds could suppress natural riparian forest renewal because the adjacent upland areas contain young, regenerating stands that are less prone to disturbances like fire, wind, and insect infestations that could otherwise cross into riparian areas. In the absence of periodic disturbance and regeneration, riparian habitat heterogeneity probably will decrease (riparian areas will become more homogeneous) and, as a result, structural and functional riparian diversity may decrease to a condition that is less than the natural range of variability (Degraaf and Yamasaki 2000, Swanson

TABLE 1. Examples of forestry best management practices (BMPs) intended to mitigate potential impacts on water resources.

BMP	Protection goals	Operational applications
Appropriate operational timing	Minimize soil disturbance, rutting, water diversion in wet areas	Harvest under frozen, snow-covered, or dry conditions
Roads and stream crossings	Minimize erosion, sediment runoff, water diversion; avoid fish movement barriers	Locate roads and crossings to avoid wet soils when possible, minimize water flow disruption; use course surfacing and back-fill materials, other sediment retention or diversion techniques; proper bridge and culvert installations, avoid culverts perched above stream beds
Landings	Minimize erosion, soil disturbance, sediment runoff	Keep number and size of landings to minimum; avoid wetlands, steep, or unstable slopes; keep out of riparian areas; use appropriate ditches, out-sloping to keep landings dry
Skidding and skid trails	Minimize erosion, soil disturbance, sediment runoff	Avoid low-lying or wet areas; skid across slopes and parallel to shorelines or stream channels; divert water by lead-off ditches; avoid concentrated and repeated skidder movement on wet or shallow or organic soils
Stabilize exposed soils on slopes	Prevent soil damage; minimize sediment and nutrient runoff	Use appropriate equipment such as low ground pressure tires; smooth out ruts on slopes to prevent channeling; promote revegetation as quickly as possible; apply stabilizing ground cover such as branches, mulch, cobble
Riparian buffer strips or management zones	Retain shoreline stability, shade, litter and wood inputs, terrestrial and semiaquatic habitats; intercept sediment and nutrient runoff; minimize near-water soil compaction or disturbance	Restricted or modified harvesting in buffers, restricted machine movement; specified tree retention
Proper equipment fuelling and maintenance	Avoid contaminating water bodies with fuels, lubricants, other contaminants	Place fuelling and maintenance areas away from water sources; provide waste containers on site; conduct regular checks for leaks or wear

et al. 2011). The unnatural landscape patterns created by these riparian buffers may not be consistent with the goals of forest management based on END patterns (Macdonald et al. 2004, Holmes et al. 2010) because natural disturbances often occur in riparian areas (Andison and McCleary 2002, Nitschke 2005).

Unnatural landscape patterns arising from the systematic retention of no-logging buffers are resulting in changes or proposed changes to riparian forest management guidelines in Canada (Morissette and Donnelly 2010, Sibley and Gordon 2010). Several recent studies or guidelines have incorporated intentional shoreline disturbance by careful near-water logging to create more natural riparian forest conditions (Kardynal et al. 2009, Kreutzweiser et al. 2010, OMNR 2010, Naylor et al. 2012). This approach seems counterintuitive for the protection of water resources and aquatic ecosystem integrity. The forest management guidelines that focus on the retention of riparian buffers are intended to prevent or mitigate disturbances to shoreline areas, thereby protecting aquatic ecosystems. The intended outcomes from these buffer guidelines often are not clearly defined by specific, quantitative targets (Richardson and Thompson

2009), but the premise generally has been protection of aquatic systems by avoiding or minimizing change from prelogging (baseline) or nearby no-logging (reference) conditions. Changes induced by forest management operations from these baseline or reference conditions usually have been construed as undesirable impacts (see the reviews on forestry impacts listed above). However, in forests that have evolved over a range of disturbances, using baseline or reference conditions defined over a short time period (often only a few years) to represent the natural range of variability in forests and their water bodies over longer periods (decades or more) may be inappropriate. If the target for management under END is “no change beyond the range of natural disturbance” (Long 2009), then short-term changes from reference conditions or sites may be acceptable. Therefore, applying END to riparian forest management (i.e., intentional disturbance of some riparian forests) may require redefining acceptable and unacceptable changes in aquatic ecosystems using specific conservation targets based on mimicking natural disturbance (Richardson and Thompson 2009, Moore and Richardson 2012).

Empirical studies are clearly needed to develop and assess relevant conservation targets for applying END to riparian forests. Studies should be done to determine the effects of intentional shoreline disturbance under END and the extent to which riparian logging emulates or modifies the effects of natural disturbances (Nitschke 2005). These studies also should establish the bounds of natural variability in forest water bodies to help determine when logging-induced changes become unacceptable impacts or simply natural (or emulated) pulses in longer-term renewal processes of a disturbance-based ecosystem. This approach differs from the historical assessments of intensive logging to shorelines that have demonstrated impacts on aquatic communities (e.g., Ely and Wallace 2010), because shoreline harvesting under END principles will follow careful harvesting practices under specific guidelines. For example, although the goal is to mimic natural disturbance patterns as closely as possible, shoreline harvesting would be constrained in some places by restrictions on machine movement or ground disturbance in wet areas, residual forest requirements for special wildlife habitats, and downed wood, inoperable terrain, cultural heritage sites, or other values (Naylor et al. 2012).

### **A Theoretical Basis for Applying END to Riparian Forest Management**

Forest ecologists have long regarded natural disturbance regimes as critical factors in structuring and sustaining diverse forest communities (reviewed by McCarthy 2001), but the role of disturbance in enhancing spatial heterogeneity and sustaining aquatic communities has been recognized increasingly in more recent years (Lake 2000, Lepori and Hjerdt 2006, Lepori and Malmqvist 2007). The emphasis in our paper is on streams, but the principles of disturbance effects would apply to other aquatic ecosystems as well. General disturbance theory postulates that spatial heterogeneity in biotic communities, patchiness, and temporal dynamics arise in large part from disturbance and that the absence of this disturbance-mediated temporal and spatial variability would preclude the existence of many species (Sousa 1984). Disturbance in ecology is variously defined, but White and Pickett's (1985, p. 7) definition of "any relatively discrete event in time that disrupts ecosystems, community, or population structure, and changes resources, substrate availability, or the physical environment" is useful. The simplified premise of disturbance theory is that disturbance increases habitat heterogeneity and complexity by destroying some patches (conditions) while creating

others, thereby changing the colonization, habitation, and succession patterns of biota and influencing community structure. The resultant different-sized patches under varying conditions support a higher biodiversity than is supported in more homogeneous systems.

Disturbance theory has been successful in explaining or predicting community structure in aquatic systems. The effects of natural disturbances on aquatic communities can be scale- and context-dependent, but disturbances promote habitat heterogeneity that underpins the continuance of diverse aquatic communities (Reice 1994, Lepori and Hjerdt 2006). The interactive effects of disturbances and patch dynamics on biotic communities are complex, but in general, major disturbances like floods and drought create new patterns of patchiness that regulate biotic community structure by varying resource availability and use (Lake 2000).

Disturbances are of particular importance to forest watersheds and riparian areas where land-water linkages connect terrestrial and aquatic ecosystems because they can play a role in establishing and sustaining those linkages. Nakamura et al. (2000) demonstrated this importance by focusing on the sequencing or cascading of geomorphological disturbances down the gravitational flow paths of stream networks and their riparian systems in mountainous landscapes. The same principles should be applicable to other landscapes. They suggest that streams and their riparian zones are networks containing a shifting mosaic of disturbance patches that are linear and parallel, linking riparian conditions to stream conditions in the context of disturbance. Swanson et al. (2011) pointed out that natural recovery processes in forest ecosystems that promote succession, renewal, complexity and stability also occur in riparian areas and influence aquatic ecosystems.

Thus, it follows that disturbances to forested watersheds are important natural determinants of aquatic habitat conditions and communities because of the strong ecological linkages between forest water bodies and their terrestrial watersheds. Natural disturbances in riparian and upland forests can influence these ecological linkages and induce changes in receiving waters. For example, insect defoliation of riparian forests can change stream water chemistry through elevated nitrification in riparian soils and leaching of nutrients and labile C from insect frass and green litterfall (Lewis and Likens 2007). Wildfire can induce large changes in aquatic ecosystems and their communities through canopy removal, increased runoff, altered biogeochemical processes, nutrient fluxes, sediment delivery, and other influences (Bisson

et al. 2003, Spencer et al. 2003, Neary et al. 2008, Malison and Baxter 2010, Moore and Richardson 2012). Forest gap generation by windthrow can influence in-stream light availability and periphyton communities via increased spatial variability in canopy structure (Stovall et al. 2009).

It appears then, that natural forest disturbances in watersheds will promote aquatic habitat heterogeneity and, therefore, natural patterns in biological diversity. Disturbance is a significant ecological process that maintains biodiversity and ecosystem integrity (Mori 2011). Therefore, forest watershed disturbances may be required for long-term aquatic ecosystem stability. One could argue that restricting natural riparian and upland disturbances through fire suppression, insect control, or no-logging buffers could conceivably constrain these natural disturbance and renewal processes and their effects on aquatic habitats and communities. No-logging buffers cannot accommodate the natural range of variability in riparian forest composition and function, and they disregard the fact that disturbance is a natural part of riparian forests (Palik et al. 2000, Sibley and Gordon 2010). This logic suggests that intentional disturbances by carefully planned and implemented forest management operations (e.g., Naylor et al. 2012) could be used to emulate natural disturbances in watersheds and riparian forests to sustain aquatic ecosystems over the long term, recognizing that some short-term alterations (potentially perceived as impacts) could be incurred. However, this suggestion is largely untested in forest watersheds.

### Conclusions

Previously published forums on END in forestry (Kuuluvainen 2002, Mitchell et al. 2002, Perera et al. 2004) have not considered implications for watershed and riparian management or for conservation of aquatic biodiversity and ecosystem integrity. Jurisdictions that have implemented some END principles for riparian forest management usually have done so to create complexity through early-successional regeneration (Swanson et al. 2011) in shoreline habitats to support terrestrial and semiaquatic biodiversity (Kreutzweiser et al. 2005, 2010, Kardynal et al. 2009, Naylor et al. 2012), while striving to avoid adverse effects on aquatic systems. We now suggest, based on disturbance ecology and the recognized land–water linkages in forest watersheds, that END in forest management can also be applied to sustain aquatic biodiversity and ecosystem integrity by mimicking natural riparian and watershed disturbances and renewal processes. This suggestion does

not imply unrestrained riparian logging in all areas, but rather carefully planned and implemented logging in some riparian forests under science-based guidelines (Morissette and Donnelly 2010, Sibley and Gordon 2010, Moore and Richardson 2012, Naylor et al. 2012, Sibley et al. 2012). However, much of this idea is untested and many uncertainties remain around the application of END to forest watersheds. The subsequent papers in this *BRIDGES* cluster explore some of the outstanding issues.

Our intent is for this cluster to promote an exchange of ideas on how, why (or why not), when, and where to apply END in riparian forests. Using intentional riparian disturbance by forest management to emulate natural disturbance patterns and processes is a new and evolving concept in forestry, and the potential implications for aquatic ecosystems have not been explored. Indeed, this paucity of information on the implications of END for aquatic ecosystems was the impetus for exploring this topic through a *BRIDGES* cluster in hope of generating advanced discussion and study. We have drawn on established disturbance theory to provide a theoretical basis for using intentional riparian disturbance as a management tool because few empirical studies are available. Our group has notionally embraced the concept of applying END to riparian forest management, but we recognize the need to move on to empirical testing. Discussions arising from this cluster should advance and direct that process.

### Acknowledgements

Many of the ideas for this *BRIDGES* cluster were spawned during discussions at the White River Riparian Harvesting Impacts Project field tour, supported in part by the Enhanced Forest Productivity Science Program Grant 010-2-R1 (DPK). Subsequent discussion and suggestions from Ashley Moerke advanced the ideas for a journal series and resulted in this cluster. Brian Naylor and anonymous referees provided useful comments on earlier manuscripts.

### Literature Cited

- ANDISON, D. W., AND K. MCCLEARY. 2002. Disturbance in riparian zones on foothills and mountain landscapes of Alberta. Alberta Foothills Disturbance Ecology Research Series Report No. 3. Foothills Model Forest, Hinton, Alberta. (Available from: [http://foothillsresearchinstitute.ca/Content\\_Files/Files/ND/ND\\_report3.pdf](http://foothillsresearchinstitute.ca/Content_Files/Files/ND/ND_report3.pdf))
- BARLING, R. W., AND I. D. MOORE. 1994. Role of buffer strips in management of waterway pollution: a review. *Environmental Management* 18:543–558.
- BISSON, P. A., B. E. RIEMAN, C. LUCE, P. F. HESSBURG, D. C. LEE, J. L. KERSHNER, G. H. REEVES, AND R. E. GRESSWELL. 2003.

- Fire and aquatic ecosystems of the western USA: current knowledge and key questions. *Forest Ecology and Management* 178:213–229.
- BROADMEADOW, S., AND T. R. NISBET. 2004. The effects of riparian forest management on the freshwater environment: a literature review of best management practices. *Hydrology and Earth System Sciences* 8:286–305.
- BUTTLE, J. M. 2002. Rethinking the donut: the case for hydrologically relevant buffer zones. *Hydrological Processes* 16:3093–3096.
- BUTTLE, J. M., I. F. CREED, AND R. D. MOORE. 2009. Advances in Canadian forest hydrology, 2003–2007. *Canadian Water Resources Journal* 34:113–126.
- DEGRAAF, R. M., AND M. YAMASAKI. 2000. Bird and mammal habitat in riparian areas. Pages 139–156 in E. S. Verry, J. W. Hornbeck, and A. Dolloff (editors). *Riparian management in forests of the continental eastern United States*. Lewis Publishers, CRC Press, Boca Raton, Florida.
- DREVER, C. R., G. PETERSON, C. MESSIER, Y. BERGERON, AND M. FLANNIGAN. 2006. Can forest management based on natural disturbances maintain ecological resilience? *Canadian Journal of Forest Research* 36:2285–2299.
- ELY, D. T., AND J. B. WALLACE. 2010. Long-term functional group recovery of lotic macroinvertebrates from logging disturbance. *Canadian Journal of Fisheries and Aquatic Sciences* 67:1126–1134.
- FORTINO, K., A. E. HERSHEY, AND K. J. GOODMAN. 2004. Utility of biological monitoring for detection of timber harvest effects on streams and evaluation of Best Management Practices: a review. *Journal of the North American Benthological Society* 23:634–646.
- GUNDERSON, L. H. 2000. Ecological resilience—in theory and application. *Annual Review of Ecology and Systematics* 31:425–439.
- HICKEY, B., AND B. DORAN. 2004. A review of the efficiency of buffer strips for the maintenance and enhancement of riparian ecosystems. *Water Quality Research Journal of Canada* 39:311–317.
- HOLMES, S. B., D. P. KREUTZWEISER, AND P. S. HAMILTON. 2010. Operational and economic feasibility of logging within forested riparian zones. *Forestry Chronicle* 86:601–607.
- HUNTER, M. L. 1993. Natural fire regimes as spatial models for managing boreal forests. *Biological Conservation* 65:115–120.
- HUNTER, M. L. (EDITOR). 1999. *Maintaining biodiversity in forest ecosystems*. Cambridge University Press, Cambridge, UK.
- HYNES, H. B. N. 1975. The stream and its valley. *Verhandlungen der Internationalen Vereinigung für theoretische und angewandte Limnologie* 19:1–15.
- ICE, G. G., E. SCHILLING, AND J. VOWELL. 2010. Trends for forestry best management practices implementation. *Journal of Forestry* 108:267–273.
- KARDYNAL, K. J., K. A. HOBSON, S. L. VAN WILGENBURG, AND J. L. MORISSETTE. 2009. Moving riparian management guidelines towards a natural disturbance model: an example using boreal riparian and shoreline forest bird communities. *Forest Ecology and Management* 257:54–65.
- KIMMINS, J. P. 2004. Emulating natural forest disturbances: what does this mean? Pages 8–28 in A. H. Perera, L. J. Buse, and M. G. Weber (editors). *Emulating natural forest landscape disturbances*. Columbia University Press, New York.
- KNEESHAW, D. D., B. D. HARVEY, G. P. REYES, M.-N. CARON, AND S. BARLOW. 2011. Spruce budworm, windthrow and partial cutting: do different partial disturbances produce different forest structures? *Forest Ecology and Management* 262:482–490.
- KREUTZWEISER, D. P., S. S. CAPELL, AND K. P. GOOD. 2005. Macroinvertebrate community responses to selection logging in riparian and upland areas of headwater catchments in a northern hardwood forest. *Journal of the North American Benthological Society* 24:208–222.
- KREUTZWEISER, D. P., K. P. GOOD, S. S. CAPELL, AND S. B. HOLMES. 2008a. Leaf litter decomposition and invertebrate communities in boreal forest streams linked to upland logging disturbance. *Journal of the North American Benthological Society* 27:1–15.
- KREUTZWEISER, D. P., P. W. HAZLETT, AND J. M. GUNN. 2008b. Logging impacts on the biogeochemistry of boreal forest soils and nutrient export to aquatic systems: a review. *Environmental Reviews* 16:157–179.
- KREUTZWEISER, D. P., E. A. MUTO, S. B. HOLMES, AND J. M. GUNN. 2010. Effects of upland clearcutting and riparian partial-harvesting on leaf pack breakdown and aquatic invertebrates in boreal forest streams. *Freshwater Biology* 55:2238–2252.
- KUULUVAINEN, T. 2002. Introduction: disturbance dynamics in boreal forests: defining the ecological basis of restoration and management of biodiversity. *Silva Fennica* 36:5–11.
- LAKE, P. S. 2000. Disturbance, patchiness, and diversity in streams. *Journal of the North American Benthological Society* 19:573–592.
- LECERF, A., AND J. S. RICHARDSON. 2010. Litter decomposition can detect effects of high and moderate levels of forest disturbance on stream condition. *Forest Ecology and Management* 259:2433–2443.
- LEE, P., C. SMYTH, AND S. BOUTIN. 2004. Quantitative review of riparian buffer width guidelines from Canada and the United States. *Journal of Environmental Management* 70:165–180.
- LEPORI, F., AND N. HJERDT. 2006. Disturbance and aquatic biodiversity: reconciling contrasting views. *BioScience* 56:809–818.
- LEPORI, F., AND B. MALMQVIST. 2007. Predictable changes in trophic community structure along a spatial disturbance gradient in streams. *Freshwater Biology* 52:2184–2195.
- LEWIS, G. P., AND G. E. LIKENS. 2007. Changes in stream chemistry associated with insect defoliation in a Pennsylvania hemlock-hardwood forest. *Forest Ecology and Management* 238:199–211.
- LINDENMAYER, D. B., J. F. FRANKLIN, AND J. FISCHER. 2006. General management principles and a checklist of strategies to guide forest biodiversity conservation. *Biological Conservation* 131:433–445.

- LONG, J. N. 2009. Emulating natural disturbance regimes as a basis for forest management: a North American view. *Forest Ecology and Management* 257:1868–1873.
- MACDONALD, E., C. J. BURGESS, G. J. SCRIMGEOUR, S. BOUTIN, S. REEDYK, AND B. KOTAK. 2004. Should riparian buffers be part of forest management based on emulation of natural disturbances? *Forest Ecology and Management* 187:185–196.
- MACLEAN, D. A. 2004. Predicting forest insect disturbance regimes for use in emulating natural disturbance. Pages 69–84 in A. H. Perera, L. J. Buse, and M. G. Weber (editors). *Emulating natural forest landscape disturbances*. Columbia University Press, New York.
- MALISON, R. L., AND C. V. BAXTER. 2010. The fire pulse: wildfire stimulates flux of aquatic prey to terrestrial habitats driving increases in riparian consumers. *Canadian Journal of Fisheries and Aquatic Sciences* 67: 570–579.
- MARTEL, N., M. A. RODRIGUEZ, AND P. BERUBE. 2007. Multi-scale analysis of responses of stream macrobenthos to forestry activities and environmental context. *Freshwater Biology* 52:85–97.
- MCCARTHY, J. 2001. Gap dynamics of forest trees: a review with particular attention to boreal forests. *Environmental Reviews* 9:1–59.
- MITCHELL, R. J., B. J. PALIK, AND M. L. HUNTER. 2002. Preface: natural disturbance as a guide to silviculture. *Forest Ecology and Management* 155:315–316.
- MOORE, R. D., AND J. S. RICHARDSON. 2012. Natural disturbance and forest management in riparian zones: comparison of effects at reach, catchment, and landscape scales. *Freshwater Science* 31:239–247.
- MORI, A. S. 2011. Ecosystem management based on natural disturbances: hierarchical context and non-equilibrium paradigm. *Journal of Applied Ecology* (online) doi: 10.1111/j.1365-2664.2010.01956.x
- MORISSETTE, J., AND M. DONNELLY. 2010. Riparian areas: challenges and opportunities for conservation and sustainable forest management. Sustainable Forest Management Network, Edmonton, Alberta. (Available from: [http://www.ales.ualberta.ca/forestry/Sustainable\\_Forest\\_Management/Publications/SynthesisReports.aspx](http://www.ales.ualberta.ca/forestry/Sustainable_Forest_Management/Publications/SynthesisReports.aspx))
- NAKAMURA, F., F. J. SWANSON, AND S. M. WONDZELL. 2000. Disturbance regimes of stream and riparian systems – a disturbance-cascade perspective. *Hydrological Processes* 14:2849–2860.
- NAYLOR, B. J., R. W. MACKERETH, D. P. KREUTZWEISER, AND P. K. SIBLEY. 2012. Merging END concepts with protection of fish habitat and water quality in new direction for riparian forests in Ontario: a case study of science guiding policy and practice. *Freshwater Science* 31: 248–257.
- NEARY, D. G., K. C. RYAN, AND L. F. DEBANO (EDITORS). 2008. *Wildland fire in ecosystems: effects of fire on soil and water*. General Technical Report RMRS-GTR-42-vol. 4. US Department of Agriculture, Forest Service, Rocky Mountain Research Station, Ogden, Utah.
- NITSCHKE, C. R. 2005. Does forest harvesting emulate fire disturbance? A comparison of effects on selected attributes in coniferous-dominated headwater systems. *Forest Ecology and Management* 214:305–319.
- OMNR (ONTARIO MINISTRY OF NATURAL RESOURCES). 2001. *Forest management guide for natural disturbance pattern emulation*. Version 3.1. Ontario Ministry of Natural Resources, Queen's Printer for Ontario, Toronto. (Available from: [http://www.web2.mnr.gov.on.ca/mnr/forests/forestdoc/ebr/guide/natural\\_dist/part%20one.pdf](http://www.web2.mnr.gov.on.ca/mnr/forests/forestdoc/ebr/guide/natural_dist/part%20one.pdf))
- OMNR (ONTARIO MINISTRY OF NATURAL RESOURCES). 2010. *Forest management guide for conserving biodiversity at the stand and site scales*. Queen's Printer for Ontario, Toronto, Ontario. (Available from: [www.publications.serviceontario.ca](http://www.publications.serviceontario.ca))
- PALIK, B., J. ZASADA, AND C. HEDMAN. 2000. Ecological considerations for riparian silviculture. Pages 233–254 in E. Verry, J. W. Hornbeck, and C. A. Dolloff (editors). *Riparian management in riparian forests of the continental eastern United States*. Lewis Publishers, CRC Press, Boca Raton, Florida.
- PERERA, A. H., AND L. J. BUSE. 2004. Emulating natural disturbance in forest management: an overview. Pages 3–7 in A. H. Perera, L. J. Buse, and M. G. Weber (editors). *Emulating natural forest landscape disturbances*. Columbia University Press, New York.
- PERERA, A. H., L. J. BUSE, AND M. G. WEBER (EDITORS). 2004. *Emulating natural forest landscape disturbances*. Columbia University Press, New York.
- PHILLIPS, M. J., L. W. SWIFT, AND C. R. BLINN. 2000. Best management practices for riparian areas. Pages 273–286 in E. S. Verry, J. W. Hornbeck, and C. A. Dolloff (editors). *Riparian management in forests of the continental eastern United States*. Lewis Publishers, Boca Raton, Florida.
- PREPAS, E. E., B. PINEL-ALLOUL, R. J. STEEDMAN, D. PLANAS, AND T. CHARETTE. 2003. Impacts of forest disturbance on boreal surface waters in Canada. Pages 369–393 in P. J. Burton, C. Messier, D. W. Smith, and W. L. Adamowicz (editors). *Towards sustainable management of the boreal forest*. NRC Research Press, Ottawa, Ontario.
- REICE, S. R. 1994. Nonequilibrium determinants of biological community structure. *American Scientist* 82:424–435.
- RICHARDSON, J. S. 2008. Aquatic arthropods and forestry: effects of large-scale land use on aquatic systems in Nearctic temperate regions. *Canadian Entomologist* 140: 495–509.
- RICHARDSON, J. S., AND R. J. DANEHY. 2007. A synthesis of the ecology of headwater streams and their riparian zones in temperate forests. *Forest Science* 53:131–147.
- RICHARDSON, J. S., R. J. NAIMAN, AND P. A. BISSON. 2012. How did fixed-width buffers become standard practice for protecting freshwaters and their riparian areas from forest harvest practices? *Freshwater Science* 31:232–238.
- RICHARDSON, J. S., AND R. M. THOMPSON. 2009. Setting conservation targets for freshwater ecosystems in forested catchments. Pages 244–263 in M. A. Villard and B. G. Jonsson (editors). *Setting conservation targets for managed forest landscapes*. Cambridge University Press, Cambridge, UK.

- ROOD, S. B. 2006. Unusual disturbance: forest change following a catastrophic debris flow in the Canadian Rocky Mountains. *Canadian Journal of Forest Research* 36:2204–2215.
- SCHILLING, E. 2009. Compendium of forestry best management practices for controlling nonpoint source pollution in North America. Technical Bulletin No. 966. National Council for Air and Stream Improvement, Research Triangle Park, North Carolina. (Available from: <http://www.ncasi.org/Publications/Detail.aspx?id=3204>)
- SEYMOUR, R. S., A. S. WHILE, AND P. G. DE MAYNADIER. 2002. Natural disturbance regimes in northeastern North America—evaluating silvicultural systems using natural scales and frequencies. *Forest Ecology and Management* 155:357–367.
- SIBLEY, P. K., AND A. M. GORDON. 2010. Managing riparian forests: a decision support system. Sustainable Forest Management Network, Edmonton, Alberta. (Available from: [http://www.ales.ualberta.ca/forestry/Sustainable\\_Forest\\_Management/Publications/~~/media/2CD412CA6C1441E2B7110CE698195A70.ashx](http://www.ales.ualberta.ca/forestry/Sustainable_Forest_Management/Publications/~~/media/2CD412CA6C1441E2B7110CE698195A70.ashx))
- SIBLEY, P. K., D. P. KREUTZWEISER, B. J. NAYLOR, J. S. RICHARDSON, AND A. M. GORDON. 2012. Emulation of natural disturbance (END) for riparian forest management: synthesis and recommendations. *Freshwater Science* 31:258–264.
- SOUSA, W. P. 1984. The role of disturbance in natural communities. *Annual Review of Ecology and Systematics* 15:353–391.
- SPENCER, C. N., K. O. GABEL, AND F. R. HAUER. 2003. Wildfire effects on stream food webs and nutrient dynamics in Glacier National Park, USA. *Forest Ecology and Management* 178:141–153.
- STEEDMAN, R. J., C. J. ALLAN, R. L. FRANCE, AND R. S. KUSHNERIUK. 2004. Land, water, and human activity in boreal watersheds. Pages 59–86 in J. M. Gunn, R. J. Steedman, and R. A. Ryder (editors). *Boreal Shield watersheds: lake trout ecosystems in a changing environment*. CRC Press, Boca Raton, Florida.
- STOVALL, J. P., W. S. KEETON, AND C. E. KRAFT. 2009. Late-successional riparian forest structure results in heterogeneous periphyton distributions in low-order streams. *Canadian Journal of Forest Research* 39:2343–2354.
- SUFFLING, R., AND A. H. PERERA. 2004. Characterizing natural forest disturbance regimes: concepts and approaches. Pages 43–54 in A. H. Perera, L. J. Buse, and M. G. Weber (editors). *Emulating natural forest landscape disturbances*. Columbia University Press, New York.
- SWANSON, M. E., J. F. FRANKLIN, R. L. BESCHTA, C. M. CRISAFULLI, D. A. DELLA SALA, R. L. HUTTO, D. B. LINDENMAYER, AND F. J. SWANSON. 2011. The forgotten stage of forest succession: early-successional ecosystems on forest sites. *Frontiers in Ecology and the Environment* 9:117–125.
- VOWELL, J. L., AND R. B. FRYDENBORG. 2004. A biological assessment of best management practice effectiveness during intensive silvicultural and forest chemical application. *Water, Air, and Soil Pollution: Focus* 4: 297–307.
- WEBSTER, J. R., S. W. GOLLADAY, E. F. BENFIELD, J. L. MEYER, W. T. SWANK, AND J. B. WALLACE. 1992. Catchment disturbance and stream responses: an overview of stream research at Coweeta Hydrologic Laboratory. Pages 231–253 in P. J. Boon, P. Calow, and G. E. Petts (editors). *River conservation and management*. John Wiley and Sons, Chichester, UK.
- WHITE, P. S., AND S. T. A. PICKETT. 1985. Natural disturbance and patch dynamics: an introduction. Pages 3–9 in S. T. A. Pickett and P. S. White (editors). *The ecology of natural disturbance and patch dynamics*. Academic Press, New York.

Received: 28 August 2011

Accepted: 10 November 2011