

Fibroblast Growth Factor May Regulate the Initiation of Oocyte Growth in the Developing Ovary of the Medaka, *Oryzias latipes*

Authors: Watanabe, Akihiko, Kobayashi, Emiko, Ogawa, Taku, and Onitake, Kazuo

Source: Zoological Science, 15(4) : 531-536

Published By: Zoological Society of Japan

URL: [https://doi.org/10.2108/0289-0003\(1998\)15\[531:FGFMRT\]2.0.CO;2](https://doi.org/10.2108/0289-0003(1998)15[531:FGFMRT]2.0.CO;2)

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Fibroblast Growth Factor May Regulate the Initiation of Oocyte Growth in the Developing Ovary of the Medaka, *Oryzias latipes*

Akihiko Watanabe*, Emiko Kobayashi, Taku Ogawa and Kazuo Onitake

Department of Biology, Faculty of Science, Yamagata University,
1-4-12 Kojirakawa-machi, Yamagata 990-8560, Japan

ABSTRACT—The distribution of fibroblast growth factor (FGF) was investigated in developing and matured ovaries of the medaka, *Oryzias latipes*. In the fry, FGF localized in the cytoplasmic region of all oocytes in the ovary at the pre-vitellogenic stage. Before the initiation of vitellogenesis, it disappeared in the cytoplasmic region and newly appeared around each oocyte, and then it localized around all oocytes in the ovary at the vitellogenic stage. Interestingly, the change in FGF distribution was orderly occurring from the posterior to anterior region of the ovary. In the adult, FGF was detected by immunofluorescence staining around the oocytes. These results suggest that FGF plays a significant role in the initiation of oocyte development through follicle cells, and the expression of FGF is rigidly regulated in the developing ovary of *O. latipes*.

INTRODUCTION

Oocyte development is rigidly regulated by the interaction between germ cells and somatic cells in vertebrates. It is initiated by the effect of gonadotropin from the pituitary gland and regulated by steroid hormone produced in the matured ovary.

In the teleostean ovary, oogenesis involves two sequential steps, the growth of the oocyte and the subsequent maturation, that are influenced by several kinds of hormones, as is true also in other vertebrates. During the growth phase, meiosis proceeds and is arrested at the diplotene stage of the first meiotic division in the oocyte. The egg membrane is formed around the oocyte and the yolk protein accumulates, causing the oocyte to increase in size. Estradiol-17 β is produced in the follicle cells in response to gonadotropin (Nagahama *et al.*, 1995), and the former hormone acts as the regulator of these events (Hamazaki *et al.*, 1989; Nagahama *et al.*, 1995).

In *Oryzias latipes*, germ cell development is initiated earlier in females than males in the developing gonad, and oocytes are seen in the ovary of the fry soon after they hatch out (Onitake, 1972; Satoh and Egami, 1972; Satoh, 1974). Oogenesis continuously occurs from oogonium even in the matured ovary. Sex-reversal can be induced in the fry of the gonad with steroid hormone (Yamamoto, 1958, 1975) as seen in many species of vertebrates (Jimenez *et al.*, 1993; Piferrer *et al.*, 1994; Ganesh and Raman, 1995). Then, the production of steroid hormone must be rigidly regulated in the developing ovary. It is reported that steroidogenesis occurs in granu-

losa cells in *O. latipes* (Onitake and Iwamatsu, 1986). However, the mechanism regulating steroidogenesis in the developing ovary is unknown.

Fibroblast growth factors (FGFs) are important regulators of growth and differentiation of the various types of cell in vertebrates. In mammals, it is known that basic FGF supports the proliferation and differentiation of the granulosa cells *in vitro* (Gospodarowicz *et al.*, 1977; Gospodarowicz and Bialecki, 1978; Lavranos *et al.*, 1994). Thus, it seems likely that basic FGF might be one of the regulators of follicular morphogenesis, acting through granulosa cells (Gospodarowicz, 1990).

In this study, to examine the participation and the mechanism of FGF in controlling oocyte development in the developing ovary of teleost, we investigated the distribution of FGF in the developing ovary of the medaka, *Oryzias latipes*, by immunofluorescence staining.

MATERIALS AND METHODS

Animals

Adult or immature specimens of the medaka, *Oryzias latipes*, were purchased from Morikawa Fish Farm (Yamato-koriyama, Nara, Japan). Immature fish were obtained from the fertilized eggs prepared in our laboratory. They were kept at 26°C in our laboratory before use in experiments.

Histology

The ovaries dissected out from the bodies of the adult and immature fish, which were 7–12 mm in body length, were fixed in Bouin's fixative. Sections of 6 μ m thickness were prepared and stained in Delafield's hematoxylin and eosin solution.

* Corresponding author: Tel. +81-23-628-4619;
FAX. +81-23-628-4617.

Immunofluorescence staining

At least five individuals were used for each experiment. The methods used were the same as those previously reported (Watanabe *et al.*, 1997). In brief, ovaries from the adult or immature fish, 7–12 mm in body length, were fixed in 2% paraformaldehyde in PBS (0.05 M phosphate buffer, pH 7.4) at 4°C for 2 hr, and then they were immersed in 30% sucrose in PBS at 4°C for 1.5 hr. They were embedded in O.C.T. compound (Miles) and frozen in liquid nitrogen. Frozen sections of 8 µm in thickness were serially prepared. They were reacted with a human FGF2-specific monoclonal antibody (described as FGF antibody) (Reilly *et al.*, 1989) at room temperature for 30 min, and then with the FITC-labeled antibodies raised in the goat against mouse IgG (diluted 1:30 in PBS that contained 3% BSA; MBL) at room temperature for 40 min.

In controls, two types of experiments were pursued. One was that 100 ng of bovine FGF2 (Austral Biological) was added to 200 µl of the solution containing the FGF2 specific antibody, and then used for the first immunoreaction. Another was that the chick-specific monoclonal antibody (Watanabe *et al.*, 1993) or PBS that did not include any antibody was used for the first immunoreaction instead of the FGF2-specific monoclonal antibody. The sections were observed with a fluorescence microscope (BH-RFK, Olympus).

RESULTS

Distribution of FGF in the developing ovary of *O. latipes*

In the immature fish, 7–8.5 mm in body length, the ovary was localized between the mesonephric tubule and gut in the dorsal-posterior region of the abdominal cavity (Fig. 1a, b). The anterior region of the ovary bound to peritoneum under

the air-bladder. Many oocytes in the ovaries were at the pachytene or early diplotene stage of meiotic prophase (Fig. 2a). The size of each oocyte was about 30 µm, which suggested that they were at pre-vitellogenic stage. Follicle cells were seen but not developed. Strong fluorescence was detected in the ovary and muscle by the immunofluorescence staining with the anti-FGF antibodies (Fig. 2b). In the ovary, it was localized in the cytoplasmic region of all oocytes.

In control, auto-fluorescence was seen in the nucleus of somatic cells including follicle cells and muscle (Fig. 2c). It was easily distinguishable from that of FITC by the morphology and the difference in wavelength for excitation. No signal was detected in any oocyte or follicle cell.

In the immature fish of 9.0 mm in body length, the morphological features of the oocytes and follicle cells in the ovary were similar to those in the younger fry. Strong fluorescence was detected in the ovary (Fig. 3a, b). It localized to the cytoplasmic regions of each oocyte in the anterior region of the ovaries (Fig. 3d). Little fluorescence was detected at the sites of follicle cells. This pattern was the same as that seen in the individuals of less than 8.5 mm in body length. However, in the posterior region of the same ovaries, strong fluorescence was seen in the follicle cell layer but not in the oocytes (Fig. 3e). The localization pattern was just inverted compared to that of the anterior region of the same ovary. In control, significant fluorescence was not detected (Fig. 3c).




Fig. 1. Sections of the ovary of *O. latipes*. Six µm paraffin sections of the body of the immature fish (a, b), and the ovary of the adult (c, d) were cut and stained with hematoxylin-eosin solution. (a) A section of the immature fish of 8.5 mm in length. Arrow indicates the ovary. The size of each oocyte was similar. (b) A section of the immature fish of 12 mm in length. Arrow indicates the ovary. The number of oocytes was increased and the size of each oocyte varied compared with (a). (c) Many oocytes at different stages were observed in the ovary. (d) High-magnification view of (c). I, II, III, and IV indicate the developmental stage of each oocyte (Iwamatsu *et al.*, 1988). g, gut; m, mesonephric tubule. Bar: 200 µm (c), and 50 µm (a, d).




Fig. 2. Immunofluorescence staining of the ovary of fry, 8.5 mm in length, of *O. latipes* with the monoclonal antibody against FGF protein. (a) A six µm transverse section was stained with hematoxylin-eosin solution. (b) An eight µm frozen section of the fry was stained. The cytoplasmic regions of each oocyte and muscle were strongly stained. (c) Control. Bovine FGF2 was added to the solution of FGF-specific antibody and then, used in the first immunoreaction. Auto-fluorescence was seen in the nucleus of somatic cells. Arrows indicate the ovary. g, gut; m, mesonephric tubule; mu, muscle. Bar: 50 µm.

In the immature fish of more than 10–12 mm in body length, many oocytes were observed in the ovary (Fig. 1b). The size of these oocytes varied (Figs. 1b, 4a), indicating that some had begun to undergo vitellogenesis. Fluorescence was detected not in the oocyte but at the site of follicle cells (Fig. 4b). No difference was seen in the localization pattern of fluorescence. In control, significant fluorescence was not detected

(Fig. 4c).

Distribution of FGF in the matured ovary of *O. latipes*

In the ovary of the adult medaka, oocytes at various stages of oogenesis were seen (Fig. 1c, d). Growing oocytes were surrounded by granulosa cells and theca cells. Strong fluorescence was detected around developing oocytes at all stages by the immunofluorescence staining with the anti-FGF antibodies (Fig. 5a). These regions corresponded to the sites of follicle cells. In control, no signal was detected in or around the oocytes (Fig. 5b).

Specificity of the antibodies against FGF

To examine the specificity of the fluorescence staining with the antibodies against human FGF2, the binding of the antibodies to the antigen was blocked with bovine FGF2 in the first immunoreaction. No fluorescence of specific staining was seen in the ovarian sections of the immature (Figs. 2, 4) and the mature fish (Fig. 5b). The results were quite similar as for control.

DISCUSSION

FGFs are regulating factors of the proliferation and/or differentiation of various types of cell in vertebrates (Gospodarowicz, 1990). In this study, we used the immunostaining technique with a monoclonal antibody against human basic FGF for the distribution of FGF in the developing ovary of the medaka. Human FGF2 protein is effective in other species (Dawid *et al.*, 1990; Niswander *et al.*, 1993). In this study, the first immunoreaction was inhibited by the addition of bovine FGF2 to the solution containing the monoclonal antibody against human FGF2 (Fig. 2b). It has been reported that the antibody can neutralize the effect of basic FGF (Reilly *et al.*, 1989). Recently a cDNA fragment of the medaka was isolated with this antibody as a probe in our laboratory. It codes a functional domain of basic and acidic FGF for the binding to the receptor that includes a cysteine residue conserved among species (unpublished data). These findings suggest that the antigen stained with anti-FGF antibody in this study was a homologue of FGF in the medaka.

The change in FGF distribution in the developing ovary of the medaka

FGF was localized in the cytoplasmic region of oocytes of the fries, 8.5 mm in body length (Fig. 2). It disappeared in the oocyte cytoplasm of fry, 9 mm in body length, and synchronously appeared in follicle cells (Fig. 3). In teleosts, follicle cells appear at the early stage of development of the ovary (Sato, 1974; Nakamura and Nagahama, 1985; Nakamura *et al.*, 1993). In the fry of less than 9 mm, the oocytes were nearly equal in diameter, as shown in the present study. This indicates that all oocytes were at the pre-vitellogenic stage. And then, some of the oocytes enlarged as the fry grew. These results suggest that FGF acts as a paracrine factor on follicle cells and regulates the initiation of vitellogenesis in the devel-




Fig. 3. Sections of the ovary of the fry, 9.0 mm in length, of *O. latipes*. An eight μ m frozen section of the immature fish was stained with the monoclonal antibody against FGF. (a) The section including the anterior portion of the ovary. An air-bladder (asterisk) and a pair of guts, the features of the anterior region, were sectioned. (b) The section including the posterior portion of the same ovary of (a). Air-bladder was not seen, and gut was sectioned alone. (c) Control. The FGF-specific antibody was not included in the immunoreaction. Auto-fluorescence was seen in mesonephric tubules and gut. No significant fluorescence was seen in the ovary and muscle. (d) and (e) High magnification view of (a) and (b). Strong fluorescence was specifically seen in the cytoplasmic region of each oocyte (d) and the somatic cells around them (e). Arrows indicate the ovary. g, gut; m, mesonephric tubule; mu, muscle. Bar: 200 μ m (a) and 50 μ m (d).

opening ovary of the medaka.

In this study, the change in FGF localization occurred along the anterior-posterior axis in the developing ovary (Fig. 3). It occurred in all oocytes in the ovary before oocytes are not developed. This result suggests that the expression of FGF is regulated independently of oocyte development in the developing ovary. However, the mechanism regulating it is unknown.

In teleosts, vitellogenesis is regulated by estradiol-17 β produced by follicle cells (Nagahama *et al.*, 1995). FGF may activate steroidogenesis for the initiation of vitellogenesis in the developing ovary of the medaka. It has been reported that sex-reversal could artificially take place in the gonad of the fry of *O. latipes* by the stimulation of steroid hormone (Yamamoto, 1958, 1975). In female, it is induced before the period when steroidogenesis is initiated (Onitake, 1972; Satoh and Egami, 1972; Yamamoto, 1975). The period of loss of the response to sex-reversal corresponds with the period when FGF disappeared in the oocytes. From these points, it is speculated that change in FGF distribution correlates with sex determination of germ cells in the developing gonad. The process of sex

determination is not well-known in teleost and FGF may be a useful marker for the analysis of it.

The possible role of FGF in follicle cells of the medaka

In the teleostean ovary, the follicle consists of the oocyte and two layers of follicle cells. During oocyte growth, the oocyte increases in size with the accumulation of yolk protein, and a thick egg membrane is formed between oocyte and follicle cells. It is known that the follicle cells, both granulosa cells and theca cells, are indispensable for the regulation of these events since estradiol-17 β is synthesized in them in response to gonadotropin (Nagahama *et al.*, 1995). In medaka, it has been reported that steroidogenesis occurs only in granulosa cells (Onitake and Iwamatsu, 1986).

In this study, FGF was found around the oocytes in the ovary of the adult medaka. The distribution of FGF corresponded to the site of follicle cells. In mammal, basic FGF can support the proliferation of granulosa cells in the bovine follicles *in vitro* (Gospodarowicz *et al.*, 1977; Lavranos *et al.*, 1994). It is also known that basic FGF can affect the differentiation of granulosa cells *in vitro* (Gospodarowicz and Bialecki,




Fig. 4. Immunofluorescence staining of the ovary of fry, 12 mm in length, of *O. latipes* with the monoclonal antibody against FGF protein. (a) A six μm section was stained with hematoxylin-eosin solution. (b) An eight μm frozen section of the fry was stained. Strong fluorescence was seen around each oocyte. (c) Control. Bovine FGF2 was added to the solution of FGF-specific antibody and then, used in the first immunoreaction. Auto-fluorescence was seen in the nucleus of somatic cells. Bar: 50 μm .

1978). Furthermore, very electron-dense lipid granules can be seen in granulosa cells cultured under the influence of FGF (Gospodarowicz *et al.*, 1977). These results suggest that FGF acts as a regulator of the proliferation and differentiation of those cells in the medaka.

Steroidogenic cells first appear in the ovaries of fish of 7–9 mm in body length (Kanamori *et al.*, 1985). In this study, FGF was first seen at the sites of follicle cells at the same

period and continued to localize to the same region (Fig. 4). This suggests that FGF is important for steroidogenesis in follicle cells. The steroidogenic activities of follicle cells increase as oocyte growth proceeds (Nagahama *et al.*, 1995). However, the distribution of FGF was not specific for the stages of developing oocytes (Fig. 5). It is reported that TGF- β is produced in the theca cells and stimulates the synthesis of estradiol-17 β in the granulosa cells *in vitro* in the bovine (Skinner *et al.*, 1987). It seems likely that some other factors are cooperate with FGF to regulate steroidogenesis in follicle cells during oocyte growth.

ACKNOWLEDGMENTS

We thank Dr. Reilly, T. and Dr. Walton, H. of E. I. du Pont de Nemours and Company for the gift of the monoclonal antibody against human FGF2 protein. We also thank Dr. Nagahama, Y. of National Institute for Basic Biology for his valuable encouragement.

REFERENCES

- Dawid IB, Sargent TD, Rosa F (1990) The role of growth factors in embryonic induction in amphibians. In "Growth Factors and Development" Ed by M Nilsen-Hamilton, Academic Press, San Diego, pp 261–288
- Ganesh S, Raman R (1995) Sex reversal by testosterone and not by estradiol or temperature in *Calotes versicolor*, the lizard lacking sex chromosomes. *J Exp Zool* 271: 139–144
- Gospodarowicz D (1990) Fibroblast growth factor and its involvement in developmental process. In "Growth Factors and Development" Ed by M Nilsen-Hamilton, Academic Press, San Diego, pp 57–93
- Gospodarowicz D, Ill CR, Birdwell CR (1977) Effects of fibroblast and Epidermal growth factors on ovarian cell proliferation *in vitro*. 1. Characterization of the response of granulosa cells to FGF and EGF. *Endocrinology* 100: 1108–1120
- Gospodarowicz D, Bialecki H (1978) The effects of epidermal and fibroblast growth factors on replicative lifespan of cultured bovine granulosa cells. *Endocrinology* 103: 854–865
- Hamazaki TS, Nagahama Y, Iuchi I, Yamazaki K (1989) A glycoprotein from the liver constitutes the inner layer of the egg envelope (zona pellucida) of the fish, *Oryzias latipes*. *Dev Biol* 133: 101–110
- Iwamatsu Y, Ohta T, Oshima E, Sakai N (1988). Oogenesis in the medaka *Oryzias latipes*—Stages of oocyte development. *Zool Sci* 5: 349–369
- Jimenez R, Burgos M, Sanchez A, Sinclair AH, Alarcon FJ, Marin JJ, Ortega E, Guardia RD (1993) Fertile female of the mole *Talpa occidentalis* are phenotypic intersexes with ovotestes. *Development* 118: 1303–1311
- Kanamori A, Nagahama Y, Egami N (1985) Development of the tissue architecture in the gonads of the medaka *Oryzias latipes*. *Zool Sci* 2: 695–706
- Lavranos TC, Rodgers HF, Bertoncello I, Rodgers RJ (1994) Anchorage-independent culture of bovine granulosa cells: the effects of basic fibroblast growth factor and dibutyryl cAMP on cell division and differentiation. *Exp Cell Res* 211: 245–251
- Nagahama Y, Yoshikuni M, Yamashita M, Tokumoto T, Katsu Y (1995) Regulation of oocyte growth and maturation in fish. In "Current Topics in Developmental Biology Vol 30" Ed by RA Pedersen, GP Schatten, Academic Press, New York, pp 103–145
- Nakamura M, Nagahama Y (1985) Steroid producing cells during ovarian differentiation of the tilapia, *Sarotherodon niloticus*. *Development Growth Differ* 27: 701–708




Fig. 5. Immunofluorescence staining of the ovary of mature fish of *O. latipes* with the monoclonal antibody against FGF protein. (a) An eight μm frozen section of the ovary of the mature fish was stained. (b) Control. The FGF-specific antibody was not included in the immunoreaction. Auto-fluorescence was seen in haemocytes and the nucleus of follicle cells. Bar: 200 μm .

- Nakamura M, Specker JL, Nagahama Y (1993) Ultrastructural analysis of the developing follicle during early vitellogenesis in the tilapia, *Oreochromis niloticus*, with special reference to the steroid-producing cells. *Cell Tissue Res* 272: 33–39
- Niswander L, Tickle C, Vogel A, Booth I, Martin GR (1993) FGF-4 replaces the apical ectodermal ridge and directs outgrowth and patterning of the limb. *Cell* 75: 579–587
- Onitake K (1972) Morphological studies of normal sex-differentiation and induced sex-reversal process of gonads in the medaka, *Oryzias latipes*. *Annot Zool Japon* 45: 159–169
- Onitake K, Iwamatsu T (1986) Immunocytochemical demonstration of steroid hormones in the granulosa cells of the medaka, *Oryzias latipes*. *J Exp Zool* 239: 97–103
- Piferrer F, Zanuy S, Carrillo M, Solar I, Devlin RH, Donaldson EM (1994) Brief treatment with an aromatase inhibitor during sex differentiation causes chromosomally female salmon to develop as normal, functional males. *J Exp Zool* 270: 255–262
- Reilly TM, Taylor DS, Herblin WF, Thoolen MJ, Chiu AT, Watson DW, Timmermans PBMW (1989) Monoclonal antibodies directed against basic fibroblast growth factor which inhibits its biological activity *in vitro* and *in vivo*. *Biochem Biophys Res Commun* 164: 736–743
- Satoh N (1974) An ultrastructural study of sex differentiation in the teleost *Oryzias latipes*. *J Embryol Exp Morph* 32: 195–215
- Satoh N, Egami N (1972) Sex differentiation of germ cells in the teleost, *Oryzias latipes*, during normal embryonic development. *J Embryol Exp Morph* 28: 385–395
- Skinner MK, Keski-Oja J, Osteen KG, Moses HL (1987) Ovarian thecal cells produce transforming growth factor-beta which can regulate granulosa cell growth. *Endocrinology* 121: 786–792
- Watanabe A, Ohsugi K, Ide H (1993) Formation of distal structures from stumps of chick wing buds at stages 24–25 following the grafting of quail tissue from X-irradiated distal limb buds. *J Exp Zool* 267: 447–453
- Watanabe A, Hatakeyama N, Yasuoka A, Onitake K (1997) The distribution of fibroblast growth factor and the mRNA for its receptor, *MFR1*, in the developing testis of the medaka, *Oryzias latipes*. *J Exp Zool* 279: 177–184
- Yamamoto T (1958) Artificial induction of functional sex-reversal in genotypic female of the medaka (*Oryzias latipes*). *J Exp Zool* 137: 227–264
- Yamamoto, T (1975) *Medaka (Killifish). Biology and Strains*. Keigaku Publ Co, Tokyo

(Received January 12, 1998 / Accepted April 13, 1998)