



Extensive Distribution of Galanin-Immunoreactive Neuronal Systems in a Fresh Water Snail, *Indoplanorbis exustus*, and Partial Coexistence with Choline Acetyltransferase-like Immunoreactivity

Authors: Yamamoto, Toshiharu, Suzuki, Hirohumi, and Uemura, Haruko

Source: Zoological Science, 13(4) : 577-585

Published By: Zoological Society of Japan

URL: <https://doi.org/10.2108/zsj.13.577>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Extensive Distribution of Galanin-Immunoreactive Neuronal Systems in a Fresh Water Snail, *Indoplanorbis exustus*, and Partial Coexistence with Choline Acetyltransferase-like Immunoreactivity

Toshiharu Yamamoto*, Hirohumi Suzuki and Haruko Uemura

Department of Biology, Kanagawa Dental College, Yokosuka 238, Japan

ABSTRACT—Immunohistochemical techniques were employed to investigate the distribution of galanin-like immunoreactivity in the central nervous system (CNS) of the snail, *Indoplanorbis exustus*, and immunoreactivity with an antibody against choline acetyltransferase (CAT) in galanin-immunoreactive somata was also examined. Galanin-immunoreactive (Gal-IR) cells were observed in all ganglia. They were classified into large (50 μm), medium (21 μm), and small (13 μm)-sized neurons and small Gal-IR cells dominated over large and medium Gal-IR cells in number. Gal-IR cells were most abundant in the cerebral ganglion. Several immunoreactive somata were seen in the parietal, visceral, and pedal ganglia, and only a few Gal-IR cells in the pleural and buccal ganglia. Densely arranged Gal-IR fiber bundles were observed in the cerebro-pleural, parieto-pleural and bucco-cerebral connectives. Of the peripheral nerves, the pallial and pharyngeal nerves contained more numerous Gal-IR fibers than other peripheral nerves. By Western blot analysis, the galanin antibody detected approximately 6.1 kDa band, suggesting a pre-form or a long form of galanin. Some Gal-IR cells in the cerebral ganglion and a few in the pleural and parietal ganglia were also immunoreactive for CAT. These results indicate that a distinct neuronal system with galanin or galanin-like peptide is present in the CNS of the snail, and suggest that a part of the galanin neuronal system is cholinergic. The peptide may function as a neurotransmitter/neuromodulator and/or neurohormone in the snail.

INTRODUCTION

Galanin is a biologically active peptide consisting of 29 amino acid residues and was originally isolated from the small intestine of pigs (Tatemoto *et al.*, 1983). Relatively minor variations in amino acid sequences are known among various species of vertebrates (rat, Vrontakis *et al.*, 1987; Kaplan *et al.*, 1988; bovine, Rökæus and Carlquist, 1988; chicken, Norberg *et al.*, 1991). Human galanin appears to be unique, comprising 30 amino acids possessing an additional nonamidated serine residue as its C-terminus (Evans and Shine, 1991; Bersani *et al.*, 1991; Schmidt *et al.*, 1991). In invertebrates, galanin was biochemically studied only in blowfly (*Phormia terraenovae*), and its galanin appears to be highly alkaline in nature but similar in size to those of vertebrates (Lundquist *et al.*, 1991).

The neuroanatomical distribution of galanin has been well documented in the rat (Skofitsch and Jacobowitz, 1985; Melander *et al.*, 1986), primates (Kordower and Mufson, 1990), opossum (Elmqvist *et al.*, 1992), amphibians (Olivereau and Olivereau, 1992), and fish (Olivereau and Olivereau, 1991; Yamamoto *et al.*, 1992). Galanin-immunoreactive (Gal-IR)

somata and fibers were widely distributed in the vertebrate central nervous system (CNS), especially in the hypothalamus, Gal-IR fibers being densely located in the median eminence. These profiles suggest that galanin may act as a neurotransmitter, a neuromodulator, or a neurohumoral peptide (Hökfelt *et al.*, 1987; Bartfai *et al.*, 1993). It is of interest that galanin coexists with various neurotransmitter markers and bioactive peptides, including cholinergic markers in the basal forebrain (Melander *et al.*, 1985; Melander and Staines, 1986), tyrosine hydroxylase in the arcuate nucleus (Everitt *et al.*, 1986), adenosine deaminase in the tuberomammillary nucleus (Staines *et al.*, 1986), luteinizing hormone releasing hormone in the diagonal band of Broca (Merchantaler *et al.*, 1991), dopamine in the median eminence (Nordström *et al.*, 1987), and vasopressin in the supraoptic and paraventricular nuclei (Rökæus *et al.*, 1988; Gaymann and Martin, 1989; Skofitsch *et al.*, 1989).

The snail CNS has been shown to be a good model for neurobiological studies because of its simplicity and readily accessible localization of neurons (Boer and van Minnen, 1988). However, only *Bulla gouldiana*, a marine gastropod, was studied in regard to the distribution of galanin-like immunoreactivity in the neuronal system (Roberts *et al.*, 1989). This prompted us to investigate in detail galanin-containing

* To whom correspondence should be addressed.

neuronal systems in a fresh water snail, *Indoplanorbis exustus*, which has long hair-like tentacles and this morphological characteristic may offer good opportunities for analyzing the tentacle-withdrawal reflex. In fact, a closely related species, *Planorbarius (Planorbis) corneus*, has been utilized to study an experimental-learning paradigm involving the tentacle-withdrawal reflex (Biral *et al.*, 1978a, b). In the present study, we examined galanin-like immunoreactivity in the CNS of the snail and demonstrated its colocalization with choline acetyltransferase (CAT)-like immunoreactivity in a subpopulation of Gal-IR cells of the cerebral ganglion. Furthermore, we analyzed extracts of the CNS by Western blots.

MATERIALS AND METHODS

Indoplanorbis exustus were anesthetized with ethyl m-aminobenzoate methanesulfonate (Nacalai Tesque, Inc., Kyoto, Japan) and the CNS were carefully dissected out with the aid of a stereoscopic microscope. The CNS of six animals were removed with a part of the pharynx and fixed for 16 hr in Bouin-Hollande solution at room temperature. After washing in distilled water, the tissue was dehydrated and wax-embedded according to the conventional method. Horizontal sections were cut at a thickness of 5 μm , and serial sections at intervals of 20 μm were mounted onto slides for galanin immunohistochemistry. Consecutive sections were used for comparison of galanin and CAT immunoreactivities.

Sections on slides were dewaxed in xylene and gradually hydrated with graded alcohol solutions. The sections were then washed overnight in 0.1 M sodium phosphate buffer containing 0.9 % saline (PBS) at pH 7.4 and incubated for 48 hr at 4°C with rabbit anti-rat galanin serum (Peptide Institute, Inc., Osaka, Japan) diluted in PBS containing 0.05% Triton X-100 (PBST) at 1:2,000. After washing in PBST, the sections were incubated for 1 hr at room temperature with biotinylated anti-rabbit IgG (Vector Laboratories, Berlingame, USA) diluted 1:100 in PBST. The sections were then washed again in PBST and incubated for 1 hr at room temperature with avidin-biotin-horseradish peroxidase complex (Vector Laboratories) diluted 1:100 in PBST. After final wash in PBST, the sections were reacted with a 0.05 M Tris-HCl buffer solution (pH 7.4) containing 0.02% 3,3'-diaminobenzidine hydrochloride (DAB) and 0.005% hydrogen peroxide, before being dehydrated, cleared in xylene, and coverslipped with Malinol (Muto Pure Chemicals Ltd., Tokyo, Japan). Alternate sections were immunohistochemically stained with anti-galanin and goat anti-human placental CAT antibodies (1:5,000; Chemicon International Inc., Temecula, USA). Immunohistochemical controls were carried out by omission of the antiserum in the first incubation, or by incubation with the normal rabbit or the goat serum, or with antiserum preabsorbed with synthetic rat galanin (10 $\mu\text{g}/\text{ml}$; Peptide Institute, Inc.) as the primary antibody. Some adjacent sections were stained with Gomori's paraldehyde-fuchsin. Some immunohistochemically-stained sections were lightly counterstained for 1 min with 0.05% thionine solution buffered with 0.02 M acetic buffer at pH 4.0 for observation of general histology.

For block-staining, the CNS was fixed for 24 hr in PB (pH 6.9) containing 4% paraformaldehyde and 0.2% picric acid and kept for 1 week in PBS containing 0.3% Triton X-100 with several changes of the solution. Then the tissue was incubated for 48 hr at 4°C with anti-galanin serum. After washing with PBST, the tissue was incubated for 24 hr at 4°C with dichlorotriazinyl aminofluorescence-conjugated anti-rabbit IgG (1:100; Chemicon International Inc.), and cleared in anti-fading medium (Valnes and Brandtzaeg, 1985).

For Western blots, CNSs from 20 snails (about 28 mg wet weight) were homogenized in 140 μl of 0.05 M Tris-HCl buffer (pH 7.4)

containing 140 units of aprotinin (a protease inhibitor) with a microhomogenizer at 4°C. Then the homogenates were centrifuged at 10,000 $\times g$ for 1 hr at 4°C. Equivalent volume of Laemmli's sample buffer (Laemmli, 1970) was added to the supernatant. These samples were analyzed by sodium dodecyl sulfate-polyacrylamid gel (15%) electrophoresis (SDS-PAGE). After SDS-PAGE, separated proteins were transferred to Clear Blot Membrane-p (Atto Corporation, Tokyo, Japan) and probed with anti-rat galanin antibody (1:2,000) after blocking non-specific protein binding with PBS containing 3% bovine serum albumin (BSA). Probed membranes were then incubated with biotinylated anti-rabbit IgG (1:1,000) and avidin-biotin-horseradish peroxidase complex (1:1,000) diluted in PBS containing 3% BSA, and finally reacted with DAB solution containing 0.005% hydrogen peroxide and 0.6% ammonium nickel sulfate for 5-10 min. Rat galanin (Peptide Institute, Inc.) and molecular marker proteins (cytochrome *c*, monomer to hexamer; Oriental East Co., Ltd., Osaka, Japan) were also electrophoresed simultaneously.

RESULTS AND DISCUSSION

General anatomy of the CNS

For general neuroanatomy of the snail, we referred to the description of *Planorbarius corneus* (Sonetti *et al.*, 1990) and *Lymnaea stagnalis* (Bonga, 1970). The CNS of the snail, *Indoplanorbis exustus*, was observed to be composed of five paired ganglia (cerebral, pleural, parietal, pedal, and buccal ganglia) and an unpaired visceral ganglion, being basically same to the CNS of *Planorbarius corneus* (Sonetti *et al.*, 1990). The dorsal body was located anterodorsally on the cerebral commissure in a single mass, not in two masses as in *Lymnaea*. The cerebral ganglion was large and lunar in shape with a small lateral lobe, while the pleural and buccal ganglia were relatively small and round. The left pleural ganglion was situated under the left parietal ganglion, and right pleural ganglion was formed into ring-shape with the cerebral, visceral, and left parietal ganglia. The left parietal ganglion was larger than the right one, in contrast with the larger right parietal ganglion in *Lymnaea*. The pedal and visceral ganglia were relatively large and slightly flattened.

Immunoreactive somata

Gal-IR cells were seen in all ganglia (Fig. 1). Their somata were round, ovoid, or sometimes triangular in shape. Almost all Gal-IR neurons were unipolar with a process directing to the center of the ganglion. They were classified into three types based on their size; i.e., small (7.5-17.5 μm ; mean, 12.7 μm in diameter), medium (18.7-25.0 μm ; mean, 21.1 μm in diameter), and large (38.4-63.7 μm ; mean, 50.3 μm in diameter) neurons. Large and medium Gal-IR cells were a minor population and were located on the peripheral region of the ganglion. On the other hand, small Gal-IR cells were abundant and tended to be situated in the inner area of the ganglion.

Gal-IR cells were seen most abundantly in the cerebral ganglion (Figs. 1 and 2A-C). The cerebral ganglion contained lateral, medial, and anterior Gal-IR cell groups. A few medium and many small Gal-IR cells were seen in each cell group, while only one large Gal-IR cell was observed in the medial Gal-IR cell group of the ganglion in each side (Fig. 1). A few




Fig. 1. Schematic diagram of the dorsal view of the CNS of the snail, *Indoplanorbis exustus* showing the localization of Gal-IR cells and fibers. Asterisks, squares, and circles represent large, medium and small Gal-IR cells, respectively. The pedal ganglion (PE) is dislocated for presentation. Abbreviations: BU, buccal ganglion; CE, cerebral ganglion; DB, dorsal body; LL, lateral lobe of the cerebral ganglion; PA, parietal ganglion; pc, pedal commissure; PL, pleural ganglion; VI, visceral ganglion; 1, tentacular nerve (n); 2, frontal lip n; 3, median lip n; 4, inferior pedal n; 5, median pedal n; 6, superior pedal n; 7, pallial n; 8, intestinal n; 9, anal n; 10, pharyngeal n; 11, gastric n; 12, radular n. Scale bar: about 200 μm .

medium Gal-IR cells were seen in the lateral lobe (Fig. 1). One large Gal-IR cell was seen in the medial portion of the left and right parietal ganglia (Figs. 1 and 3A). A few medium and small Gal-IR cells were located in the left parietal ganglion, while the right ganglion contained fewer of both medium and small Gal-IR cells (Figs. 1 and 3A, B). Just a few small Gal-IR cells were seen in the pleural ganglion. In the pedal ganglion, one large and several small Gal-IR cells were seen (Figs. 1 and 3C, D). One large, a few medium and small Gal-IR cells were observed in the visceral ganglion (Figs. 1 and 4C). In the buccal ganglion, a few medium Gal-IR cells were detected on both sides (Figs. 1 and 5C) but no large and small Gal-IR cells were found in this ganglion.

One large Gal-IR cell was observed in each cerebral, parietal, pedal, and visceral ganglion. The number of these cells in individual ganglion could not be underestimated, because the size of large Gal-IR cells was approximately 50 μm and we counted the cell number in 5 μm -thick sections at intervals of 20 μm .

Immunoreactive fibers

Fine Gal-IR fibers were seen in the centrally located neuropil in the each ganglion and in almost all peripheral nerves. In the intercerebral commissure, a few Gal-IR fiber bundles were seen (Figs. 1, 2A and 4A). The tentacular and lip nerves from the cerebral ganglion contained massive Gal-IR fibers (Figs. 1 and 4A). Most immunoreactive fibers from the medial Gal-IR cell group of the cerebral ganglion traveled to the ipsilateral parietal ganglion through the pleural ganglion (Figs. 1, 3A and 4A). Many Gal-IR fibers were observed in the

pallial nerve from the parietal ganglion (Figs. 1, 3B and 4A). Relatively abundant Gal-IR fibers were seen in the intestinal nerve from the visceral ganglion (Fig. 4C). A few Gal-IR fibers were seen in the pedal commissure, and thick fiber bundles were seen in the neuropil of the pedal ganglion (Fig. 4A). Among three main nerves from the pedal ganglion, the inferior pedal nerve contained most abundant Gal-IR fibers (Figs. 1 and 4B). A considerable number of immunoreactive fibers were seen in the bucco-cerebral connective nerve (Fig. 1), and the pharyngeal nerve from the buccal ganglion contained many Gal-IR fibers (Fig. 5).

These immunostaining profiles were abolished by preabsorption of antiserum with synthetic rat galanin, and no immunoreactivity was observed in the sections processed with normal rabbit serum or with omission of the primary antibody (data not shown).

Many bioactive peptides discovered originally in the mammalian organs were also present in the CNS of gastropods (Boer *et al.*, 1979; Schot *et al.*, 1981; Roberts *et al.*, 1989; Sonetti *et al.*, 1990; Kasuya *et al.*, 1991; Honma *et al.*, 1994). Furthermore, FMRFamide, one of bioactive peptides first isolated from a clam *Macrocallista nimbosa* (Price and Greenberg, 1977), was immunohistochemically demonstrated to be expressed in the gastropoda CNS in a significant population of cells (*Lymnaea*, Schot and Boer, 1982; Schot *et al.*, 1984; *Aplysia*, Schaefer *et al.*, 1985). Compared with the relative number of FMRFamide-immunoreactive cells in these species, the Gal-IR cell population is minor in *Indoplanorbis*, but still is a distinct neuronal system. A considerable number of Gal-IR neurons and fibers were found in the CNS of




Fig. 2. Photomicrographs showing Gal-IR somata and fibers in the cerebral ganglion. A: Photomontage showing the anterior Gal-IR cell group of cerebral ganglion (CE) and fibers in the cerebral commissure (cc). A large arrow, small arrows, and arrowheads indicate a medium Gal-IR cell, small Gal-IR cells and Gal-IR fibers, respectively. B,C: Low (B) and high (C) magnifications showing Gal-IR cells in the lateral Gal-IR cell group of the cerebral ganglion (CE). Weakly stained medium neurons (large arrows) are located in the peripheral region of the ganglion and intensely stained small neurons (small arrows) are relatively centrally situated. No large Gal-IR cell is located in this area. B and C are lightly counterstained with thionine. Abbreviation: LL, lateral lobe. Scale bars: A, 100 μm ; B, 50 μm ; C, 20 μm .

Indoplanorbis exustus. On the other hand, in the CNS of *Bulla gouldiana*, a marine gastropod, a few Gal-IR neurons, but almost no Gal-IR fibers, were detected only in the cerebral ganglion (Roberts *et al.*, 1989). This discrepancy may have been caused by different antiserum used, species difference and/or different habitats of these two species; *Indoplanorbis exustus* is a fresh water snail and *Bulla gouldiana* a marine gastropod. The antiserum used here was against rat galanin, but Roberts *et al.* (1989) used an antiserum against porcine galanin. Between these two kinds of galanin, three among 29 amino acid residues are different (Tatemoto *et al.*, 1983; Vrontakis *et al.*, 1987).

Western blot analysis

The anti-rat galanin antibody used here detected approximately 6.1 kDa band by Western blots of the supernatant from the homogenates of the snail CNS (Fig. 6).

This size was larger than that of rat galanin itself (3.2 kDa). This suggests that snail galanin may have a larger size than those of other higher animals. Alternately the main form of galanin in the snail CNS is a precursor. In the extracts of adrenal gland, the presence of large forms of galanin, as well as the usual molecular size, was reported in mammals (Bauer *et al.*, 1986).

Comparison with CAT-like immunoreactivity

CAT-immunoreactive (CAT-IR) cells and fibers were also widely distributed in the CNS. CAT-like immunoreactivity was seen in medium and small cells based on our classification, and no large CAT-IR cells were detected in the present study. CAT-IR cells were relatively abundant in the cerebral, left parietal, visceral, and pedal ganglia. Only a few CAT-IR cells were seen in the pleural and right parietal ganglia. No CAT-IR cells were found in the buccal ganglion. By comparison with




Fig. 3. Photomicrographs showing Gal-IR cells and fibers in the left pleural, parietal (A, B), and pedal (C, D) ganglia. A: Gal-IR fibers (small arrows) in the neuropil of the pleural (PL) and parietal (PA) ganglia and one large Gal-IR cell (large arrow). B: Two medium Gal-IR cells (large arrows) in the parietal ganglion (PA) and immunoreactive fiber bundles (small arrows) in the neuropil of the ganglion and the pallial nerve (pan). In these two planes, no small Gal-IR cells are seen. C: Small Gal-IR cells (arrows) in the pedal ganglion. D: One large Gal-IR cell (large arrow) in the pedal ganglion and immunoreactive fibers (small arrows) in the centrally located neuropil. Scale bars: A, B, 100 μm ; C, D, 20 μm .

Gal-IR cells, CAT-IR cells were observed more frequently than Gal-IR cells in the cerebral ganglion (Fig. 7A, B), and about 20% of CAT-IR cells were also immunopositive for galanin (Fig. 7C, D). That ratio corresponded to about 25% of Gal-IR cells in the cerebral ganglion. However, in the pedal and visceral ganglion, no CAT-IR cells showing galanin-like immunoreactivity were found. In the pleural and parietal ganglia, only a few co-existing cells were seen. In general, CAT-IR fibers were distributed more densely than Gal-IR fibers in the neuropil of all ganglia except in the buccal ganglion. Although CAT-IR fibers existed in all main peripheral nerves, especially many CAT-IR fibers were seen in the three pedal nerves. No CAT immunoreactivity was observed in the sections processed with normal goat serum or with omission of the

primary antibody (data not shown).

The antiserum used here is against human placental CAT. Although the cross-reactivity of the antiserum with rat CAT has been reported (Shiromani *et al.*, 1990), no data are available showing cross-reactivity with snail CAT. Therefore, results obtained with this serum should be cautiously interpreted. CAT and acetylcholine esterase activities have been biochemically demonstrated in the CNS of the snail *Helix aspersa* (Emson and Kerkut, 1971) and leech *Hirudo medicinalis* (Sargent, 1977; Wallace, 1981). It may be reasonable to consider that CAT-IR neurons of *Indoplanorbis* are cholinergic. Several bioactive peptides (myomodulin, buccalin, and FMRFamide) existing in the cholinergic system have been demonstrated to modulate effects of acetylcholine




Fig. 4. Fluorescent micrographs showing Gal-IR cells and fibers in the whole-mount CNS of the snail. A: Ventral view of the left cerebral (CE), pleural (PL), pedal (PE), and parietal (PA) ganglia, and main peripheral nerves (1, tentacular nerve (n); 2, frontal lip n; 3, median lip n; 7, pallial n). Abbreviations: cc, cerebral commissure; cp, cerebro-pedal connective; pp, parieto-pleural connective. B: Fluorescent micrograph showing Gal-IR fibers in the inferior (4), median (5), and superior (6) pedal nerves from the pedal ganglion (PE). C: Fluorescent micrograph showing Gal-IR fibers in the intestinal nerve (8) from the visceral ganglion (VI). Scale bars: A, 200 μ m; B, C, 50 μ m.




Fig. 5. Photomontage showing galanin-like immunoreactivity in the buccal ganglion (BU) and pharyngeal (10), gastric (11), and radular (12) nerves. Note dense Gal-IR fibers in the pharyngeal nerve. Scale bar, 50 μ m.




Fig. 6. Western blots of extracts from the snail CNS. Left numbers indicate molecular weights of marker proteins and rat galanin (3.2) by kDa. bb, the level of brilliant blue. A, rat galanin; B, supernatant stained with anti-galanin antibody, arrowhead indicates a 6.1 kDa band; C, supernatant stained with normal rabbit serum diluted 1:1,000.




Fig. 7. Photomicrographs of consecutive sections showing colocalization of galanin- (A, C) and CAT-like immunoreactivities (B, D) in the cerebral ganglion. A-D: Low (A, B) and high (C, D) magnification photographs showing that some of medium Gal-IR cells (arrows in A, C) are also immunopositive with an anti-CAT antibody (arrows in B, D). Small Gal-IR cells (arrowheads in C) are difficult to recognize in the adjacent section D. Scale bars: A, B, 100 μm ; C, D, 20 μm .

in the terminal area of the *Aplysia* CNS (Cropper *et al.*, 1987, 1988) and in the *Busycon* heart (Smith and Hill, 1987). In *Indoplanorbis*, CAT-IR cells were medium and small in size, and some of the medium-sized cells showed galanin-like immunoreactivity. Galanin in CAT-IR cells possibly modulates cholinergic functions. In this context, it has been suggested in the rat that galanin creates functional diversity of cholinergic signaling in relation to higher brain functions such as memory and learning (Melander and Staines, 1986). In the *Aplysia*, excellent analyses have been achieved about memory and learning which are composed of two different levels, namely, habituation and sensitization (Kandel, 1977). In the gill-withdrawal reflex, short-time habituation involves a transient decrease in synaptic efficacy, and long-term habituation produces a more prolonged change leading to a functional disruption between terminals of sensory neurons and motoneurons located in the abdominal ganglion. Serotonergic metacerebral cells in the head sensitize the reflex by means of an increase in cAMP. Similar mechanisms seem to underlie the feeding-behavior and heart rate control. It is very interesting to study whether neurons showing both galanin and CAT immunoreactivities in the snail cerebral

ganglion related to memory and learning.

The presence of Gal-IR neurons with and without CAT immunoreactivity suggests the possibility that some Gal-IR neurons contain other neurotransmitter markers such as tyrosine hydroxylase or adenosine deaminase as observed in the rat brain as cited in Introduction. Furthermore, the large Gal-IR cell in the parietal ganglion was stained with paraldehyde-fuchsin stain, but the large Gal-IR cells in the cerebral and pedal ganglia were not stained (data not shown). This suggests that subpopulation of Gal-IR cells belong to the classical neurosecretory cells. Such partial overlap of other bioactive peptide-containing cells with classical neurosecretory cells was also reported in other snails (Schot *et al.*, 1981; Sonetti *et al.*, 1990). Thus, the present study suggests the functional diversity of galanin in the snail. Further studies are required to clarify whether galanin in these cells acts as neurohormone or neurotransmitter/neuromodulator. Gal-IR neurons are possibly good candidates to colocalize a number of other signal molecules and to analyze physiological and pharmacological functions of galanin.

REFERENCES

- Bartfai T, Hökfelt T, Langel Ü (1993) Galanin — A neuroendocrine peptide. *Critical Reviews in Neurobiol* 7: 229–274
- Bauer FE, Adrian TE, Yanaihara N, Polak JM, Bloom SR (1986) Chromatographic evidence for high-molecular-mass galanin immunoreactivity in pig and cat adrenal glands. *FEBS Lett* 201: 327–331
- Bersani M, Johnsen AH, Hojrup P, Dunning BE, Andreasen JJ, Holst JJ (1991) Human galanin: primary structure and identification of two molecular forms. *FEBS Lett* 283: 189–194
- Biral GP, Zini I, Lombardo F, Ferri M, Sonetti D (1978a) Apparto per training di apprendimento in *Planorbis corneus*. *Boll Soc Ital Biol Sper* 54: 961–966
- Biral GP, Zini I, Lombardo F, Ferri M, Sonetti D (1978b) Test di apprendimento in *Planorbis corneus*. *Boll Soc Ital Biol Sper* 54: 967–973
- Boer HH, Schot LPC, Roubos EW, Maat At, Lodder JC, Beichelt D (1979) ACTH-like immunoreactivity in two electrotonically coupled giant neurons in the pond snail *Lymnaea stagnalis*. *Cell Tissue Res* 202: 231–240
- Boer HH, von Minnen J (1988) Immunocytochemistry of hormonal peptides in molluscs: optical and electron microscopy and the use of monoclonal antibodies. In "Neurohormones in Invertebrates" Ed by MC Thorndyke, GJ Goldsworthy, Cambridge University Press, Cambridge, pp 19–41
- Bonga SEW (1970) Ultrastructure and histochemistry of neurosecretory cells and neurohaemal areas in the pond snail *Lymnaea stagnalis* (L.). *Z Zellforsch* 108: 190–224
- Cropper EC, Tenenbaum R, Kolks MAG, Kupfermann I, Weiss KR (1987) Myomodulin: A bioactive neuropeptide present in an identified cholinergic buccal motor neuron of *Aplysia*. *Proc Natl Acad Sci USA* 84: 5483–5486
- Cropper EC, Miller MW, Tenenbaum R, Kolks MAG, Kupfermann I, Weiss KR (1988) Structure and action of buccalin: A modulatory neuropeptide localized to an identified small cardioactive peptide-containing cholinergic motor neuron of *Aplysia californica*. *Proc Natl Acad Sci USA* 85: 6177–6181
- Elmqvist JK, Fox CA, Ross LR, Jacobson CD (1992) Galanin-like immunoreactivity in the adult and developing Brazilian opossum brain. *Dev Brain Res* 67: 161–176
- Emsen PC, Kerkut GA (1971) Acetylcholinesterase in snail brain. *Comp Biochem Physiol* 39B: 879–889
- Evans HF, Shine J (1991) Human galanin: molecular cloning reveals a unique structure. *Endocrinology* 129: 1682–1684
- Everitt BJ, Meister B, Hökfelt T, Melander T, Terenius T, Rökaeus Å, Theodorsson-Norheim E, Dockray G, Edwardson J, Cuello C, Elde B, Goldstein M, Hemmings H, Ouimet C, Walaas I, Greengard P, Vale W, Weber E, Wu J-Y, Chang K-J (1986) The hypothalamic arcuate nucleus-median eminence complex: immunohistochemistry of transmitters, peptides and DARPP-32 with special reference to coexistence in dopamine neurons. *Brain Res Rev* 11: 97–155
- Gaymann W, Martin R (1989) Immunoreactive galanin-like material in magnocellular hypothalamo-neurohypophysial neurones of the rat. *Cell Tissue Res* 255: 139–147
- Hökfelt T, Millhorn D, Seroogy K, Tsuruo Y, Ceccatelli S, Lindh B, Meister B, Melander T, Schalling M, Bartfai T, Terenius L (1987) Coexistence of peptides with classical neurotransmitters. *Experientia* 43: 768–780
- Honma Y, Kawahara T, Chiba A (1994) Immunohistochemical localization of neuropeptide Y-like substance in the abdominal ganglion of the sea hare *Aplysia kurodai*: relationship with FMRFamide-like substance. *Fisheries Sci* 60: 53–58
- Kandel ER (1977) Cellular insights into behavior and learning. The Harvey Lectures, Series 73: 19–92
- Kaplan LM, Spindel ER, Isselbacher KJ, Chin WW (1988) Tissue-specific expression of the rat galanin gene. *Proc Natl Acad Sci USA* 85: 1065–1069
- Kasuya Y, Kobayashi H, Uemura H (1991) Endothelin-like immunoreactivity in the nervous system of invertebrates and fish. *J Cardiovasc Pharmacol* 17 (Suppl 7): S463–S466
- Kordower JH, Mufson EJ (1990) Galanin-like immunoreactivity within the primate basal forebrain: differential staining patterns between humans and monkeys. *J Comp Neurol* 294: 281–292
- Laemmli UK (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T₄. *Nature* 227: 680–685
- Lundquist CT, Rökaeus Å, Nässel DR (1991) Galanin immunoreactivity in the blowfly nervous system: localization and chromatographic analysis. *J Comp Neurol* 312: 77–96
- Melander T, Staines WA, Hökfelt T, Rökaeus Å, Eckenstein F, Salvaterra PM, Wainer BH (1985) Galanin-like immunoreactivity in cholinergic neurons of the septum-basal forebrain complex projecting to the hippocampus of the rat. *Brain Res* 360: 130–138
- Melander T, Hökfelt T, Rökaeus Å (1986) Distribution of galaninlike immunoreactivity in the rat central nervous system. *J Comp Neurol* 248: 475–517
- Melander T, Staines WA (1986) A galanin-like peptide coexists in putative cholinergic somata of the septum-basal forebrain complex and in acetylcholinesterase-containing fibers and varicosities within the hippocampus in the owl monkey (*Aotus trivirgatus*). *Neurosci Lett* 68: 17–22
- Merchenthaler I, Lopez J, Lennard EE, Negro-Vilar A (1991) Sexual differences in the distribution of neurons coexpressing galanin and luteinizing hormone-releasing hormone in the rat brain. *Endocrinology* 129: 1977–1986
- Norberg Å, Sillard R, Carlquist M, Jörnvall H, Mutt V (1991) Chemical detection of natural peptides by specific structures isolation of chicken galanin by monitoring for its N-terminal dipeptide, and determination of the amino acid sequence. *FEBS Lett* 288: 151–153
- Nordström Ö, Melander T, Hökfelt T, Bartfai T, Goldstein M (1987) Evidence for an inhibitory effect of the peptide galanin on dopamine release from the rat median eminence. *Neurosci Lett* 73: 21–26
- Olivereau M, Olivereau JM (1991) Immunocytochemical localization of a galanin-like peptidergic system in the brain and pituitary of some teleost fish. *Histochemistry* 96: 343–354
- Olivereau M, Olivereau JM (1992) Immunocytochemical localization of a galanin-like peptidergic system in the brain of two urodele and two anuran species (Amphibia). *Histochemistry* 98: 51–66
- Price DA, Greenberg MJ (1977) The structure of a molluscan cardioexcitatory peptide. *Science* 197: 670–671
- Roberts MH, Speh JC, Moore RY (1989) The central nervous system of *Bulla gouldiana*: peptide localization. *Peptides* 9: 1323–1334
- Rökaeus Å, Carlquist M (1988) Nucleotide sequence analysis of cDNA encoding a bovine galanin precursor protein in the adrenal medulla and chemical isolation of bovine gut galanin. *FEBS Lett* 234: 400–406
- Rökaeus Å, Young III WS, Mezey E (1988) Galanin coexists with vasopressin in the normal rat hypothalamus and galanin's synthesis is increased in the Brattleboro (diabetes insipidus) rat. *Neurosci Lett* 90: 45–50
- Sargent PB (1977) Synthesis of acetylcholine by excitatory motoneurons in central nervous system of the leech. *J Neurophysiol* 40: 453–460
- Schaefer M, Picciotto MR, Kreiner T, Kaldany R-R, Taussing R, Scheller RH (1985) *Aplysia* neurons express a gene encoding multiple FMRFamide neuropeptides. *Cell* 41: 457–467
- Schmidt WE, Kratzin H, Eckart K, Dreves D, Mundkowski G, Clemens A, Katsoulis S, Schafer H, Gallwitz B, Creutzfeldt W (1991) Isolation and primary structure of pituitary human galanin, a 30-residue nonamidated neuropeptide. *Proc Natl Acad Sci USA* 88:

11435–11439

- Shiromani PJ, Floyd C, Velázquez-Moctezuma J (1990) Pontine cholinergic neurons simultaneously innervate two thalamic targets. *Brain Res* 532: 317–322
- Schot LPC, Boer HH, Swaab DF, Noorden SV (1981) Immunocytochemical demonstration of peptidergic neurons in the central nervous system of the pond snail *Lymnaea stagnalis* with antisera raised to biologically active peptides of vertebrates. *Cell Tissue Res* 216: 273–291
- Schot LPC, Boer HH (1982) Immunocytochemical demonstration of peptidergic cells in the pond snail *Lymnaea stagnalis* with an antiserum to the molluscan cardioactive tetrapeptide FMRFamide. *Cell Tissue Res* 225: 347–354
- Schot LPC, Boer HH, Montagne-Wajer C (1984) Characterisation of multiple immunoreactive neurons in the central nervous system of the pond snail *Lymnaea stagnalis* with different fixatives and antisera adsorbed with the homologous and the heterologous antigens. *Histochemistry* 81: 373–378
- Skofitsch G, Jacobowitz DM (1985) Immunohistochemical mapping of galanin-like neurons in the rat central nervous system. *Peptides* 6: 509–546
- Skofitsch G, Jacobowitz DM, Amann R, Lembeck F (1989) Galanin and vasopressin coexist in the rat hypothalamoneurohypophyseal system. *Neuroendocrinology* 49: 419–427
- Smith PJS, Hill RB (1987) Modulation of output from an isolated gastropod heart: effects of acetylcholine and FMRFamide. *J Exp Biol* 127: 105–120
- Sonetti D, Lusvardi C, Fasolo A (1990) Immunohistochemical localization of some vertebrate-like neuropeptides (SP, NPY, CGRP, CCK) in the central nervous system of the freshwater snail *Planorbarius corneus*. *Cell Tissue Res* 260: 435–448
- Staines WA, Yamamoto T, Daddona PE, Nagy JI (1986) Neuronal colocalization of adenosine deaminase, monoamine oxidase, galanin and 5-hydroxytryptophan uptake in the tuberomammillary nucleus of the rat. *Brain Res Bull* 17: 351–365
- Tatemoto K, Rokaeus A, Jornvall H, McDonald TJ, Mutt V (1983) Galanin — a novel biologically active peptide from porcine intestine. *FEBS Lett* 164: 124–128
- Valnes K, Brandtzaeg P (1985) Retardation of immunofluorescence fading during microscopy. *J Histochem Cytochem* 33: 755–761
- Vrontakis ME, Peden LM, Duckworth ML, Friesen HG (1987) Isolation and characterization of a complementary DNA (galanin) clone from estrogen induced pituitary tumor messenger RNA. *J Biol Chem* 262: 16755–16758
- Wallace BG (1981) Distribution of AChE in cholinergic and non-cholinergic neurons. *Brain Res* 219: 190–195
- Yamamoto T, Maler L, Nagy JI (1992) Organization of galanin-like immunoreactive neuronal systems in weakly electric fish (*Apteronotus leptorhynchus*). *J Chem Neuroanat* 5: 19–38

(Received February 8, 1996 / Accepted April 16, 1996)