

Detection of Several Families of Deiminated Proteins Derived from Filaggrin and Keratins in Guinea Pig Skin

Authors: Kan, Shuhei, Asaga, Hiroaki, and Senshu, Tafsuo

Source: Zoological Science, 13(5) : 673-678

Published By: Zoological Society of Japan

URL: <https://doi.org/10.2108/zsj.13.673>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Detection of Several Families of Deiminated Proteins Derived from Filaggrin and Keratins in Guinea Pig Skin

Shuhei Kan¹, Hiroaki Asaga* and Tatsuo Senshu

*Department of Cell Chemistry, Tokyo Metropolitan Institute of Gerontology,
Itabashi-ku, Tokyo 173, Japan*

ABSTRACT—Structural proteins in the mammalian epidermis contain citrulline residues generated by enzymatic deimination of arginine residues. We analyzed deiminated proteins solubilized from sequentially stripped layers of guinea pig epidermis. Deiminated proteins were localized in the granular and cornified layers. Those in the inner layer enriched with granular cells were resolved into numerous components by two-dimensional gel electrophoresis. An arc-shaped high-molecular-weight smear and two series of charged isomers among them coincided with filaggrin immunoreactivity. Several groups of filaggrin-negative spots appeared to be generated by further deimination and proteolysis of these filaggrins. Deiminated protein spots co-migrating with type II and type I keratins were also detected. Deiminated filaggrins and their further processed derivatives disappeared in the outer layer, while deiminated keratins persisted. These data suggested that filaggrin as well as profilaggrin were deiminated during the posttranslational processing in guinea pig skin, and that some keratins were deiminated preferentially during the cornification of epidermis. Possible biological significance of protein deimination in guinea pig skin was discussed in comparison with our recent finding on deiminated proteins in rat skin.

INTRODUCTION

The process of normal epidermal differentiation is characterized by a series of morphologic and biochemical changes as keratinocytes progress from the germinative basal layer through the spinous and granular layers to the outer cornified layer. Aggregation of keratin filaments mediated by filaggrin is one of the key events underlying the process (Lynley and Dale, 1983; Manabe *et al.*, 1991; Steinert *et al.*, 1981). Filaggrin is generated from its high-molecular-weight precursor (profilaggrin), which does not aggregate keratin filaments (Haydock and Dale, 1986; Meek *et al.*, 1983; Rothnagel *et al.*, 1987; Scott and Harding, 1981) and is deposited in keratohyalin granules in highly phosphorylated forms (Lonsdale-Eccles *et al.*, 1980, 1982; Resing *et al.*, 1985; Scott and Harding, 1981). Profilaggrin is dephosphorylated during processing to filaggrin (Lonsdale-Eccles *et al.*, 1982; Resing *et al.*, 1984, 1985; Scott and Harding, 1981), and filaggrin is further degraded in the cornified layer to amino acids, which are thought to be important for retaining moisture in the cornified cells (Scott *et al.*, 1982; Scott and Harding, 1986). Therefore, both phosphorylation/dephosphorylation and proteolysis are involved in the functional role of filaggrin. A

relatively unexplored posttranslational modification, which filaggrin seems to be subjected to, is “deimination”. Deimination is catalyzed by “peptidylarginine deiminases (EC 3.5.3.15)” which deiminate arginine residues generating citrulline residues (Fujisaki and Sugawara, 1981; Kubilus *et al.*, 1980; Rogers *et al.*, 1977; Rothnagel and Rogers, 1984; Terakawa *et al.*, 1991; Watanabe *et al.*, 1988). Harding and Scott (1983) suggested that the deimination of arginine residues of filaggrin modulates its interaction with keratins and facilitates its breakdown to amino acids. However, information concerning biological significance of protein deimination is limited, partly because sensitive enough methods to detect deiminated proteins specifically were not available.

We recently showed that not only filaggrin but also keratins were deiminated in rat skin (Senshu *et al.*, 1995). These deiminated proteins were detected specifically by means of chemically modifying citrulline residues then probing them with a monospecific antibody (Senshu *et al.*, 1992). We also showed that deiminated proteins were localized in the granular and cornified layers, and suggested a possible role for protein deimination during the terminal stage of epidermal differentiation (Senshu *et al.*, 1995). However, in rat skin, not only the number of arginine residues modified in a given filaggrin molecule but also the amount of deiminated filaggrin in the total filaggrin pools appeared to be too small to argue for the functional significance of filaggrin deimination. We therefore extended our studies using guinea pig skin, in which multiple charged isomers of filaggrin were detected as

¹ Present address: Bio Signal Pathway Project, Kanagawa Academy of Science and Technology in Meiji Institute of Health Science, Odawara 250, Japan.

* To whom correspondence should be addressed.

presumptive products of deimination (Scott, 1986; Harding and Scott, 1983). We found numbers of spots of deiminated components at various stages of proteolytic processing of profilaggrin in the granular layer, that disappeared in the cornified layer. We also found the preferential deimination of some keratins that persisted in the cornified layer.

MATERIALS AND METHODS

Animals and tissue samples

Five-week-old female albino guinea pigs (Hartley strain) weighing 300-350 g were obtained from Japan SLC Co. Ltd. (Shizuoka, Japan). All operations were performed under pentobarbital anesthesia or after decapitation under the anesthesia. Back skin was clipped and depilated using depilatory cream (Epilat; Kanebo, Japan). Epidermal layers were stripped using cyanoacrylate glue. The glue was spread as a thin film over a delineated area, and then removed with the attached cell layers. This operation was repeated three times. Histological examination showed relative enrichment of the upper cornified cells in the first layer, that of the lower cornified cells in the second layer, and that of the granular cells in the third layer.

Extraction of proteins

The stripped layers were minced in 9.5 M urea (Ultra Pure, ICN Biochemicals, Inc., Ohio), 2% Nonidet P-40, 2% Ampholines (pH 3.5-10, Pharmacia LKB), and 5% 2-mercaptoethanol, and insoluble materials were removed by brief centrifugal filtration through a 200-mesh nylon screen. Protein concentrations in the extracts were estimated according to Lowry *et al.* (1951) after precipitation with 20% trichloroacetic acid. Bovine serum albumin was used as the standard.

SDS-PAGE and immunoblotting

The extracts were resolved by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) or two-dimensional gel electrophoresis using a nonequilibrium pH gradient gel electrophoresis (pH 3.5-10) in the first dimension as described (Senshu *et al.*, 1995). Samples for SDS-PAGE were diluted at least 5-fold to reduce the concentration of Nonidet P-40. They were not heated to avoid artificial carbamylation of proteins. Resolved proteins were Western blotted to a BA-S 85 membrane (S & S, Dassel, Germany). Total proteins were visualized by staining with Amido Black 10B. The blot for detecting deiminated proteins was pretreated with ovalbumin and paraformaldehyde to improve the retention of deiminated filaggrins on the membrane (Senshu *et al.*, 1995). Citrulline residues on the blot were modified by an overnight incubation at 37°C in 0.0125% FeCl₃, 2.3 M H₂SO₄, 1.5 M H₃PO₄, 0.25% diacetyl monoxime, 0.125% antipyrine, and 0.25 M acetic acid (modification medium), and the resulting blot was incubated with anti-modified citrulline IgG (0.125 µg/ml) (Senshu *et al.*, 1992, 1995). The specificity of the immunochemical detection was confirmed by probing equivalent blots incubated in modification medium free of diacetyl monoxime, antipyrine, and acetic acid. Filaggrin was immunoblotted using a polyclonal antibody purified from an anti-rat filaggrin serum (Senshu *et al.*, 1995) by adsorption to rat filaggrin on the Western blot followed by elution with 3 M sodium thiocyanate. It was used at an IgG concentration of 13 ng/ml estimated by dot immunoassay using normal rabbit IgG as the reference. Keratins were detected using a mixture of the monoclonal antibodies AE1 and AE3 (provided by Dr. T.-T. Sun, New York University School of Medicine) at 500- and 1000-fold dilutions, respectively. The blots were incubated with appropriate peroxidase-labeled second IgGs (Bio-Rad, used at 5000-fold dilution), and the bound second antibodies were detected by the enhanced luminol reaction using Renaissance (Dupont NEN). The intensity of the chemiluminescence signal was increased by adding 0.015% hydrogen peroxide to the reagent cocktail. The data were recorded

by a Macintosh computer using a chilled charge-coupled device camera and displayed on a video screen at a suitable intensity. The immunoblot was post-stained with Amido Black 10B to correlate the chemiluminescence spots with stains in the total protein profile. Visible images were recorded by a Macintosh computer using a high performance charge-coupled device camera. Hard copies were obtained using a video printer.

Immunohistochemistry

Tissues were fixed in a Bouin Hollande Sublimate solution, then cryosections were prepared and mounted as described (Asaga and Senshu, 1993; Senshu *et al.*, 1995). Sections for staining deiminated proteins were post-fixed as described to ensure tight adherence to the slides (Asaga and Senshu, 1993). Endogenous peroxidases were inactivated by an incubation in 1% hydrogen peroxide in methanol for 15 min. Sections were then incubated with 0.25% trypsin (GIBCO/BRL) for 15 min, followed by modification medium at 37°C for 3 hr (Asaga and Senshu, 1993; Senshu *et al.*, 1995). Control sections were incubated in medium free of diacetyl monoxime, antipyrine, and acetic acid. The resulting sections were incubated with anti-modified citrulline IgG (0.125 µg/ml). Sections for filaggrin immunostaining were incubated with the affinity purified antibody to rat filaggrin (13 ng/ml). Bound antibodies were stained by means of a VECTASTAIN ABC Elite kit (Vector Laboratories), using 3,3'-diaminobenzidine as the chromogenic substrate. Stained sections were counterstained with hematoxylin.

RESULTS

Detection of deiminated proteins in stripped epidermal layers

We first compared SDS-PAGE profiles of proteins extracted from the upper three layers of stripped epidermis. Three major bands (195, 61, and 56 kDa) were detected in the total protein profile (Fig. 1a). These bands were detected with the mixture of monoclonal antibodies AE1 and AE3, although the 61-kDa band was under-represented probably due to its lower affinity to the monoclonal antibodies (Fig. 1b). Two minor bands (66 and 46 kDa) were detectable in the total protein profiles of the inner two layers. The former but not the latter was detected with the monoclonal antibody mixture. Immunoblotting for filaggrin gave only a weak 195-kDa band in the first layer (Fig. 1c). The band intensity increased in the inner layers along with a leading smear. These signals probably represented profilaggrin and its high-molecular-weight processing intermediates. Two additional bands co-migrating with the 66- and 46-kDa bands found in the total protein profile were detected in the second layer, and their signal intensity increased further in the third layer. Detection of deiminated proteins on equivalent blots showed four major bands (206, 165, 61, and 56 kDa) in the first layer (Fig. 1d). The signal intensity of the two smaller bands, which co-migrated with those detected by the AE1/AE3 mixture, decreased in the inner layers, whereas two additional bands (66 and 46 kDa) co-migrating with those detected by anti-filaggrin became detectable. In addition, broad bands (about 34, 30, and 23 kDa) were discernible above the background smear. The signal intensity of the 46-, 34-, 30-, and 23-kDa bands appeared to be stronger in the second layer than in the third layer, while that of the 66-kDa band remained about equal. This and the relative abundance of 66- and 46-kDa filaggrins




Fig. 1. SDS-PAGE profiles of various epidermal proteins. Samples were resolved by SDS-PAGE and Western blotted onto nitrocellulose membranes to stain total proteins with Amido Black 10B (a), as well as to detect keratins (b), filaggrin (c), and deiminated proteins (d) as described in the text. Lanes 1-3, first, second, and third layers stripped from the epidermis, respectively. The amount applied was 5 μ g. The gel was calibrated with the Molecular Weight Markers "DAIICHI" II (Daiichi Pure Chemicals Co., Ltd., Tokyo, Japan).

in these layers suggested that these filaggrins in the second layer were deiminated at larger numbers of arginine residues than those in the third layer.

Identification of deiminated proteins on two-dimensional blots

The above one-dimensional analyses allowed quantitative comparison of the band intensity between different samples. We next performed two-dimensional Western blotting of the first and the third layer fractions to identify deiminated proteins. Keratins, filaggrin, and deiminated proteins were detected on equivalent blots and results then correlated with the stained total protein profiles. It should be mentioned that each immunoblot was displayed at an appropriate intensity to show its protein composition. This does not necessarily mean that the data are suitable for quantitative comparison between different samples. The first layer showed two diffuse major spots of deiminated proteins (Fig. 2d). Overlapping spots were visible in the total protein (Fig. 2a). The less anodic spots co-migrated with a diffuse type II keratin spot corresponding to the 61-kDa band in the one-dimensional analysis (Fig. 2b). The other spot co-migrated with a 56-kDa type I keratin spot. These keratins are probably the products of various posttranslational modifications as measured from the rather diffuse appearance of the spots. A few curved smears of deiminated proteins were found as minor signals near the upper anodic corner. Similar signals were detected in infant rat skin (Senshu *et al.*, 1995), which were thought to be derived from trichohyalin (Rogers *et al.*, 1977). These deiminated proteins did not coincide with weak signals detected by anti-

filaggrin (Fig. 2c). A streak in the upper anodic region was detected by either anti-filaggrin or anti-keratins. It probably represented high-molecular-weight aggregates containing both filaggrin and keratins.

The total protein profile of the third layer showed an arc-shaped smear in the upper cathodic region, multiple spots of keratins in the neutral to acidic region, and two series of charged isomers (at about 66 and 46 kDa) scattered in the cathodic region (Fig. 2e). The mobility of these charged isomers in the second dimension appeared to decrease slowly with the decreasing first dimensional mobility. An additional series of charged isomers (about 34 kDa) was also discernible in the neutral region. Immunoblotting for filaggrin showed a major signal co-migrating with the arc-shaped smear representing high-molecular-weight processing intermediates, and minor signals co-migrating with basic components of the 66- and 46-kDa charged isomers (Fig. 2g). Detection of deiminated proteins on equivalent blots visualized a prominent signal co-migrating with the arc-shaped smear. This suggests deimination of high-molecular-weight processing intermediates of profilaggrin (Fig. 2h). Two series of charged isomers (66 and 44 kDa) co-migrating with those in the total protein profile were also detected. The arrays of these charged isomers extended further towards the neutral region than those detected with anti-filaggrin, suggesting that less basic components were deiminated at larger numbers of arginine residues. In addition, weak signals of 34-kDa charged isomers were detected in the neutral region. Series of smaller charged isomers could be detected when the data were displayed at increased sensitivity. These isomers appeared to be vertically or near vertically linked with trailing smears, giving the overall impression of a flowing stream. This layer also contained a series of intense signals that co-migrated with a series of 61-kDa type II keratins and minor signals migrating in the 56-kDa type I keratin region (Fig. 2f). These spots of deiminated proteins accounted for all the bands found in the one-dimensional analysis. It should be noted that only limited numbers of keratin spots were deiminated.

Immunolocalization of deiminated proteins

Figure 3 shows micrographs of immunohistochemically stained sections in comparison with that stained with hematoxylin and eosin (Fig. 3a). Filaggrin immunoreactivity was localized in the granular and lower cornified layers (Fig. 3b). Deiminated proteins were localized in the whole cornified layer as well as in the granular layer (Fig. 3c). Control sections incubated in modification medium without diacetyl monoxime, antipyrine, and acetic acid were not stained (Fig. 3d).

DISCUSSION

Filaggrin is the product of the dephosphorylation and proteolytic processing of the precursor profilaggrin (Haydock and Dale, 1986; Lonsdale-Eccles *et al.*, 1980, 1982; Meek *et al.*, 1983; Resing *et al.*, 1984, 1985; Rothenagel *et al.*, 1987; Scott and Harding, 1981). Filaggrin can be separated by two-




Fig. 2. Characterization of the deiminated proteins by two-dimensional gel electrophoresis. Samples were resolved by nonequilibrium pH gradient gel electrophoresis and SDS-PAGE in the first and second dimensions, respectively. Total proteins (a, e), keratins (b, f), filaggrin (c, g), and deiminated proteins (d, h) were visualized as described. Panels a-d, proteins extracted from the first layer; Panels e-h, proteins extracted from the third layer. The amount loaded was about 30 μ g. Acidic type I and neutral-basic type II keratins were distinguished by separately probing equivalent blots with AE1 and AE3. Reference points are shown by numbered arrowheads.




Fig. 3. Micrographs of stained skin sections. Cryosections were processed as described in the text. (a) Stained with hematoxylin and eosin; (b) anti-filaggrin; (c) post-fixed, incubated in the modification medium, and stained with anti-modified citrulline IgG; (d) treated similarly to b except that diacetyl monoxime, antipyrine, and acetic acid were omitted from the modification medium. (C) Stratum corneum; (G) stratum granulosum; (S) stratum spinosum. Dashed line, epidermal-dermal junction. Bar, 50 μ m.

dimensional gel electrophoresis into families differing in molecular weight and/or charge, depending on the species (Harding and Scott, 1983). Guinea pig filaggrin has been shown to be highly complex, including high-molecular-weight variants (200 kDa or above) (Harding and Scott, 1983; Scott, 1986). We used an affinity purified polyclonal antibody to rat filaggrin for immunoblotting. The two-dimensional immunoblot reported here resembled those of guinea pig filaggrin reported by other investigators (Harding and Scott, 1983; Scott, 1986). In addition, the antibody specifically stained the lower cornified and granular layers. These results indicated the acceptable cross-reactivity and specificity of our antibody for probing guinea pig filaggrin.

Harding and Scott (1983) suggested the deimination of guinea pig filaggrin based on amino acid analysis and on the detection of [3 H]citrulline in filaggrin derived from guinea pig skin pre-labeled with [3 H]arginine. They further proposed that deimination progressively lowers the affinity of filaggrin for keratins based on their observation that less basic variants of filaggrins interacted poorly with keratins *in vitro*. However, direct identification of deiminated filaggrins on the two-dimensional map was not conducted by these authors. Here we used a highly sensitive and specific method for the detection of deiminated proteins on the two-dimensional Western blot, and detected several families of deiminated filaggrins at various stages of processing, ranging from the arc-shaped smear in the high-molecular-weight region to multiple charged isomers in the low-molecular-weight region. Furthermore, it should be noted that the present method enabled us to detect multiply modified or extensively processed components that were not detectable with anti-filaggrin. The enrichment of deiminated filaggrins in the second layer and their almost complete deprivation in the first layer probably reflected increased deimination of filaggrins in the lower cornified layer followed by their degradation in the upper cornified layer. This is in accord with the proposal of Harding and Scott (1983) suggesting that the deimination of filaggrin facilitates its breakdown to amino acids. However, the biological significance of filaggrin deimination or even filaggrin itself in guinea pig skin is still unclear. We do not know which are functional filaggrin units to aggregate keratins among various filaggrin-positive components detected on the Western blots. The 66-kDa component may be regarded to be functional from its relative abundance. However, if so, why was the high-molecular-weight smear already deiminated? If they were deiminated prior to binding to keratins, did it reflect the regulated production of non-deiminated, functional filaggrin and deiminated, non-functional filaggrin? Furthermore, it is puzzling that deimination of filaggrin appeared to occur at only few arginine residues in a minor fraction of the total filaggrin in rat skin (Senshu *et al.*, 1995). Nevertheless, filaggrin was thought to be efficiently degraded as measured from its deprivation in the upper cornified layer. These data suggest that deimination of filaggrin may not be an obligatory step for its breakdown in rat skin. In addition, deimination of high-molecular-weight processing intermediates were not detected

in rat skin. Therefore, the biological significance of filaggrin deimination might differ between different species.

It is well-known that a basic type II keratin K5 and its acidic type I keratin partner K14 are expressed in the basal layer of epidermis forming fine cytoskeletal networks (Steinert and Freedberg, 1991). Another basic type II keratin K1 and its acidic type I keratin partner K10 are expressed in the suprabasal cells and incorporated into the pre-existing networks of K5 and K14 (Steinert and Freedberg, 1991). The resulting keratin networks are converted to a densely packed filamentous structure mediated by filaggrin during the transition of the granular cells to the cornified cells (Lynley and Dale, 1983; Manabe *et al.*, 1991; Steinert *et al.*, 1981). The presence of deiminated keratins and filaggrins in these upper epidermal layers is consistent with our proposal that the protein deimination was playing a possible role during the terminal stages of epidermal differentiation (Senshu *et al.*, 1995). It should be noted that only a few of keratin spots were deiminated, as has been observed in rat skin. This suggests a preferential action of peptidylarginine deiminases towards these keratins. The major deiminated products are type II keratins. The deimination of specific arginine residues of such keratins might play a role in modulating interactions with their direct partners, other keratins in the cytoskeletal network, or keratin-associated protein filaggrin. Detailed identification of deiminated keratins was not attempted because antibodies specific enough to identify guinea pig keratins were not available. We are in the process of identifying the deiminated residues in defined keratins.

ACKNOWLEDGMENTS

We thank Dr. T.-T. Sun (New York University School of Medicine) for supplying the monoclonal antibodies AE1 and AE3. We also thank Prof. Seiichiro Kawashima for his encouragement. Supported in part by a Grant-in-Aid #07680703 from the Ministry of Education, Science, Sports and Culture, Japan.

REFERENCES

- Asaga H, Senshu T (1993) Combined biochemical and immunocytochemical analyses of postmortem protein deimination in the rat spinal cord. *Cell Biol Int* 17: 525–532
- Fujisaki M, Sugawara K (1981) Properties of peptidylarginine deiminase from the epidermis of new born rat. *J Biochem (Tokyo)* 89: 257–263
- Harding CR, Scott IR (1983) Histidine-rich proteins (filaggrins): Structural and functional heterogeneity during epidermal differentiation. *J Mol Biol* 170: 651–673
- Haydock PV, Dale BA (1986) The repetitive structure of the profilaggrin gene as demonstrated using epidermal profilaggrin cDNA. *J Biol Chem* 261: 12520–12525
- Kubilus J, Waitkus RW, Baden HP (1980) Partial purification and specificity of an arginine-converting enzyme from bovine epidermis. *Biochim Biophys Acta* 615: 246–251
- Lonsdale-Eccles JD, Haugen JA, Dale BA (1980) A phosphorylated keratohyalin-derived precursor of epidermal stratum corneum basic protein. *J Biol Chem* 255: 2235–2238
- Lonsdale-Eccles JD, Teller DC, Dale BA (1982) Characterization of a phosphorylated form of the intermediate filament-aggregating protein filaggrin. *Biochemistry* 21: 5940–5948
- Lowry OH, Rosebrough NJ, Farr AL, Randall RJ (1951) Protein measurement with the Folin phenol reagent. *J Biol Chem* 193: 265–275
- Lynley AM, Dale BA (1983) Characterization of human epidermal filaggrin, a histidine-rich, keratin filament aggregating protein. *Biochim Biophys Acta* 744: 28–45
- Manabe M, Sanchez M, Sun T-T, Dale BA (1991) Interaction of filaggrin with keratin filaments during advanced stages of normal human epidermal differentiation and in ichthyosis vulgaris. *Differentiation* 48: 43–50
- Meek RL, Lonsdale-Eccles JD, Dale BA (1983) Epidermal filaggrin is synthesized on a large messenger ribonucleic acid as a high-molecular-weight precursor. *Biochemistry* 22: 4867–4871
- Resing KA, Walsh KA, Dale BA (1984) Identification of two intermediates during processing of profilaggrin to filaggrin in neonatal mouse epidermis. *J Cell Biol* 99: 1372–1378
- Resing KA, Dale BA, Walsh KA (1985) Multiple copies of phosphorylated filaggrin in epidermal profilaggrin demonstrated by analysis of tryptic peptides. *Biochemistry* 24: 4167–4175
- Rogers GE, Harding HWJ, Llewellyn-Smith IJ (1977) The origin of citrulline-containing proteins in the hair follicle and the chemical nature of trichohyalin, an intracellular precursor. *Biochim Biophys Acta* 495: 159–175
- Rothnagel JA, Mehrel T, Idler WW, Roop DR, Steinert PM (1987) The gene for mouse epidermal filaggrin precursor. Its partial characterization, expression, and sequence of a repeating filaggrin unit. *J Biol Chem* 262: 15643–15648
- Scott IR (1986) Alterations in the metabolism of filaggrin in the skin after chemical- and ultraviolet-induced erythema. *J Invest Dermatol* 87: 460–465
- Scott IR, Harding CR (1981) Studies on the synthesis and degradation of a high molecular weight histidine-rich phosphoprotein from mammalian epidermis. *Biochim Biophys Acta* 669: 65–78
- Scott IR, Harding CR (1986) Filaggrin breakdown to water binding compounds during development of the rat stratum corneum is controlled by the water activity of the environment. *Dev Biol* 115: 84–92
- Scott IR, Harding CR, Barrett JG (1982) Histidine-rich protein of the keratohyalin granules. Source of the free amino acids, urocanic acid and pyrrolidone carboxylic acid in the stratum corneum. *Biochim Biophys Acta* 719: 110–117
- Senshu T, Sato T, Inoue T, Akiyama K, Asaga H (1992) Detection of citrulline residues in deiminated proteins on polyvinylidene difluoride membrane. *Anal Biochem* 203: 94–100
- Senshu T, Akiyama K, Kan S, Asaga H, Ishigami A, Manabe M (1995) Detection of deiminated proteins in rat skin: Probing with a monospecific antibody after modification of citrulline residues. *J Invest Dermatol* 105: 163–169
- Steinert, PM, Freedberg, IM (1991) Molecular and cellular biology of keratins. In "Physiology, Biochemistry, and Molecular Biology of the Skin Vol 1" Ed by LA Goldsmith, Oxford University Press, New York, pp 113–147
- Steinert PM, Cantieri JS, Teller DC, Lonsdale-Eccles JD, Dale BA (1981) Characterization of a class of cationic proteins that specifically interact with intermediate filaments. *Proc Natl Acad Sci USA* 78: 4097–4101
- Terakawa H, Takahara H, Sugawara K (1991) Three types of peptidylarginine deiminase: Characterization and tissue distribution. *J Biochem (Tokyo)* 110: 661–666
- Watanabe K, Akiyama K, Hikichi K, Ohtsuka R, Okuyama A, Senshu T (1988) Combined biochemical and immunocytochemical comparison of peptidylarginine deiminases present in various tissues. *Biochim Biophys Acta* 966: 375–383

(Received March 28, 1996 / Accepted May 29, 1996)