



## **Relative Potency of Three Homologous Natriuretic Peptides (ANP, CNP and VNP) in Eel Osmoregulation**

Authors: Tsukada, Takehiro, and Takei, Yoshio

Source: Zoological Science, 18(9) : 1253-1258

Published By: Zoological Society of Japan

URL: <https://doi.org/10.2108/zsj.18.1253>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Relative Potency of Three Homologous Natriuretic Peptides (ANP, CNP and VNP) in Eel Osmoregulation

Takehiro Tsukada\* and Yoshio Takei

Ocean Research Institute, the University of Tokyo,  
Tokyo 164-8639, Japan

**ABSTRACT**—Evidence has accumulated that atrial natriuretic peptide (ANP) plays important roles in sea-water adaptation in eels. However, the roles of the other two natriuretic peptides (CNP and VNP) in osmoregulation have not been examined yet. In the present study, the effects of homologous ANP, CNP and VNP were compared on plasma  $\text{Na}^+$  concentration (an indicator of plasma osmolality), hematocrit (an approximate indicator of blood volume) and drinking rate in freshwater- and seawater-adapted eels. In seawater eels, ANP and VNP, but not CNP, infused at 5 pmol/kg/min decreased plasma  $\text{Na}^+$  concentration and drinking rate and increased hematocrit. In freshwater eels, ANP and VNP failed to decrease plasma  $\text{Na}^+$  concentration but increased hematocrit to the same extent as in seawater eels. Inhibition of drinking was not detectable in freshwater eels because of little drinking before NP infusions. These results show that the effects of NPs on plasma  $\text{Na}^+$  concentration, drinking rate and hematocrit are mediated by NPR-A, since only ANP and VNP that bind with higher affinity to NPR-A are effective in seawater eels. The mechanisms of regulation of plasma  $\text{Na}^+$  concentration and hematocrit are unknown, but NPR-A is present in the responsible tissues for regulation of hematocrit in both freshwater and seawater eels. However, NPR-A may be absent in the tissues of freshwater eels that are responsible for regulation of plasma  $\text{Na}^+$  concentration.

**Key words:** atrial natriuretic peptide, ventricular natriuretic peptide, C-type natriuretic peptide, drinking, plasma sodium concentration

## INTRODUCTION

Accumulating data indicate that natriuretic peptides (NPs) are involved in cardiovascular and body fluid homeostasis in vertebrates (Farrell and Olson, 2000; Loretz and Pollina, 2000; Takei, 2000a). The NP family consists of atrial natriuretic peptide (ANP), brain natriuretic peptide (BNP), and C-type natriuretic peptide (CNP) in tetrapods. ANP and BNP are circulating hormones secreted mainly from the heart, while CNP is basically a paracrine factor in the brain and periphery (Fowkes and McArdle, 2000; Takei, 2000a). Three NP receptors have been identified in mammals, of which NPR-A and NPR-B are members of the membrane-bound guanylyl cyclases and are involved in biological actions of the NP family. Another type, NPR-C, lacks the enzyme domain and may be involved in the clearance of NPs (Maack, 1992). ANP and BNP bind to NPR-A preferentially and CNP is specific for NPR-B, while all three NPs have high affinity to NPR-C.

In the eel, ANP, ventricular natriuretic peptide (VNP), CNP and four NP receptors (NPR-A, -B, -C, and -D) have been

identified (Takei, 2000a). VNP is a teleost-specific NP with a long C-terminal sequence extending from the intramolecular ring (Takei *et al.*, 1991). VNP may be unique in its function because it has high affinity not only to NPR-A but also to NPR-B equivalent to 1/10 of CNP (Katafuchi *et al.*, 1994). NPR-D is a new type of guanylyl cyclase-deficient receptor and has been identified only in the eel thus far (Kashiwagi *et al.*, 1995).

Previous studies showed that ANP is secreted soon after transfer of freshwater eels to seawater (Kaiya and Takei, 1996), and acts on various osmoregulatory organs to decrease NaCl in the body, thereby promoting adaptation to seawater (Takei, 2000a). In contrast, expressions of CNP gene as well as plasma CNP level are greater in freshwater eels than in seawater eels (Takei *et al.*, 2001). NPR-B messages are more abundantly expressed in freshwater eels than in seawater eels (Katafuchi *et al.*, 1994) and CNP increases cyclic guanosine 3', 5'-monophosphate (cGMP) accumulation in the membrane fraction of freshwater eel gills but not of seawater eel gills (Mishina and Takei, 1997). These results indicate a role of CNP in freshwater adaptation. However, nothing is known about the role of VNP in osmoregulation except for the vascular action (Takei *et al.*, 1991).

In the present study, the relative potency of three eel NPs: ANP, CNP and VNP, were compared in water and electrolyte

\* Corresponding author: Tel. 03-5351-6464;  
FAX. 03-5351-6463.  
E-mail: tsuka@ori.u-tokyo.ac.jp

metabolism in freshwater and seawater eels using an *in vivo* infusion system. The parameters measured were drinking rate, plasma  $\text{Na}^+$  concentration and hematocrit. Drinking was measured because it is essential for adaptation of eels to hyperosmotic environments (Takei *et al.*, 1998). Plasma  $\text{Na}^+$  concentration and hematocrit were used to assess the changes in two major parameters of body fluid balance: plasma osmolality and blood volume, respectively. The differences in potencies of three NPs in different actions and in different osmotic environments are discussed in relation to the involvement of different NP receptors.


## MATERIALS AND METHODS

**Animals:** Cultured, immature eels, *Anguilla japonica*, of mixed sexes were purchased from a local dealer. They were maintained without feeding in seawater or freshwater tanks for at least two weeks before use. Water in the tank was continuously circulated, aerated, and regulated at 18°C. Eels weighed  $198.3 \pm 7.6$  g ( $n=5$ ; seawater) and  $200.7 \pm 8.2$  g ( $n=6$ ; freshwater) at the time of surgery.


**Drugs:** Three homologous NPs were dissolved in distilled water, aliquoted and frozen at  $-20^\circ\text{C}$  until use. On the day of experimentation, the stock solution ( $10^{-4}$  M) was diluted with isotonic saline (0.9% NaCl) containing 0.01% Triton X-100 (Wako pure chemicals industries, Osaka).

**Surgical procedures:** Eels were anesthetized by immersion in 0.1% (w/v) tricaine methanesulfonate (Sigma, St. Louis, MO, USA) for 15 min. Vinyl tubes (o.d.: 1.5 mm, Becton Dickinson Co., Sparks, MD, USA) were inserted into the esophagus and stomach of eels as described previously (Takei *et al.*, 1998). In addition, a polyethylene tube (o.d.: 0.8 mm, SP31, Natsume Seisakusho Co., Tokyo) was inserted into the ventral aorta for infusion of NPs and blood collection. After surgery, eels were placed in a plastic trough through which aerated seawater or fresh water continuously circulated at 18°C. The troughs were covered with a black vinyl sheet to minimize visual stress during the experiments.

**Synchronized drop counter and pulse injector system:** The catheter placed in the esophagus was connected to a drop counter for continuous measurement of drinking rate. It has been shown that seawater dropped from the catheter is diluted to 80% by esophageal desalination in seawater eels (Tsuchida and Takei, 1998). Thus 80% seawater of the same volume as a drop was injected into the stom-



**Fig. 1.** The time course of changes in plasma  $\text{Na}^+$  concentration during 30-min infusions of homologous ANP (○), CNP (●) and VNP (□) (5 pmol/kg/min) in seawater eels (A;  $n=5$ ) and freshwater eels (B;  $n=6$ ). Saline was infused before and after NP infusions. Data were expressed as percentage to the values before NP infusion. Values represent means  $\pm$  SE. \* $p < 0.05$ , \*\* $p < 0.01$  compared with the value before NP infusion by Fisher's PLSD test.



**Fig. 2.** The time course of changes in hematocrit during 30-min infusions of homologous ANP (○), CNP (●) and VNP (□) (5 pmol/kg/min) in seawater eels (A;  $n=5$ ) and freshwater eels (B;  $n=6$ ). Saline was infused before and after NP infusions. Data were expressed as percentage to the values before NP infusion. Values represent means  $\pm$  SE. \* $p < 0.05$ , \*\* $p < 0.01$  compared with the value before NP infusion by Fisher's PLSD test.


ach by a pulse injector synchronized with the drop counter.

**Experimental protocol:** On the day following surgery (more than 12 hr later), eels were infused with one of three homologous NPs (ANP, CNP or VNP) at 5 pmol/kg/min for 30 min successively in random order. This dose was shown to be sufficient to ensure the renal and vascular NP actions (Takei and Kaiya, 1998). Before and after each NP infusion, a control infusion of saline was made for 30 min and 1 hr, respectively. A period of 1 hr without infusion was placed between different NP infusions to exclude the influence of previous infusion. Infusion rate was 100  $\mu$ l/30min, whereas blood was collected into a capillary (40  $\mu$ l) every 30 min during infusion for measurements of hematocrit and plasma Na<sup>+</sup> concentration. At the time of exchange of infusates, 50  $\mu$ l of the next infusate was injected as a prime to obtain stable plasma NP levels quickly. The Na<sup>+</sup> concentration was determined in an atomic absorption spectrophotometer (Z5300 Hitachi, Tokyo). All measurements were made in triplicate.


**Statistical analysis:** All data were expressed in terms of percentages from the value of initial saline infusion. The ANOVA was used to determine the changes in time course data, which was followed by the Fisher's Protected Least Significant Difference (Fisher's PLSD) test at each time point. The Mann-Whitney's U-test was used for comparison between different groups. Significance was set at  $p < 0.05$ . All results were expressed as means  $\pm$  SE.

## RESULTS

**Plasma Na<sup>+</sup> concentration:** Plasma Na<sup>+</sup> concentration was 171.0  $\pm$  4.9 mM in seawater eels and 153.1  $\pm$  1.3 mM in fresh-



**Fig. 3.** The time course of changes in drinking rate during 30-min infusions of homologous ANP (○), CNP (●) and VNP (□) (5 pmol/kg/min) in seawater eels (n=5). Saline was infused before and after NP infusions. Data were expressed as percentage to the values before NP infusion. Freshwater eels did not drink before NP infusion. Values represent means  $\pm$  SE. \* $p < 0.05$ , \*\* $p < 0.01$  compared with the value before NP infusion by Fisher's PLSD test.



**Fig. 4.** Comparison of the effects of ANP (open bar), CNP (filled bar) and VNP (hatched bar) on plasma Na<sup>+</sup> concentration, hematocrit and drinking rate in seawater eels (A; n=5) and freshwater eels (B; n=6). Data were expressed as percentages from the value before NP infusions. Data on drinking rate in freshwater eels are not included because they did not drink before NP infusion. Values represent means  $\pm$  SE. \* $p < 0.05$ , \*\* $p < 0.01$  compared among NPs by Mann-Whitney's U-test. The asterisks on the shoulder of the bar of freshwater eels show significant differences from the values of seawater eels.

water eels before infusion of NPs. Plasma  $\text{Na}^+$  concentration decreased significantly during ANP or VNP infusion in seawater eels compared with the value of saline-infused controls before NP infusion (Fig. 1A). The levels were restored when infusate was changed to saline, but they were lower than the initial levels for 1 hr. By contrast, CNP infusion did not alter plasma  $\text{Na}^+$  concentration. In freshwater eels, all NPs did not alter plasma  $\text{Na}^+$  concentration except for VNP that increased it slightly after infusion (Fig. 1B).

**Hematocrit:** Hematocrit was  $23.8 \pm 1.0\%$  in seawater eels and  $24.2 \pm 0.8\%$  in freshwater eels before NP infusion. Hematocrit increased significantly after infusion of ANP or VNP in seawater eels and the increase lasted for 30–60 min (Fig. 2A). However, CNP did not change hematocrit. Similarly, only ANP and VNP increased hematocrit in freshwater eels (Fig. 2B).

**Drinking rate:** Drinking rate was  $46.2 \pm 4.5 \mu\text{l}/\text{min}$  in seawater eels and  $0.8 \pm 0.4 \mu\text{l}/\text{min}$  in freshwater eels. Drinking rate was profoundly inhibited by ANP and VNP, but CNP had no effect in seawater eels (Fig. 3). The antidipsogenic effect continued for 1 hr after termination of ANP or VNP infusion. Since no appreciable drinking was observed in freshwater eels, it was impossible to detect any inhibition of drinking rate in these fish. In fact, 5 of 6 freshwater eels did not drink at all throughout the experiment (data not shown).

**Comparison of ANP, CNP and VNP:** In seawater eels, CNP was less potent than ANP and VNP in all effects examined except for the effect on hematocrit (Fig. 4A). There were no differences in the effects of ANP and VNP on plasma  $\text{Na}^+$  concentration and drinking rate. However, the effect of VNP on hematocrit was greater than that of ANP. In freshwater eels, no differences were detected in the effects of three NPs on plasma  $\text{Na}^+$  concentration. However, CNP was less effective than ANP and VNP on increasing hematocrit (Fig. 4B). The antidipsogenic effect could not be expressed as percent changes because of no appreciable drinking before NP infusions. Comparing the effects between freshwater and seawater eels, the effects of ANP and VNP on decreasing plasma  $\text{Na}^+$  concentration is stronger in seawater eels than in freshwater eels, whereas no difference was detected for the CNP effects (Fig. 4A, B). The effects of three NPs on hematocrit did not differ between freshwater and seawater eels.

## DISCUSSION

Three members of the NP family (ANP, CNP and VNP) have been identified in the eel, but only the effect of ANP on seawater adaptation has been well established (Takei, 2000a). CNP is principally a local paracrine factor in mammals (Forkes and McArdle, 2000). However, recent study showed that CNP circulates in the blood of freshwater eels possibly by secretion from the heart and intestine (Takei *et al.*, 2001). Therefore, it is possible that not only ANP and VNP but also CNP is a circulating hormone to regulate water and electrolyte balance in eels. However, effects of the three NPs have not been compared in detail with respect to the body fluid regulation except for the hemodynamic effects (Takei *et al.*, 1991).

## Effects on plasma $\text{Na}^+$ concentration

In seawater eels, ANP and VNP infused into the circulation decreased plasma  $\text{Na}^+$  concentration, but CNP was without effect. It is known that ANP and VNP bind with higher affinity to NPR-A and increase greater cGMP accumulation compared with CNP (Kashiwagi *et al.*, 1999). Thus, a decrease in plasma  $\text{Na}^+$  concentration seems to be mediated by NPR-A. One possible route for the decrease is via  $\text{Na}^+$  excretion by the gill since ANP and VNP increase the  $\text{Na}^+, \text{K}^+$ -ATPase activity in the gill cells isolated from seawater eels (Takei *et al.*, 1999). Another possible route is through decreased oral intake of environmental seawater and decreased absorption of NaCl by the intestine. As shown in the present study, ANP and VNP greatly decreased drinking rate in seawater eels. Furthermore, ANP and VNP inhibit NaCl absorption by the intestine as shown in the Ussing's chamber *in vitro* (Ando and Hara, 1994; Loretz and Takei, 1997). In fact, we recently found that the hyponatremic effect is suppressed if eels are forced to drink environmental seawater at a normal rate during ANP infusion (T Tsukada and Y Takei, unpublished data). The hyponatremic effect of ANP is not due to increased  $\text{Na}^+$  excretion by the kidney since ANP did not cause natriuresis in seawater eels (Takei and Kaiya, 1998).

In contrast to the result in seawater eels, ANP and VNP did not decrease plasma  $\text{Na}^+$  concentration in freshwater eels. This may be due to the less important role of drinking and the subsequent intestinal absorption of NaCl and water for body fluid regulation in freshwater eels than in seawater eels. On the other hand, CNP did not alter plasma  $\text{Na}^+$  concentration in this study, although the messages of CNP-specific receptor, NPR-B (Koller *et al.*, 1991), are more abundantly expressed in the gills and other osmoregulatory organs of freshwater eels than in those of seawater eels (Katafuchi *et al.*, 1994),

## Effects on drinking

Drinking rate is generally high in seawater fish in order to compensate for water lost osmotically across the body surfaces (Hirano, 1974; Evans, 1993). Consistently, the seawater eels used in this study drank copiously and continuously throughout the experiment. Inhibition of drinking was apparent during ANP or VNP infusion, but CNP was without effect on drinking. Therefore, the antidipsogenic effect of NP is most probably mediated by NPR-A. This result is consistent with that obtained in the rat in which ANP inhibits drinking (Antunes-Rodrigues *et al.*, 1985; Katsuura *et al.*, 1986) but CNP slightly enhanced drinking (Samson *et al.*, 1991).

In contrast to seawater eels, freshwater eels used in the current study scarcely drank during the experiment. Thus, it was impossible to detect an inhibitory effect of NPs on their drinking. It is generally thought that fish in fresh water drink little to cope with an osmotic influx of water across the gills (Evans, 1993). However, ANP and VNP are potently antidipsogenic in freshwater eels when infused at low doses into the circulation (Tuchida and Takei, 1998). The low drinking rate in freshwater eels in the present study may be due to little bleeding during surgery because eels are highly responsive

to hypovolemia by drinking (Hirano, 1974; Takei, 2000b).

### Effects on hematocrit

In both seawater and freshwater eels, ANP and VNP increased hematocrit, but CNP had no effect. Therefore, NPR-A may be present in the target tissues of eels that are responsible for regulation of water permeability across the vascular wall. The effects of ANP, BNP and CNP on hematocrit have been compared in the dog (Woods and Jones, 1999). The results show that ANP and BNP increased hematocrit but CNP was without effect, showing the involvement of NPR-A. Thus, similar mechanisms may be in function for regulation of capillary permeability in eels.

The hemoconcentrating effects of ANP have been extensively studied in mammals (Renkin and Tucker, 1996). It is possible that an increase in hematocrit, i.e., the extravasation of plasma fluid, is caused by an increase in the hydraulic conductivity across the endothelial cells and/or by enlargement of the functional capillary surface area by recruitment of non-perfused capillaries. However, Valentin *et al.* (1997) reported that sodium nitroprusside, which causes similar changes in the vasculature as does ANP through increased accumulation of cGMP, has no effect on hematocrit in the nephrectomized rat. Furthermore, if only water is exuded into the interstitial space, resultant increase in plasma oncotic pressure would re-withdraw interstitial water into plasma immediately as observed after administration of furosemide (Renkin and Tucker, 1996). It is also shown that ANP contributes to the development of pulmonary edema in the rat (Tamura *et al.*, 2000). It is more likely, therefore, that ANP and VNP increase permeability of plasma proteins through the intercellular pathway toward interstitial compartment. In addition, increases in hematocrit are achieved by an increased supply of red cells from blood-storing organs such as spleen and liver, and by expansion of red cell volume (Nikinmaa and Huestis, 1984). The exact mechanisms are not fully understood yet, but the hemoconcentrating effects of NPs occur from fish to mammals.

### Comparison of ANP, CNP and VNP

In mammals, comparative studies of ANP, CNP and BNP have been performed with respect to cardiovascular and renal actions. The results show that ANP and BNP are generally equipotent but CNP is much less potent in all actions examined except for the effects on the mesenteric vasoconstriction in the dog (Woods and Jones, 1999) and on the inhibition of aldosterone secretion in the sheep (Charles *et al.*, 1996). The hypotensive actions of ANP, CNP and VNP have also been compared in the eel, which showed the potency order of VNP=ANP>CNP (Takei *et al.*, 1991). This result is similar to the effects on drinking, plasma Na<sup>+</sup> concentration and hematocrit as observed in this study. Since the potency order is similar to the order of affinity to NPR-A (Kashiwagi *et al.*, 1999), the effects on drinking, plasma Na<sup>+</sup> concentration, and hematocrit may be mediated by NPR-A.

### Comparison of seawater and freshwater eels

It is highly probable that ANP is an important hormone for promoting seawater adaptation (Takei, 2000a). In fact, our present study showed that ANP and VNP decreased plasma Na<sup>+</sup> concentration, which allows eels to survive in the high NaCl environment. Although the precise mechanisms for the hyponatremia are not understood yet, the close relationship between decreased drinking rate and hyponatremia in seawater eels suggests a role of drinking in regulation of plasma Na<sup>+</sup> concentration as mentioned above. Although the effects on drinking, plasma Na<sup>+</sup> concentration and hematocrit share the same receptor, NPR-A, the effects differ between freshwater and seawater eels probably because of the difference in the target tissues or in the density of the receptor in the same target tissue.

### ACKNOWLEDGMENTS

The authors thank Dr. Howard A. Bern of University of California at Berkeley for critical reading of the manuscript. We also thank Dr. Koji Inoue and Ms. Sanae Hasegawa of Ocean Research Institute for advice. This work was supported by Grant-in-Aid for Specially Promoted Grant (01902008) and for Creative Basic Research (12NP0201) from the Ministry of Education, Science, Sports and Culture of Japan.

### REFERENCES

- Ando M, Hara I (1994) Alteration of sensitivity to various regulators in the intestine of the eel following seawater acclimation. *J Comp Physiol A* 109: 447–453
- Antunes-Rodrigues J, McCann SM, Rogers LC, Samson WK (1985) Atrial natriuretic factor inhibits dehydration- and angiotensin II-induced water intake in the conscious, unrestrained rat. *Proc Natl Acad Sci USA* 82: 8720–8723
- Charles CJ, Espiner EA, Richards AM, Nicholls MG, Yandle TG (1996) Comparative bioactivity of atrial, brain, and C-type natriuretic peptides in conscious sheep. *Am J Physiol* 270: R1324–R1331
- Evans DH (1993) Osmotic and ionic regulation. In "The Physiology of Fishes" Ed by DH Evans, CRC Press, Boca Raton, pp 315–341
- Farrell AP, Olson KR (2000) Cardiac natriuretic peptides: a physiological lineage of cardioprotective hormone? *Physiol Biochem Zool* 73: 1–11
- Fowkes RC, McArdle CA (2000) C-type natriuretic peptide: an important neuroendocrine regulator? *Trends Endocrinol Metab* 11: 333–338
- Hirano T (1974) Some factors regulating water intake by the eel, *Anguilla japonica*. *J Exp Biol* 61: 737–747
- Kaiya H, Takei Y (1996) Changes in plasma atrial and ventricular natriuretic peptide concentrations after transfer of eels from freshwater to seawater or vice versa. *Gen Comp Endocrinol* 104: 337–345
- Kashiwagi M, Katafuchi T, Kato A, Inuyama H, Ito T, Hagiwara H, Takei Y, Hirose S (1995) Cloning and properties of a novel natriuretic peptide receptor, NPR-D. *Eur J Biochem* 233: 102–109
- Kashiwagi M, Miyamoto K, Takei Y, Hirose S (1999) Cloning, properties and tissue distribution of natriuretic peptide receptor-A of euryhaline eel, *Anguilla japonica*. *Eur J Biochem* 259: 204–211
- Katafuchi T, Takashima A, Kashiwagi M, Hagiwara H, Takei Y, Hirose S (1994) Cloning and expression of eel natriuretic-peptide receptor B and comparison with its mammalian counterparts. *Eur J Biochem* 222: 835–842
- Katsuura G, Nakamura M, Inouye K, Kono M, Nakao K, and Imura H (1986) Regulatory role of atrial natriuretic polypeptide in water

- drinking in rats. *Eur J Pharmacol* 121: 285–287
- Koller KJ, Lowe DG, Bennett GL, Minamino N, Kangawa K, Matsuo H, Goeddel DV (1991) Selective activation of the B natriuretic peptide receptor by C-type natriuretic peptide (CNP). *Science* 252: 120–123
- Loretz CA, Pollina C (2000) Natriuretic peptides in fish physiology. *Comp Biochem Physiol A* 125: 169–187
- Loretz CA, Takei Y (1997) Natriuretic peptide inhibition of intestinal salt absorption in the Japanese eel: physiological significance. *Fish Physiol Biochem* 17: 319–324
- Maack T (1992) Receptors of atrial natriuretic factor. *Annu Rev Physiol* 54: 11–27
- Mishina S, Takei Y (1997) Characterization of natriuretic peptide receptors in eel gill. *J Endocrinol* 154: 415–422
- Nikinmaa M, Huestis WH (1984) Adrenergic swelling of nucleated erythrocytes: cellular mechanisms in a bird, domestic goose, and two teleosts, striped bass and rainbow trout. *J Exp Biol* 113: 215–224
- Renkin EM, Tucker VL (1996) Atrial natriuretic peptide as a regulator of transvascular fluid balance. *News Physiol Sci* 11: 138–143
- Samson WK, Skala KD, Huang FL (1991) CNP-22 stimulates, rather than inhibits, water drinking in the rat: evidence for a unique biological action of the C-type natriuretic peptides. *Brain Res* 568: 285–288
- Takei Y (2000a) Structural and functional evolution of the natriuretic peptide system in vertebrates. *Int Rev Cytol* 194: 1–66
- Takei Y (2000b) Comparative physiology of body fluid regulation in vertebrates with special reference to thirst regulation. *Jap J Physiol* 50: 171–186
- Takei Y, Balment RJ (1993) Biochemistry and physiology of a family of eel natriuretic peptides. *Fish Physiol Biochem* 11: 183–188
- Takei Y, Flik G, Hirose S (1999) Interaction of the natriuretic peptide system and the rennin-angiotensin system in eel osmoregulation. In "Recent Developments in Comparative Endocrinology and Neurobiology" Ed by Roubos EW, Wendelaar-Bonga SE, Vaudry H, de Loof A, Shaker Publishers, Maastricht 144–148
- Takei Y, Inoue K, Ando K, Ihara T, Katafuchi T, Kashiwagi M, Hirose S (2001) Enhanced expression and release of C-type natriuretic peptide in freshwater eels. *Am J Physiol* 280: R1727–1735
- Takei Y, Kaiya H (1998) Antidiuretic effect of eel ANP infused at physiological doses in conscious, seawater-adapted eels, *Anguilla japonica*. *Zool Sci* 15: 399–404
- Takei Y, Takahashi A, Watanabe TX, Nakajima K, Sakakibara S (1991) A novel natriuretic peptide isolated from eel cardiac ventricles. *FEBS Lett* 282: 317–320
- Takei Y, Tsuchida T, Tanakadate A (1998) Evaluation of water intake in seawater adaptation in eels using synchronized drop counter and pulse injector system. *Zool Sci* 15: 677–682
- Tamura K, Takamori S, Mifune H, Hayashi A and Shirouzu K (2000) Changes in atrial natriuretic peptide concentration and expression of its receptors after pneumonectomy in the rat. *Clin Sci* 99: 343–348
- Tsuchida T, Takei Y (1998) Effects of homologous atrial natriuretic peptide on drinking and plasma ANG II level in eels. *Am J Physiol* 275: R1605–R1610
- Valentin JP, Ribstein J, Nafrialdi N, Mimran A (1997) Influence of dopamine receptor and adrenoceptor blockade on the hemoregulating and hypotensive actions of atrial natriuretic peptide. *Can J Physiol Pharmacol* 75: 1142–1147
- Woods RL, Jones MJ (1999) Atrial, B-type, and C-type natriuretic peptides cause mesenteric vasoconstriction in conscious dogs. *Am J Physiol* 276: R1443–R1452

(Received July 24, 2001 / Accepted September 22, 2001)