



## **Arginine Kinase from the Tardigrade, *Macrobiotus occidentalis*: Molecular Cloning, Phylogenetic Analysis and Enzymatic Properties**

Authors: Uda, Kouji, Ishida, Mikako, Matsui, Tohru, and Suzuki, Tomohiko

Source: Zoological Science, 27(10) : 796-803

Published By: Zoological Society of Japan

URL: <https://doi.org/10.2108/zsj.27.796>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Arginine Kinase from the Tardigrade, *Macrobiotus occidentalis*: Molecular Cloning, Phylogenetic Analysis and Enzymatic Properties

Kouji Uda<sup>1\*</sup>, Mikako Ishida<sup>2</sup>, Tohru Matsui<sup>2</sup> and Tomohiko Suzuki<sup>1</sup>

<sup>1</sup>Laboratories of Biochemistry, Faculty of Science, Kochi University, Kochi 780-8520, Japan

<sup>2</sup>Plant Taxonomy, Faculty of Science, Kochi University,  
Kochi 780-8520, Japan

Arginine kinase (AK), which catalyzes the reversible transfer of phosphate from ATP to arginine to yield phosphoarginine and ADP, is widely distributed throughout the invertebrates. We determined the cDNA sequence of AK from the tardigrade (water bear) *Macrobiotus occidentalis*, cloned the sequence into pET30b plasmid, and expressed it in *Escherichia coli* as a 6x His-tag-fused protein. The cDNA is 1377 bp, has an open reading frame of 1080 bp, and has 5'- and 3'-untranslated regions of 116 and 297 bp, respectively. The open reading frame encodes a 359-amino acid protein containing the 12 residues considered necessary for substrate binding in *Limulus* AK. This is the first AK sequence from a tardigrade. From fragmented and non-annotated sequences available from DNA databases, we assembled 46 complete AK sequences: 26 from arthropods (including 19 from Insecta), 11 from nematodes, 4 from mollusks, 2 from cnidarians and 2 from onychophorans. No onychophoran sequences have been reported previously. The phylogenetic trees of 104 AKs indicated clearly that *Macrobiotus* AK (from the phylum Tardigrada) shows close affinity with *Epiperipatus* and *Euperipatoides* AKs (from the phylum Onychophora), and therefore forms a sister group with the arthropod AKs. Recombinant 6x His-tagged *Macrobiotus* AK was successfully expressed as a soluble protein, and the kinetic constants ( $K_m$ ,  $K_d$ ,  $V_{max}$  and  $k_{cat}$ ) were determined for the forward reaction. Comparison of these kinetic constants with those of AKs from other sources (arthropods, mollusks and nematodes) indicated that *Macrobiotus* AK is unique in that it has the highest values for  $k_{cat}$  and  $K_d/K_m$  (indicative of synergistic substrate binding) of all characterized AKs.

**Key words:** guanidino kinase, phosphagen kinase, arginine kinase, creatine kinase, water bear, *Macrobiotus occidentalis*

## INTRODUCTION

Phosphagen (guanidino) kinases catalyze the reversible transfer of the high-energy phosphoryl group of ATP to naturally occurring guanidine compounds. Members of this enzyme family play a key role in animals as ATP-buffering systems in cells that display high and variable rates of ATP turnover. Phosphorylated high-energy guanidines are referred to as phosphagens. In vertebrates, phosphocreatine is the only phosphagen, and the corresponding phosphagen kinase is creatine kinase (CK). In contrast, invertebrates have various phosphagens in addition to phosphocreatine: phosphoglycocyamine (catalyzed by glycocyamine kinase: GK), phosphotaurocyamine (taurocyamine kinase: TK), phosphohypotaurocyamine (hypotaurocyamine kinase: HTK), phospholombricine (lombricine kinase: LK) and phosphoarginine (arginine kinase: AK). Phosphagen kinases are phylogenetically separated into two distinct groups: the AK group, which includes AK and HTK, and the

CK group, which includes CK, GK, LK and TK (Ellington, 2001; Wyss et al., 1992; Schlattner et al., 2005; McLeish and Kenyon, 2005; Ellington and Suzuki, 2006; Uda et al., 2005a). Interestingly, several AKs such as those from the echinoderm *Stichopus* and the annelid *Sabellastarte* are clustered in the CK group, indicating that they have evolved secondarily from CK (Suzuki et al., 1999; Uda and Suzuki, 2007).

Most AKs are monomers of 40 kDa, but in some species they exist as dimers (Seals and Grossman, 1988; Suzuki et al., 1999) or contiguous dimers (two-domain AKs), presumably as a result of gene duplication and subsequent fusion (Suzuki et al., 1997; Suzuki et al., 1998).

Typical AKs are most widely distributed among organisms such as arthropods, mollusks, nematodes, cnidarians, poriferae, protozoans (ciliates and choanoflagellates), and bacteria, indicating their ancient origin (Andrews et al., 2008; Uda et al., 2006). In three major invertebrate groups (arthropods, nematodes, and mollusks), AK is the only phosphagen

\* Corresponding author. Phone: +81-88-844-8488;  
Fax : +81-88-844-8359;  
E-mail: k-uda@cc.kochi-u.ac.jp  
doi:10.2108/zsj.27.796

## ABBREVIATIONS

AK, arginine kinase; CK, creatine kinase; GK, glycocyamine kinase; GS region, guanidine specificity region; LK, lombricine kinase; TK, taurocyamine kinase; EST, expressed sequence tag.



**Table 1.** AKs used for the phylogenetic analysis.

Phylum	Class	Order	Genus/Species/Isoform	Accession number/Database <sup>a</sup>			
Alveolata	Oligohymenophorea	Hymenostomatida	<i>Tetrahymena thermophila</i> AK1 <i>Tetrahymena thermophila</i> AK2	EAS01428 EAS01429			
Arthropoda	Arachnida	Arachnida	<i>Aleuroglyphus ovatus</i> AK <i>Loxosceles laeta</i> AK	ABU97463 EY188599			
		Araneae	<i>Aphonopelma</i> sp. AK	Genbank EST : FC823446, FC824317			
		Astigmata	<i>Dermatophagoides farinae</i> AK1 <i>Dermatophagoides farinae</i> AK2	AAP57094 ABU97470			
		Ixodida	<i>Ixodes scapularis</i> AK	Genbank EST : EW821872, EW873512			
		Prostigmata	<i>Tetranychus urticae</i> AK	Trace Archive : 2267574886, 2267695435			
		Anostraca	<i>Artemia franciscana</i> AK	AAL25092			
		Diplostraca	<i>Daphnia pulex</i> AK	Trace Archive : 895565747, 897280293, 895554084			
		Insecta	Blattaria	<i>Blattella germanica</i> AK	ABC86902		
			Blattaria	<i>Periplaneta americana</i> AK	AAT77152		
				Coleoptera	<i>Tribolium castaneum</i> AK	Trace Archive : 569305708, 580631152	
				Diptera	<i>Drosophila melanogaster</i> AK <i>Anopheles gambiae</i> AK <i>Aedes aegypti</i> AK <i>Ceratitis capitata</i> AK	AAN11983 EAA44056 ABF18260 Genbank EST : FG083307, FG075954	
					<i>Drosophila pseudoobscura</i> AK	Genbank EST : DR124999, DR145664	
<i>Glossina morsitans</i> AK	Genbank EST : DV618298, FM982907						
<i>Lutzomyia longipalpis</i> AK	Genbank EST : AM109228, AM109239						
<i>Phlebotomus papatasi</i> AK	Genbank EST : EY204603, EY214760						
<i>Cochliomyia hominivorax</i> AK	Genbank EST : FG300496, FG296874						
<i>Teleopsis dalmanni</i> AK	Genbank EST : GO297058, GO298184						
<i>Homalodisca vitripennis</i> AK	AAT01074						
<i>Oncometopia nigricans</i> AK	AAU95198						
<i>Nilaparvata lugens</i> AK	Genbank EST : DB840416, DB826716						
<i>Rhodnius prolixus</i> AK	Genbank EST : EH114777, FG544166						
		Hymenoptera	<i>Solenopsis invicta</i> AK	ACF04198			
			<i>Apis mellifera</i> AK	AAC39040			
			<i>Nasonia vitripennis</i> AK	Trace Archive : 1081135584, 1076813375, 1068958665, 1105139233			
			<i>Lysiphlebus testaceipes</i> AK	Genbank EST : EH010491, EH015342, EH010390			
			Lepidoptera	<i>Plodia interpunctella</i> AK	CAC85911		
				<i>Bombyx mori</i> AK	ABD36282		
						<i>Danaus plexippus</i> AK	Genbank EST : EY260080, EY271098
						<i>Spodoptera frugiperda</i> AK	Genbank EST : DV076460, DY898274
						<i>Manduca sexta</i> AK	Genbank EST : BF046795, BE015379, BE015528
						<i>Trichoplusia ni</i> AK	Genbank EST : CF259256, FF370292
						<i>Ostrinia nubilalis</i> AK	Genbank EST : GH997366, GH989259
						Orthoptera	<i>Schistocerca americana</i> AK
<i>Locusta migratoria</i> AK	ABF68036						
		Phthiraptera				<i>Gryllus bimaculatus</i> AK	Genbank EST : DC443130, DC446501
						<i>Pediculus humanus</i> AK	Trace Archive : 1382191351, 1379696849, 1386063845
		Malacostraca				<i>Gammarus pulex</i> AK	Genbank EST : EH275731, EH275602
			<i>Pachygrapsus marmoratus</i> AK	AAG01175			
			<i>Litopenaeus vannamei</i> AK	ABI98020			
			<i>Fenneropenaeus chinensis</i> AK	AAV83993			
			<i>Neohelice granulata</i> AK	AAF43438			
			<i>Callinectes sapidus</i> AK	AAF43436			
			<i>Marsupenaeus japonicus</i> AK	AAB31477			
			<i>Homarus gammarus</i> AK	CAA48654			
			<i>Procambarus clarkii</i> AK	2020435A			
			<i>Neocaridina denticulata</i> AK	BAH56609			
			<i>Penaeus monodon</i> AK	AAO15713			
			<i>Eriocheir sinensis</i> AK	AAF43437			
			<i>Petrolisthes cinctipes</i> AK	Genbank EST : FE756031, FE750140			
			<i>Carcinus maenas</i> AK	AAD48470			
			<i>Eurydice pulchra</i> AK	Genbank EST : CO869027, CO868808, CO868911			
			<i>Limulus polyphemus</i> AK	P51541			
			<i>Homo sapiens</i> MCK <sup>b</sup>	AAA96609			
			<i>Anthopleura japonica</i> 2DAK	O15992			
			<i>Aiptasia pallida</i> AK	Genbank EST : GH579704, GH574852, GH575418			
			<i>Acropora millepora</i> 2DAK	Genbank EST : DY586394, EZ016454, EH038119, EH037125			
		Chordata	<i>Scapharca broughtonii</i> AK	BAD11949			
			<i>Crassostrea gigas</i> AK	BAD11950			
			Cephalopoda	<i>Nautilus pompilius</i> AK	BAA95594		
				<i>Octopus vulgaris</i> AK	BAA95609		
			Gastropoda	<i>Septoteuthis lessoniana</i> AK	BAA95610		
				<i>Aplysia kurodai</i> AK	BAB41095		
			Dacrydium	<i>Cellana grata</i> AK	BAB41096		
				<i>Haliotis madaka</i> AK	P51544		
			Polyplacophora	<i>Batillius cornutus</i> AK	BAA22870		
				<i>Liolophura japonica</i> AK	BAA22871		
		Cephalopoda	<i>Euprymna scolopes</i> AK	Genbank EST : DW282592, DW279554			
			<i>Idiosepius paradoxus</i> AK	Genbank EST : DB918583, DB916072, DB919901			
			<i>Aplysia californica</i> AK	Trace Archive : 1161815795, 1809265942, 1182066208, 1162368191			
			<i>Biomphalaria glabrata</i> AK	Genbank EST : ES491406, FC856201			
			Nematoda	<i>Trichinella spiralis</i> AK	Trace Archive : 1724989270, 1724991545		
				<i>Toxocara canis</i> AK	ABK76312		
						<i>Pristionchus pacificus</i> AK1	Trace Archive : 989893386, 987437388, 760524991
						<i>Pristionchus pacificus</i> AK2	Genbank EST : FG097924, B1500767, A1988904
						<i>Caenorhabditis elegans</i> AK1	AAO21426
						<i>Caenorhabditis elegans</i> AK2	CAB00062
<i>Caenorhabditis elegans</i> AK3	CAB05517						
<i>Caenorhabditis elegans</i> MiAK	AAK21503						
<i>Heterorhabditis bacteriophora</i> AK	Trace Archive : 1877615891, 1949656867						
<i>Haemonchus contortus</i> AK	Genbank EST : CB015139, BM139164						
<i>Strongyloides ratti</i> AK1	Genbank EST : BI073820, FC816131, FC816421						
<i>Strongyloides ratti</i> AK2	Genbank EST : FC812688, FC818348, BI742298						
			<i>Heterodera glycines</i> AK1	AAO49799			
			<i>Heterodera glycines</i> AK2	AAP41028			
			<i>Globodera rostochiensis</i> AK	Genbank EST : BM355956, BM354963			
			<i>Meloidogyne hapla</i> AK	Genbank EST : CA997516, CA997485			
			<i>Xiphinema index</i> AK	Genbank EST : CV568581, CV509691, CV581377			
			<i>Dictyocaulus viviparus</i> AK	Genbank EST : EV853193, EV851844			
			<i>Epiperipatus</i> sp. AK	Genbank EST : AM499754, AM500583			
			<i>Euperipatoides kanangrensis</i> AK	Trace Archive : 1987166188, 1987167250			
			<i>Paragonimus westermani</i> TK <sup>c</sup>	ACT37385			
			<i>Siphonosoma cumanense</i> HTK <sup>c</sup>	BAE16970			

<sup>a</sup>For sequences obtained from GenBank, accession numbers are shown. For the assembled sequences in this study, the database name used and accession numbers are shown.

<sup>b</sup>*Homo sapiens* MCK is used as an outgroup.

<sup>c</sup>Recent phylogenetic analyses of *Paragonimus* TK and *Siphonosoma* HTK indicate that they evolved from AK genes (Uda et al., 2005; Jarilla et al., 2009).

\*The 46 newly assembled sequences.

tion. The Maximum-Likelihood (ML) analysis with the approximate likelihood-ratio test for branches (aLRT; Anisimova and Gascuel, 2006) was performed in the program PhyML v3.0 (Guindon and Gascuel, 2003) using the LG amino acid replacement matrix.

## RESULTS AND DISCUSSION

cDNA for AK from *Macrobiotus occidentalis* was amplified by PCR and cloned into the plasmids pGEM-T Easy and pET30b. Fig. 1 shows the nucleotide and derived amino acid sequences of *Macrobiotus* AK. The nucleotide sequence consists of 1377 bp, with an open reading frame (ORF) of 1080 bp, and 5'- and 3'-untranslated regions of 116 and 297 bp, respectively. The sequence was deposited into the DDBJ database (accession number: AB537977). This is the first reported AK sequence from a tardigrade.

The ORF codes were consistent with a protein of 359 amino acid residues, with a calculated molecular mass of 40,060 Da and an estimated pI of 6.81. When the amino acid sequence was compared with *Limulus* AK, for which the crystal structure has been determined (Zhou et al., 1998), it was found that *Macrobiotus* AK completely conserved all key residues believed necessary for AK function (underlined in Fig. 1). Conserved residues include seven that interact with the substrate arginine in *Limulus* AK (S63, G64, V65, Y68, E228, C274 and E317) and five residues that interact with the substrate ADP (R127, R129, R232, R283 and R312). The results show that *Macrobiotus* AK and *Limulus* AK may have very similar substrate recognition systems.

At present, at least 60 complete sequences of invertebrate AKs have been deposited in protein or DNA databases. We also know that many EST or genomic DNA databases contain fragmented and non-annotated AK sequences. We performed a comprehensive search for AK fragments across multiple databases using known AK

sequences as references, and assembled the fragments into complete cDNA sequences. As a result, we obtained 46 complete AK sequences: 26 from arthropods (including 19 from Insecta (Coleoptera: *Tribolium castaneum*, Diptera: *Ceratitis capitata*, *Drosophila pseudoobscura*, *Glossina morsitans*, *Lutzomyia longipalpis*, *Phlebotomus papatasi*, *Cochliomyia hominivorax*, *Teleopsis dalmanni*, Hemiptera: *Nilaparvata lugens*, *Rhodnius prolixus*, Hymenoptera: *Nasonia vitripennis*, *Lysiphlebus testaceipes*, Lepidoptera: *Danaus plexippus*, *Spodoptera frugiperda*, *Manduca sexta*, *Trichoplusia ni*, *Ostrinia nubilalis*, Orthoptera: *Gryllus bimaculatus*, Phthiraptera: *Pediculus humanus*)), three from cnidarians, four from mollusks, 11 from nematodes and two from onychophorans (see Table 1). These onychophoran AK sequences are the first to be reported for that taxon.

The amino acid sequences of 104 invertebrate AKs, including *Macrobiotus* AK, the 46 AKs obtained by our in silico analyses (Table 1), and *Paragonimus* TK and *Siphonosoma* HTK (both of which evolved from AK genes; Uda et al., 2005; Jarilla et al., 2009), were aligned using the ClustalW program (data not shown). The sequence of *Macrobiotus* AK showed the highest identity (75%) with AK from the onychophorans *Epiperipatus* and *Euperipatoides*, 62–74% with arthropod AKs, 59–65% with nematode AKs, and 49–55% with mollusk AKs.


A phylogenetic tree was constructed from the above alignments using the ML (Fig. 2) and NJ (data not shown) methods. The two trees show similar topology, and the protostome AK sequences are separated into two distinct groups: lophotrochozoans (mollusks, platyhelminths and sipunculids) and ecdysozoans (arthropods, nematodes, onychophorans and tardigrades). Recent molecular phylogenetic studies suggest three possibilities for the phylogeny of ecdysozoans: (a) Tardigrada and Onychophora are included within Arthropoda (Colgan et al., 2008), (b) Tardigrada has

```

1 gaaocggactggaagaagcogagacatttatttgcactcttggttttttcagtcgcaaaacgactatcagccgcccctttttgtcgtttctt 90
91 cttcttttggtaatacgcaggttttgcattggccgctctgtgacacgctcagaaaattctccgaggctcctggcattcttgcgaaggagaccaaa 180
1 M A A V D H A Q K I S E A P G I L Q G D Q K 22
181 AGGGACACTCCCTGCTCAAGAAATACCTGTGCGAAAGATGTCGCGAGAAAAGTTGAAGAAGCACAACCTGGCATGGGTGCCAGCCTTTGGG 270
23 G H S L L K K Y L S K D V A E K L K N D K T G M G A S L L W D 52
271 ACTGCATCCAGTCTGGTGTGGCCAATCTGGACAGCGGTGTGGCATCTACGCCCTGATGCGGAATCCTACACCAAATTCGGATGTCT 360
53 C I Q S G V A N L D S G V G I Y A P D A E S Y T K F S D V F 82
361 TCTATCCCATCATCCAGGATTACCACATGGATTGACACTGAAGCTGGAGCCAAACCCACCGGCTGACTTCGGTCTGGACAACTCA 450
83 Y P I I Q D Y H I G F D L K A G A K H P P A D F G L D K L N 112
451 ATTTCCCAATCCGACCCGACTGGCGAATACATATTTGCACTCGCGTCCGATGGTTCGCTCGCTGGCTGGATATCCGTTCAACCCCG 540
113 F P N P D P T G E Y I I S T R V R C G R S L A G Y P F N P L 142
541 TCTTAAACGAAACCAATATAAAGAAATGGAAGAGAAAGTGAAGAGCGCACTCACTGGATTGACCGGAGAACTAGCCGGCACTTACTACC 630
143 L N E Q Q Y K E M E E K V K S A L T G L T G E L A G T Y Y P 172
631 CACTTACCGCATGGACAAGGCCACCCAAAACCACTCATCGAGGACATTTCTTGTTCAGGAGGGAGATCGTTCTTGAAGCTGCCA 720
173 L T G M D K A T Q N Q L I E D H F L F K E G D R F L Q A A N 202
Phos.con
721 ACGTAGCCGTTTCTGGCCACTGGTCTGGAATCTTCCACAACAAGGACAAGACTTTCCTGGTCTGGGTCAACGAGGAGGACCATCTCC 810
203 A S R F W P T G R G I F H N K D K T F L V W V N E E D H L R 232
kuma AK R2
811 GCATCATCAGCATGCAAAAGGGCGCGATTGTTGGCAGTCTTCAAGCCTCTGATTGAGGGTGTGAAACAAATCGAGGCGCAAACTGCCCT 900
233 I I S M Q K G G D L L A V F K R L I E G V K Q I E A K L P F 262
kuma AK R1
901 TCTCCCGTGTGACCCGCTGGGTTATTGACTTTTGCCTCCACCAACCTGGGCACCACCATCCGCGCCAGTGTGCATATCAAGTACCCCA 990
263 S R D D R L G Y L T F C P T N L G T T I R A S V H I K L P K 292
991 AGATCAGCAAAACCTCGACGAGTTCCACAAGATTGCCGCTAAATATAACCTCCAAGTTCGTGGTACATCTGGAGAACACTCAGAATCCA 1080
293 I S K N L D E F H K I A A K Y N L Q V R G T S G E H S E S I 322
1081 TCGCGGAGTTTACGACGCTCCCAACAAGCGTCGCATGGGTCTGACCGAATACGATGCCGTCAGGAAATGTACGCGGTATTGTCGAAT 1170
323 G G V Y D V S N K R R M G L T E Y D A V K E M Y D G I V E L 352
1171 TGATCAAGCTGGAGAAAGCTTCTGAgctctggattttgtgcaaatgattgtctgagactcctctactagcgaacaccttggcggctgcaaa 1260
353 I K L E K A S * 359
1261 tgcctgacctgaccctgttttttctttgacacattgtgacctgctagaatgcaattggccgagaacgaacgaactttaaattgttcogcgagaat 1350
1351 tttgacctataaaqcaactaqaacqcc 1377

```

**Fig. 1.** Nucleotide and derived amino acid sequence of cDNA of *Macrobiotus* AK. Primers used to amplify the cDNA are shown by arrows. The key residues interacting with the substrates, arginine and ADP, are underlined.




**Fig. 2.** Maximum-likelihood (ML) tree for amino acid sequences of invertebrate AKs. The tree was constructed using the PhyML program. The approximate likelihood-ratio test (aLRT) values are shown at the branching points. *Homo* muscle-type creatine kinase was used as an out-group. Accession numbers of the sequences are listed in Table 1. *Macrobiotus* AK is boxed, and the 46 newly assembled sequences are marked by asterisks.

close affinity with Onychophora, and they form a sister group with Arthropoda (Mallatt and Giribet, 2006), and (c) Onychophora has close affinity with Arthropoda, and they form a sister group with Tardigrada (Dunn et al., 2008). Our

phylogenetic tree (Fig. 2) indicates that AK from the tardigrade *Macrobiotus* has very close affinity with onychophoran AKs, and forms a sister group with the arthropod AKs. Thus, our analyses support possibility (b), which was originally deduced

from 28S and 18S rRNA analyses using the ML method (Mallatt and Giribet, 2006; Mallatt et al., 2004).

Recombinant 6x His-tagged *Macrobiotus* AK was successfully expressed as a soluble protein, and purified by affinity chromatography. Fig. 3 shows the result of SDS-



**Fig. 3.** SDS-PAGE of His-tagged *Macrobiotus* AK. Lane 1, marker proteins (Precision Plus Protein Standards, Bio Rad). Lane 2, soluble proteins from the *E. coli* crude extract. Lane 3, His-tagged *Macrobiotus* AK enzyme purified by affinity chromatography.

PAGE of the purified recombinant enzyme. The recombinant enzyme gave a major single band with a molecular mass of 40 kDa (lane 3), suggesting that the enzyme is sufficiently pure to allow determination of its kinetic constants.

The kinetic constants for *Macrobiotus* AK were obtained using software written by R. Viola (Enzyme Kinetics Programs, ver. 2.0); the results are summarized in Table 2. The kinetic constants were compared with those of AKs from other sources: the arthropods *Locusta* (Wu et al., 2007; Li et al., 2006), *Neocaridina* (Iwanami et al., 2009), *Cissites* (Tanaka et al., 2007), and *Periplaneta* (Brown and Grossman, 2004), the nematode *Toxocara* (Wickramasinghe et al., 2007), the mollusks *Nautilus* (Uda and Suzuki, 2004; Matsumoto and Suzuki, unpublished data), *Scapharca* (Takeuchi et al., 2004), *Octopus* (Takeuchi et al., 2004), and *Crassostrea* (Fujimoto et al., 2005), and the sea anemone *Anthopleura* (Tada et al., 2008; Tada et al., 2010) (Table 2).

The values for  $K_m^{\text{arg}}$  (0.68 mM) and  $K_m^{\text{ATP}}$  (0.86 mM) from *Macrobiotus* AK are in the range found for other AKs: 0.12–1.44 mM for  $K_m^{\text{arg}}$  and 0.14–2.17 mM for  $K_m^{\text{ATP}}$ .

The  $K_d/K_m$  and  $k_{\text{cat}}$  values for *Macrobiotus* AK appear to be unique. In many phosphagen kinase reactions, two substrates, arginine (or phosphoarginine) and MgATP (or MgADP) in AK reaction, typically exhibit synergistic binding to AK. That is, binding of the first substrate facilitates binding of the second substrate. In terms of kinetic constants, this means that  $K_d$ , the dissociation constant in the absence of the second substrate, is higher than  $K_m$  ( $K_d/K_m > 1$ ). This synergism may be associated with substrate-induced conformational changes within the tertiary complex. In previous works, we showed that the amino acid residues at positions 62 and 193 (positions relative to *Limulus* AK), which are conserved in normal AKs, including *Macrobiotus* AK, as Asp and Arg, respectively, form a hydrogen bond in the transition state analogue complex in *Limulus* AK (Zhou et al., 1998) and are key residues for synergism (Suzuki et al., 2000; Takeuchi et al., 2004; Fujimoto et al., 2005). Interestingly, *Macrobiotus* AK exhibits higher synergism in substrate binding ( $K_d/K_m = 5.78$ ) than do other AKs ( $K_d/K_m = 0.9$ –3.99; Table 2). In addition, the  $k_{\text{cat}}$  value ( $291 \text{ s}^{-1}$ ) of *Macrobiotus*

**Table 2.** Comparison of kinetic constants of invertebrate AKs at 25°C for the forward reaction (phosphagen synthesis).

Source	Enzyme state	Reference	$K_m^{\text{arg}}$ (mM)	$K_d^{\text{arg}}$ (mM)	$K_m^{\text{ATP}}$ (mM)	$K_d^{\text{ATP}}$ (mM)	$k_{\text{cat}}$ (1/s)	$K_d/K_m$
Tardigrada								
<i>Macrobiotus</i>	His-tag	This work	$0.683 \pm 0.15$	$3.95 \pm 0.70$	$0.858 \pm 0.119$	$4.96 \pm 1.16$	$291 \pm 27$	5.78
Arthropoda								
<i>Locusta</i>	Native	Li et al. (2006)	0.94		1.29		163	
	no tag	Wu et al. (2007)	$0.951 \pm 0.08$	$2.67 \pm 0.22$	$1.27 \pm 0.23$	$3.56 \pm 0.32$	$159 \pm 6.2$	3.2
<i>Neocaridina</i>	His-tag	Iwanami et al. (2009)	$0.376 \pm 0.039$	$0.466 \pm 0.078$	$0.989 \pm 0.064$	$1.23 \pm 0.23$	$200 \pm 5.2$	1.24
<i>Cissites</i>	MBP-tag	Tanaka et al. (2007)	$1.01 \pm 0.07$	$0.99 \pm 0.03$	$0.95 \pm 0.16$	$0.92 \pm 0.16$	$2.02 \pm 0.05$	0.99
<i>Periplaneta</i>	Native	Brown and Grossman (2004)	0.49	0.45	0.14	0.17	1.30	0.92
Nematoda								
<i>Toxocara</i>	MBP-tag	Wickramasinghe et al. (2007)	$0.12 \pm 0.003$	$0.23 \pm 0.03$	$0.30 \pm 0.04$	$0.60 \pm 0.07$	$29.2 \pm 0.19$	1.96
Mollusca								
<i>Nautilus</i>	MBP-tag	Uda and Suzuki (2004)	$0.67 \pm 0.11$	$2.26 \pm 0.07$	$1.40 \pm 0.11$	$4.72 \pm 0.36$	$2.51 \pm 0.16$	3.37
	His-tag	Matsumoto and Suzuki (unpublished data)	$0.56 \pm 0.01$				$33.0 \pm 0.60$	
<i>Crassostrea</i>	MBP-tag	Fujimoto et al. (2005)	$0.35 \pm 0.01$	$0.82 \pm 0.37$	$0.97 \pm 0.25$	$2.26 \pm 0.59$	$79.7 \pm 3.44$	2.34
<i>Scapharca</i>	MBP-tag	Takeuchi et al. (2004)	$1.44 \pm 0.28$	$2.57 \pm 0.29$	$0.65 \pm 0.15$	$1.16 \pm 0.25$	$72.1 \pm 7.5$	1.78
<i>Octopus</i>	MBP-tag	Takeuchi et al. (2004)	$0.95 \pm 0.033$	$3.78 \pm 0.05$	$0.75 \pm 0.121$	$4.72 \pm 0.36$	$29.4 \pm 0.72$	3.99
Cnidaria								
<i>Anthopleura</i>	MBP-tag	Tada et al. (2008)	$0.25 \pm 0.04$	$0.33 \pm 0.07$	$2.17 \pm 0.20$	$2.83 \pm 0.83$	$129 \pm 5.26$	1.32
	His-tag	Tada and Suzuki (2010)	$0.28 \pm 0.05$	$0.30 \pm 0.08$	$1.52 \pm 0.16$	$1.61 \pm 0.55$	$678 \pm 33$	1.07

AK is also higher than other AKs (1.3–200 s<sup>-1</sup>; Table 2), except for that (678 s<sup>-1</sup>) of *Anthopleura* His-tagged AK, which exhibits an unusual two-domain structure (Tada and Suzuki, 2010). These results indicate that *Macrobiotus* AK is distinguished from other AKs by its high  $k_{\text{cat}}$  and  $K_d/K_m$  values.

We determined preliminary temperature/activity profiles at pH 8.0 for His-tagged recombinant *Macrobiotus* AK and *Nautilus* AK, a well-characterized AK (Fig. 4). Comparison of the profiles indicates that the optimum temperature of *Macrobiotus* AK appears to be shifted about 10°C to the high temperature region, and maintains higher activity over 35°C, compared with *Nautilus* AK.


These characteristics of *Macrobiotus* AK (high  $k_{\text{cat}}$  and  $K_d/K_m$  values, and differences in temperature-dependent activity) may be related to the survival of *Macrobiotus occidentalis* under extreme conditions.

#### ACKNOWLEDGEMENTS

This work was supported by a Grant-in-Aid for Scientific Research in Japan to KU (21770080) and to TS (17570062 and 20570072).

#### REFERENCES

- Andrews LD, Graham J, Snider MJ, Fraga D (2008) Characterization of a novel bacterial arginine kinase from *Desulfotalea psychrophila*. *Comp Biochem Physiol B* 150: 312–319
- Anisimova M, Gascuel O (2006) Approximate likelihood-ratio test for branches: A fast, accurate, and powerful alternative. *Syst Biol* 55: 539–552
- Borson ND, Salo WL, Drewes LR (1992) A lock-docking oligo(dT) primer for 5' and 3' RACE PCR. *PCR Meth Appl* 2: 144–148
- Brown A, Grossman SH (2004) The mechanism and modes of inhibition of arginine kinase from the cockroach (*Periplaneta americana*). *Arch Insect Biochem Physiol* 57: 166–177
- Chomczynski P, Sacchi N (1987) Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal Biochem* 162: 156–159
- Cleland WW (1979) Statistical analysis of enzyme kinetic data. *Method Enzymol* 63: 103–138
- Colgan DJ, Hutchings PA, Beacham E (2008) Multi-gene analyses of the phylogenetic relationships among the Mollusca, Annelida, and Arthropoda. *Zool Studies* 47: 338–351
- Dunn CW, Hejnal A, Matus DQ, Pang K, Browne WE, Smith SA, et al. (2008) Broad phylogenomic sampling improves the resolution of the animal tree of life. *Nature* 452: 745–749
- Ellington WR (2001) Evolution and physiological roles of phosphagen systems. *Ann Rev Physiol* 63: 289–325
- Ellington WR, Suzuki T (2006) Evolution and divergence of creatine kinase genes. In "Molecular Anatomy and Physiology of Proteins: Creatine Kinase" Ed by C Vial, Nova Science, New York pp 1–26
- Fujimoto N, Tanaka K, Suzuki T (2005) Amino acid residues 62 and 193 play the key role in regulating the synergism of substrate binding in oyster arginine kinase. *FEBS Lett* 579: 1688–1692
- Guindon S, Gascuel O (2003) A simple fast and accurate algorithm to estimate large phylogenies by maximum likelihood. *Syst Biol* 52: 696–704
- Horikawa DD, Sakashita T, Katagiri C, Watanabe M, Kikawada T, et al. (2006) Radiation tolerance in the tardigrade *Milnesium tardigradum*. *Int J Radiat Biol* 82: 843–848
- Iwanami K, Iseno S, Uda K, Suzuki T (2009) A novel arginine kinase from the shrimp *Neocaridina denticulata*: The fourth arginine kinase gene lineage. *Gene* 437: 80–87
- Jarilla BR, Tokuhiro S, Nagataki M, Hong SJ, Uda K, Suzuki T, Agatsuma T (2009) Molecular characterization and kinetic properties of a novel two-domain taurocyamine kinase from the lung fluke *Paragonimus westermani*. *FEBS Lett* 583: 2218–2224
- Jonsson KI, Rabbow E, Schill RO, Harms-Ringdahl M, Rettberg P (2008) Tardigrades survive exposure to space in low Earth orbit. *Curr Biol* 18: R729–R731
- Li M, Wang X, Bai J (2006) Purification and characterization of arginine kinase from locust. *Protein Pept Lett* 13: 405–410
- Mallatt J, Giribet G (2006) Further use of nearly complete 28S and 18S rRNA genes to classify Ecdysozoa: 37 more arthropods and a kinorhynch. *Mol Phylogenet Evol* 40: 772–794
- Mallatt JM, Garey JR, Shultz JW (2004) Ecdysozoan phylogeny and Bayesian inference: first use of nearly complete 28S and 18S rRNA gene sequences to classify the arthropods and their kin. *Mol Phylogenet Evol* 31: 178–191
- McLeish MJ, Kenyon GL (2005) Relating structure to mechanism in creatine kinase. *Crit Rev Biochem Mol Biol* 40: 1–20
- Morrison JF, James E (1965) The mechanism of the reaction catalysed by adenosine triphosphate-creatine phosphotransferase. *Biochem J* 97: 37–52
- Nelson RD (2002) Current status of the tardigrada: evolution and ecology. *Integr Comp Biol* 42: 652–659
- Ramlov H, Westh P (2001) Cryptobiosis in the eutardigrade *Adorybiotus (Richtersius) coronifer*: tolerance to alcohols, temperature and *de novo* protein synthesis. *Zool Anz* 240: 517–523
- Schlattner U, Tokarska-Schlattner M, Wallimann T (2006) Mitochondrial creatine kinase in human health and disease. *Biochim Biophys Acta* 1762: 164–180
- Seals DJ, Grossman SH (1988) Purification and characterization of arginine kinase from the sea cucumber *Caudina arenicola*. *Comp Biochem Physiol B* 89: 701–707
- Seki K, Toyoshima M (1998) Preserving tardigrades under pressure. *Nature* 395: 853–854
- Suzuki T, Furukohri T (1994) Evolution of phosphagen kinase Primary structure of glycoamine kinase and arginine kinase from invertebrates. *J Mol Biol* 237: 353–357



**Fig. 4.** Temperature/activity profiles of *Macrobiotus* AK and *Nautilus* AK. Profiles represent activity relative to each maximum activity. Activities at pH 8.0 were measured between 10 and 45°C under substrate concentrations of 9.52 mM arginine and 4.76 mM ATP, using His-tagged recombinant enzymes.



- Suzuki T, Kawasaki Y, Furukohri T (1997) Evolution of phosphagen kinase: Isolation characterization and cDNA-derived amino acid sequence of two-domain arginine kinase from the sea anemone *Anthopleura japonicus*. *Biochem J* 328: 301–306
- Suzuki T, Kawasaki Y, Unemi Y, Nishimura Y, Soga T, Kamidochi M, Yazawa Y, Furukohri T (1998) Gene duplication and fusion have occurred frequently in the evolution of phosphagen kinases—a two-domain arginine kinase from the clam *Pseudocardium sachalinensis*. *Biochim Biophys Acta* 1388: 253–259
- Suzuki T, Kamidochi M, Inoue N, Kawamichi H, Yazawa Y, Furukohri T, Ellington WR (1999) Arginine kinase evolved twice: evidence that echinoderm arginine kinase originated from creatine kinase. *Biochem J* 340: 671–675
- Suzuki T, Fukuta H, Nagato H, Umekawa M (2000) Arginine kinase from *Nautilus pompilius*, a living fossil: Site-directed mutagenesis studies on the role of amino acid residues in the Guanidino specificity region. *J Biol Chem* 275: 23884–23890
- Tada H, Suzuki T (2010) Cooperativity in the two-domain arginine kinase from the sea anemone *Anthopleura japonicus*. II. Evidence from site-directed mutagenesis studies. *Int J Biol Macromol* 47: 250–254
- Tada H, Nishimura Y, Suzuki T (2008) Cooperativity in the two-domain arginine kinase from the sea anemone *Anthopleura japonicas*. *Int J Biol Macromol* 42: 46–51
- Takeuchi M, Mizuta C, Uda K, Fujimoto N, Okamoto M, Suzuki T (2004) Unique evolution of Bivalvia arginine kinases. *Cell Mol Life Sci* 61: 110–117
- Tanaka K, Ichinari S, Iwanami K, Yoshimatsu S, Suzuki T (2007) Arginine kinase from the beetle *Cissites cephalotes* (Olivier). Molecular cloning, phylogenetic analysis and enzymatic properties. *Insect Biochem Mol Biol* 37: 338–345
- Uda K, Suzuki T (2004) Role of amino acid residues on the GS region of *Stichopus* arginine kinase and *Danio* creatine kinase. *Protein J* 23: 53–64
- Uda K, Suzuki T (2007) A novel arginine kinase with substrate specificity towards D-arginine. *Protein J* 26: 281–291
- Uda K, Iwai A, Suzuki T (2005) Hypotaurocyamine kinase evolved from a gene for arginine kinase. *FEBS Lett* 579: 6756–6762
- Uda K, Fujimoto N, Akiyama Y, Mizuta K, Tanaka K, Ellington WR, Suzuki T (2006) Evolution of the arginine kinase gene family. *Comp Biochem Physiol D* 1: 209–218
- Wickramasinghe S, Uda K, Nagataki M, Yatawara L, Rajapakse RPVJ, Watanabe Y, Suzuki T, Agatsuma T (2007) *Toxocara canis*: molecular cloning, characterization, expression and comparison of the kinetics of cDNA-derived arginine kinase. *Exp Parasitol* 117: 124–132
- Wickramasinghe S, Yatawara L, Nagataki M, Takamoto M, Watanabe Y, Rajapakse RPVJ, Uda K, Suzuki T, Agatsuma T (2008) Development of a highly sensitive IgG-ELISA based on recombinant arginine kinase of *Toxocara canis* for serodiagnosis of visceral larva migrans in the murine model. *Parasitol Res* 103: 853–858
- Wu QY, Li F, Zhu WJ, Wang XY (2007) Cloning, expression, purification, and characterization of arginine kinase from *Locusta migratoria manilensis*. *Comp Biochem Physiol B Biochem Mol Biol* 148: 355–362
- Wyss M, Smeitink J, Wevers R, Wallimann T (1992) Mitochondrial creatine kinase: a key enzyme of aerobic energy metabolism. *Biochim Biophys Acta* 1102: 119–166
- Zhou G, Somasundaram T, Blanc E, Parthasarathy G, Ellington WR, Chapman MS (1998) Transition state structure of arginine kinase: implications for catalysis of bimolecular reactions. *Proc Natl Acad Sci USA* 95: 8449–8454

(Received March 2, 2010 / Accepted May 4, 2010)