

Two new cytotypes and additional karyological records for blind mole rats, *Nannospalax xanthodon* and *N. ehrenbergi* (Mammalia, Rodentia) in Turkey

Authors: Sözen, Mustafa, Çolak, Faruk, Sevindik, Murat, and Matur, Ferhat

Source: *Folia Zoologica*, 64(2) : 167-172

Published By: Institute of Vertebrate Biology, Czech Academy of Sciences

URL: <https://doi.org/10.25225/fozo.v64.i2.a10.2015>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Two new cytotypes and additional karyological records for blind mole rats, *Nannospalax xanthodon* and *N. ehrenbergi* (Mammalia, Rodentia) in Turkey

Mustafa SÖZEN, Faruk ÇOLAK, Murat SEVİNDİK and Ferhat MATUR*

Bülent Ecevit University, Faculty of Arts and Science, Department of Biology, 67100 Zonguldak, Turkey;
e-mail: spalaxtr@hotmail.com, farukcolak@gmail.com, muratsevindik@hotmail.com, ferhat.matur@gmail.com

Received 1 April 2015; Accepted 16 July 2015

Abstract. Blind mole rats are model organisms for studies of chromosomal evolution, and have a variety of chromosomal forms in Turkey. This study was performed on subterranean mole rats of *Nannospalax xanthodon* (Satunin, 1898) and *N. ehrenbergi* (Nehring, 1898) in Turkey. Karyotypes of 63 specimens originating from 30 localities were analysed. Two new cytotypes with $2n = 54$ from Adana and $2n = 56$ from Karaman, two different populations of the cytotypes $2n = 54C$ and $2n = 58S$, and four different chromosomal arm numbers of the $2n = 60$ cytotype (NF = 74, 76, 78 and 80) were determined in *N. xanthodon*. The cytotypes characterized by $2n = 54$, NF = 74 from Tufanbeyli and Saimbeyli in the Adana province (54S), and by $2n = 56$, NF = 70 from Karaman (56K) are new for *N. xanthodon* in Turkey. A population of *N. ehrenbergi* from the Osmaniye province in southern Anatolia had a complement with $2n = 56$, NF = 70. Additional karyological records for other cytotypes have extended their known distribution areas, and filled most karyological gaps in Turkey.

Key words: rodents, karyotype, Anatolia

Introduction

According to recent taxonomic checklists, three mole rat species, *Nannospalax xanthodon*, *N. leucodon* and *N. ehrenbergi*, are distributed in Turkey (Wilson & Reeder 2005, Yiğit et al. 2006, Kryštufek & Vohralík 2009). Within these three recognized species, more than 30 cytotypes have currently been recognized in blind mole rats in Turkey (Sözen & Kıvanç 1998a, b, Sözen et al. 1999, 2006a, b, 2013, Coşkun 2003, Matur & Sözen 2005, Kankılıç et al. 2007a, b, 2013, 2015, Ivanitskaya et al. 2008, Arslan & Bölükbaş 2010, Arslan & Zima 2013, 2014, 2015, Matur et al. 2013, Arslan et al. 2014a). The diploid number of chromosomes varies between $2n = 36$ and 60 in Turkish blind mole rats, while the fundamental number of chromosomal arms ranges between NF = 66 and 92 (e.g. Sözen et al. 2006b, 2011, 2013, Ivanitskaya et al. 2008, Kankılıç et al. 2009, 2010, 2013, Arslan & Zima 2014, Arslan et al. 2014a). The cytotype with 62 chromosomes was eliminated from the list of Turkish *N. xanthodon* cytotypes (Ivanitskaya et al. 2008).

Nevo et al. (1994) indicated a correlation between $2n$ value and aridity stress. They showed that trends in chromosomal evolution revealed in Israeli blind mole rats by comparison of $2n$ and heterozygosity, were positively correlated with aridity stress and climatic unpredictability. Later Sözen et al. (2000a) proposed that NF values were also correlated with aridity stress and climatic unpredictability. However, Matur et al. (2011) rejected this hypothesis in a study of G- and C- banded chromosomes of four different $2n = 50$ population, and concluded that such a correlation is useless in Turkish mole rats. Geographic structures act as an isolation mechanism and they separate the cytotypes (Matur & Sözen 2005). However, most cytotypes seem to be differentiated despite the lack of a barrier. The ranges of individual cytotypes are geographically close to each other but almost no hybrids have been found in Turkey (Sözen et al. 2006b, 2013, Ivanitskaya et al. 2008, Kankılıç et al. 2009, 2013, Matur et al. 2013).

Though extensive karyological studies have been made since 1978 in Turkey and new cytotypes have

* Corresponding Author

been recently discovered (Arslan et al. 2014a), some karyological gaps still exist in some areas that should be filled. Determination of all cytotypes and their distribution areas will open new ways for general evaluation and reaching of significant results about their real taxonomic status. The purpose of this study is to describe the karyotypic characteristics of several populations in Anatolia to fill the gaps in our information.

Material and Methods

In this study, 63 specimens (25 males, 38 females) of blind mole rats belonging to *Nannospalax xanthodon* and *N. ehrenbergi* were studied from 30 localities in Anatolia. The sampled localities, the number of individuals analysed and karyological results are presented in Table 1, and the distribution of cytotypes and collection localities in Fig. 1.

Karyotypes were prepared from bone marrow according to Ford & Hamerton (1956), and about 25–30 metaphase cells were examined in each animal. The diploid number of chromosomes ($2n$), the number of autosomal arms (NFa), the total number of chromosomal arms (NF) and the sex chromosomes were determined from photos of metaphase plates. The cytotypes were named according to their chromosome diploid number, the geographic position of their occurrence, or species pertinence (e.g. S – southern, C – central, K – Karaman, *eh* – *N. ehrenbergi*).

The karyotype preparations and animals examined were deposited in the Department of Biology, the




Fig. 1. Map of study area in Turkey, and distribution of cytotypes determined. The numbers of localities are the same as in Table 1.

Faculty of Art and Sciences, Bülent Ecevit University.
Results

Two new cytotypes of *Nannospalax xanthodon* with



Fig. 2. Karyotype of a female of $2n = 54S$ cytotype from Saimbeyli.




Fig. 3. Karyotype of a male of $2n = 56K$ cytotype from Çandır Plateau.

$2n = 54S$ from Adana and $56K$ from Karaman, two different populations for *Nannospalax xanthodon* possessing $2n = 54C$ and $2n = 58S$, $NF = 72$, four different chromosomal arm numbers of $2n = 60$ cytotype ($NF = 74, 76, 78$ and 80), and one cytotype of *N. ehrenbergi* ($2n = 56eh$) were revealed.

$2n = 54S$: This cytotype was found in Tufanbeyli and Saimbeyli in the Adana province. The X chromosome is medium-sized metacentric, and the Y could not be determined since all animals of this cytotype were females. Autosomal set consists of nine pairs of bi-armed and 17 pairs of acrocentric chromosomes ($2n = 54$, $NF = 74$; Fig. 2).

$2n = 54C$: This cytotype was recorded from Sorgun (Yozgat province), Kozaklı (Nevşehir), Kırşehir centre, Kaman (Kırşehir). The X chromosome is medium-sized metacentric and the Y small acrocentric. Autosomal set contains nine pairs of bi-armed and 17 pairs of acrocentric chromosomes ($2n = 54$, $NF = 74$, $NFa = 70$).

$2n = 56K$: This karyotype was found in samples from the Barjin high-plateau (Karaman), and the Çandır high-plateau (Mersin) populations. The X chromosome is medium-sized metacentric, and the

Table 1. Synopsis of the specimens examined and basic results. See text for details.

Locality	Province	no. ♂♂	no. ♀♀	2n	NF	X	Y
1 Tufanbeyli 1 km S	Adana	0	2	54S	74	sm	-
2 Saimbeyli 10 km N	Adana	0	2	54S	74	sm	-
3 Sorgun	Yozgat	2	0	54C	74	sm	a
4 Kozaklı 10km N	Nevşehir	0	3	54C	74	sm	a
5 Kırşehir 10 km N	Kırşehir	0	2	54C	74	sm	a
6 Kaman	Kırşehir	0	1	54C	74	sm	a
7 Çukurköprü 1 km E, Kadirli	Osmaniye	1	1	56eh	70	sm	a
8 Karaman 20 km S	Karaman	1	0	56K	70	sm	a
9 Ermenek 40 km S, Çandır plateau	Mersin	1	1	56K	70	sm	a
10 Barjin plateau	Karaman	1	2	56K	70	sm	a
11 Ulukışla 30 km N, Kolsuz village	Niğde	1	2	58S	72	sm	a
12 Çamardı 15 km W	Niğde	1	1	58S	72	sm	a
13 Demirkazık village, Çamardı	Niğde	0	1	58S	72	sm	a
14 Ovacık 2 km W	Karabük	1	1	60	78	sm	a
15 Fındıkpınarı plateau	Mersin	1	2	60	78	sm	a
16 Çamiçi plateau, Niksar	Tokat	2	1	60	78	sm	a
17 Altunhisar 2 km SE	Niğde	0	1	60	80	sm	a
18 Hüyük village 2 km SW	Niğde	0	1	60	80	sm	a
19 Yeşilhisar 9 km S	Kayseri	1	1	60	80	sm	a
20 Bakırdağ 6 km W	Kayseri	1	1	60	80	sm	a
21 Bakırdağ 10 km E	Kayseri	1	0	60	80	sm	a
22 Bakırdağ 15 km E	Kayseri	1	1	60	80	sm	a
23 Mansurlu village	Adana	1	3	60	74	sm	a
24 Ayrancı town	Karaman	1	0	60	76	sm	a
25 Ladik 20 km S	Samsun	2	2	60	78	sm	a
26 Gediksaray 10 km N	Amasya	0	2	60	78	sm	a
27 Turhal 10 km S	Tokat	0	1	60	76	sm	a
28 Çekerek 5 km E	Yozgat	1	1	60	76	sm	a
29 Çayıralan	Yozgat	1	2	60	76	sm	a
30 Karataş plateau, Koru graveyard, Şebinkarahisar	Giresun	3	0	60	78	sm	a
		25	38				

Y chromosome small acrocentric. The autosomal set consists of six pairs of bi-armed and 21 pairs of acrocentric chromosomes ($2n = 56$, $NF = 70$, $NFa = 66$; Fig. 3).

$2n = 56eh$: The karyotype was found in the sample from Kadirli (Osmaniye). The X chromosome is medium-sized submetacentric and the Y chromosome small acrocentric. The autosomal set contains six pairs of bi-armed and 21 pairs of acrocentric chromosomes ($2n = 56$, $NF = 70$, $NFa = 66$).

$2n = 58S$: The karyotype was found in samples from Kolsuz and Demirkazık villages, and from vicinity of Çamardı. The X chromosome is medium-sized submetacentric; Y chromosome small acrocentric. The autosomal set contains six pairs of bi-armed

and 22 pairs of acrocentric chromosomes ($2n = 58$, $NF = 72$, $NFa = 68$).

$2n = 60$: The karyotype with 60 chromosomes was found in most localities studied. Four different NF values ($NF = 74, 76, 78$ and 80) were determined in the complement, and the differences in the NF value were due to varying number of bi-armed and acrocentric chromosomes. The X chromosome is medium-sized submetacentric, and Y chromosome small acrocentric.

Discussion

The $2n = 54$ cytotype of *N. xanthodon* was previously recorded from central, northern and eastern Anatolia (Nevo et al. 1994, Yüksel & Gülkaç 2001, Coşkun 2004, Sözen 2004, Aşan & Yağcı 2008, Arslan et al.

2011b, Kankılıç et al. 2013, 2015). The additional distribution records for the 54C cytotype from central Anatolia (Yozgat, Nevşehir and Kırşehir) fill the gaps in the distribution in the Kızılırmak basin in Central Anatolia. The distribution records of the populations with 54 and 60 chromosomes show that the 54C cytotype is surrounded by populations with $2n = 60$ in Central Anatolia (Sözen et al. 2006a, 2013, Arslan et al. 2011b, Kankılıç et al. 2013). This distribution pattern also suggests that the River Kızılırmak forms a barrier between the $2n = 60$ and $2n = 54$ cytotypes in the west, however, the river is not a barrier for the former cytotype in the southern part of the basin. The 54S cytotype from Saimbeyli and Tufanbeyli was recorded for the first time. The geographic distance between the assumed ranges of the 54C and 54S cytotypes is considerable, and the latter cytotype is accepted as a new one here. The River Kızılırmak separates the two $2n = 54$ cytotypes and the area between their ranges is occupied by populations with 60 chromosomes (Fig. 1).

The $2n = 56$ cytotype was previously recorded from Gülek, Pozantı and Aksu in southern Anatolia (Sözen & Kıvanç 1998, Sözen et al. 2006b, Kandemir et al. 2012, Arslan et al. 2014b, Kankılıç & Gürpınar 2014). Arslan et al. (2014b) compared four different cytotypes with $2n = 56$ according to the C-banding pattern and the NOR distribution. The localities studied in this paper are new records for this cytotype in southern Anatolia. Considering the geographic distance between the Pozantı and Karaman regions as well as the specific chromosome morphologies found in karyotypes of the specimens from Karaman, we propose to recognize these populations as a new cytotype designated as 56K to indicate that this cytotype is distributed around Karaman region.

Records of populations with $2n = 56$ belonging to *N. ehrenbergi* were previously reported from Tarsus (Nevo et al. 1994, 1995, Ivanitskaya et al. 1997), Kadirli (Osmaniye), Ceyhan and Kozan (Adana) (Coşkun et al. 2006), and Adana and Şeyfımurat (Sözen et al. 2006a). In all these localities, the number of chromosomal arms in the karyotype was $NF = 70$. We

recorded the same cytotype near Kadirli, and extended the distribution area of this cytotype northward.

The $2n = 58S$ cytotype was recorded in *N. xanthodon* from Ereğli, Ulukışla and Pozantı in southern Anatolia by Sözen & Kıvanç (1998), Sözen et al. (2000b, 2006b), and Arslan et al. (2011a). We found the same karyotype in additional three localities and extended the distribution of the cytotype to the north by more than 50 km.

The karyotype with $2n = 60$ was mostly recorded from central Anatolia (Yüksel 1984, Gülkaç & Yüksel 1989, Nevo et al. 1994, 1995, Sözen et al. 1999, 2000b, 2006a, b, 2011, 2013, Tez et al. 2001, Yüksel & Gülkaç 2001, Sözen 2004, Matur & Sözen 2005, Kankılıç et al. 2007a, b, 2009, 2010, Ivanitskaya et al. 2008, Arslan & Bölükbaş 2010, Arslan et al. 2011a, Kandemir et al. 2012, Aşan et al. 2013, Matur et al. 2013). In the karyotype with 60 chromosomes, several different NF values were determined in various parts of its occurrence (reviewed in Sözen et al. 2006a and Arslan & Zima 2014). This cytotype was recorded from 18 additional localities here and these new localities generally determine its northern and southern distribution borders, and fill most karyological gaps in central Anatolia.

Chromosome studies have provided rather comprehensive picture of karyotypic variation of blind mole rat populations in Anatolia. Cytogenetic investigations should be supplemented now by analyses of other phylogenetic markers. Molecular systematic studies on chromosomal forms of blind mole rats have been recently performed (e.g. Arslan et al. 2010, Kandemir et al. 2012, Kryštufek et al. 2012, Kankılıç et al. 2013, 2015, Kankılıç & Gürpınar 2014), and we can expect that new species will be described based on such data in the near future.

Acknowledgements

We would like to thank Prof. Dr. Ahmet Karataş (Niğde University) for drawing the map in Figure 1, to Burak Çoban, Meltem Gölgeci Matur and Selcen Zorlu for their carefully linguistic edit on the manuscript. This study was supported by TUBITAK (101T084 and 106T225) and by Zonguldak Karaelmas University (2000-13-02-08 and 2004-13-06-08).

Literature

- Arslan A., Akan Ş. & Zima J. 2011a: Variation in C-heterochromatin and NORs distribution among chromosomal races of mole rats (Spalacidae) from Central Anatolia, Turkey. *Mamm. Biol.* 76: 28–35.
- Arslan A., Arisoy A. & Zima J. 2014b: Comparison of the chromosome banding pattern in the $2n = 56$ cytotypes of *Nannospalax leucodon* and *N. xanthodon* from Turkey. *Sci. World J.* 2014: 121690.
- Arslan A. & Bölükbaş F. 2010: C-heterochromatin and NORs distribution of mole rat, *Nannospalax xanthodon* from Aksaray, Turkey. *Caryologia* 63: 398–404.
- Arslan A., Toyran K., Gözütok S. & Yorulmaz T. 2011b: C- and NOR stained karyotypes of mole rat, *Nannospalax xanthodon* ($2n = 54$) from Kırıkkale, Turkey. *Turk. J. Biol.* 35: 655–661.

- Arslan A. & Zima J. 2013: The banded karyotype of the $2n = 58$ chromosomal race of mole rat from Erzincan, Turkey. *Folia Zool.* 62: 19–23.
- Arslan A. & Zima J. 2014: Karyotypes of the mammals of Turkey and neighbouring regions: a review. *Folia Zool.* 63: 1–62.
- Arslan A. & Zima J. 2015: Chromosome banding pattern retrieves an independent origin of $2n = 50$ chromosome populations of *Nannospalax xanthodon* from Turkey. *Mamm. Biol.* 80: 440–445.
- Arslan A., Zima J., Yorulmaz T. & Arslan E. 2014a: A new cytotype ($2n = 46$) of *Nannospalax xanthodon* from Turkey. *Zool. Middle East* 60: 283–287.
- Arslan E., Gülbahçe E., Arıkoğlu H., Arslan A., Bužan E.V. & Kryštufek B. 2010: Mitochondrial divergence between three cytotypes of the Anatolian mole rat, *Nannospalax xanthodon* (Nordmann, 1849). *Zool. Middle East* 50: 27–34.
- Aşan N. & Yağcı T. 2008: Karyotype and hair scale structure of *Nannospalax leucodon* (Nordmann, 1840) from Central Anatolia (Rodentia: Spalacidae). *Turk. J. Zool.* 32: 125–130.
- Aşan Baydemir N., Yağcı T. & Çakır Ş. 2013: Karyological studies of the $2n = 60$ cytotype of *Nannospalax nehringi* from Central Anatolia. *Caryologia* 66: 49–53.
- Coşkun Y. 2003: A study on the morphology and karyology of *Nannospalax nehringi* (Satunin, 1898) (Rodentia: Spalacidae) from North-eastern Anatolia, Turkey. *Turk. J. Zool.* 27: 171–176.
- Coşkun Y. 2004: A new species of mole rat, *Nannospalax munzuri* sp. n., and karyotype of *Nannospalax tuncelicus* (Coşkun, 1996) (Rodentia: Spalacidae) in eastern Anatolia. *Zool. Middle East* 33: 153–162.
- Coşkun Y., Ulutürk S. & Yürümez G. 2006: Chromosomal diversity in mole-rats of the species *Nannospalax ehrenbergi* (Mammalia: Rodentia) from South Anatolia, Turkey. *Mamm. Biol.* 71: 244–250.
- Ford C.E. & Hamerton J.L. 1956: A colchicine hypotonic citrate, squash for mammalian chromosomes. *Stain Technol.* 31: 247–251.
- Gülkaç M.D. & Yüksel E. 1989: A cytogenetical study on blind mole rats around Malatya province. *Doğa Tü Biyol Dergisi (Ankara)* 13: 63–71. (in Turkish with English summary)
- Ivanitskaya E., Coşkun Y. & Nevo E. 1997: Banded karyotypes of mole rats (*Spalax*, Spalacidae, Rodentia) from Turkey: a comparative analysis. *J. Zool. Syst. Evol. Res.* 35: 171–177.
- Ivanitskaya E., Sözen M., Rashkovetsky L., Matur F. & Nevo E. 2008: Discrimination of $2n = 60$ *Spalax leucodon* cytotypes (Spalacidae, Rodentia) in Turkey by means of classical and molecular cytogenetic techniques. *Cytogenet. Genome Res.* 122: 139–149.
- Kandemir I., Sözen M., Matur F., Kankılıç Te., Çolak F., Özkurt S. & Çolak E. 2012: Phylogeny of species and cytotypes of mole rats (Spalacidae) in Turkey inferred from mitochondrial cytochrome *b* gene sequences. *Folia Zool.* 61: 25–33.
- Kankılıç Te., Çolak R., Kankılıç To. & Çolak E. 2007a: On the morphology and karyology of *Spalax leucodon armeniacus* Mehely, 1909, and *Spalax leucodon cilicicus* Mehely, 1909 (Mammalia: Rodentia) in Turkey. *Acta Zool. Bulg.* 59: 41–46.
- Kankılıç Te., Çolak E. & Kankılıç To. 2009: Macro-anatomical and karyological features of two blind mole rat subspecies (Rodentia: Spalacidae) from Turkey. *Anat. Histol. Embryol.* 38: 145–153.
- Kankılıç Te. & Gürpınar C. 2014: Revised classification design of the Anatolian species of *Nannospalax* (Rodentia: Spalacidae) using RFLP analysis. *Turk. J. Zool.* 38: 68–78.
- Kankılıç Te., Kankılıç To., Çolak R., Çolak E. & Karataş A. 2007b: Karyological comparison of populations of the *Spalax leucodon* Nordmann, 1840 superspecies (Rodentia: Spalacidae) in Turkey. *Zool. Middle East* 42: 15–24.
- Kankılıç Te., Kankılıç To., Seker P.S., Çolak R., Selvi E. & Çolak E. 2010: Contributions to the karyology and distribution areas of cytotypes of *Nannospalax leucodon* (Rodentia: Spalacidae) in Western Anatolia. *Acta Zool. Bulg.* 62: 161–167.
- Kankılıç Te., Kankılıç To., Sözen M. & Çolak E. 2013: Genetic diversities and geographic variations of chromosomal forms of *Nannospalax nehringi* (Satunin, 1898) and *Nannospalax ehrenbergi* (Nehring, 1898) from Turkey, revealed by RAPD Analysis. *Acta Zool. Bulg.* 65: 45–58.
- Kankılıç Te., Kankılıç To., Sözen M. & Çolak E. 2015: Allozyme variations in Anatolian populations and cytotypes of the blind mole rats (*Nannospalax*). *Biochem. Syst. Ecol.* 59: 126–134.
- Kryštufek B., Ivanitskaya E., Arslan A., Arslan E. & Bužan E. 2012: Evolutionary history of mole rats (genus: *Nannospalax*) inferred from mitochondrial cytochrome *b* sequences. *Biol. J. Linn. Soc.* 105: 446–455.
- Kryštufek B. & Vohralík V. 2009: Mammals of Turkey and Cyprus. Rodentia II. Cricetinae, Muridae, Spalacidae, Calomyscidae, Hystricidae, Castoridae. *Univerzita za Primorskem, Koper*.
- Matur F., Çolak F., Sevindik M. & Sözen M. 2011: Chromosome differentiation of four $2n = 50$ chromosomal forms of Turkish mole rat, *Nannospalax nehringi*. *Zool. Sci.* 28: 61–67.
- Matur F., Çolak F., Ceylan T., Sevindik M. & Sözen M. 2013: Chromosomal evolution of the genus *Nannospalax* (Palmer, 1903) (Rodentia: Muridae) from western Turkey. *Turk. J. Zool.* 37: 470–487.
- Matur F. & Sözen M. 2005: A karyological study on subterranean mole rats of the *Spalax leucodon* Nordmann, 1840 (Mammalia: Rodentia) superspecies around Bilecik province in Turkey. *Zool. Middle East* 36: 5–10.
- Nevo E., Filippucci M.G., Redi C., Korol A. & Beiles A. 1994: Chromosomal speciation and adaptive radiation of mole rats in Asia Minor correlated with increased ecological stress. *Proc. Natl. Acad. Sci. U. S. A.* 91: 8160–8164.
- Nevo E., Filippucci M.G., Redi C., Simson S., Heth G. & Beiles A. 1995: Karyotype and genetic evolution in speciation of subterranean mole rats of the genus *Spalax* in Turkey. *Biol. J. Linn. Soc.* 54: 203–229.
- Sözen M. 2004: A karyological study on subterranean mole rats of the *Spalax leucodon* Nordmann, 1840 superspecies in Turkey. *Mamm. Biol.* 64: 420–429.
- Sözen M., Çataklı K., Eroğlu F., Matur F. & Sevindik M. 2011: Distribution of chromosomal forms of *Nannospalax nehringi* (Satunin, 1898) (Rodentia: Spalacidae) in Çankırı and Çorum provinces, Turkey. *Turk. J. Zool.* 35: 367–374.
- Sözen M., Çolak E., Yiğit N., Özkurt Ş. & Verimli R. 1999: Contributions to the karyology and taxonomy of the genus *Spalax* Guldensstaedt, 1770 (Mammalia: Rodentia) in Turkey. *Z. Säugetierkd.* 64: 210–219.

- Sözen M., Çolak E. & Yiğit N. 2000a: Contributions to the karyology and taxonomy of *Spalax leucodon nehringi* Satunin, 1898 and *Spalax leucodon armeniacus* Mehely, 1909 (Mammalia: Rodentia) in Turkey. *Z. Säugetierkd.* 65: 309–312.
- Sözen M., Çolak E. & Yiğit N. 2000b: A study on karyotypic evolution of the genus *Spalax* Gldenstaedt, 1770 (Mammalia: Rodentia) in Turkey. *Isr. J. Zool.* 46: 239–242.
- Sözen M., Çolak F., Sevindik M. & Matur F. 2013: Cytotypes of *Nannospalax xanthodon* (Satunin, 1898) (Rodentia: Spalacidae) from Western Anatolia. *Turk. J. Zool.* 37: 462–469.
- Sözen M. & Kıvanç E. 1998a: Two new karyotypic forms of *Spalax leucodon* (Nordmann, 1840) (Mammalia: Rodentia) from Turkey. *Z. Säugetierkd.* 63: 307–310.
- Sözen M. & Kıvanç E. 1998b: A new karyotype of *Spalax leucodon cilicicus* Mehely, 1909 (Mammalia: Rodentia) from type locality in Turkey. *Isr. J. Zool.* 44: 53–56.
- Sözen M., Matur F., Çolak E., Özkurt Ş. & Karataş A. 2006a: Some karyological records and a new chromosomal form for *Spalax* (Mammalia: Rodentia) in Turkey. *Folia Zool.* 55: 247–256.
- Sözen M., Sevindik M. & Matur F. 2006b: Karyological and some morphological characteristics of *Spalax leucodon* Nordmann, 1840 (Mammalia: Rodentia) superspecies around Kastamonu Province, Turkey. *Turk. J. Zool.* 30: 205–219.
- Tez C., Gndz İ. & Kefelioğlu H. 2001: Karyological study of *Spalax leucodon* Nordmann, 1840 in central Anatolia. *Pak. J. Biol. Sci.* 4: 869–871.
- Wilson E.O. & Reeder M. 2005: Mammal species of the world: a taxonomic and geographic reference, 3rd ed. *The Johns Hopkins University Press, Baltimore.*
- Yiğit N., Çolak E., Sözen M. & Karataş A. 2006: Rodents of Trkiye. *Meteksan, Ankara.*
- Yksel E. 1984: Cytogenetic study in *Spalax* (Rodentia: Spalacidae) from Turkey. *Communications, Serie C: Biologie* 2: 1–12.
- Yksel E. & Glkaç M.D. 2001: The cytogenetical comparison of *Spalax* (Rodentia: Spalacidae) populations from middle Kızılırmak basin, Turkey. *Turk. J. Biol.* 25: 17–24.