

## **First record of Crested (or Crested-type) Honey Buzzard *Pernis ptilorhynchus* for Greece**

Authors: Zannetos, Stylianos P., Zevgolis, Yiannis, and Akriotis, Triantafyllos

Source: Bulletin of the British Ornithologists' Club, 138(4) : 386-388

Published By: British Ornithologists' Club

URL: <https://doi.org/10.25226/bboc.v138i4.2018.a10>

---

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# First record of Crested (or Crested-type) Honey Buzzard *Pernis ptilorhynchus* for Greece

by Stylianios P. Zannetos, Yiannis Zevgolis & Triantafyllos Akriotis

Received 11 September 2018, revised 12 October 2018, published 14 December 2018

<http://zoobank.org/urn:lsid:zoobank.org:pub:BEE11831-B795-4FFB-9D66-8D75ACFA1FB2>

Crested Honey Buzzard *Pernis ptilorhynchus orientalis* is a long-distance migrant that breeds across southern Siberia to Sakhalin and Japan, and winters mainly in South-East Asia, Indonesia and the Philippines (Higuchi *et al.* 2005, Wells 2010, Orta *et al.* 2018). In southern Siberia, at its western limit, the breeding range partially overlaps with that of European Honey Buzzard *P. apivorus* (Stepanyan 1983, Ferguson-Lees & Christie 2001). Six subspecies of Crested Honey Buzzard are recognised, but only *orientalis* is a migrant (Orta *et al.* 2018). The species was recorded for the first time in the Western Palearctic at Borçka, north-east Turkey, in September 1979 (Laine 1996) and then at Eilat, Israel, in May 1994 (Shirihai 1994). *P. ptilorhynchus* is now considered regular in small numbers on passage through Israel, mainly at Eilat, which is a major passage bottleneck for European Honey Buzzards (Shirihai 1994). Fifteen to 20 individuals are recorded every spring, mainly in May, with 5–12 in autumn, mainly in mid September (Babbington & Campbell 2016). There have also been many recent records of *P. ptilorhynchus* at Batumi (Georgia) where the first to be officially accepted was in autumn 2007 (Abuladze 2013). Since then, the species has been identified annually at Batumi, with a total of 163 records until 2018 and a max. 51 birds in 2013 (<https://www.batumiraptorcount.org/migration-count-data#annual-totals>).

It is presumed that those Crested Honey Buzzards recorded in Israel, and elsewhere in the Middle East in spring, joined flocks of *P. apivorus* wintering in Africa (Ferguson-Lees & Christie 2001). Many records have been suspected to be potential hybrids with *P. apivorus* (Babbington & Campbell 2016).

In Europe, Crested Honey Buzzard has been fully documented just twice: on Cyprus in October 2012 (Harrison 2014) and in Italy on 18 May 2011, at the Strait of Messina between Sicily and the mainland (Scuderi & Corso 2011).

On 2 May 2018, near the village of Alyfanta (39°06'04"N, 26°31'45"E) on Lesbos, 4 km from the largest urban centre on the island (Mytilini), we observed an adult male Crested (or Crested-type) Honey Buzzard. It was watched as it soared, gradually gaining height, for c.3 minutes at a distance of c.150 m from the observers (SPZ, YZ). SPZ managed to take 12 photographs of the bird (Figs. 1–3). Subsequently, it headed south-west and was not seen again. Identification was made by the authors, following the observation, based on the photographs. This is the first documented record for Greece and the third to be accepted for Europe.

*P. ptilorhynchus* can be easily confused with *P. apivorus*. In this case, the bird's structure was obviously different: heavier bodied, slightly larger and bulkier (more eagle-like) compared with *P. apivorus*. Furthermore, its wings appeared broader and the tail shorter than that of European Honey Buzzard. The absence of the diagnostic carpal patch of *P. apivorus*, the six clearly fingered primaries protruding from the trailing edge of the wing, and the dark tail with a broad white bar in the centre of the undertail, are diagnostic features of *P. ptilorhynchus* (Ferguson-Lees & Christie 2001, Svensson *et al.* 2009, Forsman 2016). The inner secondaries show two well-defined bars while a third bar is visible on the inner primaries and outer secondaries. The head is grey with a pale throat, bordered by a



Figures 1–3. Crested (or Crested-type) Honey Buzzard *Pernis ptilorhynchus*, Alyfanta, Lesvos, Greece, 2 May 2018 (S. P. Zannetos)

dark ‘gorget’ that contrasts with the pale sandy-ochre underparts and underwings. This plumage is commonest in adult males according to Forsman (2016).

Alternatively, the possibility of hybridisation between *P. apivorus* and *P. ptilorhynchus* (Faveyts 2011, Forsman 2016) and some structural and plumage features that do not match perfectly with *P. ptilorhynchus* made us consider the possibility that the bird was a potential hybrid. Specifically, the bird shows a quite rounded wingtip, rather than the blunt tip of Crested Honey Buzzard (p5 is not clearly longer). Furthermore, the wing is typically more rectangular in Crested Honey Buzzard, but in the Greek bird appears broadest at the carpal joint, tapering towards the body and tip. The intermediate underwing and tail barring, and possible hint of a darker carpal area, typical of supposed hybrids, reinforce this hypothesis (D. Forsman *in litt.* 2018).

However, the lack of genetic research into the hybridisation question, in parallel with the fact that *P. apivorus* and *P. ptilorhynchus* are (a) not known to form mixed pairs in the region of overlap (Mosquitin 1973, Kislenko 1974, Stepanyan 1983), and (b) are not even each other’s closest relatives (Gamauf & Haring 2004), raises doubts as to whether it is justifiable to discuss hybrids between the two species. The unquestionable similarity of *P. ptilorhynchus* to *P. apivorus* and, for most European observers, the lack of understanding of their distinguishing features and especially their morphological variability, lead us to suspect that *P. ptilorhynchus* may be a more frequent vagrant to parts of south-east Europe than is currently perceived. More attention should be paid along the major raptor passage flyways in the Western Palearctic to better understand the western limit of Crested Honey Buzzard’s migration route. Furthermore, genetic analysis is critical to provide a more solid basis for discussing hybridisation between these two *Pernis* species.

#### Acknowledgements

We thank Killian Mullarney and Dick Forsman for their help in identifying the bird, Rob Bijlsma and Andrea Corso for their very useful suggestions and comments on the submitted draft, and Apostolis Christopoulos, Eleni Galinou, Fanis Theophanopoulos and the members of the Hellenic Rarities Committee, headed by George Handrinos, for accepting the record.

#### References:

- Abuladze A. 2013. *Birds of prey of Georgia*. Institute of Zoology, Ilia State Univ., Tbilisi.
- Babbington, J. & Campbell, O. 2016. Recent status and occurrence of Crested Honey Buzzards *Pernis ptilorhynchus* in the Arabian Peninsula, with emphasis on Saudi Arabia and the United Arab Emirates. *Sandgrouse* 38: 12–22.

- Faveyts, W., Valkenburg, M. & Granit, B. 2011. Crested Honey Buzzard: identification, western occurrence and hybridisation with European Honey Buzzard. *Dutch Birding* 33: 149–162.
- Ferguson-Lees, J. & Christie, D. A. 2001. *Raptors of the world*. Christopher Helm, London.
- Forsman, D. 2016. *Flight identification of raptors of Europe, North Africa and the Middle East*. Bloomsbury, London.
- Gamauf, A. & Haring, E. 2004. Molecular phylogeny and biogeography of honey-buzzards (genera *Pernis* and *Henicopernis*). *J. Zool. Syst. & Evol. Res.* 42: 145–153.
- Harrison, I. 2014. From the rarities committees. *Sandgrouse* 36: 110–116.
- Higuchi, H., Shiu, H. J., Nakamura, H., Uematsu, A., Kuno, K., Saeki, M., Hotta, M., Tokita, K., Moriya, E., Morishita, E. & Tamura, M. 2005. Migration of Honey-buzzards *Pernis ptilorhynchus* based on satellite tracking. *Orn. Sci.* 4: 109–115.
- Kislenko, G. S. 1974. [Comparative ecology of the Oriental Honey buzzard and European Honey buzzard]. Pp. 65–66 in *Sixth All-Union Orn. Conf., Moscow*, pt. 2. Moscow Univ. Press. (In Russian.)
- Laine, L. J. 1996. The 'Borçka puzzle' – the first Western Palearctic Crested Honey Buzzard. *Birding World* 9: 324–325.
- Mosquitin, S. S. 1973. Materials on distribution and mode of life of some Siberian birds. *Proc. Biol. Inst. Siberian Acad. Sci.* 16: 263–268.
- Orta, J., Marks, J. S. & Kirwan, G. M. 2018. Oriental Honey-buzzard (*Pernis ptilorhynchus*). In del Hoyo, J., Elliott, A., Sargatal, J., Christie, D. A. & de Juana, E. (eds.) *Handbook of the birds of the world Alive*. Lynx Edicions, Barcelona (retrieved from <https://www.hbw.com/node/52959> on 16 May 2018).
- Scuderi, A. & Corso, A. 2011. Crested Honey Buzzard in Europe. *Birding World* 24: 252–256.
- Shirihai, H. 1994. The Crested Honey Buzzard in Israel – a new Western Palearctic bird. *Birding World* 7: 404–406.
- Stepanyan, L. S. 1983. *Superspecies and sibling species in avifauna of the USSR*. Nauka, Moscow.
- Svensson, L., Mullarney, K. & Zetterström, D. 2009. *Collins bird guide*. Second edn. HarperCollins, London.
- Wells, D. R. 2010. *The birds of the Thai-Malay Peninsula*, vol. 2. Bloomsbury, London.

*Addresses:* Stylianios P. Zannetos, Biodiversity Conservation Laboratory, Dept. of Environment, University Hill, Mytilene, Lesvos, 81100, Greece, e-mail: [stylianoszannetos@gmail.com](mailto:stylianoszannetos@gmail.com). Yiannis Zevgolis, Biodiversity Conservation Laboratory, Dept. of Environment, University Hill, Mytilene, Lesvos, 81100, Greece, e-mail: [zevgolis@env.aegean.gr](mailto:zevgolis@env.aegean.gr). Triantafyllos Akriotis, Biodiversity Conservation Laboratory, Dept. of Environment, University Hill, Mytilene, Lesvos, 81100, Greece, e-mail: [takr@aegean.gr](mailto:takr@aegean.gr)