

## **A new subfamily taxon for *Sasia* and *Verreauxia* (Picidae)**

Authors: Sangster, George, Gaudin, Jimmy, and Fuchs, Jérôme

Source: Bulletin of the British Ornithologists' Club, 142(4) : 478-479

Published By: British Ornithologists' Club

URL: <https://doi.org/10.25226/bboc.v142i4.2022.a6>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# A new subfamily taxon for *Sasia* and *Verreauxia* (Picidae)

by George Sangster, Jimmy Gaudin & Jérôme Fuchs

Received 16 June 2022; revised 28 September 2022; published 7 December 2022

<http://zoobank.org/urn:lsid:zoobank.org:pub:90B82FEC-8BA0-4C7C-A12F-917E999E8ED8>

**SUMMARY.**—A review of eight molecular phylogenetic studies supports the distinctiveness of the genera *Sasia*/*Verreauxia* from *Picumnus* and casts doubt on the monophyly of Picumninae. We propose to restrict Picumninae to *Picumnus* and to place *Sasia* and *Verreauxia* in a new subfamily, Sasiinae.

The piculets have long been classified as four genera: *Sasia* Hodgson, 1837; *Verreauxia* Hartlaub, 1857; *Picumnus* Temminck, 1825 (including *Vivia* Hodgson, 1837); and *Nesocittes* Hargitt, 1890. They have typically been grouped in the subfamily Picumninae (e.g. Peters 1948, Winkler *et al.* 1995, Dickinson 2003), although most authors have removed *Nesocittes* from this subfamily and placed it either in Picinae (Dickinson & Remsen 2013) or in its own subfamily, Nesocitinae (Wolters 1976, Benz *et al.* 2006, Gaudin 2022). In contrast, *Sasia*, *Verreauxia* and *Picumnus* have always been placed in Picumninae (e.g. Peters 1948, Wolters 1976, Winkler *et al.* 1995, Dickinson 2003, Dickinson & Remsen 2013).

In the first multilocus molecular phylogenetic study by Fuchs *et al.* (2006) *Sasia* and *Verreauxia* formed a separate clade from *Picumnus* and these two groups represented an unresolved trichotomy with Picinae. *Nesocittes* was not included in this analysis. Benz *et al.* (2006) found *Sasia* and *Verreauxia* sister to *Picumnus*, albeit with no support, but the two groups were separated by a deep divergence exceeding that found between Nesocitinae and Picinae. In another multilocus phylogeny (Fuchs *et al.* 2007), *Sasia* and *Verreauxia* were not sister to *Picumnus*; instead, *Picumnus* was more closely related to Picinae than to *Sasia* and *Verreauxia*. In another study (Fuchs *et al.* 2013), phylogenies based on mitochondrial DNA, nine autosomal and three Z-linked loci did not place *Sasia* and *Picumnus* as sister taxa; rather, analyses of mitochondrial DNA and the Z-linked loci placed *Sasia* closer to Picinae than to *Picumnus*, whereas the autosomal loci placed *Picumnus* closer to Picinae than to *Sasia*. A further study based on mitochondrial DNA sequences placed *Sasia* closer to Picinae than to *Picumnus* (Winkler *et al.* 2014). A sixth multilocus study placed *Sasia* and *Verreauxia* sister to *Picumnus*, but again separated by a deep divergence (Dufort 2016). More recently, a multilocus study by Shakya *et al.* (2017) placed *Sasia* and *Verreauxia* closer to Picinae than to *Picumnus*. Analyses of the two mitochondrial rRNA and 13 protein-coding genes again recovered a deep split between *Sasia*/*Verreauxia* and *Picumnus* but failed to unambiguously resolve the relationships between these taxa and *Nesocittes*/Picinae (JF *et al.* unpubl. data). All phylogenetic studies that included *Nesocittes* placed it as the sister of (other) Picinae, consistent with its removal from Picumninae.

The phylogenetic evidence clearly supports the distinctiveness of *Sasia*/*Verreauxia* from *Picumnus*. Importantly, it is unclear if Picumninae (excluding *Nesocittes*) is a monophyletic group. This suggests that the piculets (excluding *Nesocittes*) may be best treated as two separate taxonomic groups: Picumninae is restricted to *Picumnus* (and *Vivia*, if that genus is recognised for *P. innominatus*, e.g. Gaudin 2022); and another taxon is recognised for *Sasia* and *Verreauxia*. 'Sasiinae' was recently used as a group name for the latter two taxa but this represents a *nomen nudum*, as was indicated by the author by his use of square brackets (Gaudin 2022). A Google search (24 September 2022) showed that the name 'Sasiidae' is

sometimes used in online publications for a group of moths but this is clearly a misspelling of the name Sesiidae Boisduval, 1828 (Lepidoptera). As it appears that no family-group name is available for *Sasia* and *Verreauxia* (Bock 1994), and that the names 'Sasiidae' and 'Sasiinae' have not been made available for any other taxonomic group in zoological nomenclature, we propose:

### Sasiinae, new subfamily

**Type genus.**—*Sasia* Hodgson, 1837

**Diagnosis.**—Very small woodpeckers (body size 8–10 cm). All three species differ from *Picumnus* and *Nesocittes* by the distinctly shorter tail, red legs, presence of a bare area around the eye, and lack of stripes, bars or spots on the crown and underparts. In addition, the three species differ from *Picumnus* by the absence of white tail stripes.

**Included taxa.**—*Sasia abnormis* (Temminck, 1825), *Sasia ochracea* Hodgson, 1837, and *Verreauxia africana* (Verreaux & Verreaux, 1855).

#### Acknowledgements

We are grateful for the constructive comments of the referee.

#### References:

- Benz, B. W., Robbins, M. B. & Peterson, A. T. 2006. Evolutionary history of woodpeckers and allies (Aves: Picidae): placing key taxa on the phylogenetic tree. *Mol. Phylo. & Evol.* 40: 389–399.
- Bock, W. J. 1994. History and nomenclature of avian family-group names. *Bull. Amer. Mus. Nat. Hist.* 222: 1–281.
- Dickinson, E. C. (ed.) 2003. *The Howard and Moore complete checklist of the birds of the world*. Third edn. Christopher Helm, London.
- Dickinson, E. C. & Remsen, J. V. (eds.) 2013. *The Howard and Moore complete checklist of the birds of the world*, vol. 1. Fourth edn. Aves Press, Eastbourne.
- Dufort, M. J. 2016. An augmented supermatrix phylogeny of the avian family Picidae reveals uncertainty deep in the family tree. *Mol. Phylo. & Evol.* 94A: 313–326.
- Fuchs, J., Ohlson, J. I., Ericson, P. G. P. & Pasquet, E. 2006. Molecular phylogeny and biogeographic history of the piculets (Piciformes: Picumninae). *J. Avian Biol.* 37: 487–496.
- Fuchs, J., Ohlson, J. I., Ericson, P. G. P. & Pasquet, E. 2007. Synchronous intercontinental splits between assemblages of woodpeckers suggested by molecular data. *Zool. Scripta* 36: 11–25.
- Fuchs, J., Pons, J.-M., Liu, L., Ericson, P. G. P., Couloux, A. & Pasquet, E. 2013. A multi-locus phylogeny suggests an ancient hybridization event between *Campophilus* and melanerpine woodpeckers (Aves: Picidae). *Mol. Phylo. & Evol.* 67: 578–588.
- Gaudin, J. 2022. *Noms français normalisés des oiseaux du monde*. Privately published, La Rochelle.
- Peters, J. L. 1948. *Check-list of birds of the world*, vol. 6. Mus. Comp. Zool., Cambridge, MA.
- Shakya, S. B., Fuchs J., Pons J.-M. & Sheldon F. H. 2017. Tapping the woodpecker tree for evolutionary insight. *Mol. Phylo. & Evol.* 116: 182–191.
- Winkler, H., Christie, D. A. & Nurney, D. 1995. *Woodpeckers: a guide to the woodpeckers of the world*. Pica Press, Robertsbridge.
- Winkler, H., Gamauf, A., Nittinger, F. & Haring, E. 2014. Relationships of Old World woodpeckers (Aves: Picidae) – new insights and taxonomic implications. *Ann. Naturhist. Mus. Wien, B* 116: 69–86.
- Wolters, H. E. 1976. *Die Vogelarten der Erde*. Lief. II. Paul Parey, Hamburg.
- Addresses:** George Sangster, Naturalis Biodiversity Center, Darwinweg 2, PO Box 9517, 2300 RA Leiden, the Netherlands, e-mail: g.sangster@planet.nl. Jimmy Gaudin, 34, avenue Antoine de Saint-Exupéry, 17 000 La Rochelle, France, e-mail: jimmy.gdn.lr.taxo@gmail.com. Jérôme Fuchs, Institut de Systématique, Evolution, Biodiversité (ISYEB), Muséum National d'Histoire Naturelle, CNRS, SU, EPHE, UA, CP 51, 57 rue Cuvier, Paris Cedex 05, F-75231, France, e-mail: jerome.fuchs@mnhn.fr