



## **Arctic fox *Vulpes lagopus* den use in relation to altitude and human infrastructure**

Authors: Selås, Vidar, Johnsen, Børge Steinmo, and Eide, Nina E.

Source: *Wildlife Biology*, 16(1) : 107-112

Published By: Nordic Board for Wildlife Research

URL: <https://doi.org/10.2981/09-023>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## Arctic fox *Vulpes lagopus* den use in relation to altitude and human infrastructure

Vidar Selås, Børge Steinmo Johnsen & Nina E. Eide

One obvious threat to the endangered arctic fox *Vulpes lagopus* population in Fennoscandia is competition with the larger red fox *Vulpes vulpes*, which may have expanded its range towards the alpine tundra because of increased food availability in the low-alpine and subalpine region. The steady increase in the number of vacation cabins and roads, and thus also in human garbage and road-killed animals, may subsidise easily available food resources and improve red fox survival in these otherwise marginal areas. In Børgefjell National Park, Norway, 14 of 27 known arctic fox dens were used by arctic fox during 2001-2005. The dens that were used were situated at higher altitudes, farther from natural red fox habitats, than unused dens. In the best of our logistic regression models, there was also a statistical negative effect of the number of cabins within  $7 \times 7$  km squares around the den sites. Hence, our results support the prediction that the arctic fox is less likely to use areas where human activity might be benefiting red foxes. A successful conservation strategy for the arctic fox will probably require a reduction of the driving forces behind the red fox expansion in the alpine areas.

*Key words:* Arctic fox, competition, disturbance, ecosystem change, infrastructure, red fox, *Vulpes lagopus*, *Vulpes vulpes*

Vidar Selås & Børge Steinmo Johnsen, Department of Ecology and Natural Resource Management, Norwegian University of Life Sciences, PO Box 5003, NO-1432 Ås, Norway - e-mail addresses: vidar.selas@umb.no (Vidar Selås), borgesjohnsen@hotmail.com (Børge Steinmo Johnsen).

Nina E. Eide, Norwegian Institute for Nature Research, Tungasletta 2, NO-7485 Trondheim, Norway - e-mail: nina.eide@nina.no.

Corresponding author: Vidar Selås

Received 19 March 2009, accepted 11 September 2009

Associate Editor: Shyamala Ratnayake

The arctic fox *Vulpes lagopus* population in Fennoscandia (Norway, Sweden and Finland) decreased dramatically from 1850 to the 1930s, has since remained small (Lönnerberg 1927, Linnell et al. 1999), and was recently estimated to  $< 120$  adult individuals (Elmhagen et al. 2004, Linnell et al. 2004). Competition and predation from the larger red fox *Vulpes vulpes* is probably the major threat to the endangered arctic fox (Østbye et al. 1978, Frafjord et al. 1989, Angerbjörn et al. 2002, Tannerfeldt et al. 2002a, Elmhagen 2003, Selås & Vik 2007). In fact, sterilised red foxes have successfully been used to eradicate arctic foxes on islands in Alaska (Bailey 1992). The altitudinal segregation between the two

fox species is probably caused by the higher nutritional need of the red fox (Hersteinsson & Macdonald 1992, Tannerfeldt et al. 2002b), preventing it from occupying the high-alpine areas, where the amount of prey is generally low and more unstable in availability (Oksanen et al. 1999).

Improved availability of winter food, i.e. ungulate carcasses and human garbage, has been suggested as the main cause for the rapid expansion of red fox populations during the 20th century (Hjeljord 1980, Selås & Vik 2006). Human infrastructure such as roads and power lines has also increased the availability of accidentally killed animals (Forman & Alexander 1998, Bevanger & Brøseth 2004). In

both Norway and Sweden, garbage, domestic animals and carcasses can constitute a significant part of the red fox winter diet (Lund 1962, Lindström 1982). Back-tracking of red foxes in low-alpine habitats has revealed that red foxes use areas close to cabins more than other areas in search of food (Røhnebæk 2004).

In Norway and Sweden, arctic fox dens with recorded breeding after 1980 were situated at higher altitudes and farther from red fox dens located above the tree line than those without breeding (Linnell et al. 1999, Dalerum et al. 2002, Frafjord 2003b, Tannerfeldt et al. 2002a). Based on the hypothesis that human-induced increase in food availability favours red fox populations, we predict that arctic fox den use depends on altitude and on the degree of human activities in the surrounding landscape. We tested this prediction by analysing arctic fox den use in Børgfjell National Park, Norway, during 2001-2005, in relation to both altitude and human infrastructure.

## Material and methods

### Study area

Our study area is Børgfjell/Byrkije National Park in Norway (1,447 km<sup>2</sup>, 65°4'N, 13°49'E), which together with the adjacent Swedish Borgafjäll has the most vital arctic fox population in Fennoscandia (Eide et al. 2009). Most of the area is situated above the tree line (500-600 m a.s.l.). The landscape is characterised by deep valleys and high mountains (up to 1,699 m a.s.l.), with many rivers and lakes. The summer is short with snowmelt around June and first snowfall in September-October. The national park has a low degree of development for tourism, with few trails and only 10 rentable/open cabins. However, there are several vacation cabins, frequently used both in summer and winter, relatively close to the park. The mean number of cabins counted within 7 × 7 and 15 × 15 km squares around each arctic fox den (N = 27, see below) was 0-12 (mean: 1.4) and 1-63 (mean: 11.0), respectively. During 1998-2007, the number of vacation cabins within the four municipalities in which the national park is situated increased by 26%, and during our study period (2001-2005) by 10% (Central Bureau of Statistics of Norway; available at: <http://www.ssb.no>).

### Den use (response variable)

We used national monitoring data from 27 known arctic fox dens in Børgfjell (see Andersen et al. 2005). All known arctic fox dens in Børgfjell were monitored for breeding activity during 2001-2005. None of the dens were used in 2003, when there was a decline in the population of small rodents, a food source that is important for arctic fox reproduction in this area (Frafjord et al. 1998).

During our study period litters were recorded in nine dens. In addition, there were five instances where adult(s) were observed at a den site in the breeding season, but we could not confirm litters. We assumed that these were dens with unsuccessful breeding attempts. The remaining dens showed no signs of use. The first two groups were combined into one group of dens in use (N = 14), and tested against dens not in use (N = 13). Because interference with red fox is a possible cause of unsuccessful breeding, we also tested dens used with litters (N = 9) against the combined latter two groups, i.e. dens with no recorded litters (N = 18).

### Predictors

Frafjord (2003b) found that den altitude was a good predictor when testing for differences between arctic fox dens with and without litters in Børgfjell. We therefore used den altitude (m a.s.l.) as the main explanatory variable in our study. Because of the small sample size, we did not use additional predictors that were significantly correlated with altitude.

Frafjord (2003b) did not find any significant relationships between arctic fox den use and distances to different types of infrastructure, assumed to reflect human activities. However, in Børgfjell, with a few cabins situated within the National Park and several aggregated outside, we regarded the total load of infrastructure within a certain distance from the den to be a much better predictor for the presumed human influence on fox interactions than the distance to infrastructure. The predictors used in our analyses were 1) kilometres of trails, 2) kilometres of roads (public and private) and 3) number of cabins. We expected all three variables to be correlated with the availability of garbage. Furthermore, trails often contain remains from freshwater fishing, and roads the carcasses of road-killed animals.

### Scales of analysis

We examined predictor variables at two different scales, 7 × 7 km and 15 × 15 km quadrates with dens

at the centre. Arctic foxes are territorial and the fairly uniform distribution of dens suggested that most were located near the centre of arctic fox home ranges. Therefore, we expected that the scale  $7 \times 7$  km ( $49 \text{ km}^2$ ) quantified most infrastructure within an arctic fox's home range, which in Fennoscandia varies from 17 to  $62 \text{ km}^2$  (Angerbjörn et al. 1997, Landa et al. 1998). The larger  $15 \times 15$  km scale was included because arctic foxes may not establish or use dens within 8 km or more from breeding dens of red foxes (Tannerfeldt et al. 2002a, Elmhagen 2003).

The different scales were measured on M711-maps (1:50,000) with an electronic telemeter (Model DM-138). Seven maps compose the study area, of which five were produced during 2001-2005 and two in 1985. Infrastructure outside the national park was possibly underestimated at the  $15 \times 15$  km scale for one den in use, whereas two dens, which were not in use, were affected at the  $7 \times 7$  km scale.

## Statistical analyses

We used the likelihood ratio tests in logistic regression models to compare predictor variables associated with used and unused arctic fox dens, and dens with and without litters. Because of the limited number of dens, we used a maximum of two predictor variables in each model, i.e. altitude and one infrastructure variable. The selected models were tested for overdispersion. We compared models using Akaike's information criterion corrected for small sample sizes ( $AIC_c$ ; Akaike 1973). We ranked models according to  $\Delta AIC_c$ , i.e. the difference in  $AIC_c$  (between each candidate model and the model with the lowest  $AIC_c$ ). Statistical analyses were carried out in R, version 2.6.1 (available at: <http://www.r-project.org>) and JMP, version 4 (© 1989-2000).

## Results

Dens used by arctic foxes in Børgefjell during 2001-2005 were on average situated at higher altitudes than dens not used (Fig. 1, Table 1). A similar difference was found between dens with recorded




Figure 1. Box plots, showing median, lower and upper quartiles (box), 10 and 90% quartiles (intervals) and sample minimum and maximum, for the relationship between arctic fox den use in Børgefjell and A) den altitude (in m a.s.l.), B) kilometres of trails, C) kilometres of roads and D) number of cabins. The three infrastructure variables were measured within squares of  $15 \times 15$  km with the den in the centre. For each predictor, the box to the left gives the values for dens used ( $N=14$ ) whereas the box to the right gives the values for dens not used ( $N=13$ ) during 2001-2005.

Table 1. Likelihood ratio tests in logistic regression models with the response variables arctic fox dens used ( $N=14$ ) and not used ( $N=13$ ) during 2002-2005. The predictors are kilometres of trails and roads and number of cabins within  $7 \times 7$  and  $15 \times 15$  km with the den in the centre. +/- signs after the variable names denote the effect in the statistical model.

Model	Used - not used			
	$\chi^2$	P	$R^2$	$\Delta AIC_c$
Den altitude +	28.43	< 0.001		
Cabins ( $7 \times 7$ ) -	12.84	< 0.001	0.83	0.00
Den altitude +	27.23	< 0.001		
Roads ( $7 \times 7$ ) -	10.62	0.001	0.77	2.21
Den altitude +	21.59	< 0.001		
Roads ( $15 \times 15$ ) -	7.81	0.005	0.69	5.80
Den altitude +	21.47	< 0.001		
Trails ( $15 \times 15$ ) -	6.77	0.009	0.67	6.06
Den altitude +	18.11	< 0.001	0.48	10.29
Den altitude +	18.95	< 0.001		
Trails ( $7 \times 7$ ) -	0.86	0.353	0.51	11.97
Cabins ( $15 \times 15$ ) -	14.18	< 0.001	0.38	14.23
Roads ( $15 \times 15$ ) -	4.33	0.038	0.12	24.08
Den altitude +	Unstable parameter estimates			
Cabins ( $15 \times 15$ ) -	Unstable parameter estimates			

Table 2. Likelihood ratio tests in logistic regression models with the response variables arctic fox dens with (N = 9) and without recorded litters (N = 18) during 2002-2005. See legend to Table 1 for explanations.

Model	Litters - no litters			
	$\chi^2$	P	R <sup>2</sup>	$\Delta AIC_c$
Den altitude + Roads (15 × 15) -	6.17	0.013	0.39	0.00
Den altitude + Cabins (15 × 15) -	5.00	0.025	0.32	2.29
Den altitude + Trails (7 × 7) -	8.16	0.004	0.31	2.61
Den altitude + Roads (15 × 15) -	3.48	0.062	0.21	3.55
Den altitude + Roads (15 × 15) -	7.29	0.007	0.21	3.62
Den altitude + Roads (7 × 7) -	7.21	0.007	0.28	3.89
Den altitude + Cabins (7 × 7) -	8.66	0.003	0.26	4.57
Den altitude + Cabins (15 × 15) -	1.52	0.218	0.18	4.75
Den altitude + Trails (15 × 15) -	6.09	0.014	0.24	5.02
Den altitude + Trails (15 × 15) -	6.83	0.009		
Den altitude + Trails (15 × 15) -	1.07	0.301		

litters and dens without litters (Table 2). Within the 7 × 7 km squares, the mean length of trails or roads and the number of cabins did not differ significantly between dens used and dens not used, or between dens with and without litters. Within the 15 × 15 km squares, the mean length of roads and the number of cabins was lower for dens in use than for dens not in use (see Fig. 1 and Table 1), and a similar difference was found between dens with and without litters (see Table 2). Also the mean length of trails tended to be lower within the 15 × 15 m squares for dens in use than for dens not in use (see Fig. 1).

For dens used/not used, the best (lowest AIC<sub>c</sub>) model included den altitude and number of cabins within 7 × 7 km squares, whereas the second best model included altitude and length of roads within 7 × 7 km (see Table 1). For dens with and without litters, the best model included altitude and length of roads within the 15 × 15 km squares, and the second best altitude and number of cabins within the same scale (see Table 2).

## Discussion

As predicted, arctic fox den use in Børgfjell during 2001-2005 was related not only to altitude, as pre-

viously reported (Frafjord 2003b), but also to human infrastructure, i.e. number of cabins and kilometres of roads. For dens used, the best model included fewer cabins within 7 × 7 km squares, whereas dens with litters were best explained by less roads within 15 × 15 km squares. One possible interpretation is that a low degree of infrastructure within the 7 × 7 km scale is sufficient for the arctic fox to use a den, but not sufficient to allow successful breeding if there is a high load of infrastructure within the 15 × 15 km scale.

We find the most likely explanation for the negative relationship between arctic fox den use and infrastructure to be increased interference with red foxes that might have expanded their distribution following human activity and associated increases in garbage, remains from fishing and road-killed animals. Such food resources should be more stable in abundance compared to the very cyclic availability of small rodents, and may be especially important for the red fox in periods when natural food resources are scarce (Nielsen 1990, Lucherini & Crema 1994, Ferrari & Weber 1995). We suggest that increased food availability at the interface of the two fox species has affected the total red fox population positively, resulting in an influx of non-breeding red foxes to high-alpine areas, where the summer may be too short for the red fox to breed (Frafjord 2003a).

Alternative hypotheses for a negative relationship between arctic fox den use and infrastructure are that arctic foxes avoid breeding close to humans, or that populations of other arctic fox predators have responded positively to human infrastructure. We exclude the first alternative hypothesis because arctic foxes are usually not very shy; in fact, the Lappish name of the species means the fearless and foolhardy (Østbye & Pedersen 1990). We exclude the second alternative hypothesis because the two potential predators of arctic fox cubs in Børgfjell, the golden eagle *Aquila chrysaetos* and the wolverine *Gulo gulo* (Frafjord et al. 1989, Tannerfeldt & Angerbjörn 1996), avoid areas with human activity (May et al. 2006, Kaisanlahti-Jokimaki et al. 2008). Besides, the wolverine population in Børgfjell is low, with 0-2 recorded reproductions each year during our study period (Brøseth & Andersen 2007).

Børgfjell currently supports the most successful population of arctic fox in Fennoscandia, both with regard to numbers of breeding animals and reproductive performance. Winter feeding of arctic foxes

and red fox control in Swedish Borgafjäll (Elmhagen et al. 2004) may be one contributing factor, but it may also be important that Børgfjell historically has had a relatively low natural abundance of red fox compared to other alpine areas. This might be because winters are severe, but also because Børgfjell has been less affected by human activities than most other alpine regions. Hence, recent increases in the number of vacation cabins around the park may have increased the negative impact on the arctic fox. As interference with the red fox appears to be a likely explanation for the observed patterns in arctic fox den use, further research should focus on factors that are of importance for the red fox expansion to the alpine habitats. Conservation efforts should hence also be focused on reducing the driving forces behind red fox expansion.

*Acknowledgements* - we thank the Norwegian Directorate for Nature Management for giving us access to data from the national arctic fox database. We also thank Per A. Lorentzen, Statskog, for practical help and for valuable discussions in the field, and Karl Frafjord, Shyamala Ratnayake and three anonymous referees for comments on earlier versions of the manuscript. Special thanks to Christian Nellemann for inspiration to explore the influence of human development in alpine landscapes.

## References

- Akaike, H. 1973: Information theory and an extension of the maximum likelihood principle. - In: Petran, B.N. & Csaki, F. (Eds.); International symposium on information theory. 2nd edition. Akademiai Kiado, Budapest, Hungary, pp. 267-281
- Andersen, R., Linnell, J.D.C., Eide, N.E. & Landa, A. 2005: Fjellrev i Norge 2005. Overvåkingsrapport. (In Norwegian with an English summary: Arctic foxes in Norway 2005. Monitoring report). - NINA Rapport 101: 1-21.
- Angerbjörn, A., Ströman, J. & Becker, D. 1997: Home range patterns in arctic foxes in Sweden. - *Journal of Wildlife Research* 2: 9-14.
- Angerbjörn, A., Tannerfeldt, M., Henttonen, H., Elmhagen, B. & Dalén, L. 2002: SEFALO Bevarande av fjällräv *Alopex lagopus* i Sverige och Finland. - Sluttrapport July 1998 - December 2002, 37 pp.
- Bailey, E.P. 1992: Red foxes, *Vulpes vulpes*, as biological control agents for introduced arctic foxes, *Alopex lagopus*, on Alaskan islands. - *Canadian Field-Naturalist* 106: 200-205.
- Bevanger, K. & Brøseth, H. 2004: Impact of power lines on bird mortality in a subalpine area. - *Animal Biodiversity and Conservation* 27: 67-77.
- Brøseth, H. & Andersen, R. 2007: Yngleregistreringer av jerv i Norge i 2007. (In Norwegian with an English summary: Monitoring of active wolverine dens in Norway in 2007). - NINA Rapport 295: 1-20.
- Dalerum, F., Tannerfeldt, M., Elmhagen, B., Becker, D. & Angerbjörn, A. 2002: Distribution, morphology and use of arctic fox *Alopex lagopus* dens in Sweden. - *Wildlife Biology* 8(3): 185-192.
- Eide, N.E., Landa, A., Flagstad, Ø., Andersen, R., van Dijk, J., Meås, R., Berntsen, F. & Bruteig, I.E. 2009: Bevaringsbiologi fjellrev. Framdriftsrapport 2007-2008. (In Norwegian with an English summary: Conservation biology Arctic fox. Progress report 2007-2008). - NINA Rapport 390: 1-57.
- Elmhagen, B. 2003: Interference competition between arctic and red foxes. - PhD thesis, Department of Zoology, University of Stockholm, 34 pp.
- Elmhagen, B., Angerbjörn, A., Henttonen, H., Eide, N. & Landa, A. 2004: Saving the endangered Fennoscandian *Alopex lagopus*. - SEFALO+, Stockholm, 35 pp.
- Forman, R.T.T. & Alexander, L.E. 1998: Roads and their major ecological effects. - *Annual Review of Ecology and Systematics* 29: 207-231.
- Ferrari, N. & Weber, J-M 1995: Influence of the abundance of food resources on the feeding habits of the red fox, *Vulpes vulpes*, in western Switzerland. - *Journal of Zoology (London)* 236: 117-129.
- Frafjord, K. 2003a: Den røde fare: kan fjellreven overleve rødreven? (In Norwegian with an English summary: Threats of the reds: can arctic foxes in Norway survive the red fox?). - *Fauna* 56: 150-159.
- Frafjord, K. 2003b: Ecology and use of arctic fox *Alopex lagopus* dens in Norway: tradition overtaken by interspecific competition? - *Biological Conservation* 111: 445-453.
- Frafjord, K., Becker, D. & Angerbjörn, A. 1989: Interactions between arctic and red foxes in Scandinavia - Predation and aggression. - *Arctic* 42: 354-356.
- Frafjord, K., Spjøtvoll, Ø., Lorentzen, P. & Grøneng, T. 1998: Reproduksjon hos fjellrev i Børgfjell. (In Norwegian with an English summary: Dynamics of arctic fox reproduction in Børgfjell, Norway). - *Fauna* 51: 128-135.
- Hersteinsson, P. & Macdonald, D.W. 1992: Interspecific competition and the geographical distribution of red and arctic foxes *Vulpes vulpes* and *Alopex lagopus*. - *Oikos* 64: 505-515.
- Hjeljord, O. 1980: Viltbiologi. - Landbruksforlaget, Oslo, 318 pp. (In Norwegian).
- Kaisanlahti-Jokimaki, M-L, Jokimaki, J., Huhta, E., Ukkola, M., Helle, P. & Ollila, T. 2008: Territory occupancy and breeding success of the Golden Eagle (*Aquila chrysaetos*) around tourist destinations in northern Finland. - *Ornis Fennica* 85: 2-12.
- Landa, A., Strand, O., Linnell, J.D.C. & Skogland, T. 1998: Home-range sizes and altitude selection for arctic foxes

- and wolverines in an alpine environment. - Canadian Journal of Zoology 76: 448-457.
- Lindström, E. 1982: Population ecology of the red fox (*Vulpes vulpes L.*) in relation to food supply. - PhD thesis, University of Stockholm, Sweden, 24 pp.
- Linnell, J.D.C., Strand, O. & Landa, A. 1999: Use of dens by red *Vulpes vulpes* and arctic *Alopex lagopus* foxes in alpine environments: Can inter-specific competition explain the non-recovery of Norwegian arctic fox populations? - Wildlife Biology 5(3): 167-176.
- Linnell, J.D.C., Landa, A., Andersen, R., Strand, O., Eide, N., van Dijk, J. & May, R. 2004: Captive-breeding, population supplementation and reintroduction as tools to conserve endangered arctic fox populations in Norway: Detailed proposal and progress 2001-2004. - NINA Oppdragsmelding 825: 1-22.
- Lönnberg, E. 1927: Fjellrevstammen i Sverige 1926. - Svenska Vetenskapsakademiens Skrifter i Naturskydds-ärenden 7: 1-23. (In Swedish).
- Lucherini, M. & Crema, G. 1994: Seasonal variation in diet and trophic niche of the red fox in an alpine habitat. - Zeitschrift für Säugetierkunde 59: 1-8.
- Lund, H.M-K. 1962: The red fox in Norway II. The feeding habits of the red fox in Norway. - Papers of the Norwegian State Game Research Institute, 2. series, no. 12: 1-79.
- May, R., Landa, A., van Dijk, J., Linnell, J.D.C. & Andersen, R. 2006: Impact of infrastructure on habitat selection of wolverine *Gulo gulo*. - Wildlife Biology 12(3): 285-295.
- Nielsen, S.M. 1990: The food of rural and suburban woodland foxes *Vulpes vulpes* in Denmark. - Jutlandica 23: 25-32.
- Oksanen, T., Schneider, M., Rammul, Ü, Hambäck, P. & Aunapuu, M. 1999: Population fluctuations of voles in North Fennoscandian tundra: contrasting dynamics in adjacent areas with different habitat composition. - Oikos 86: 463-478.
- Østbye, E. & Pedersen, Ø. 1990: Fjellreven. - In: Semb-Johansson, A. & Frislid, R. (Eds.); Norges Dyr, Pattedyrene 1. Cappelen, Oslo, pp. 48-59. (In Norwegian).
- Østbye, E., Skar, H.J., Svalastog, D. & Westby, K. 1978: Fjellrev og rødrev på Hardangervidda; hiøkologi, utbredelse og bestandsstatus. (In Norwegian with an English summary: Arctic fox (*Alopex lagopus*) and red fox (*Vulpes vulpes*) on Hardangervidda; den ecology, distribution and population status). - Meddelelser fra norsk viltforskning 3. serie, nr. 4. Direktoratet for vilt og ferskvannsfisk, Trondheim, 66 pp.
- Røhnbæk, E. 2004: Rødrevens aktivitet i forhold til hyttefelt - Sporfordeling av rødrev og byttedyr i og rundt fem hyttefelt i Ringsakfjellet. - Prosjektoppgave ved 3-årig utmarksforvaltning, Høgskolen i Hedmark, Evenstad, 31 pp. (In Norwegian).
- Selås, V. & Vik, J.O. 2006: Possible impact of snow depth and ungulate carcasses on red fox *Vulpes vulpes* populations in Norway 1897-1976. - Journal of Zoology (London) 269: 299-308.
- Selås, V. & Vik, J.O. 2007: The arctic fox *Alopex lagopus* in Fennoscandia: a victim of human-induced changes in interspecific competition and predation? - Biodiversity and Conservation 16: 3575-3583.
- Tannerfeldt, M. & Angerbjörn, A. 1996: Life history strategies in fluctuating environment: establishment and reproductive success in the arctic fox. - Ecography 19: 209-220.
- Tannerfeldt, M., Elmhagen, B. & Angerbjörn, A. 2002a: Exclusion by interference competition? The relationship between red and arctic foxes. - Oecologia 132: 213-220.
- Tannerfeldt, M., Elmhagen, B. & Angerbjörn, A. 2002b: Food-niche overlap between arctic and red foxes. - Canadian Journal of Zoology 80: 1274-1285.