

Suitability of NIRS analysis for estimating diet quality of free-living red deer *Cervus elaphus* and roe deer *Capreolus capreolus*

Authors: Kamler, Jiří, Homolka, Miloslav, and Čižmár, David

Source: Wildlife Biology, 10(3) : 235-240

Published By: Nordic Board for Wildlife Research

URL: <https://doi.org/10.2981/wlb.2004.021>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Suitability of NIRS analysis for estimating diet quality of free-living red deer *Cervus elaphus* and roe deer *Capreolus capreolus*

Jiří Kamler, Miloslav Homolka & David Čížmár

Kamler, J., Homolka, M. & Čížmár, D. 2004: Suitability of NIRS analysis for estimating diet quality of free-living red deer *Cervus elaphus* and roe deer *Capreolus capreolus*. - Wildl. Biol. 10: 235-240.

In this study, we tested the efficiency of near infrared reflectance spectroscopy (NIRS) to assess nitrogen content in faeces of free-living ruminants. Faecal nitrogen (FN) content was analysed in 168 pellet groups from red deer *Cervus elaphus* and roe deer *Capreolus capreolus* in the growing season and in winter using both the standard Kjeldahl method and NIRS analysis. Estimates of nitrogen content obtained by the two methods did not differ ($P > 0.1$), and the correlation between FN values was significant ($P < 0.001$). FN content ranged within 1.10-4.58% of dry matter and, as anticipated, it was higher in the growing season than in winter in both species ($P < 0.01$). Faecal nitrogen values were also higher for roe deer than for red deer, although the difference was only significant in the growing season ($P = 0.007$). Our study confirmed that faecal nitrogen of free-living ruminants can be accurately determined with NIRS analysis. NIRS represents a low-cost analytical technique, which could replace conventional labourious methods and is highly promising for analyses of diet quality in free-living ruminants.

Key words: deer, diet quality, faecal nitrogen, near infrared reflectance spectroscopy, NIRS

Jiří Kamler & Miloslav Homolka, Institute of Vertebrate Biology, Academy of Sciences of the Czech Republic, 603 65 Brno, the Czech Republic - e-mail addresses: kamler@brno.cas.cz (J. Kamler); homolka@brno.cas.cz (M. Homolka)

David Čížmár, Central Institute for Supervising and Testing in Agriculture, Hroznová 2, 65606 Brno, the Czech Republic - e-mail: david.cizmar@lo.zeus.cz

Corresponding author: Jiří Kamler

Received 12 April 2002, accepted 22 September 2003

Associate Editor: Anne Loison

Information on the botanical composition and quality of herbivore diet is an essential component of studies of their feeding ecology. Studies of the diet of free-liv-

ing ungulates are usually based on samples of faeces or rumen contents (Holeček, Vavra & Pieper 1982, de Jong, Gill, van Wieren & Burlton 1995, Cornelis,

Casaer & Hermy 1999, Latham, Staines & Gorman 1999). Collection of rumen contents from killed animals is often limited to certain periods of the year and can be impossible in protected areas or in species that are not hunted. Therefore, it is important to develop reliable non-invasive techniques such as faecal analysis.

Faecal samples collected for studies of the botanical composition of diets may also be used to estimate diet quality (Sinclair, Krebs & Smith 1982, Putman & Hemmings 1986). This is due to the fact that there seems to be good correlation between diet quality (e.g. dietary nitrogen, dietary digestible energy, dry matter intake, and dry matter digestibility) and faecal chemical constituents (e.g. faecal nitrogen, faecal 2,6-diaminopimelic acid, and faecal fibre; Leslie, Starkey & Vavra 1984, Leslie & Starkey 1985, Jenks, Soper, Lochmiller & Leslie 1990).

Recent studies indicate that near infrared reflectance spectroscopy (NIRS) can be a valuable method to estimate the variety of chemical components in different materials like soil, plant or animal tissues (Büning-Pfaue, Hartman, Harder, Kehraus & Urban 1998, Foley, McIlwee, Lawler, Aragones, Woolnough & Berding 1998, Ludwig, Khanna, Bauhus & Hopmans 2002). NIRS has been successfully applied in agriculture to determine the chemical composition (quality) of pastures and other food (Offer, Percival, Dewhurst & Thomas 1998, Kays, Barton & Windham 2000). Some studies also used NIRS to estimate diet quality of ruminants from faecal samples. Diet crude protein, diet digestibility and also energetic values of the diet have been successfully estimated from faeces using this method (Lyons & Stuth 1992, Leite & Stuth 1994, Leite & Stuth 1995, Purnomoadi, Kurihara, Nishida, Shibata, Abe & Kameoka 1996). NIRS can also be used to estimate other characteristics of diet such as mineral content (Windham, Hill & Stuedemann 1991), energy content of the diet of non-ruminants (van Barneveld, Nuttall, Flinn & Osborne 1999) and botanical composition of the diet as well (Petersen, Barton, Windham & Hoveland 1987, Volesky & Coleman 1996, Walker, Clark & McCoy 1998, Walker, McCoy, Launchbaugh, Fraker & Powel 2002).

The potential for using NIRS in this way has been tested mostly on domestic animals so far; wider verification of the method for use in studies of diet quality in free-living ruminants has not previously been undertaken. In this study, we compare FN content obtained by the traditional Kjeldahl method and by NIRS to test the applicability of NIRS for this analysis in free-living ruminants. To verify the method on various types of material we used dung pellets sampled from red deer *Cervus elaphus* and roe deer *Capreolus capreolus* in the grow-

ing and winter seasons. Differences in feeding ecology between these two species exist, particularly when availability and quality of food is high, thus allowing a greater degree of selection; the diets of the same two species are more similar in localities (or seasons) where food resources are more limited (Homolka 1996). Red and roe deer are ruminants belonging to distinct feeding types (Hofmann 1989). 'Concentrate-selectors' such as roe deer, are generally more selective in their choice of diet as they require diet of higher quality (higher nitrogen, energy and lower fibre content) than do 'intermediate-feeders' such as red deer. The diet of the roe deer thus contains higher quality components like browse and forbs (de Jong et al. 1995, Homolka 1996, Raymond, Servello, Griffith & Eschholz 1996) and smaller proportions of grasses of low quality than does the diet of red deer (Homolka 1995, Merrill, Callahan-Olson, Rae-deke, Taber & Anderson 1995, Gebert & Verheyden 2001).

In winter, free-living ruminants in the temperate zone face great limitation in food resources (Worden & Perkins 1995) and the quality of their diet is generally lower than in the growing season (Risenhoover 1989, Kucera 1997).

We therefore expected to find higher FN content in roe deer than in red deer and higher FN values for both species in summer than in winter.

Material and methods

Material for our study was collected in the Jeseniky Mountains in the Czech Republic. We collected 168 fresh faecal pellet groups in two seasons: the vegetation period running from May through September and the winter running from November through February (see Table 1). Samples were air-dried at 60°C for 48 hours, ground in a mill to pass a 1-mm screen, and divided into two subsamples. The first group of subsamples (reference values) were analysed for nitrogen using the standard Kjeldahl procedures (AOAC 1980) and the second group of subsamples were analysed using NIRS.

NIRS is based on the development of a calibration equation that reflects the relationship between the constituents in the sample and NIRS spectral information. Measurements were carried out using a FOSS NIR-System 6500, a near infrared reflectance spectrophotometer in the 1,100–2,500 nm wavelength range. The spectra produced by the NIRS instrument represent the total chemical and physical properties of a sample. Chemical information appears at a specific location in the spectrum. Physical properties of a sample, such as

particle size, are eliminated by mathematical corrections. We used standard normal variate correction (SNV) which corrects each ith spectrum (row) in the data matrix separately by subtraction of the row mean and normalising in the row direction (Barnes, Dhanoa & Lister 1989). Data were analysed by linear regression using software NIR Calibration 1.0 (EFFICHEM, CZ). After defining the calibration data set, the software computes the calibration equation. Calibration equations quantify the relationship between NIR absorption and laboratory reference method. Partial least square (PLS) was chosen as the regression because it is generally considered to be the method of choice in multivariate calibration (Martens & Næs 1991). To examine the relationship between the results obtained using the two tested methods, we used the correlation coefficient and standard error of cross validation (SECV). SECV was used to estimate the error of prediction for unknown samples by simulating the prediction process by leaving part of the data set out, developing the calibration model on the rest of the data matrix, and making predictions for samples left out. This process was repeated several times so that each sample was left out once.

Before comparing the values obtained by the two methods, we tested the differences of reference faecal nitrogen (FN) values in subsamples (species, season). Both season and species affected the FN values (ANOVA: $F_{3,164} = 32.76$; $P < 0.001$); therefore, we tested the differences between the two methods used separately for seasons and species (t-test). To test if NIRS performed well for both feeding specialists, we calculated the equation of the regression curve separately for red and roe deer. Differences in the x-y regression slope were examined according to Armitage, Berry & Matthews (2001).

Finally, we tested the significance of differences in the reference FN values between species and seasons (two species in one season and one species in two seasons) using a t-test.




Figure 1. Reference nitrogen (%; obtained using the Kjeldahl method) and NIRS predicted nitrogen (%) in faeces of red deer (CE) and roe deer (CC) in the growing season (g = spring-summer) and winter (w) in Jeseníky Mountains ($y = 0.9964x$, $r^2 = 0.968$, $SECV = 0.15$). The thin line represents the ideal equation $Y = X$.

Results

In all samples, faecal nitrogen ranged within 1.10–4.58% of dry matter. We did not find any significant differences between the two methods when applied to paired samples whatever the species or season ($P > 0.1$ in all cases; Table 1). Values of the nitrogen content obtained by the two methods were closely correlated in pooled data sets ($r = 0.985$, $N = 168$, $P < 0.001$, $SECV = 0.15$; Fig. 1).

The calibration values determined separately for each of the species were also significantly correlated (red deer: $r = 0.985$, $P < 0.001$, $SECV = 0.147$; roe deer: $r = 0.986$, $P < 0.001$, $SECV = 0.12$), and the equations of the two regression curves were nearly identical (red deer: $y = 0.9642x + 0.0811$; roe deer: $y = 0.9509x + 0.1405$) and their slopes did not differ significantly ($t = 0.500$; $P = 0.597$).

In accordance with our hypothesis, the reference FN values were higher in roe deer than in red deer in sum-

Table 1. Number of samples (N) and nitrogen volume (%; $\bar{x} \pm SD$) in red deer (CE) and roe deer (CC) faeces during the growing season (g = spring-summer) and winter (w) in our study (Jeseníky) found using the Kjeldahl method (lab) and NIRS analysis (NIR), significance of their differences (t-test) and comparison with literature data from Georgia (Leslie & Starkey 1985) and Nevada (Osborn & Jenks 1998), Sweden (Wahlström & Kjellander 1995) and Hampshire (Putman & Hemmings 1986).

| | N | Jeseníky-lab | Jeseníky-NIR | Differences | | Georgia | Nevada | Sweden | Hampshire |
|------|----|---------------|---------------|-------------|-------|------------------|------------------|--------|-----------|
| | | | | t | P | | | | |
| CE-g | 96 | 2.5 ± 0.7 | 2.5 ± 0.7 | 0.001 | 0.999 | 3.1 ± 0.03 | | | 4.1** |
| CE-w | 15 | 1.4 ± 0.1 | 1.5 ± 0.2 | 0.363 | 0.720 | 1.4 ± 0.04 | | | 2.4** |
| CC-g | 30 | 2.9 ± 0.6 | 2.9 ± 0.6 | 0.018 | 0.986 | $3.4 \pm 0.12^*$ | | | 3.5 |
| CC-w | 23 | 1.5 ± 0.5 | 1.5 ± 0.4 | 0.552 | 0.584 | $2.1 \pm 0.12^*$ | $1.7 \pm 0.04^*$ | 2.43 | 2.6 |

* *Odocoileus* spp.

** *Cervus nippon*

mer ($t = 2.737$, $P = 0.007$), but they did not differ in winter ($t = 0.994$; $P = 0.330$; see Table 1). Values of faecal nitrogen were significantly higher in the growing season than in winter for both red and roe deer ($t = 13.557$, $P < 0.001$ and $t = 9.445$, $P < 0.001$ in red and roe deer, respectively).

Discussion

NIRS has been successfully used for analysis of the chemical composition of different materials. Several studies have used NIRS to estimate diet quality (crude protein, digestible organic matter or energy) in domestic and also free-living ruminants (Lyons & Stuth 1992, Purnomoadi et al. 1996). Our study verified the applicability of NIRS for estimation of faecal nitrogen in free-living ruminants. We did not find any differences in reliability of NIRS between deer species and seasons, and we confirmed advantages of NIRS.

We applied NIRS in estimation of faecal nitrogen, which is generally accepted as a reliable indicator of diet quality of free-living herbivores (e.g. Sinclair et al. 1982, Putman & Hemmings 1986, Hodgman, Davitt & Nelson 1996, Kamler, Homolka & Kráčmar 2003). Despite some doubts about its reliability (Hobbs 1987) NIRS is now widely used. Our results corresponded well with the expected differences in diet quality between seasons and deer species. We expected higher FN values in browsers (roe deer) than in intermediate feeders (red deer), and in the growing season than in winter. Roe deer usually consume a diet of less grasses and of a better quality than do red deer (Kuen & Bubenik 1980, Hearney & Jennings 1983, Latham et al. 1999). The fact that we found no differences in FN contents between the two species in winter may be due to the close similarity of red and roe deer diets in the period with snow cover (Homolka 1993).

The differences between the content of FN found in our samples and those reported by other authors (see Table 1) may have been caused by different quality of forage supply. A lower content of FN (2.43% DM) was found in roe deer from a forest environment in Sweden during summer (Wahlström & Kjellander 1995). On the other hand, the content of nitrogen detected in black-tailed deer *Odocoileus hemionus columbianus* (a feeding specialist like roe deer) in Georgia, USA, in a region with broadleaved forests was higher (3.4% DM; Leslie & Starkey 1985).

Our results showed the potential in using NIRS to estimate diet quality in free-living ruminants from faeces. We would also emphasise the advantages of NIRS, for

instance speed and low cost of analysis. Furthermore, the number of samples that can be processed is not limited and the validity of the estimated parameters is not influenced by sample freshness (Pearce, Lyons & Stuth 1993, Leite & Stuth 1994). Our findings agree with the results of similar studies showing the potential of NIRS to estimate diet quality in ruminants (Leite & Stuth 1995, Cozzolino, La Manna & Martins 2002). We find the application of NIRS highly promising for faecal analyses of different characteristics of the diet of free-living ruminants. The real advantages of NIRS are its ability to estimate diet quality (digestible energy and protein) from faeces and also that it makes it possible to determine the botanical composition of the diet in wild ruminants from faecal material. Nevertheless, differences in botanical composition of the diet may influence the accuracy of NIRS, thus we recommend that further studies be carried out, especially in different feeding specialists.

Acknowledgements - we thank Anne Loison and two referees who provided constructive remarks that improved this article. Rory Putman substantially improved the structure and the English of the final version. The study was supported by the Grant Agency of the Czech Republic, Grant No. 206/99/D053.

References

- AOAC 1980: Official methods of analysis. - Association of Official Analytical Chemists, Washington, D.C., 1081 pp.
- Armitage, P., Berry, G. & Matthews, J.N.S. 2001: Statistical methods in medical research. - Blackwell Science, Oxford, 832 pp.
- Barnes, R.J., Dhanoa, M.S. & Lister, S.J. 1989: Standard normal variate transformation and de-trending of near-infrared diffuse reflectance spectra. - *Applied Spectroscopy* 43: 772-777.
- Büning-Pfaue, H., Hartman, R., Harder, J., Kehraus, S. & Urban, C. 1998: NIR-spectrometric analysis of food. Methodical development and achievable performance values. - *Fresenius Journal of Analytical Chemistry* 360: 832-835.
- Cornelis, J., Casaer, J. & Hermly, M. 1999: Impact of season, habitat and research techniques on diet composition of roe deer (*Capreolus capreolus*): a review. - *Journal of Zoology* 248: 195-207.
- Cozzolino, D., La Manna, A. & Martins, D.V. 2002: Use of near infrared reflectance spectroscopy to analyse bovine faecal samples. - *Journal of Near Infrared Spectroscopy* 10: 309-314.
- de Jong, C.B., Gill, R.M.A., Van Wieren, S.E. & Burlton, F.W.E. 1995: Diet selection by roe deer *Capreolus capre-*

- olus in Kielder Forest in relation to plant cover. - *Forest Ecology and Management* 79: 91-97.
- Foley, W.J., McIlwee, A., Lawler, I., Aragones, L., Woolnough, A.P. & Berding, N. 1998: Ecological applications of near infrared reflectance spectroscopy a tool for rapid, cost-effective prediction of the composition of plant and animal tissues and aspects of animal performance. - *Oecologia* 116: 293-305.
- Gebert, C. & Verheyden, H. 2001: Variations of diet composition of red deer (*Cervus elaphus* L.) in Europe. - *Mammalian Review* 31: 189-201.
- Hearney, A.W. & Jennings, T.J. 1983: Annual foods of the red deer (*Cervus elaphus*) and the roe deer (*Capreolus capreolus*) in the east of England. - *Journal of Zoology* 201: 565-570.
- Hobbs, N.T. 1987: Fecal indices to dietary quality: a critique. - *Journal of Wildlife Management* 51: 317-320.
- Hodgman, T.P., Davitt, B.B. & Nelson, J.R. 1996: Monitoring mule deer diet quality and intake with fecal indices. - *Journal of Range Management* 49: 215-222.
- Hofmann, R.R. 1989: Evolutionary step of ecophysiological adaptation and diversification of ruminants: a comparative view of their digestive system. - *Oecologia* 78: 443-457.
- Holechek, J.L., Vavra, M. & Pieper, R.D. 1982: Methods for determining the nutritive quality of range ruminant diets: a review. - *Journal of Animal Science* 54: 363-376.
- Homolka, M., 1993: The food niches of three ungulate species in a woodland complex. - *Folia Zoologica* 4: 193-203.
- Homolka, M. 1995: The diet of *Cervus elaphus* and *Capreolus capreolus* in deforested areas of the Moravskoslezské Beskydy Mountains. - *Folia Zoologica* 44: 227-236.
- Homolka, M. 1996: Foraging strategy of large herbivores in forest habitats. - *Folia Zoologica* 45: 127-136.
- Jenks, J.A., Soper, R.B., Lochmiller, R.L. & Leslie, J.R. 1990: Effect of exposure on nitrogen and fiber characteristic of white-tailed deer feces. - *Journal of Wildlife Management* 54: 389-391.
- Kamler, J., Homolka, M. & Kráčmar, J. 2003: Nitrogen characteristics of ungulates faeces: effect of time of exposure and storage. - *Folia Zoologica* 52: 31-35.
- Kays, S.E., Barton, F.E. & Windham, W.R. 2000: Predicting protein content by near infrared reflectance spectroscopy in diverse cereal food products. - *Journal of Near Infrared Spectroscopy* 8: 35-43.
- Kucera, T.E. 1997: Fecal indicators, diet, and population parameters in mule deer. - *Journal of Wildlife Management* 61: 550-560.
- Kuen, H. & Bubenik, A.B. 1980: Botanische Pansenanalysen bei Rotwild (*Cervus elaphus hippelaphus*), Rehwild (*Capreolus capreolus*) und Gamswild (*Rupicapra rupicapra*). - In: *Alpine Umweltprobleme*. Achenkirch. T. V-IX; A 37, Beiträge zur Umweltgestaltung: 41-60. (In German).
- Latham, J., Staines, B.W. & Gorman, M.L. 1999: Comparative feeding ecology of red (*Cervus elaphus*) and roe deer (*Capreolus capreolus*) in Scottish plantation forests. - *Journal of Zoology* 247: 409-418.
- Leite, E.R. & Stuth, J.W. 1994: Influence of duration of exposure to field condition on viability of fecal samples for NIRS analysis. - *Journal of Range Management* 47: 312-314.
- Leite, E.R. & Stuth, J.W. 1995: Fecal NIRS equations to assess diet quality of free-ranging goats. - *Small Ruminant Research* 15: 223-230.
- Leslie, D.M. & Starkey, E.E. 1985: Fecal indices to dietary quality of cervids in old-growth forests. - *Journal of Wildlife Management* 49: 142-146.
- Leslie, D.M. & Starkey, E.E. 1987: Fecal indices to dietary quality: a reply. - *Journal of Wildlife Management* 51: 321-325.
- Leslie, D.M., Starkey, E.E. & Vavra, M. 1984: Elk and deer diets in old-growth forests in western Washington. - *Journal of Wildlife Management* 48: 762-775.
- Ludwig, K., Khanna, P.K., Bauhus, J. & Hopmans, P. 2002: Near infrared spectroscopy of forest soils to determine chemical and biological properties related to soil sustainability. - *Forest Ecology and Management* 171: 121-132.
- Lyons, R.K. & Stuth, J.W. 1992: Fecal NIRS equations for predicting diet quality of free-ranging cattle. - *Journal of Range Management* 45: 238-244.
- Martens, H. & Næs, T. 1991: *Multivariate Calibration*. - Wiley, Chichester, 438 pp.
- Merrill, E.H., Callahan-Olson, A.C., Raedeke, K.J., Taber, R.D. & Anderson, R.J. 1995: Elk (*Cervus elaphus roosevelti*) Dietary Composition and Quality in the Mount St. Helens Blast Zone. - *Northwest Science* 69: 9-18.
- Offer, N.W., Percival, D.S., Dewhurst, R.J. & Thomas, C. 1998: Prediction of the voluntary intake potential of grass silage by sheep and dairy cows from laboratory silage measurements. - *Animal Science* 66: 357-367.
- Osborn, F.G. & Jenks, J.A. 1998: Assessing dietary quality of white-tailed deer using fecal indices: effects of supplemental feeding and area. - *Journal of Mammalogy* 79: 437-447.
- Pearce, R.A., Lyons, R.K. & Stuth, J.W. 1993: Influence of handling methods on fecal NIRS evaluations. - *Journal of Range Management* 46: 274-276.
- Petersen, J.C., Barton, F.E., Windham, W.R. & Hoveland, C.S. 1987: Botanical composition definition of tall fescue-white clover mixtures by near infrared reflectance spectroscopy. - *Crop science* 27: 1077-1080.
- Purnomoadi, A., Kurihara, M., Nishida, T., Shibata, M., Abe, A. & Kameoka, K.I. 1996: Application of near infrared reflectance spectroscopy to predict fecal composition and its use for digestibility estimation. - *Animal Science & Technology* 67: 851-861.
- Putman, R.J. & Hemmings, G.J. 1986: Can dietary quality of free-ranging ungulates be simply determined from faecal chemistry? - *Acta Theriologica* 31: 257-270.
- Raymond, K.S., Servello, F.A., Griffith, B. & Eschholz, W.E. 1996: Winter foraging ecology of moose on glyphosate-treated clearcuts in Maine. - *Journal of Wildlife Management* 60: 753-763.
- Risenhoover, K.L. 1989: Composition and quality of moose winter diets in interior Alaska. - *Journal of Wildlife Management* 53: 568-577.

- Sinclair, A.R.E., Krebs, C.J. & Smith, J.N.M. 1982: Diet quality and food limitation in herbivores - the case of the snowshoe hare. - *Canadian Journal of Zoology* 60: 889-897.
- van Barneveld, R.J., Nuttall, J.D., Flinn, P.C. & Osborne, B.G. 1999: Near infrared reflectance measurement of the digestible energy content of cereals for growing pigs. - *Journal of Near Infrared Spectroscopy* 7: 1-7.
- Volesky, J.D. & Coleman, S.W. 1996: Estimation of botanical composition of esophageal extrusa samples using near infrared reflectance spectroscopy. - *Journal of Range Management* 49: 163-166.
- Wahlström, L.K. & Kjellander, F. 1995: Ideal free distribution and natal dispersal in female roe deer. - *Oecologia* 103: 302-308.
- Walker, J.W., Clark, D.G. & McCoy, S.D. 1998: Fecal NIRS for predicting percent leafy spurge in diets. - *Journal of Range Management* 51: 450-455.
- Walker, J.W., McCoy, S.D., Launchbaugh, K.L., Fraker, M.J. & Powel, J. 2002: Calibrating fecal NIRS equations for predicting botanical composition of diets. - *Journal of Range Management* 55: 374-382.
- Windham, W.R., Hill, N.S. & Stuedemann, J.A. 1991: Ash in forage esophageal and fecal samples analyzed using near-infrared reflectance spectroscopy. - *Crop Science* 31: 1345-1349.
- Worden, K.A. & Pekins, P.J. 1995: Seasonal change in feed-intake, body-composition, and metabolic-rate of white-tailed deer. - *Canadian Journal of Zoology* 73: 452-457.