

Index to Volume 124

Source: Rhodora, 124(1000) : 468-471

Published By: New England Botanical Society

URL: <https://doi.org/10.3119/0035-4902-124.1000.468>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

INDEX TO VOLUME 124

New scientific names are in bold.

A

Acanthaceae of the Nueva Galicia region of west-central Mexico: key to genera, species list with distributions by state, and a new species—*Justicia wilburii*. Thomas F. Daniel 228–253

acidic swamps 17–70

agaricomycetes 1–16

Aldrovanda (Droseraceae) 103–105; *A. vesiculosa* 103–105; *A. vesiculosa* var. *vesiculosa* 103

alga, red 98–102

Alford, Mac H. 141–145, 361–373

Almeda, Frank 133–146, 147–157

Andes Mountains 158–227

ANNOUNCEMENTS

Graduate Student Research Awards 465

Les Mehrhoff Botanical Research Awards 466

Merritt Lyndon Fernald Award 464

Antithamnion (Ceramiaceae) 98–102; *A. pectinatum* 98–102

Asher, Olivia A. 1–16

Ashrafi, Hamid 301–332

Asteraceae 361–373

B

Bahia, Brazil 147–157

Basidiomycota 1–16

Bertin, Robert I. 375–412

biosystematics 301–332

blueberries 301–332

BOOK REVIEWS

Ecological Guide to the Mosses & Common Liverworts of the Northeast by Sue Alix Williams 453–455

Boletes in the Bronx and beyond: a study of Boletales (Agaricomycetes, Basidiomycota, Fungi) specimen records of New York City. Olivia A. Asher and Naveed Davoodian 1–16

botanical history 71–90, 158–227

Botanical history and geographical distribution of *Carex waponahkikensis* (Cyperaceae). Marilee Lovit 71–90

Brazil 147–157

Bronx, New York 1–16

bryophytes 333–360

Button sedge fen: a newly described natural community

in three peatlands in New Hampshire. Daniel D. Sperduto and William F. Nichols (New England Note) 445–450

C

Cadeia do Espinhaço 147–157

Cameron, Kree 361–373

campo rupestre 147–157

Carex (Cyperaceae) 71–90; *C. albolutescens* 80; *C. crawfordii* 78; *C. longii* 76–77; *C. scoparia* 73–83; *C. scoparia* var. *tessellate* 71–90; *C. waponahkikensis* 71–90

Carter, Richard J. 361–373

cedar glade 287–300

Central America 228–253, 254–278

Chapada Diamantina, Bahia, Brazil 147–157

checklist 413–435

Connecticut 17–70

common garden 287–300

Crowl, Andrew A. 301–332

cudweeds 361–373

Cyperus (Cyperaceae) 279–286; *C. almensis* 284; *C. altonii* 284; *C. anisistii* 284; *C. conservator-davidii* 284; *C. consors* 285; *C. dichromeniformis* 284; *C. fedoniae* 284; *C. grandisimplex* 282, 284; *C. inops* 284; *C. lacustris* 284–285; *C. lundellii* 284; *C. miliifolius* 284; *C. odoratus* 279; *C. pearcei* 284; *C. simplex* 284; *C. squarrosus* 279; *C. subcastaneus* 284; *C. verlooeei* sect. *Incurvi* 279–286

Cyperus verlooeei sect. *Incurvi*, a new species from the Guiana Shield region of South America. Gordon C. Tucker and Mark T. Strong 279–286

D

Daniel, Thomas F. 139–140, 128–253

Davoodian, Naveed 1–16

dendrochronology 17–70

digitized herbarium specimens 1–16

distribution records 228–253, 413–435

diversity, *Sphagnum* 413–435

DNA barcoding 361–373

documented occurrences, Acanthaceae 228–253

E

early flowering 287–300

ectomycorrhizal fungi 1–16

edaphic endemic 287–300
 endemism 147–157, 228–253

F

Factors associated with the transition to weediness in the granitic outcrop endemic, *Phacelia dubia* var. *georgiana* (Hydrophyllaceae), and dating the time of the transition. Stacy Taylor-Bennetts and Foster Levy 287–300

Fernald, Merritt L. 71–90

Fidanza, Karina 147–157

floristic change 375–412

floristics 228–253

Franklin County, Massachusetts 375–412

Fritsch, Peter W. 301–332

fungi 1–16

G

Gamochaeta (Compositae) 361–373; *G. americana* 364; *G. antillana* 364; *G. argyrinea* 363–364, 367; *G. calviceps* 364, 368; *G. chionesthes* 364, 367; *G. coarctata* 363–364; *G. impatiens* 363–364, 367; *G. pensylvanica* 364, 368; *G. purpurea* 363–364, 367; *G. simplicicaulis* 364, 367; *G. ustulata* 363–364, 367–368

Gilman, Arthur V. 91–97

Globally rare *Aldrovanda vesiculosa* (Droseraceae) new to New Hampshire, U.S.A., The. Donald J. Padgett, Emmi Kurosawa, and Michael P. Graziano (New England Note) 103–105

Gnaphalieae 361–373

Gnaphalium 361–373

Graziano, Michael P. 103–105

Guyana 279–286

H

Haines, Arthur 91–97

hemlock-northern hardwoods 17–70

herbarium 287–300, 333–360

herbarium data 375–412

Hickler, Matthew G. 375–412

I

illustrations, botanical 158–227

Imperforate St. John's-wort (*Hypericum maculatum* subsp. *obtusiusculum*): a new overlooked nonnative forb for Eastern North America. Étienne Lacroix-Carignan, Colin J. Chapman-Lam, C. Sean Blaney, and Étienne Léveillé-Bourret (Note) 336–444

inoscultation 17–70

invasive plants 17–70, 375–412

Inventory of Connecticut's primeval forests, An. Jack Ruddat 17–70

J

Justicia (Acanthaceae) 228–253; *J. aurea* 232; *J. brandegeana* 232; *J. candicans* 231–232; *J. carnea* 232; *J. cuixmalensis* 232; *J. hilsenbeckii* 231; *J. Pacifica* 231; *J. pringlei* 239, 243–244; *J. salviiflora* 239, 243–244; *J. stellata* 232; *J. spicigera* 232; *J. wilburii* 132, 232, 239–244

K

Karlin, Eric F. 413–435

Kurosawa, Emmi 103–105

L

late seral 17–70

lectotype 91–97, 158–227

Lectotypification and an emended description of *Rosa* × *fernaldiorum*. Arthur V. Gilman and Arthur Haines 91–97

Levy, Foster 115–132, 140–141, 287–300

lichens 333–360

Lovit, Marilee 71–90

Luteyn, James L. 136–139, 158–227

M

Mackenzie, Kenneth K. 71–90

Macleania (Ericaceae: Vaccinieae) 158–227; *M. alata* 172; *M. alpicola* 173; *M. amplexicaulis* 173; *M. angulata* 163–170, 173–177; *M. antioquiae* 177; *M. arcuata* 177; *M. attenuata* 178; *M. benthamiana* 168, 178–179; *M. bullata* 179; *M. cardiophylla* 179–180; *M. coccinea* 165, 167–169, 180, 191; *M. coccoloboides* 180; *M. compacta* 181; *M. cordata* 165, 167–169, 181–182; *M. cordifolia* 168, 172, 182–183, 198–200; *M. costaricensis* 183; *M. costeroides* 183; *M. crassa* 183; *M. crenulata* 184–185; *M. dodsonii* 185; *M. ecuadorensis* 185; *M. elliptica* 185–186; *M. epiphytica* 186; *M. ericae* 186; *M. euryphylla* 186; *M. farinosa* 186; *M. floribunda* 159, 165, 167, 174, 178–179, 187; *M. glabra* 187–189; *M. hirtiflora* 189; *M. insignis* 163–172, 180, 190–191; *M. irazuensis* 191; *M. kalbreyeri* 191; *M. laurina* 191; *M. loeseneriana* 191; *M. longiflora* 164–169, 192; *M. macrantha* 179, 192–193; *M. maldonadensis* 193; *M. megabraceolata* 193; *M. mollis* 193; *M. multibracteata* 193; *M. nervosa* 193; *M. nitida* 194; *M. ovata* 194–195; *M. penduliflora* 195; *M. pentaptera* 186, 195–197; *M. pilgeriana* 197; *M. poortmannii* 197–198; *M. popenoei* 168, 197; *M. pubiflora* 198; *M. pulchra* 165, 167–169, 198–199; *M. punctata* 164, 167–168, 199–200; *M. racemosa* 200; *M. recumbens* 201; *M. recurvata* 177, 201; *M. reducta* 201; *M. robusta* 177, 201–202; *M. rotundifolia* 202; *M. rupestris* 168, 173, 177, 202–207; *M. salapa* 168–170, 207–208; *M. sleumeriana* 208;

- M. smithiana* 208–209; *M. sodiroi* 209; *M. speciosissima* 164–165, 167–169, 209–210; *M. spectabilis* 210; *M. stricta* 210; *M. subracemosa* 210; *M. subsessilis* 211; *M. talamancensis* 211–212; *M. tenuifolia* 212; *M. tovarensis* 212; *M. trianae* 212–213; *M. tropica* 213; *M. tuberosa* 172; *M. turrialbana* 213
- Maine 91–97
- Manos, Paul S. 301–332
- Marcetia* (Melastomataceae) 147–157; *M. acerosa* 153, 155; *M. ericoides* 147, 153, 155; *M. luetzelburgii* 153, 155; *M. lychnophoroides* 153, 155; *M. shepherdii* 147, 153, 155; *M. taxifolia* 147, 153; *M. wilburiana* 132, 147–157
- Marcetieae 147–157
- Massachusetts 375–412, 413–435
- Mathieson, Arthur C. 98–102
- Mexico 228–253
- Moore, Gerry 333–360
- morphology 301–332
- Motzkin, Glenn 375–41
- N**
- natural rose hybrid 91–97
- NEBS Meeting News 106–114, 456–463
- Neotropics 158–227
- neotype 158–227
- New England 17–70, 91–97, 98–102, 103–105, 375–412, 413–435
- New Hampshire 98–102, 103–105
- New *Marcetia* (Melastomataceae) from Chapada Diamantina, Bahia, Brazil, A. Frank Almeda, Ricardo Pacifico, and Karina Fidanza 147–157
- New York 1–16
- Nomenclator of Neotropical *Macleania* (Ericaceae: Vacciniaceae) with historical notes about early introductions into cultivation, A. James L. Luteyn 158–227
- nomenclature 91–97, 158–227, 254–278, 333–360
- North America 301–332, 333–360
- Nueva Galicia region, Mexico 228–253
- O**
- Occurrence of the introduced red alga *Antithamnion pectinatum* (Ceramiaceae) in New Hampshire, U.S.A. Arthur C. Mathieson and Kimberly W. Payne (New England Note) 98–102
- old growth 17–70
- P**
- Pacifico, Ricardo 147–157
- Padgett, Donald J. 103–105
- Palicourea* (Rubiaceae: Palicoureaeae) 254–278; sect. *Corymbiferae* 275; sect. *Montanae* 273, 275; sect. *Pseudocephaelis* 263, 268; *P. albaniana* 274–275; *P. carruthersii* 255–257; *P. clarkia* 273; *P. drewiana* 271, 273; *P. furcata* 257–260, 268; *P. geminata* 258, 260–264; *P. harlingii* 268, 271; *P. hispidula* 259; *P. hoffmannseggiana* 258–260, 263, 268; *P. iodotricha* 264–267; *P. japurensis* 267–268; *P. monsalvaeae* 269–271; *P. wilburiana* 132, 271–273; *P. winkleri* 258, 262; *P. woynkowskii* 273–274
- Patterns and changes in the nonnative flora of Franklin County, Massachusetts. Robert I. Bertin, Karen B. Searcy, Glenn Motzkin, and Matthew G. Hickler 375–412
- Payne, Kimberly W. 98–102
- peat mosses 413–435
- Phacelia dubia* (Hydrophyllaceae) 287–300; *P. dubia* var. *georgiana* 287–300; *P. dubia* var. *interior* 287–300; *P. dubia* var. *ronensis* 287–300
- phenotypic plasticity 287–300
- pollen 228–253
- polyploidy 301–332
- Preliminary review of North American material in Rafinesque’s herbarium, A. Gerry Moore 333–360
- primary forests 17–70
- Psychotria 254–278
- R**
- rapid evolution 287–300
- Remembering Robert L. Wilbur. Frank Almeda 133–146
- Report of the 2023 NEBS collecting foray in Bristol County, Rhode Island. Lisa A. Standley (The Botanists’ Corner) 451–452
- restingas 147–157
- Reviewers of Manuscripts 467
- Robert Lynch Wilbur: botanist, mentor, family man. Foster Levy and Charlotte M. Taylor 115–132
- Rosa* (Rosaceae) 91–97; *R. carolina* 91–97; *R. ×fernaldiorum* 91–97; *R. humilis* 92; *R. lucida* 92; *R. nitida* 91–97; *R. palustris* 91–97
- Rubiacearum Americanarum Magna Hama Pars LI: new species and nomenclatural clarifications in *Palicourea* (Rubiaceae: Palicoureaeae) in honor of Dr. Robert L. Wilbur. Charlotte M. Taylor 254–278
- Ruddat, Jack 17–70
- S**
- Searcy, Karen B. 375–412
- secondary old-growth 17–70
- sedge 279–286
- South America 147–157, 254–278, 279–286
- species delimitation 361–373
- Sphagnum* species of Massachusetts, The. Eric F. Karlin 413–435

- Sphagnum* (Sphagnopsida) 413–435; *S. affine* 422, 427–428; *S. andersonianum* 419, 426; *S. angermanicum* 427; *S. ×atlanticum* 421, 426; *S. austinii* 422, 427; *S. bartlettianum* 419, 426; *S. Beothuk* 429; *S. brevifolium* 426, 429; *S. ×carolinianum* 423–427; *S. capillifolium* 419–421; *S. ×centrale* 426; *S. contortum* 415, 428; *S. cuspidatum* 421–422, 426; *S. cyclophyllum* 419; *S. divinum* 419, 422, 427, 429; *S. diabolicum* 419, 422, 427, 429; *S. fallax* 426, 429; *S. fimbriatum* 427; *S. flavicomans* 427–428; *S. flexuosum* 414, 426; *S. ×henryense* 426; *S. imbricatum* 419, 422; *S. inundatum* 419, 423; *S. isoviitae* 426; *S. lescurii* 423–426, 428; *S. macrophyllum* 414, 426–427; *S. magellanicum* 419, 422, 426–427, 429; *S. majus* subsp. *Norvegicum* 419, 422, 426–427; *S. medium* 422, 427; *S. ×mississippiense* 419, 421, 426–427; *S. ×missouricum* 419, 423–426; *S. ×palustre* 426–427; *S. portoricense* 419; *S. pylaesii* 427; *S. quinquefarium* 427; *S. recurvum* 426–427; *S. rubellum* 419; *S. subsecundum* 422–427; *S. subnitens* 419; *S. tenellum* 414; 419, 426, 428–429; *S. tenerum* 414, 427; *S. teres* 428; *S. ×torreyanum* 421; *S. trinitense* 421, 427; *S. viride* 421, 426; *S. warnstorffii* 414
- Strong, Mark T. 279–286
- Systematics and evolution of *Vaccinium* sect. *Cyanococcus* (Ericaceae): progress and prospects. Peter W. Fritsch, Andrew A. Crowl, Hamid Ashrafi, and Paul S. Manos 301–332
- T**
- Taylor, Charlotte M. 115–132, 254–278
- Taylor-Bennetts, Stacy 287–300
- taxonomy 254–278, 301–332
- traprock ridges 17–70
- tree rings 17–70
- Tucker, Gordon C. 279–286
- typification 158–227, 333–360
- U**
- United States 1–16, 17–70, 91–97, 98–102, 103–105, 361–373, 375–412, 413–435
- urban ecology 1–16
- Utility of DNA markers for delimiting and identifying species of *Gamochoaeta* (Compositae) in the eastern United States, The. Mac H. Alford, Kree Cameron, and J. Richard Carter 361–373
- V**
- Vaccinium* (Ericaceae) 301–332; *V. altomontanum* 312, 315, 325; *V. amoenum* 313–314, 324–327; *V. angustifolium* 311–318, 325; *V. arboretum* 326; *V. arkansanum* 314, 324, 327–328; *V. ashei* 309, 313, 316, 324, 327; *V. atrococcum* 311–318, 324–328; *V. australe* 313–314; *V. boreale* 311–318; *V. brittonii* 311–312, 325; *V. caesariense* 311–318, 327–328; *V. constablaei* 309, 314–315, 327; *V. corymbosum* 302–303, 309–318, 327–328; *V. darrowii* 311–318, 325; *V. elliottii* 315–318, 324–327; *V. formosum* 314–315, 327; *V. fuscum* 313–318, 324, 327; *V. hirsutum* 306, 315; *V. holophyllum* 315; *V. lamarckii* 311–312, 325; *V. pallidum* 311–319, 325, 327; *V. ×marianum* 315, 327–328; *V. myrsinites* 303, 313, 315, 325–326; *V. myrtilloides* 311–318; *V.* sect. *Cyanococcus* 301–332; *V. simulatum* 314–315, 327–328; *V. stamineum* 326; *V. tenellum* 311–318, 325–326; *V. uliginosum* 303; *V. vacillans* 315–318, 325; *V. virgatum* 313–316, 324–327
- Venezuela 279–286
- virgin woodlands 17–70
- W**
- Wilbur, Robert Lynch 115–132, 133–146, 254–278, 333–360