



## **Presence of *Myotis alcathoe* Helversen & Heller, 2001 (Chiroptera: Vespertilionidae) in the Iberian Peninsula**

Authors: Agirre-Mendi, Pablo T., García-mudarra, Juan L., Juste, Javier, and Ibáñez, Carlos

Source: Acta Chiropterologica, 6(1) : 49-57

Published By: Museum and Institute of Zoology, Polish Academy of Sciences

URL: <https://doi.org/10.3161/001.006.0104>

---

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## Presence of *Myotis alcathoe* Helversen & Heller, 2001 (Chiroptera: Vespertilionidae) in the Iberian Peninsula

PABLO T. AGIRRE-MENDI<sup>1</sup>, JUAN L. GARCÍA-MUDARRA<sup>2</sup>, JAVIER JUSTE<sup>2,3</sup>,  
and CARLOS IBÁÑEZ<sup>2</sup>

<sup>1</sup>*c/ Paseo del Río Grande, nº 7, 3º I (Edificio Donosti), 01320 Oyón (Álava), Spain*

<sup>2</sup>*Estación Biológica de Doñana (CSIC), P.O. Box 1056, 41080 Sevilla, Spain*

<sup>3</sup>*Corresponding author: E-mail: juste@ebd.csic.es*

In a survey of bats from La Rioja (Spain), several specimens of the *mystacinus* group were captured at different mountain localities. Genetic and morphologic analyses have revealed the presence of two lineages within this group in La Rioja. The lineages have been identified as corresponding to two different species: *Myotis mystacinus* sensu stricto and the recently described *M. alcathoe*. Both species were found using the same nocturnal refugia (caves) and the same forest habitats. This study extends the distribution of *M. alcathoe* west and southwards and adds a new mammal species to the Iberian fauna.

*Key words:* bats, *Myotis mystacinus*, *M. alcathoe*, distribution, Iberia, mtDNA

### INTRODUCTION

In the last few years, a surprisingly high number of cryptic species have been found among European bats (Mayer and Helversen, 2001), particularly within the vespertilionid genera *Plecotus* (Kiefer *et al.*, 2002; Juste *et al.*, 2004) and *Myotis* (Castella *et al.*, 2000; Helversen *et al.*, 2001). *Myotis mystacinus* (Kuhl, 1817) was considered a single species until recently, but it has turned out to be a species group with as many as four different taxa recognised at present in the Western Palaearctic. *Myotis brandtii* (Eversmann, 1845) was historically the first species to be reinstated based on morphological differences from *M. mystacinus* (Hanák, 1965). Later, Volleth (1987) distinguished two new lineages: *Myotis* sp.

A and *Myotis* sp. B, based on the different distribution of active nucleolus organiser regions (NORs). In a thorough morphological revision, Benda and Tsytsulina (2000) assigned the *Myotis* sp. A lineage to the species *Myotis aurascens* Kuszakin, 1935, whereas Helversen *et al.* (2001) almost simultaneously described the lineage *Myotis* sp. B as *Myotis alcathoe* Helversen and Heller, 2001, based on differences in morphology, ecological preferences and mitochondrial DNA sequences (ND1 and 12S rRNA). In a recent phylogenetic study, Ruedi and Mayer (2001) have shown that despite morphological similarity, *M. mystacinus* and *M. brandtii* split a long time ago. The latter species is the only Palaearctic member of an otherwise monophyletic American clade. Similarly, Mayer and

Helversen (2001) found that *M. mystacinus* and *M. alcathoe* also split long ago and that they are not sister clades. The evolutionary relationships between *M. mystacinus* and *M. aurascens*, remain unclear.

Regarding the distribution of these forms occurring in the West Palaearctic, *M. brandtii* is found mainly across central and northern Europe, being generally absent in the Mediterranean basin, although it has been reported from Italy and the Balkans (Gerell, 1999a; Benda and Tsytsulina, 2000). The species *M. alcathoe* was first described as endemic from the Balkans and Hungary (Helversen *et al.*, 2001), and later reported from France (Ruedi *et al.*, 2002) and Slovakia (Benda *et al.*, 2003). *Myotis aurascens* seems to be distributed from the Balkans eastwards to Anatolia, Crimea and the Caucasus until Kazakhstan, whereas *M. mystacinus* is spread across most of Europe except in the east where it does not seem to overlap with *M. aurascens* (Benda and Tsytsulina, 2000). Of the *mystacinus* group only the species *M. mystacinus* has been reported so far from the Iberian Peninsula and shows a fragmented distribution, restricted

to mountain ranges in the northern half of the Iberian Peninsula (Fig. 1). Indeed, it is considered among the rarest bats in Iberia (Fernández and Ibáñez, 1989; Ibáñez *et al.*, 1992; Agirre-Mendi, 2002). Recently, Benda and Tsytsulina (2000) have suggested that the Iberian *M. mystacinus* is significantly larger than other European bats and have proposed their taxonomic distinction at subspecific level as *M. m. occidentalis* Benda, 2000.

In an ongoing faunal study of the ‘La Rioja’ autonomous region (Northern Spain), several specimens of bats of the *mystacinus* group were captured. The results of their morphological and genetic examination are described here and show the presence of *M. alcathoe* in Iberia, extending considerably the known distribution of this recently described species.

## MATERIAL AND METHODS

### Study area

The ‘La Rioja’ region is a small area (5,045 km<sup>2</sup>) located in central northern Iberia (42°39′–41°55′N;




FIG. 1. Left: distribution of bats of the *mystacinus* group in Iberia (Aihartza, 2001; Fernández-Gutiérrez, 2002; Gerell, 1999b; Woutersen and Bafaluy-Zoriguel, 2001). Right: sampling localities in La Rioja (Spain). Shade area > 800 m a.s.l.

3°08'–1°41'W) (Fig. 1). Despite its small size, La Rioja shows a diverse mosaic of landscapes and habitats. Climate and vegetation types are strongly correlated with altitude, which ranges from 260 m at the Ebro river in the north of the region to over 2,200 m in the mountains of the 'Sistema Ibérico' range in the south-western corner. In turns the lowlands, which are mainly under agricultural use, show strong Mediterranean influences. The mountain slopes have high humidity (rainfall over 1,000 mm) and were originally covered by a mixture of broad-leaved (*Fagus sylvatica* and *Quercus* sp.) and pine (*Pinus sylvestris*) forests (Fernández *et al.*, 1989; Martínez-Abaigar *et al.*, 1994).

### Sampling

From 1984 to 2003 a total of 197 localities were sampled for bats in La Rioja (Fig. 1). The sampling was carried out by setting up mist-nets over or near water surfaces (on 195 nights) or at the entrance of nocturnal refugia, caves and mines (on 174 nights).

A total of 25 bats of the *mystacinus* group were captured during the survey. Two older additional bibliographic records from the area (Ibáñez *et al.*, 1992) were also included in the study. Five museum specimens (preserved at the Estación Biológica de Doñana's scientific collections — EBD) were used for morphological and molecular comparisons and five additional samples of DNA of released bats were analysed (Appendix). Bats in the field were studied externally, measured (forearm length), identified and released. Wing punches were taken from some specimens, following procedures set out in Worthington Wilmer and Barratt (1996), to permit genetic study.

### Genetic Analysis

DNA was extracted from tissue samples following standard phenol/chloroform protocols (Maniatis *et al.*, 1989) or from wing biopsies preserved in ethanol following Higuchi *et al.* (1988). A fragment of subunit 1 of the mitochondrial NADH dehydrogenase (ND1) gene was amplified using primers ER65 and ER66 (Mayer and Helversen, 2001). The PCR cocktail (20 µl final reaction volume) included 2 µl of DNA extract, 1 µl of each primer (10 µM), 0.8 µl of MgCl<sub>2</sub> (50 mM), 0.16 µl dNTP (25 mM), 0.5 unit of taq-polymerase with appropriate buffer and H<sub>2</sub>O. Thermocycling consisted of 5' initial denaturation at 94°C, followed by 40 cycles at 94°C (30''), 63°C (30'') and 72°C (1'30''), with a final extension at 72°C (5'). The fragment was sequenced directly from purified PCR product using an ABI 3100 automated sequencer (Applied Biosystems, Warrington, UK)

following manufacturer's protocols. The partial sequences were aligned and edited using the program Sequencher 4.1 (Gene Codes Corp.) and inspected by eye. For species identification purposes, orthologous sequences of the European species of the *mystacinus* group (*Myotis mystacinus*, *M. alcathoe*, *M. aurascens* and *M. brandtii*) were obtained from GenBank (Helversen *et al.*, 2001; Mayer and Helversen, 2001) and included in the alignment. The best fitting substitution model was selected with hierarchical likelihood ratio tests (LRT — Modeltest, Posada and Crandall, 1998). Then, absolute differences and corrected distances between sequences were estimated using PAUP\* 4.0b10 software (Swofford, 2000) and MEGA v. 2.1 (Kumar *et al.*, 2001).

## RESULTS

### Genetic Analysis

A 565 bp fragment of the mitochondrial ND1 gene was sequenced from eight specimens from La Rioja. Sequences showed no evidence of heteroplasmy and the alignment, including the reference sequences from GenBank, showed 124 variable positions (Table 1), most of them being transitions and located in third positions. The sample consisted of four different haplotypes, two of them were new haplotypes and are deposited in the GenBank (accession numbers are given in Appendix).

Two clearly distinct lineages were found in the Iberian ND1 sequences that showed 16.3% corrected distance between them and over 16% with respect to *M. brandtii*. One of the lineages (Mmy 936, 937, 945, and 946) showed corrected distances < 0.3% and 0.8%, respectively, compared to the sequences of *M. mystacinus* and *M. aurascens* from the GenBank (Tables 1 and 2). Moreover, two individuals of this lineage shared the same haplotype with specimens of *M. mystacinus* from Germany (AY027848) and Spain (AY027847). Two other sequences (Mmy 104 and 968) clearly belonged to this lineage, but did not reach the 565 bp length in their sequences and were not included in further analyses.

TABLE 1. Polymorphic sites identified in the alignment of mitochondrial ND1 sequences from 8 bats of the *M. mystacinus* group from La Rioja, Spain (see Appendix) and orthologous sequences of *M. mystacinus* (AY027848Mmy), *M. alcaethoe* (AY027836Mal), *M. aurascens* (AY027844Mau), and *M. brandtii* (AY027851Mbr) deposited in the GenBank by Helversen et al. (2001) and Mayer and Helversen (2001)

|                     |       |             |               |             |             |
|---------------------|-------|-------------|---------------|-------------|-------------|
| Specimen            | Haplo | 11111       | 111111111     | 1111112222  | 2222222222  |
| AY027836Mal Hungary | 1     | TTTTGGAGGCA | TCAACGCTGC    | ACATAATACC  | CTCCTGTCTA  |
| Mal935 La Rioja     | 1     | .....       | .....         | .....       | .....       |
| Mal944 La Rioja     | 1     | .....       | .....         | .....       | .....       |
| Mal947 La Rioja     | 1     | .....       | .....         | .....       | .....       |
| Mal105 La Rioja     | 1     | .....       | .....         | .....       | .....       |
| AY027832Mal Greece  | 2     | .....A..... | .....G.....AT | .....T..... | .....       |
| AY027851Mbr Germany | 4     | C.CAAGAAAG  | C.TT.C.GC     | .....TCA.   | TT.TC.CC..  |
| AY027844Mau Greece  | 3     | CCCAA.AAA.  | C...T.GG.C    | ..TG.T...TT | ATC.CTC.TT  |
| AY027848Mmy Germany | 5     | CCCAA.AAA.  | C...T.GG.C    | GTG.T...TT  | ATC.CT...TT |
| Mmy945 La Rioja     | 5     | CCCAA.AAA.  | C...T.GG.C    | GTG.T...TT  | ATC.CT...TT |
| Mmy946 La Rioja     | 5     | CCCAA.AAA.  | C...T.GG.C    | GTG.T...TT  | ATC.CT...TT |
| Mmy936 La Rioja     | 6     | CCCAA.AAA.  | C...T.GG.C    | ..TG.T...TT | ATC.CT...TT |
| Mmy937 La Rioja     | 7     | CCCAA.AAA.  | C...T.GG.C    | GTG.T...TT  | ATC.CT...TT |
| AY027836Mal         | 1     | 222222222   | 2333333333    | 3333333333  | 3444444444  |
| Mal935 La Rioja     | 1     | .....       | .....         | .....       | .....       |
| Mal944 La Rioja     | 1     | .....       | .....         | .....       | .....       |
| Mal947 La Rioja     | 1     | .....       | .....         | .....       | .....       |
| Mal105 La Rioja     | 1     | .....       | .....         | .....       | .....       |
| AY027832Mal Greece  | 2     | .....       | .....         | .....       | .....       |
| AY027851Mbr Germany | 4     | CAAA.CC...  | C...GTC.TCT   | ..CATTCFA.T | ..CATTC..CC |
| AY027844Mau Greece  | 3     | .A.AG..TCT  | CTG.TCGTC.    | TCA.TCTAT.  | GCA..CGA..  |
| AY027848Mmy Germany | 5     | .A.AG..TCT  | CTG.TCGTC.    | TCA.TCTAT.  | GCA..CGA..  |
| Mmy945 La Rioja     | 5     | .A.AG..TCT  | CTG.TCGTC.    | TCA.TCTAT.  | GCA..CGA..  |
| Mmy946 La Rioja     | 5     | .A.AG..TCT  | CTG.TCGTC.    | TCA.TCTAT.  | GCA..CGA..  |
| Mmy936 La Rioja     | 6     | .A.AG..TCT  | CTG.TCGTC.    | TCA.TCTA..  | GCA..CGA..  |
| Mmy937 La Rioja     | 7     | .A.AG..TCT  | CTG.TCGTC.    | TCA.TCTA..  | GCA..CGA..  |
| AY027836Mal         | 1     | TCGCAATCTC  | TCAACTACAC    | CTGCCCTCGCC | ATGCCTAGTT  |
| Mal935 La Rioja     | 1     | .....       | .....         | .....       | .....       |
| Mal944 La Rioja     | 1     | .....       | .....         | .....       | .....       |
| Mal947 La Rioja     | 1     | .....       | .....         | .....       | .....       |
| Mal105 La Rioja     | 1     | .....       | .....         | .....       | .....       |
| AY027832Mal Greece  | 2     | .....       | .....         | .....       | .....       |
| AY027851Mbr Germany | 4     | CAAA.CC...  | C...GTC.TCT   | ..CATTCFA.T | ..CATTC..CC |
| AY027844Mau Greece  | 3     | .A.AG..TCT  | CTG.TCGTC.    | TCA.TCTAT.  | GCA..CGA..  |
| AY027848Mmy Germany | 5     | .A.AG..TCT  | CTG.TCGTC.    | TCA.TCTAT.  | GCA..CGA..  |
| Mmy945 La Rioja     | 5     | .A.AG..TCT  | CTG.TCGTC.    | TCA.TCTAT.  | GCA..CGA..  |
| Mmy946 La Rioja     | 5     | .A.AG..TCT  | CTG.TCGTC.    | TCA.TCTAT.  | GCA..CGA..  |
| Mmy936 La Rioja     | 6     | .A.AG..TCT  | CTG.TCGTC.    | TCA.TCTA..  | GCA..CGA..  |
| Mmy937 La Rioja     | 7     | .A.AG..TCT  | CTG.TCGTC.    | TCA.TCTA..  | GCA..CGA..  |
| AY027836Mal         | 1     | 4555555555  | 4444444444    | 4444444444  | 4444444444  |
| Mal935 La Rioja     | 1     | .....       | .....         | .....       | .....       |
| Mal944 La Rioja     | 1     | .....       | .....         | .....       | .....       |
| Mal947 La Rioja     | 1     | .....       | .....         | .....       | .....       |
| Mal105 La Rioja     | 1     | .....       | .....         | .....       | .....       |
| AY027832Mal Greece  | 2     | .....       | .....         | .....       | .....       |
| AY027851Mbr Germany | 4     | .....       | .....         | .....       | .....       |
| AY027844Mau Greece  | 3     | .....       | .....         | .....       | .....       |
| AY027848Mmy Germany | 5     | .....       | .....         | .....       | .....       |
| Mmy945 La Rioja     | 5     | .....       | .....         | .....       | .....       |
| Mmy946 La Rioja     | 5     | .....       | .....         | .....       | .....       |
| Mmy936 La Rioja     | 6     | .....       | .....         | .....       | .....       |
| Mmy937 La Rioja     | 7     | .....       | .....         | .....       | .....       |
| AY027836Mal         | 1     | 4555555555  | 4444444444    | 4444444444  | 4555555555  |
| Mal935 La Rioja     | 1     | .....       | .....         | .....       | .....       |
| Mal944 La Rioja     | 1     | .....       | .....         | .....       | .....       |
| Mal947 La Rioja     | 1     | .....       | .....         | .....       | .....       |
| Mal105 La Rioja     | 1     | .....       | .....         | .....       | .....       |
| AY027832Mal Greece  | 2     | .....       | .....         | .....       | .....       |
| AY027851Mbr Germany | 4     | .....       | .....         | .....       | .....       |
| AY027844Mau Greece  | 3     | .....       | .....         | .....       | .....       |
| AY027848Mmy Germany | 5     | .....       | .....         | .....       | .....       |
| Mmy945 La Rioja     | 5     | .....       | .....         | .....       | .....       |
| Mmy946 La Rioja     | 5     | .....       | .....         | .....       | .....       |
| Mmy936 La Rioja     | 6     | .....       | .....         | .....       | .....       |
| Mmy937 La Rioja     | 7     | .....       | .....         | .....       | .....       |
| AY027836Mal         | 1     | 8360581230  | 7013901793    | 7013901793  | 8360581230  |
| Mal935 La Rioja     | 1     | .....       | .....         | .....       | .....       |
| Mal944 La Rioja     | 1     | .....       | .....         | .....       | .....       |
| Mal947 La Rioja     | 1     | .....       | .....         | .....       | .....       |
| Mal105 La Rioja     | 1     | .....       | .....         | .....       | .....       |
| AY027832Mal Greece  | 2     | .....       | .....         | .....       | .....       |
| AY027851Mbr Germany | 4     | .....       | .....         | .....       | .....       |
| AY027844Mau Greece  | 3     | .....       | .....         | .....       | .....       |
| AY027848Mmy Germany | 5     | .....       | .....         | .....       | .....       |
| Mmy945 La Rioja     | 5     | .....       | .....         | .....       | .....       |
| Mmy946 La Rioja     | 5     | .....       | .....         | .....       | .....       |
| Mmy936 La Rioja     | 6     | .....       | .....         | .....       | .....       |
| Mmy937 La Rioja     | 7     | .....       | .....         | .....       | .....       |
| AY027836Mal         | 1     | CTTCTTTTCT  | CTTCTTTTCT    | CTTCTTTTCT  | CTTCTTTTCT  |
| Mal935 La Rioja     | 1     | .....       | .....         | .....       | .....       |
| Mal944 La Rioja     | 1     | .....       | .....         | .....       | .....       |
| Mal947 La Rioja     | 1     | .....       | .....         | .....       | .....       |
| Mal105 La Rioja     | 1     | .....       | .....         | .....       | .....       |
| AY027832Mal Greece  | 2     | .....       | .....         | .....       | .....       |
| AY027851Mbr Germany | 4     | .....       | .....         | .....       | .....       |
| AY027844Mau Greece  | 3     | .....       | .....         | .....       | .....       |
| AY027848Mmy Germany | 5     | .....       | .....         | .....       | .....       |
| Mmy945 La Rioja     | 5     | .....       | .....         | .....       | .....       |
| Mmy946 La Rioja     | 5     | .....       | .....         | .....       | .....       |
| Mmy936 La Rioja     | 6     | .....       | .....         | .....       | .....       |
| Mmy937 La Rioja     | 7     | .....       | .....         | .....       | .....       |

TABLE 2. Differences among haplotypes within the *mystacinus* group across a 561 bp fragment of the mitochondrial ND1 gene. The lower half of the matrix shows absolute differences and the upper half corrected distances according to a HKY85 model. Haplotypes information in Table 1

| Haplotype | 1 (Mal) | 2 (Mal) | 3 (Mau) | 4 (Mbr) | 5 (Mmy) | 6 (Mmy) | 7 (Mmy) |
|-----------|---------|---------|---------|---------|---------|---------|---------|
| 1 (Mal)   | –       | 0.011   | 0.172   | 0.165   | 0.167   | 0.162   | 0.164   |
| 2 (Mal)   | 6       | –       | 0.165   | 0.171   | 0.159   | 0.155   | 0.157   |
| 3 (Mau)   | 82      | 79      | –       | 0.167   | 0.007   | 0.007   | 0.008   |
| 4 (Mbr)   | 80      | 82      | 80      | –       | 0.176   | 0.171   | 0.174   |
| 5 (Mmy)   | 80      | 77      | 4       | 84      | –       | 0.004   | 0.002   |
| 6 (Mmy)   | 78      | 75      | 4       | 82      | 2       | –       | 0.002   |
| 7 (Mmy)   | 79      | 76      | 5       | 83      | 1       | 1       | –       |

The other lineage (Mal 135, 935, 944 and 947) shared a unique haplotype that showed a genetic distance at the ND1 fragment of 0 and 1.1% with respect to the two known sequences of *M. alcaethoe* (Tables 1 and 2).

*Morphological Comparisons*

The specimens identified as belonging to the *M. mystacinus/aurascens* lineage, showed dorsally greyish pelage, a relatively broad and rather rounded base of the upper canine. This dental character can tell the two species apart (P. Benda, in litt.). The four specimens identified as belonging to the *M. alcaethoe* lineage were clearly smaller in external measurements than the six

*M. mystacinus/aurascens* for both males (Table 3) and females (forearm length 32.5 and 33.0 mm in *M. alcaethoe* and 34.9 mm in *M. mystacinus/aurascens*). Two specimens identified as *M. alcaethoe* showed reddish brown dorsal colouration but the other two *M. alcaethoe* showed a greyish coloration similar to the *M. mystacinus/aurascens* specimens. Both greyish and reddish brown forms of *M. alcaethoe* had pinkish on the internal basal surface of the ears and hairless areas of the face. The only skull of the Spanish *M. alcaethoe* showed the anterior upper premolar located slightly off the tooth-row and displaced internally, which also showed a more developed cingulum cusp than the specimens of the *M. mystacinus/aurascens* lineage.

TABLE 3. Measurements ( $\bar{x} \pm 1SD$ ) of male specimens of *M. alcaethoe* and *M. mystacinus* from La Rioja (Spain) and conspecifics from Greece (Helversen *et al.*, 2001) and Slovakia (Benda *et al.*, 2003). Acronyms: FA: forearm length; GSL: greatest skull length; CBL: condylobasal length; ZW: zygomatic width; M<sup>3</sup>–M<sup>3</sup>: width across upper third molars; C–M<sup>3</sup>: length of the upper tooth-row; BCW: braincase width; ML: mandibular length; C–M<sub>3</sub>: length of the lower tooth-row. Sample size in parentheses

| Trait                          | <i>M. alcaethoe</i> |           |           | <i>M. mystacinus</i> |
|--------------------------------|---------------------|-----------|-----------|----------------------|
|                                | La Rioja            | Greece    | Slovakia  | La Rioja             |
| FA                             | 31.5 (2)            | 30.5 (1)  | 31.3 (1)  | 33.5 ± 0.19 (6)      |
| GSL                            | 12.9 (1)            | –         | 12.66 (1) | 13.25 (2)            |
| CBL                            | 12.3 (1)            | 11.70 (1) | 12.20 (1) | 13.15 (2)            |
| ZW                             | 8.0 (1)             | 7.87 (1)  | 8.22 (1)  | 8.30 (1)             |
| M <sup>3</sup> –M <sup>3</sup> | 5.0 (1)             | 5.15 (1)  | 5.26 (1)  | 5.20 (1)             |
| C–M <sup>3</sup>               | 4.8 (1)             | 4.80 (1)  | 4.90 (1)  | 5.35 (2)             |
| BCW                            | 6.4 (1)             | –         | 6.41 (1)  | 6.75 (2)             |
| ML                             | 9.2 (1)             | 9.01 (1)  | 9.31 (1)  | 9.85 (2)             |
| C–M <sub>3</sub>               | 5.2 (1)             | 5.05 (1)  | 5.20 (1)  | 5.70 (2)             |

### Habitat Preference

Despite intensive sampling, bats of the *mystacinus* group were found only in the humid mountain areas of the southwest of La Rioja (Fig. 1). Both species were captured in the same habitats and in fact, *M. mystacinus/aurascens* was also caught in two out of the three known localities for *M. alcaethoe* (Fig. 1). Altitudinal range varied from 790 to 1,390 m for *M. alcaethoe* and from 914 to 1,460 m for *M. mystacinus/M. aurascens*. Both species were captured at the entrance of the same nocturnal refugia, and also using the same beech forest. Besides, *M. alcaethoe* was captured in riparian forest and *M. mystacinus* in Scottish pine forests.

### DISCUSSION

The presence of two lineages in the mitochondrial DNA sequences of ten bats from La Rioja indicates that two species within the *mystacinus* group coexist in the Iberian Peninsula. The shape of the base of the upper canine identifies provisionally the specimens within the *M. mystacinus/M. aurascens* lineage as *M. mystacinus*. This was expected based on the exclusively eastern distribution of *M. aurascens* in Europe (Benda and Tsytulina, 2000). It was not possible to distinguish between *M. aurascens* and *M. mystacinus* using mitochondrial DNA (Mayer and Helversen, 2001). Interestingly, Castella *et al.* (2000) and Mayer and Helversen (2001) found also that mtDNA was unreliable to distinguish *M. myotis* and *M. blythii*. The location of the NORs in the chromosomes remains the main diagnostic characteristic between the former two species and needs to be checked for the Iberian *M. mystacinus*.

The other lineage is identified, without doubt, as *M. alcaethoe* since all specimens within the lineage showed the molecular

diagnostic sequence of this species (Helversen *et al.*, 2001; Ruedi *et al.*, 2002; Benda *et al.*, 2003). The genetic distance found between the two species in Iberia (> 16%) is similar to the value reported for this mitochondrial fragment in other studies (Helversen *et al.*, 2001). The two lineages seem to show important differences in the level of polymorphism in Iberia. The four *M. alcaethoe* from Iberia are monomorphic and share a unique haplotype with individuals from Hungary (Helversen *et al.*, 2001) and with two other known specimens from France (Ruedi *et al.*, 2002) and Slovakia (Benda *et al.*, 2003). However three haplotypes are found in the orthologous fragment of ND1 in the four Iberian *M. mystacinus*. One of these haplotypes is shared with specimens from Germany and was already reported for Spain by Helversen *et al.* (2001). Although more detailed analyses with larger sample sizes are needed, this high polymorphism could possibly indicate a role of the Iberian Peninsula as a quaternary refugium for *M. mystacinus* in recent cold periods, as is well known for other species (Hewitt, 1999; Ruedi and Castella, 2003).

At morphological level, the Iberian *M. alcaethoe* show two colour phases: the dorsal reddish brown pelage, similar to other European co-specifics and a greyer one, similar to *M. mystacinus* and already reported in the original description of the species by Helversen *et al.* (2001). The range of colour variation within *M. alcaethoe* is still unknown but the lack of pigmentation along the base of the ears and hairless areas of the face in *M. alcaethoe* appears to be a way of distinguishing between both species from external examination. The length of the forearm also seems to be a useful characteristic for distinguishing *M. alcaethoe* from *M. mystacinus*, since their measurements, at least in Iberia, do not overlap (31.4–33.0 mm in *M. alcaethoe* and 33.0–34.9 mm in *M. mystacinus*). The validity of other proposed

diagnostic characters such as relative length of tragus, length of hind feet, length of thumb and length of the claw (Helvesen *et al.*, 2001; Ruedi *et al.*, 2002; Benda *et al.*, 2003) could not be checked since most of the bats were captured before the new species was identified.

Helvesen *et al.* (2001) suggest habitat segregation between these species in Greece; *M. alcaethoe* was more commonly found in small valleys with closed vegetation, whereas *M. mystacinus* was found in forests surrounding large water bodies. Besides, Ruedi *et al.* (2002) found *M. alcaethoe* hunting far away from any watercourse. In La Rioja the habitat preferences of both species were found to strongly overlap. In fact, the two species were captured twice in the same net the same night. Sibling species found in Europe are generally sympatric and/or syntopic across large areas (e.g., *Pipistrellus pipistrellus*/*P. pygmaeus*, *M. myotis*/*M. blythii*, etc.). In Iberia, the *M. mystacinus*/*M. alcaethoe* species-complex complies with this pattern even for a very restricted distribution area. In fact, the *mystacinus* group is only found in Iberia in the humid mountain ranges in the northern regions (Fig. 1). *M. mystacinus* sensu stricto is also currently known, apart from La Rioja, from the western mountains of the 'Sistema Central' range. In fact, individuals from these mountains (Gredos and Linares de Riofrío) were molecularly identified by Helvesen *et al.* (2001). Nevertheless, more intensive studies are necessary to revise the distribution of this species group in Iberia and to define the ecological requirements of the two species.

The description of *M. alcaethoe* and its finding in Iberia, might challenge the subspecies *M. m. occidentalis* described for Iberia by Benda and Tsytsulyna (2000) in their revision of the *mystacinus* group. The little genetic differentiation found between the Iberian and German haplotypes of

*M. mystacinus* does not give support to such subspecific recognition of the Iberian population. Nevertheless, as discussed above, mitochondrial DNA seems of little use in clarifying systematics and evolutionary relationships within the *M. mystacinus*/*M. aurascens* lineage. The measurements of the molecular identified *M. mystacinus* from La Rioja are still larger than the values reported for typical specimens from Central Europe (Benda *et al.*, 2003) and this would support the differentiation suggested by Benda *et al.* (2003). Given the overlapping distributions of *M. mystacinus* and *M. alcaethoe*, more detailed studies on the genetic and morphological variation patterns across the Western Palaearctic seem necessary to reconstruct the systematics and evolutionary relationships within the group.

The finding of *M. alcaethoe* in La Rioja implies the first record of the species for the Iberian Peninsula. This record, along with the recent additions of *Pipistrellus pygmaeus* and *Plecotus macrobullaris* (Barratt *et al.*, 1997; Garin *et al.*, 2003), gives a total of 28 bat species for the Iberian Peninsula. It also extends the southwestern limit of the range of *M. alcaethoe*, a species that was originally proposed as endemic to the Balkans (Helvesen *et al.*, 2001). Finally, this, and the recent records from France and Slovakia, indicate that this species is far more widely spread across southern Europe than previously thought, although most probably in a discontinuous distribution.

#### ACKNOWLEDGEMENTS

The institutional Gobierno de La Rioja provided collecting permits. Petr Benda kindly provided useful information and Jane Orr kindly revised the English. This research is part of the projects REN2000-1639 and REN2002-01372 /GLO funded by the Dirección General de Investigación of the Spanish Ministry of Science and Technology.



## LITERATURE CITED

- AGIRRE-MENDI, P. T. 2002. *Myotis mystacina* (Kuhl, 1817), Murciélago ratonero bigotudo. Pp. 162–165, in Atlas de los Mamíferos terrestres de España (L. J. PALOMO and J. GISBERT, eds.). Dirección General de la Conservación de la Naturaleza, SECEM-SECEMU, Madrid, 564 pp.
- AIHARTZA, J. R. 2001. Quirópteros de Araba, Bizkaia y Guipuzkoa: distribución, ecología y conservación. Ph.D. Thesis, Basque Country University, Leioa, 336 pp.
- BARRATT, E. M., R. DEAVILLE, T. M. BURLAND, M. W. BRUFORD, G. JONES, P. A. RACEY, and R. K. WAYNE. 1997. DNA answers the call of pipistrelle bats species. *Nature*, 387: 138–139.
- BENDA, P., and K. A. TSYLSULINA. 2000. Taxonomic revision of *Myotis mystacinus* group (Mammalia: Chiroptera) in the western Palearctic. *Acta Societas Zoologica Bohemica*, 64: 331–398.
- BENDA, P., M. RUEDI, and M. UHRIN. 2003. First record of *Myotis alcaethoe* (Chiroptera: Vespertilionidae) in Slovakia. *Folia Zoologica*, 52: 359–365.
- CASTELLA, V., M. RUEDI, L. EXCOFFIER, C. IBÁÑEZ, R. ARLETTAZ, and J. HAUSSER. 2000. Is the Gibraltar Strait a barrier to gene flow for the bat *Myotis myotis* (Chiroptera: Vespertilionidae)? *Molecular Ecology*, 9: 1761–1772.
- FERNÁNDEZ, R., and C. IBÁÑEZ. 1989. Patterns of distribution of bats in the Iberian peninsula. Pp. 357–361, in European bat research 1987 (V. HANÁK, I. HORÁČEK and J. GAISLER, eds.). Charles University Press, Praha, 718 pp.
- FERNÁNDEZ, R., L. LOPO, and R. RODRÍGUEZ. 1989. Mapa forestal de La Rioja. Estudios Monográficos Instituto de Estudios Riojanos, 18: 1–72.
- FERNÁNDEZ-GUTIÉRREZ, J. 2002. Los murciélagos en Castilla y León. Atlas de distribución y tamaño de poblaciones. Junta de Castilla y León, Consejería de Medio Ambiente, Valladolid, 343 pp.
- GARIN, I., J. L. GARCÍA-MUDARRA, J. R. AIHARTZA, U. GOITI, and J. JUSTE. 2003. Presence of *Plecotus macbullaris* (Chiroptera: Vespertilionidae) in the Pyrenees. *Acta Chiropterologica*, 5: 243–250.
- GERELL, R. 1999a. *Myotis brandtii* (Eversmann, 1845). Pp. 104–105, in The atlas of European mammals (A. J. MITCHELL-JONES, G. AMORI, W. BOGDANOWICZ, B. KRYŠTUFEK, P. J. H. REIJNDERS, F. SPITZENBERGER, M. STUBBE, J. B. M. THISSEN, V. VOHRALÍK and J. ZIMA, eds.). Academic Press, London, xi + 484 pp.
- GERELL, R. 1999b. *Myotis mystacinus* (Kuhl, 1817). Pp. 116–117, in The atlas of European mammals (A. J. MITCHELL-JONES, G. AMORI, W. BOGDANOWICZ, B. KRYŠTUFEK, P. J. H. REIJNDERS, F. SPITZENBERGER, M. STUBBE, J. B. M. THISSEN, V. VOHRALÍK and J. ZIMA, eds.). Academic Press, London, xi + 484 pp.
- HANÁK, V. 1965. Zur Systematic der Bartfledermaus *Myotis mystacinus* Kuhl, 1819 und über das Vorkommen von *Myotis ikonikovi* Ognev, 1912 in Europa. *Věstník Československé Společnosti Zoologické*, 29: 353–367.
- HELVERSEN, O. VON, K.-G. HELLER, F. MAYER, A. NEMETH, M. VOLLETH, and P. GOMBKÖTÖ. 2001. Cryptic mammalian species: a new species of whiskered bat (*Myotis alcaethoe* n. sp.) in Europe. *Naturwissenschaften*, 88: 217–223.
- HEWITT, G. M. 1999. Post-glacial re-colonization of European biota. *Biological Journal of the Linnean Society*, 68: 87–112.
- HIGUCHI, R., C. H. VON BEROLDINGEN, G. F. SENSA-BAUGH, and H. A. ERLICH. 1988. DNA typing from hairs. *Nature*, 332: 543–546.
- IBÁÑEZ, C., A. GUILLÉN, R. FERNÁNDEZ, J. L. PÉREZ, and S. I. GUERRERO. 1992. Iberian distribution of some little known bat species. *Mammalia*, 56: 433–444.
- JUSTE, J., C. IBÁÑEZ, J. MUÑOZ, D. TRUJILLO, P. BENDA, A. KARATAŞ and M. RUEDI. 2004. Mitochondrial phylogeography of the long-eared bats (*Plecotus*) in the Mediterranean Palearctic and Atlantic Islands. *Molecular Phylogenetics and Evolution*, 31: 1114–1126.
- KIEFER, A., F. MAYER, J. KOSUCH, O. VON HELVERSEN, and M. VEITH. 2002. Conflicting molecular phylogenies of European long-eared bats (*Plecotus*) can be explained by cryptic diversity. *Molecular Phylogenetics and Evolution*, 25: 557–566.
- KUMAR, S., K. TAMURA, I. B. JAKOBSEN, and M. NEI. 2001. MEGA2: Molecular evolutionary genetics analysis software. Arizona State University, Tempe, Arizona, USA.
- MANIATIS, T., E. F. FRITSH, and J. SAMBROOK. 1989. *Molecular cloning: a laboratory manual*, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, USA.
- MARTÍNEZ ABAIGAR, J., E. NÚÑEZ OLIVERA, and R. T. LAS HERAS. 1994. *Agroclimatología de La Rioja*. Biblioteca de Temas Riojanos, IER Logroño, Spain, 155 pp.
- MAYER, F., and O. VON HELVERSEN. 2001. Cryptic diversity in European bats. *Proceedings of the Royal Society of London B*, 268: 1825–1832.
- POSADA, D., and K. A. CRANDALL. 1998. MODELTEST: testing the model of DNA substitution. *Bioinformatics*, 14: 817–818.
- RUEDI, M., and V. CASTELLA. 2003. Genetic

consequences of the ice ages on nurseries of the bat *Myotis myotis*: a mitochondrial and nuclear survey. *Molecular Ecology*, 12: 1527–1540.

RUEDI, M., and F. MAYER. 2001. Molecular systematics of bats of the genus *Myotis* (Vespertilionidae) suggests deterministic ecomorphological convergences. *Molecular Phylogenetics and Evolution*, 21: 436–448.

RUEDI, M., P. JOURDE, P. GIOSA, M. BARATAUD, and S. Y. ROUÉ. 2002. DNA reveals the existence of *Myotis alcaethoe* in France (Chiroptera: Vespertilionidae). *Revue suisse de Zoologie*, 109: 643–652.

SWOFFORD, D. L. 2000. PAUP\*. Phylogenetic analysis using parsimony. Version 4.0b10. Sinauer Associates, Sunderland, Massachusetts, USA.

VOLLETH, M. 1987. Differences in the location of nucleolus organizer regions in European vespertilionid bats. *Cytogenetics and Cell Genetics*, 44: 186–197.

WORTHINGTON WILMER, J., and E. BARRATT. 1996. A non-lethal method of tissue sampling for genetics studies of chiropterans. *Bat Research News*, 37: 1–3.

WOUTERSEN, K., and J. J. BAFALUY-ZORIGUEL. 2001. Murciélagos del Alto Aragón. K. Woutersen Publicaciones, Huesca, Spain, 144 pp.

Received 14 March 2004, accepted 06 May 2004

APPENDIX

Bats of the *mystacinus* group from La Rioja (Spain) used in this study. The specimens that could not be molecularly identified are referred as *M. mystacinus* s.l. Haplotypes AY027835 and AY027847 are from Helversen *et al.* (2001)

| Taxon                  | Locality (date of capture or number of individuals) | Accession No. | DNA ID  | Voucher  |
|------------------------|---|---------------|---------|----------|
| <i>alcaethoe</i>       | El Rasillo (07.08.87)                               | AY027835      | Mal 105 | EBD16000 |
| <i>alcaethoe</i>       | Anguiano (01.08.03)                                 | AY027835      | Mal 947 | Biopsy   |
| <i>alcaethoe</i>       | Ojacastro (19.06.01)                                | AY027835      | Mal 935 | Biopsy   |
| <i>alcaethoe</i>       | Ojacastro (29.07.03)                                | AY027835      | Mal 944 | Biopsy   |
| <i>mystacinus</i>      | El Rasillo (07.08.87)                               | AY552330      | Mmy 936 | EBD16001 |
| <i>mystacinus</i>      | El Rasillo (24.08.88)                               | AY552331      | Mmy 937 | EBD17801 |
| <i>mystacinus</i>      | El Rasillo (22.08.88)                               | –             | Mmy 104 | EBD18289 |
| <i>mystacinus</i>      | El Rasillo (02.09.88)                               | –             | Mmy 968 | EBD18290 |
| <i>mystacinus</i>      | Anguiano (01.08.03)                                 | AY027847      | Mmy 945 | Biopsy   |
| <i>mystacinus</i>      | Anguiano (01.08.03)                                 | AY027847      | Mmy 946 | Biopsy   |
| <i>mystacinus</i> s.l. | Anguiano (1)  | –             | –       | –        |
| <i>mystacinus</i> s.l. | Castroviejo (1)                                     | –             | –       | –        |
| <i>mystacinus</i> s.l. | Ezcaray (3)   | –             | –       | –        |
| <i>mystacinus</i> s.l. | Lumbreras (1)                                       | –             | –       | –        |
| <i>mystacinus</i> s.l. | Ortigosa de C. (2)                                  | –             | –       | –        |
| <i>mystacinus</i> s.l. | Villanueva de C. (1)                                | –             | –       | –        |
| <i>mystacinus</i> s.l. | Villavelayo (5)                                     | –             | –       | –        |
| <i>mystacinus</i> s.l. | Villoslada de C. (2)                                | –             | –       | –        |
| <i>mystacinus</i> s.l. | Zarzosa (1)   | –             | –       | –        |