

The First Record of Ingestion and Inhalation of Micro- and Mesoplastics by Neotropical Bats from the Brazilian Amazon

Authors: Correia, Letícia L., Ribeiro-Brasil, Danielle R. G., Garcia, Magali G., Silva, Daniela de Melo e, Alencastre-Santos, Ana B., et al.

Source: *Acta Chiropterologica*, 25(2) : 371-383

Published By: Museum and Institute of Zoology, Polish Academy of Sciences

URL: <https://doi.org/10.3161/15081109ACC2023.25.2.015>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

The first record of ingestion and inhalation of micro- and mesoplastics by Neotropical bats from the Brazilian Amazon

LETÍCIA L. CORREIA¹, DANIELLE R. G. RIBEIRO-BRASIL², MAGALI G. GARCIA³,
DANIELA DE MELO E SILVA⁴, ANA B. ALENCASTRE-SANTOS¹, and THIAGO B. VIEIRA^{1, 3, 5}

¹Laboratório de Ecologia, Faculdade de Ciências Biológicas, Universidade Federal do Pará, Campus Altamira, Altamira, 68371-155, Brazil

²Laboratório de Ecologia e Conservação de Ecossistemas Aquáticos (LECEA), Universidade Federal de Mato Grosso, Campus I Araguaia, Pontal do Araguaia, 78698-000, Brazil

³Laboratório de Microbiologia, Universidade Federal do Pará, Altamira, 68371-155, Brazil

⁴Laboratory of Mutagenesis, Institute of Biological Sciences, ICB I, Federal University of Goiás, Campus Samambaia, Goiânia, Goiás, 74690-900, Brazil

⁵Corresponding author: E-mail: vieiratb@ufpa.br

This study shows the abundance of contamination by microplastics (MPs) and the first record of contamination by MPs in bats. Additionally, we tried to understand the mechanism of the environmental contamination of bats. Therefore, the digestive and respiratory tracts from 81 adult bats belonging to 25 species were extracted for analysis. Bats were captured in different locations in the Brazilian Amazon (Altamira, Bragança, Brasil Novo, Medicilândia, Nova Timboteua, Placas, São Félix do Xingu, Uruará and Vitória do Xingu, all in the state of Pará). The results showed that all species were contaminated with MPs in at least one of the analyzed systems. For the digestive system, the form of contamination occurs through bioaccumulation and biomagnification by the ingestion of contaminated food or water. In the case of the respiratory system, contamination occurs through the inhalation of MPs suspended in the atmospheric air. The different foraging characteristics of bats, the type of capture strategy for this food, and the type of habitat reinforce the idea that plastic contaminants are present in all environments.

Key words: plastic waste, flying mammals, feeding, breathing

Este estudio muestra la abundancia de contaminación por microplásticos (MP) y el primer registro de contaminación por PM en murciélagos. Además, intentamos comprender el mecanismo de contaminación ambiental de los murciélagos. Por lo tanto, se extrajeron para su análisis los tractos digestivo y respiratorio de 81 murciélagos adultos de 25 especies. Los murciélagos fueron capturados en diferentes localidades de la Amazonía brasileña (Altamira, Bragança, Brasil Novo, Medicilândia, Nova Timboteua, Placas, São Félix do Xingu, Uruará y Vitória do Xingu, todos en el estado de Pará). Los resultados mostraron que todas las especies estaban contaminadas con PM en al menos uno de los sistemas analizados. Para el sistema digestivo, la forma de contaminación se produce por bioacumulación y biomagnificación por la ingestión de alimentos o agua contaminados. Mientras que para el sistema respiratorio, la contaminación se produce por la inhalación de PM suspendidas en el aire atmosférico. Las diferentes características de alimentación de los murciélagos, el tipo de estrategia de captura de este alimento y el tipo de hábitat refuerzan la idea de que los contaminantes plásticos están presentes en todos los entornos.

Palabras clave: residuos plásticos, mamíferos voladores, alimentación, respiración

INTRODUCTION

Plastics are polymers derived from petroleum and of anthropogenic origin, considered contaminants of emerging concern, and have gained global attention due to their abundance, durability, and persistence (Thompson *et al.*, 2009; Anderson *et al.*, 2016; Wilson *et al.*, 2021). The properties of

plastics, such as low cost and versatility, make their use widespread in society, being used in the domestic, automotive, and textile industries until they become a serious environmental problem due to improper disposal (Napper and Thompson, 2020). Plastic that is no longer useful enters the environment as plastic waste that can be divided into macroplastics (MCPs) and microplastics (MPs)

(Steensgaard *et al.*, 2017). Macroplastics comprise macro (> 25 cm) and mesoplastics (5 mm to 25 cm) sizes, while microplastics comprise micro (1 to 5 mm) and nano (< 1 mm) sizes (Wagner *et al.*, 2014). MPs can be of primary origin, produced in micro size, or secondary due to the degradation product of larger pieces (Wright *et al.*, 2013; Boucher and Friot, 2017). In terms of shape, particles can be spherical, like pellets, have irregular shapes like fragments and films, or elongated and thin like fibers (McCormick *et al.*, 2016). Plastics have already been found in fish (Ribeiro-Brasil *et al.*, 2020), birds (Tokunaga *et al.*, 2023), bottled water (Li *et al.*, 2023) and human breast milk (Ragusa *et al.*, 2022). In the Americas, studies on this topic are still scarce (Ayala *et al.*, 2023).

The degradation of plastic textile fibers produces, for example, microplastics called microfibrils; this degradation product has been observed in atmospheric precipitation, becoming breathable microfibrils (Gasperi *et al.*, 2018). This fact suggests potential exposure of microfibrils to organisms that present pulmonary respiration, such as humans and bats (Zhang *et al.*, 2020). Microfibrils can also be deposited on the surfaces of fruits and terrestrial organisms (Rillig *et al.*, 2017). In this way, bats can absorb food and inhale aerosolized microfibrils, allowing contamination by ingestion and inhalation, respectively.

MPs are the plastic waste most commonly found in the environment and of the most significant concern, as they are considered the easiest to spread and the most assimilated by organisms (Dris *et al.*, 2016; Duis and Coors, 2016; Horton *et al.*, 2017; Windsor *et al.*, 2019; Dahms *et al.*, 2020; He *et al.*, 2020; Miller *et al.*, 2020; Akhbarizadeh *et al.*, 2021; Baho *et al.*, 2021; Kumari *et al.*, 2022). The effects of MP absorption are not yet fully elucidated, but there is already knowledge about ingestion by various organisms, ranging from aquatic (Ribeiro-Brasil *et al.*, 2020; Jawad *et al.*, 2021) to terrestrial environments (Lahive *et al.*, 2019).

Bats exhibit a significant morphological and behavioral diversification, feeding on fruits, nectar, blood, insects, and vertebrates (Kalko, 1998). They play critical ecological roles in the ecosystem, such as pollination, seed dispersal, and insect control, including agricultural pest control (Fenton *et al.*, 1999; Estrada and Coates-Estrada, 2002; Castro-Luna and Galindo-González, 2012; Kasso and Balakrishnan, 2013; Rodríguez-San Pedro *et al.*, 2020; Aguiar *et al.*, 2021; de Jong *et al.*, 2021; Surtipito, 2021). In addition, some species of pioneer plants,

necessary for regeneration and ecological succession of degraded areas, are pollinated and dispersed exclusively by bats (Passos and Passamani, 2003). Because of the important services they provide, bats are considered key species in tropical forests (Fleming and Heithaus, 1981).

Studies on the ingestion or inhalation of microplastics by bats do not exist, and in situ observations of MPs contamination have only been reported in other organisms, such as marine organisms (Miller *et al.*, 2020), land plants (Kumari *et al.*, 2022), and freshwater fish (Ribeiro-Brasil *et al.*, 2022). Plastic residues have already been found in fish from farms in Rondônia (Dantas Filho *et al.*, 2023), and in other streams in the municipalities of Pará, such as Barcarena, Ipixuna do Pará, Concórdia do Pará, and Tomé-Açu (Ribeiro-Brasil *et al.*, 2020), and in the Xingu River (Andrade *et al.*, 2019) that passes through the city of Altamira. Moreover, the contamination of MPs in human organs has been demonstrated. This contamination is possibly through ingestion and inhalation; it is likely that bats are absorbing plastic waste from the environment, either directly, by inhaling particles from the air (Pauly *et al.*, 1998; Gasperi *et al.*, 2018), or indirectly, by ingesting contaminated food (Gross, 2015; da Costa Araújo and Malafaia, 2021). Bats, like humans, because both have similar respiratory systems, may be susceptible to similar contamination. Thus, the objective of this study was to identify how the ingestion and inhalation of microplastics by bats in the eastern Brazilian Amazon occurs and to confirm the presence of plastic waste through visual analysis and inspection of materials collected from bats.

MATERIALS AND METHODS

Sample Collection and Species Studied

The samplings were carried out at 26 points, between 2017 and 2021, four urban points and 22 rural point, all located in the following municipalities: Altamira, Brasil Novo, Placas, Nova Timboteua, Bragança, São Félix do Xingu, Uruará, Vitória do Xingu and Medicilândia, all located in the state of Pará (Fig. 1 and Table 1). Some rural points were carried out in cocoa plantations and/or natural vegetation, the urban points were collected within cities or in nearby places. The region has a tropical climate of type Am, according to the Köppen climate classification (Peel *et al.*, 2007), with an average temperature of 26.1°C and an average annual rainfall of 2,000 millimeters.

Bats were sampled using ten mist nets (9 × 2.5 m), open at sunset and remaining for six hours, inspected every half hour. The bats were placed in 100% cotton fabric bags and taken to the Laboratory of Ecology of Altamira — LABECO at the Federal University of Pará, Altamira campus. Afterwards,




FIG. 1. Geographical location of the collection points where bats were sampled from 2017 to 2021 by ChiroXingu: Center for Studies in Ecology and Conservation of Bats

TABLE 1. Geographical coordinates of the points and locations where bats were sampled during the years 2017 to 2021 by ChiroXingu

Point	Municipality	Geographical coordinates (decimal degrees)		Urban	Rural
		Latitude (W)	Longitude (S)		
P1	Altamira	-52.253990	-3.156140	X	
P2		-52.192110	-3.198030	X	
P3		-52.150320	-3.515619		X
P4		-52.287420	-3.341350		X
P5		-52.268830	-3.359210		X
P6		-52.181760	-3.189400		X
P7		-52.274694	-3.322750		X
P8		-52.235370	-3.117135		X
P9		-52.618720	-5.775530		X
P10		-52.202389	-3.199694	X	
P11	-52.212472	-3.210500	X		
P12	-52.070760	-3.192790		X	
P13	-52.098180	-3.151360		X	
P14	Bragança	-46.738750	-1.080000		X
P15	Brasil Novo	-52.274667	-3.322778		X
P16	Medicilândia	-53.082780	-3.583540		X
P17		-53.045560	-3.508920		X
P18		-53.262880	-3.488000		X
P19		-53.045560	-3.508920		X
P20	Nova Timboteua	-47.368556	-1.078889		X
P21	Placas	-54.355590	-3.798080		X
P22	Uruará	-53.258080	-3.494170		X
P23	Vitória do Xingu	-51.940028	-3.375472		X
P24		-51.928670	-3.352950		X
P25		-51.851060	-3.211300		X
P26		-51.505997	-3.123956		X

collected individuals were euthanized by cervical dislocation, and morphometric data (total length, foot length, ear length, tragus length, forearm length, and body mass) were measured. Subsequently, bats were fixed with 10% formaldehyde and stored in glass jars with 70% alcohol in the ChiroXingu Bat Collection: Center for studies in ecology and conservation of bats. The ChiroXingu research group collected the bats from April to August 2017, September and October 2018, January to June 2020 and July 2021. Each sampling, transport and preservation of the sampled specimens were carried out in accordance with the relevant guidelines and regulations of the Sistema de Autorização e Informação em Biodiversidade, Instituto Chico Mendes de Conservação da Biodiversidade, Ministerio do Meio Ambiente (license No. 57294-2 granted by the last author). All methods were performed according to ARRIVE guidelines as study design, sample size, statistical methods, and experimental animals. However, the protocols referring to the following works were also followed to avoid contamination of the sample in the laboratory (Nuelle *et al.*, 2014; Wagner *et al.*, 2014; Devriese *et al.*, 2015; Ribeiro-Brasil *et al.*, 2020).

Material Analysis

Extraction of the biological tissues

Organs of the digestive and respiratory systems were removed completely from each fluid-preserved specimen. The digestive system was removed from esophagus to anus, and respiratory system from the trachea to lungs. The entire process was carried out inside a laminar flow hood to avoid contamination of the samples.

Digestion of the biological tissues

The samples were put into sanitized glass vials containing potassium hydroxide (KOH; 10%, V/V) to dissolve the tissues (Ghosal *et al.*, 2018). They were placed in an oven with a temperature of 60°C for seven days with modified protocol by (Lavoy and Crossman, 2021) to accelerate the sample digestion process. After tissue digestion, samples were filtered through a 0.2 µm porosity membrane with a vacuum pump. Membranes were stored in Petri dishes, protected by aluminum foil envelopes, and returned to the oven for 24 h at 60°C for drying the membranes. Aluminum was used to avoid contamination of samples in the work environment.

Visual analysis of plastic waste

The samples were analyzed under a stereo microscope with a magnification of 100 times (Digilab-microscope Stereo Trinocular DI-106T zoom). The membranes were scanned from left to right, top to bottom. Each item was photographed and identified and placed into two categories: microplastics (MPs) length range from 1 to 5 mm) and mesoplastics (MSPs) range from 5 mm to 25 cm — Wagner *et al.*, 2014).

Quality assurance and quality control (QA/QC)

For material classification, the following criteria were followed: a) residues considered as fibers that had a structure like animal joints were disregarded; b) only plastic waste that had the same pattern from one end to the other was considered; c) confirmation of plastic for smaller particles was done through the hot needle test (Devriese *et al.*, 2015). The hot needle test is placing the hot needle over the sample, if the sample

changes shape or shrinks, it is because this sample is considered plastic.

All the necessary precautions were taken for laboratory procedures, such as wearing clothes and gowns made of only 100% cotton. All analysis material was previously washed with distilled water and filtered before use. The membranes used in the filtration process were covered with aluminum foil. In addition, we followed an analytical approach for monitoring microplastics in marine sediments where plastics were counted and removed from the samples collected from bats (we counted only the plastic residues that came from the bats, not sediment or other environmental substrates — Nuelle *et al.*, 2014).

Statistical Analysis

A *t*-test for separate variances was performed to compare the amount of plastic waste between respiratory and digestive systems. The analysis was done using the software provided by R Core Team (2021).

RESULTS

We analyzed 81 individuals from 25 species and three families; the most abundant species was *Carollia perspicillata* with nine individuals, while the rarest species, with a single record, were: *Saccopteryx bilineata*, *Lophostoma carrikeri*, *Phyllostomus elongatus* and *Artibeus gnomus* (Table 2). Seventy-eight individuals (96.3%) were contaminated by plastic residues in at least one of the analyzed organs (lung, stomach, and/or intestine).

One hundred fifty-eight samples were analyzed, representing 77 respiratory and 81 digestive systems (Table 2). There was a significant difference between the systems ($t = 4.33$, $d.f. = 98.6$, $P < 0.001$), with the digestive system being more affected (\bar{x} , SD: 4.59, 5.62) than the respiratory system (1.73, 1.90 — Fig. 2). Only the respiratory system of *Pteronotus gymnonotus*, *A. gnomus* and *Sturnira giannae* and the digestive system of *L. carrikeri* were not contaminated by plastic residues (Table 2). All plastic waste found in bats was of the fiber type (Fig. 3).

DISCUSSION

MPs' Paths

This study is the first report of contamination by microplastics (MPs) in bats and expands the list of organisms capable of absorbing MPs. In this way, we confirm that MPs can contaminate bats, and this contamination can be through the airways and digestive tracts. Only fiber-type MPs were found in analyzed organs and investigated systems. The digestive system showed higher contamination.

The ingestion and/or inhalation of plastic waste, whether through the digestive or respiratory route, are two possibilities in which bats, and other taxonomic groups, including humans, are exposed (Pauly *et al.*, 1998; Galloway, 2015; Ragusa *et al.*, 2021).

The forms of contamination by plastic waste can be through the atmospheric air or the ingestion of contaminated food or water (Revel *et al.*, 2018). The presence of many fibers can be explained by the fact that they are the lightest and most easily dispersed (Covernton *et al.*, 2019). Some studies pointed out that plastic debris is commonly found in the oceanic food web and more recently in the terrestrial food web (He *et al.*, 2020; Miller *et al.*, 2020; Baho *et al.*, 2021; Kumari *et al.*, 2022) with wind appearing to be the primary disperser of plastic, catalyzed by rain (Dris *et al.*, 2016; Akhbarizadeh *et al.*, 2021). In terrestrial environments, the main pathway of exposure is atmospheric fallout (Evangelidou *et al.*, 2020) and different agricultural practices, for example, plastic mulching (Büks and Kaupenjohann, 2020; Crossman *et al.*, 2020; Baho *et al.*, 2021).

Fiber-type microplastics are the most abundant in the environment and come mainly from clothes. Fibers are released into the environment when pieces are washed or when they come loose from wear (Hernandez *et al.*, 2017; Liu *et al.*, 2019). Another vital factor in microfiber contamination is face masks and wet wipes, which have seen a considerable increase in consumption and improper disposal during the COVID-19 pandemic (Fadare and Okoffo, 2020; Shruti *et al.*, 2021). Due to its shape, fibers tend to be retained in the lungs and digestive tract. When inhaled and ingested, the fiber-type




FIG. 2. Boxplot of *t*-test showing the difference in the concentration of plastic waste between respiratory and digestive systems

TABLE 2. Bat species collected in the Brazilian Amazon, showing the number of micro- and mesoplastic particles recorded for each species. The total (Σ) is followed by the mean (\bar{x}) \pm standard deviation (SD). N — total number of specimens in the study, N^* — number of analyzed systems

Family/Subfamily	Species	Trophic guild	N	Respiratory system			Digestive system		
				N^*	Micro Σ ($\bar{x} \pm SD$)	Meso Σ ($\bar{x} \pm SD$)	N^*	Micro Σ ($\bar{x} \pm SD$)	Meso Σ ($\bar{x} \pm SD$)
Emballonuridae Gervais, 1856	<i>Rhynchonycteris naso</i> (Wied, 1820)	Insectivorous	3	1	2	0	3	4 (1.33 \pm 1.53)	0
Emballonurinae Gervais, 1856	<i>Saccopteryx bilineata</i> (Temminck, 1838)	Insectivorous	1	1	4	0	1	2	0
	<i>S. leptura</i> (Schreber, 1774)	Insectivorous	4	4	5 (1.25 \pm 1.26)	1 (0.25 \pm 0.50)	4	17 (4.25 \pm 7.18)	1 (0.25 \pm 0.50)
	<i>Peropteryx trinitatis</i> Miller, 1899	Insectivorous	3	3	1 (0.34 \pm 0.58)	0	3	23 (7.67 \pm 12.42)	0
Mormoopidae Saussure, 1860	<i>Pteronotus gymnotus</i> (Wagner, 1843)	Insectivorous	3	3	0	0	3	20 (6.67 \pm 4.93)	0
	<i>P. rubiginosus</i> (Wagner, 1843)	Insectivorous	2	2	2 (1.00 \pm 0.00)	0	2	10 (5.00 \pm 4.24)	0
	<i>P. alitonus</i> Pavan, Bobrowiec e Percequillo, 2018	Insectivorous	5	5	5 (1.00 \pm 0.71)	0	5	5 (1.00 \pm 0.71)	0
Phyllostomidae Gray, 1825	<i>Carollia brevicauda</i> (Schinz, 1821)	Frugivorous	3	3	3 (1.00 \pm 1.00)	0	3	13 (4.34 \pm 3.51)	0
Carollinae Miller, 1924	<i>C. perspicillata</i> (Linnaeus, 1758)	Frugivorous	9	9	35 (3.89 \pm 2.37)	3 (0.33 \pm 0.50)	9	84 (9.34 \pm 7.30)	8 (0.88 \pm 1.16)
Lonchorhinae Gray, 1866	<i>Lonchorhina aurita</i> (Tomes, 1863)	Insectivorous	3	3	3 (1.00 \pm 0.00)	0	3	7 (2.34 \pm 1.53)	0
Phyllostominae Gray, 1825	<i>Lophostoma carrikeri</i> (Allen, 1910)	Insectivorous	1	1	2	0	1	0	0
	<i>L. silvicola</i> (d'Orbigny, 1836)	Insectivorous	5	4	3 (0.75 \pm 0.50)	0	5	09 (1.80 \pm 0.84)	0
	<i>Phyllostomus discolor</i> (Wagner, 1843)	Omnivorous	2	2	4 (2.00 \pm 1.41)	0	2	11 (5.50 \pm 6.37)	1 (0.5 \pm 0.70)
	<i>P. elongatus</i> (E. Geoffroy, 1810)	Omnivorous	1	1	2	0	1	2	0
	<i>P. hastatus</i> (Pallas, 1767)	Omnivorous	3	2	4 (2.00 \pm 1.41)	0	3	1 (0.34 \pm 0.58)	0
	<i>Tonatia maresi</i> Williams, Willig e Reid, 1995	Insectivorous	3	3	8 (2.67 \pm 1.53)	1 (0.33 \pm 0.57)	3	19 (6.34 \pm 6.11)	2 (0.66 \pm 1.15)
Rhinophyllinae Baker <i>et al.</i> , 2016	<i>Rhinophylla fischeriae</i> (Carter, 1966)	Frugivorous	5	5	4 (0.80 \pm 0.48)	0	5	43 (8.60 \pm 8.56)	0
Rhinodermatinae Gervais, 1856	<i>Artibeus fimbriatus</i> Gray, 1838	Frugivorous	4	4	6 (1.50 \pm 1.29)	0	4	19 (4.75 \pm 1.50)	3 (0.75 \pm 1.5)
	<i>A. gnomus</i> Handley, 1987	Frugivorous	1	1	0	0	1	1	0
	<i>A. obscurus</i> (Schinz, 1821)	Frugivorous	5	5	17 (3.40 \pm 3.43)	1 (0.2 \pm 0.44)	5	17 (3.40 \pm 3.58)	2 (0.4 \pm 0.89)
	<i>A. cinereus</i> (Gervais, 1856)	Frugivorous	5	5	9 (1.80 \pm 0.84)	0	5	23 (4.60 \pm 2.89)	4 (0.8 \pm 1.30)
	<i>Sturnira giannae</i> Velazco e Patterson, 2019	Frugivorous	3	3	0	0	3	15 (5.00 \pm 1.00)	0
	<i>S. tildae</i> (de la Torre, 1959)	Frugivorous	2	2	7 (3.50 \pm 2.12)	0	2	15 (7.50 \pm 2.12)	3 (1.5 \pm 0.70)
	<i>Uroderma bilobatum</i> (Peters, 1866)	Frugivorous	3	3	3 (1.00 \pm 1.73)	0	3	9 (3.00 \pm 2.00)	0
	<i>U. maguirostrum</i> (Davis, 1968)	Frugivorous	2	2	1 (0.50 \pm 0.71)	0	2	3 (1.50 \pm 0.71)	0
Sum			81	77			81		

plastic waste is the easiest to release by the digestive system (Suran, 2018; Saborowski *et al.*, 2019).

Plants can intercept MPs carried by winds and rains, where rough surfaces, such as stems, leaves, flowers, and fruits, can absorb microplastics (estimated 0.13 trillion MPs/cm²) (Liu *et al.*, 2020) and internalize them (Yin *et al.*, 2021). Thus, frugivorous species, such as bats of the subfamilies Caroliniae, Rhinophyllinae, and Stenodermatinae, feed on fruits that may be contaminated (Fenton *et al.*, 1999). For example, bats will consume particles that have become trapped in floral exudates and particles that have been internalized. Contamination of the digestive system of fruit bats by MPs suggests bioaccumulation.

The accumulation of plastic contaminants in secondary consumers and at levels above these are well documented in the literature (Dris *et al.*, 2016; Hocking *et al.*, 2017; Horton *et al.*, 2017; de Souza Machado *et al.*, 2018; He *et al.*, 2020; Miller *et al.*, 2020; Kumari *et al.*, 2022), especially for marine environments. For aquatic food chains, evidence of the accumulation of plastic waste was observed in producers, secondary consumers, and even quaternary consumers, thus showing the accumulation and transfer of these contaminants along at least five trophic levels in the marine food chain (Miller *et al.*, 2020).

Despite the growing number of publications in terrestrial environments, no studies described the path of plastic waste from primary producers to tertiary or quaternary consumers (He *et al.*, 2020b). However, the transport of these plastic wastes by the

vascular systems of plants has already been reported (Crossman *et al.*, 2020; Li *et al.*, 2020), leading to the presence and consequent accumulation of these residues in roots, leaves, seeds, and fruits (Dietz and Herth, 2011; Kumari *et al.*, 2022). This plastic waste can be primarily from the atmospheric air (Truong *et al.*, 2021), which is subsequently carried to the soil and bodies of water (Dris *et al.*, 2016; Truong *et al.*, 2021). Soil contamination, and consequent contamination of plant vascular systems, can be intensified by the deposition of plastic waste in river waters and areas bordering water bodies. It is estimated that large rivers can be abundant sources of plastic waste, given the urbanization around rivers, in addition to the dendritic and accumulative nature of the drainage basins (Mani *et al.*, 2015; de Souza Machado *et al.*, 2018).

The transfer of plastic waste between different trophic levels may explain the contamination of insectivorous and omnivorous bats and suggests biomagnification of MPs (Horton *et al.*, 2017; Lusher *et al.*, 2017; de Souza Machado *et al.*, 2018; Miller *et al.*, 2020; Kumari *et al.*, 2022). Biomagnification has been observed for aquatic and terrestrial trophic chains, including small and medium-sized vertebrates and invertebrates, such as annelids and arthropods (Horton *et al.*, 2017; de Souza Machado *et al.*, 2018). Prey contamination can occur in the same way as for plants, through the deposition of residues present in ambient air on the surface of the body. Another route of prey contamination is direct or indirect ingestion of plastic (Al-Jaibachi *et al.*, 2018; Windsor *et al.*, 2019b; Immerschitt and




FIG. 3. Microplastics found in the respiratory and digestive systems of bats. Images A and B refer to mesoplastic fibers. Images C, D, E, and F are microplastic-sized fibers

Martens, 2020). In this way, when preyed upon by bats, they contaminate bats (Dris *et al.*, 2016; Horton *et al.*, 2017; de Souza Machado *et al.*, 2018; Kumari *et al.*, 2022).

MPs have already been found in insects of some orders such as Coleoptera and Diptera (Heinlaan *et al.*, 2020) in addition to Ephemeroptera and Trichoptera (Ziccardi *et al.*, 2016; Lusher *et al.*, 2017) all of which are considered food resources by insectivorous bats. In this case, MPs contamination in insects can be either by the accumulation of plastic residues in their exoskeletons or their external structure (Ehlers *et al.*, 2020) since insects have a body surface with bristles that can serve as a substrate for adhesion of MPs. Thus, contamination of terrestrial vertebrates, such as birds (Boucher and Friot, 2017) and bats, occurs through inhalation of air contaminated by MPs, consumption of contaminated water (Carlin *et al.*, 2020), and bioaccumulation and biomagnification through interactions with the environment and food webs (Waite *et al.*, 2018).

Contamination of Respiratory and Digestive Systems

Contamination of the entire respiratory system of bats, demonstrated in our study, is consistent with findings in the literature that the atmospheric air is contaminated in all environments, whether urban, rural, or even those considered pristine (Dris *et al.*, 2016; He *et al.*, 2020; Akhbarizadeh *et al.*, 2021; Truong *et al.*, 2021). Plastic debris present in atmospheric air is considered the initial and primary source of contamination of all other systems (continental or oceanic) (Dris *et al.*, 2016; Akhbarizadeh *et al.*, 2021) and direct inhalation (during breathing) of atmospheric air with plastic debris (Sridharan *et al.*, 2021) is a simple and consistent mechanism for bat respiratory contamination.

Members of the subfamily Phyllostominae in our sample include species of the genera *Lophostoma*, *Phyllostomus*, and *Tonatia* captured in our study, are considered indicators of preserved habitats with primary or secondary vegetation in an advanced state of regeneration (Faria, 2006; Oliveira and Aguiar, 2015; Palheta *et al.*, 2020; Vieira *et al.*, 2021; Weier *et al.*, 2021). This corroborates that plastic waste tends to be a contaminant of emerging concern and is distributed in all environments, either in atmospheric air or in food of bats through consumption of fruits, insects and small vertebrates. Two other examples include insectivorous bats occurring in gaps and edges of vegetation, such as members of

Embalonuridae and Mormoopidae that forage in the forest, usually between the canopy and sub-canopy. Additionally, gleaner insectivores, such as Phyllostominae bats that use the interior of the vegetation to forage use sit-and-wait strategy, were found with respiratory and digestive systems contaminated with plastic debris.

Contamination by MPs in terrestrial vertebrates has already been observed in other groups and with concentrations similar to that in bats. A high frequency of MPs (94.1% of individuals) was reported in birds (Zhao *et al.*, 2016). Equal concentration to the frequency of occurrence of MPs observed in the analyzed bats (96.3%). However, the level of contamination of individuals can vary depending on the habitat and behavior of species. In studies by Carlin *et al.* (2020), where they evaluated the ingestion of MPs in eight species of birds of prey and noted that 100% of the birds were contaminated with MPs in the digestive system. Other studies show less contamination (less than 85%) in stomach samples from two species of bird residing in coastal swamps in Mississippi, USA (Weitzel *et al.*, 2021). In both studies, the dominance of fiber-type MPs was recorded, in agreement with what we observed in our study.

Unlike other authors, we analyzed both the digestive system (from esophagus to anus), with 98.8% MP contamination, and the respiratory system (from trachea to lungs), with 96.1% MP contamination. We obtained a mean (\pm SD) food intake of 14.9 ± 17.37 MPs and air intake of 5.2 ± 7.17 MPs. Generally, the previous works analyzed only one system or organ, usually stomach or intestines. Analyzing only one system or organ limits the comparison of contamination among flying terrestrial organisms.

In general, studies of MP contamination for terrestrial organisms are scarce compared to aquatic environments. MPs have already been detected in human lungs (Pauly *et al.*, 1998); and 19.5% of samples contained fibers (Jenner *et al.*, 2022). However, other studies show that MP contamination varies from 24% (Chen *et al.*, 2022) to 84.6% (Jenner *et al.*, 2022), depending on the habits and location where the person lives.

Bats showed a higher average lung contamination than humans. However, this result is likely to be false, as the whole lungs were analyzed in bats, unlike in humans, where only a small portion of the tissue was used. In addition, data available in the literature does not allow us to make a better comparison of contaminated tissue area of human lungs with the

bat lungs. This difference in the concentration of inhaled plastic particles is possibly due to the high exposure in urban centers where people live, with direct contamination inside homes, on the street, and in the workplace. In addition, other factors, such as lifespan, must be taken into account. Humans are expected to live longer (expectation 70 years) than bats (expectation 40 years) (Podlutzky *et al.*, 2005), and exposure increases over time. Due to longer life expectancy, humans are expected to inhale more plastic particles than bats. It is essential to point out that in both bats and humans, there is a lack of knowledge about the adverse effects these particles leave on the respiratory system.

Consequences of Contamination

The ingestion of MPs is not directly linked to the risk of rapid death and survival of species, but the ingestion influences, over time, ontogenetic development and fitness. Ingested MPs will cause adverse effects, such as altered endocrine functions, decreased pup body mass, and tissue inflammation (Roman *et al.*, 2019). In birds, the ingestion of MPs causes a deceleration in sexual development (Roman *et al.*, 2019), evidenced by the transfer of MPs between adults and nestling (Carey, 2011). MPs can cause a false sense of satiety in high concentrations, leading individuals to starvation (Carbery *et al.*, 2018; Fossi *et al.*, 2018). In addition, microorganisms (Amaral-Zettler *et al.*, 2020; Chang *et al.*, 2022; Wang *et al.*, 2022) and metals (Kutralam-Muniasamy *et al.*, 2021; Zong *et al.*, 2021) may adhere to the surfaces of MPs and be additional contaminants (Kutralam-Muniasamy *et al.*, 2021). Ingestion and inhalation of MPs can cause adverse effects on bat species, including local extinction of species, which can affect ecosystem functions, such as pollination, seed dispersal, and insect control performed by bats.

All bats analyzed show plastic debris in the respiratory and digestive systems. The different foraging characteristics of bats, considering both the type of food (flowers, fruits, invertebrates, and vertebrates) and capture strategy (open areas, clearings, edges of vegetation) and habitat (urban or pristine), reinforce the idea that plastic contaminants are present in all environments (especially the terrestrial environment). We also support the need to analyze various organs/tissues, mainly when referring to the direct routes of contamination (respiratory and digestive), to estimate MP contamination and determine possible sources of contamination. The effect of the environment, type of foraging, and even the kind of food

on the accumulation of plastic waste in bats remain to be understood. Thus, research is still needed to identify differences in level of contamination by plastic waste in bats and its relationship with the type of environment, foraging strategy, and food consumed. In addition, analysis of feces of terrestrial organisms makes it possible to make inferences about how much of these ingested MPs are being eliminated from the organism in question.

ACKNOWLEDGEMENTS

This research benefited from resources from Vale SA's environmental compensation administered by the Centro Nacional de Pesquisa e Conservação de Cavernas (Cecav/ICMBio) and services to the Brazilian Society for the Study of Chiropterans — SBEQ, as part of the DD Program — The Species More Unknown in Brazil and with resources from the Termo de Compromisso de Compensação Espeleológica — TCCE VALE 1/2018 — Edital Ferruginosas 01/2021 under the administration of the Instituto Brasileiro de Desenvolvimento e Sustentabilidade — IABS — IABS.

AUTHOR CONTRIBUTION STATEMENT

LLC: collection and/or assembly of data, data analysis and interpretation, and writing the article; DRGR-B: data analysis and interpretation; MGG: critical revision of the article; DMS: data analysis and interpretation, and critical revision of the article; ABAS: collection and/or assembly of data; TBV: research concept and design, collection and/or assembly of data, critical revision and final approval of the article.

LITERATURE CITED

- AGUIAR, L. M. S., I. D. BUENO-ROCHA, G. OLIVEIRA, E. S. PIRES, S. VASCONCELOS, G. L. NUNES, M. R. FRIZZAS, and P. H. B. TOGNI. 2021. Going out for dinner. The consumption of agriculture pests by bats in urban areas. *PLOS ONE*, 16: e0258066.
- AKHBARIZADEH, R., S. DOBARADARAN, M. AMOUEI TORKMAHALLEH, R. SAEEDI, R. AIBAGHI, and F. FARAJI GHASEMI. 2021. Suspended fine particulate matter (PM_{2.5}), microplastics (MPs), and polycyclic aromatic hydrocarbons (PAHs) in air: their possible relationships and health implications. *Environmental Research*, 192: 110339.
- AL-JAIBACHI, R., R. N. CUTHBERT, and A. CALLAGHAN. 2018. Up and away: ontogenic transference as a pathway for aerial dispersal of microplastics. *Biology Letters*, 14: 20180479.
- AMARAL-ZETTLER, L. A., E. R. ZETTLER, and T. J. MINCER. 2020. Ecology of the plastisphere. *Nature Reviews Microbiology*, 18: 139–151.
- ANDERSON, J. C., B. J. PARK, and V. P. PALACE. 2016. Microplastics in aquatic environments: Implications for Canadian ecosystems. *Environmental Pollution*, 218: 269–280.
- ANDRADE, M. C., D. B. FITZGERALD, K. O. WINEMILLER, P. S. BARBOSA, and T. GIARRIZZO. 2019. Trophic niche segregation among herbivorous serrasalmids from rapids of the lower Xingu River, Brazilian Amazon. *Hydrobiologia*, 829: 265–280.

- AYALA, F., M. ZETA-FLORES, S. RAMOS-BALDÁRRAGO, J. TUME-RUIZ, A. RANGEL-VEGA, E. REYES, E. QUINDE, G. E. DE-LA-TORRE, L. LAJO-SALAZAR, and S. CÁRDENAS-ALAYZA. 2023. Terrestrial mammals of the Americas and their interactions with plastic waste. *Environmental Science and Pollution Research*, 30: 57759–57770.
- BAHO, D. L., M. BUNDSCHUH, and M. N. FUTTER. 2021. Microplastics in terrestrial ecosystems: Moving beyond the state of the art to minimize the risk of ecological surprise. *Global Change Biology*, 27: 3969–3986.
- BOUCHER, J., and D. FRIOT. 2017. Primary microplastics in the oceans: a global evaluation of sources. IUCN, Gland, Switzerland, 44 pp.
- BÜKS, F., and M. KAUPENJOHANN. 2020. Global concentrations of microplastics in soils — a review. *Soil*, 6: 649–662.
- CARBERRY, M., W. O’CONNOR, and T. PALANISAMI. 2018. Trophic transfer of microplastics and mixed contaminants in the marine food web and implications for human health. *Environmental International*, 115: 400–409.
- CAREY, M. J. 2011. Intergenerational transfer of plastic debris by short-tailed shearwaters (*Ardenna tenuirostris*). *Emu — Austral Ornithology*, 111: 229–234.
- CARLIN, J., C. CRAIG, S. LITTLE, M. DONNELLY, D. FOX, L. ZHAI, and L. WALTERS. 2020. Microplastic accumulation in the gastrointestinal tracts in birds of prey in central Florida, USA. *Environmental Pollution*, 264: 114633.
- CASTRO-LUNA, A. A., and J. GALINDO-GONZÁLEZ. 2012. Seed dispersal by phyllostomid bats in two contrasting vegetation types in a Mesoamerican reserve. *Acta Chiropterologica*, 14: 133–142.
- CHANG, J., W. FANG, J. LIANG, P. ZHANG, and G. ZHANG. 2022. A critical review on interaction of microplastics with organic contaminants in soil and their ecological risks on soil organisms. *Chemosphere*, 306: 135573.
- CHEN, Q., J. GAO, H. YU, H. SU, Y. YANG, Y. CAO, Q. ZHANG, Y. REN, H. HOLLERT, H. SHI, C. CHEN, and H. LIU. 2022. An emerging role of microplastics in the etiology of lung ground glass nodules. *Environmental Sciences Europe*, 34: 25.
- COVERNTON, G. A., C. M. PEARCE, H. J. GURNEY-SMITH, S. G. CHASTAIN, P. S. ROSS, J. F. DOWER, and S. E. DUDAS. 2019. Size and shape matter: a preliminary analysis of microplastic sampling technique in seawater studies with implications for ecological risk assessment. *Science of the Total Environment*, 667: 124–132.
- CROSSMAN, J., R. R. HURLEY, M. FUTTER, and L. NIZZETTO. 2020. Transfer and transport of microplastics from biosolids to agricultural soils and the wider environment. *Science of the Total Environment*, 724: 138334.
- DA COSTA ARAÚJO, A. P., and G. MALAFAIA. 2021. Microplastic ingestion induces behavioral disorders in mice: a preliminary study on the trophic transfer effects via tadpoles and fish. *Journal of Hazardous Materials*, 401: 123263.
- DAHMS, H. T. J., G. J. VAN RENSBURG, and R. GREENFIELD. 2020. The microplastic profile of an urban African stream. *Science of the Total Environment*, 731: 138893.
- DANTAS FILHO, J. V., V. PEREZ PEDROTI, B. L. TEMPONI SANTOS, M. M. DE LIMA PINHEIRO, Á. BEZERRA DE MIRA, F. CARLOS DA SILVA, E. C. SOARES E SILVA, J. CAVALI, E. A. CECILIA GUEDES, and S. DE VARGAS SCHONS. 2023. First evidence of microplastics in freshwater from fish farms in Rondônia state, Brazil. *Heliyon*, 9: e15066
- DE JONG, J., L. MILLON, O. HASTAD, and J. VICTORSSON. 2021. Activity pattern and correlation between bat and insect abundance at wind turbines in South Sweden. *Animals*, 11: 3269.
- DE SOUZA MACHADO, A. A., W. KLOAS, C. ZARFL, S. HEMPEL, and M. C. RILLIG. 2018. Microplastics as an emerging threat to terrestrial ecosystems. *Global Change Biology*, 24: 1405–1416.
- DEVRIESE, L. I., M. D. VAN DER MEULEN, T. MAES, K. BEKAERT, I. PAUL-PONT, L. FRÈRE, J. ROBBENS, and A. D. VETHAAK. 2015. Microplastic contamination in brown shrimp (*Crangon crangon*, Linnaeus 1758) from coastal waters of the Southern North Sea and Channel area. *Marine Pollution Bulletin*, 98: 179–187.
- DIETZ, K. J., and S. HERTH. 2011. Plant nanotoxicology. *Trends in Plant Science*, 16: 582–589.
- DRIS, R., J. GASPERI, M. SAAD, C. MIRANDE, and B. TASSIN. 2016. Synthetic fibers in atmospheric fallout: a source of microplastics in the environment? *Marine Pollution Bulletin*, 104: 290–293.
- DUIS, K., and A. COORS. 2016. Microplastics in the aquatic and terrestrial environment: sources (with a specific focus on personal care products), fate and effects. *Environmental Sciences Europe*, 28: 1–25.
- EHLERS, S. M., T. AL NAJJAR, T. TAUPP, and J. H. E. KOOP. 2020. PVC and PET microplastics in caddisfly (*Lepidostoma basale*) cases reduce case stability. *Environmental Science and Pollution Research*, 27: 22380–22389.
- ESTRADA, A., and R. COATES-ESTRADA. 2002. Bats in continuous forest, forest fragments and in an agricultural mosaic habitat-island at Los Tuxtlas, Mexico. *Biological Conservation*, 103: 237–245.
- EVANGELIOU, N., H. GRYPHE, Z. KLIMONT, C. HEYES, S. ECKHARDT, S. LOPEZ-APARICIO, and A. STOHL. 2020. Atmospheric transport is a major pathway of microplastics to remote regions. *Nature Communications*, 11: 3381.
- FADARE, O. O., and E. D. OKOFFO. 2020. Covid-19 face masks: a potential source of microplastic fibers in the environment. *Science of the Total Environment*, 737: 140279.
- FARIA, D. 2006. Phyllostomid bats of a fragmented landscape in the north-eastern Atlantic forest, Brazil. *Journal of Tropical Ecology*, 22: 531–542.
- FENTON, M. B., J. O. WHITAKER, JR., M. J. VONHOF, J. M. WATERMAN, W. A. PEDRO, L. M. AGUIAR, J. E. BAUMGARTEN, S. BOUCHARD, D. M. FARIA, C. V. PORTFORS, *et al.* 1999. The diet of bats from Southeastern Brazil: the relation to echolocation and foraging behaviour. *Revista Brasileira de Zoologia*, 16: 1081–1085.
- FLEMING, T. H., and E. R. HEITHAUS. 1981. Frugivorous bats, seed shadows, and the structure of tropical forests. *Biotropica*, 13: 45–53.
- FOSSI, M. C., C. PANTI, M. BAINI, and J. L. LAVERS. 2018. A review of plastic-associated pressures: Cetaceans of the Mediterranean Sea and Eastern Australian Shearwaters as case studies. *Frontiers in Marine Science*, 5: 173.
- GALLOWAY, T. S. 2015. Micro- and nano-plastics and human health. Pp. 343–366, *in* *Marine Anthropogenic Litter* (M. BERGMAN, L. GUTOW, and M. KLAGES, eds.). Springer Open, 447 pp.
- GASPERI, J., S. L. WRIGHT, R. DRIS, F. COLLARD, C. MANDIN, M. GUERROUACHE, V. LANGLOIS, F. J. KELLY, and B. TASSIN. 2018. Microplastics in air: are we breathing it in? *Current Opinion in Environmental Science and Health*, 1: 1–5.
- GHOSAL, S., M. CHEN, J. WAGNER, Z. M. WANG, and S. WALL. 2018. Molecular identification of polymers and

- anthropogenic particles extracted from oceanic water and fish stomach — a Raman micro-spectroscopy study. *Environmental Pollution*, 233: 1113–1124.
- GROSS, M. 2015. Oceans of plastic waste. *Current Biology*, 25: 93–96.
- HE, D., K. BRISTOW, V. FILIPOVIĆ, J. LV, and H. HE. 2020. Microplastics in terrestrial ecosystems: a scientometric analysis. *Sustainability (Switzerland)*, 12: 1–15.
- HEINLAAN, M., K. KASEMETS, V. ARUOJA, I. BLINOVA, O. BONDARENKO, A. LUKJANOVA, A. KHOSROVYAN, I. KURVET, M. PULLERITS, M. SIHTMÄE, *et al.* 2020. Hazard evaluation of polystyrene nanoplastic with nine bioassays did not show particle-specific acute toxicity. *Science of the Total Environment*, 707: 136073.
- HERNANDEZ, E., B. NOWACK, and D. M. MITRANO. 2017. Polyester textiles as a source of microplastics from households: a mechanistic study to understand microfiber release during washing. *Environmental Science and Technology*, 51: 7036–7046.
- HOCKING, D. P., F. G. MARX, T. PARK, E. M. G. FITZGERALD, and A. R. EVANS. 2017. A behavioural framework for the evolution of feeding in predatory aquatic mammals. *Proceedings of the Royal Society*, 284B: 20162750.
- HORTON, A. A., A. WALTON, D. J. SPURGEON, E. LAHIVE, and C. SVENDSEN. 2017. Microplastics in freshwater and terrestrial environments: evaluating the current understanding to identify the knowledge gaps and future research priorities. *Science of the Total Environment*, 586: 127–141.
- IMMERSCHITT, I., and A. MARTENS. 2020. Ejection, ingestion and fragmentation of mesoplastic fibres to microplastics by *Anax imperator* larvae (Odonata: Aeshnidae). *Odonatologica*, 49: 57–66.
- JAWAD, L. A., N. J. ADAMS, and M. K. NIEUWOUDT. 2021. Ingestion of microplastics and mesoplastics by *Trachurus declivis* (Jenyns, 1841) retrieved from the food of the Australasian gannet *Morus serrator*: first documented report from New Zealand. *Marine Pollution Bulletin*, 170: 112652.
- JENNER, L. C., J. M. ROTCHELL, R. T. BENNETT, M. COWEN, V. TENTZERIS, and L. R. SADOFSKY. 2022. Detection of microplastics in human lung tissue using μ FTIR spectroscopy. *Science of the Total Environment*, 831: 154907.
- KALKO, E. K. V. 1998. Organisation and diversity of tropical bat communities through space and time. *Zoology*, 101: 281–297.
- KASSO, M., and M. BALAKRISHNAN. 2013. Ecological and economic importance of bats (Order Chiroptera). *ISRN Biodiversity*, 2013: 1–9.
- KUMARI, A., V. D. RAJPUT, S. S. MANDZHIEVA, S. RAJPUT, T. MINKINA, R. KAUR, S. SUSHKOVA, P. KUMARI, A. RANJAN, V. P. KALINITCHENKO, *et al.* 2022. Microplastic pollution: an emerging threat to terrestrial plants and insights into its remediation strategies. *Plants*, 11: 340.
- KUTRALAM-MUNIASAMY, G., F. PÉREZ-GUEVARA, I. E. MARTÍNEZ, and V. C. SHRUTI. 2021. Overview of microplastics pollution with heavy metals: analytical methods, occurrence, transfer risks and call for standardization. *Journal of Hazardous Materials*, 415: 125755.
- LAHIVE, E., A. WALTON, A. A. HORTON, D. J. SPURGEON, and C. SVENDSEN. 2019. Microplastic particles reduce reproduction in the terrestrial worm *Enchytraeus crypticus* in a soil exposure. *Environmental Pollution*, 255: 113174.
- LAVOY, M., and J. CROSSMAN. 2021. A novel method for organic matter removal from samples containing microplastics. *Environmental Pollution*, 286: 117357.
- LI, H., L. ZHU, M. MA, H. WU, L. AN, and Z. YANG. 2023. Occurrence of microplastics in commercially sold bottled water. *Science of the Total Environment*, 867: 161553.
- LI, L., Y. LUO, R. LI, Q. ZHOU, W. J. G. M. PEIJNENBURG, N. YIN, J. YANG, C. TU, and Y. ZHANG. 2020. Effective uptake of submicrometre plastics by crop plants via a crack-entry mode. *Nature Sustainability*, 3: 929–937.
- LIU, J., Y. YANG, J. DING, B. ZHU, and W. GAO. 2019. Microfibers: a preliminary discussion on their definition and sources. *Environmental Science and Pollution Research*, 26: 29497–29501.
- LIU, K., W. COURTENE-JONES, X. WANG, Z. SONG, N. WEI, and D. LI. 2020. Elucidating the vertical transport of microplastics in the water column: a review of sampling methodologies and distributions. *Water Research*, 186: 116403.
- LUSHER, A. L., N. A. WELDEN, P. SOBRAL, and M. COLE. 2017. Sampling, isolating and identifying microplastics ingested by fish and invertebrates. *Analytical Methods*, 9: 1346–1360.
- MANI, T., A. HAUKE, U. WALTER, and P. BURKHARDT-HOLM. 2015. Microplastics profile along the Rhine River. *Scientific Reports*, 5: 17988.
- MCCORMICK, A. R., T. J. HOELLEIN, M. G. LONDON, J. HITTIE, J. W. SCOTT, and J. J. KELLY. 2016. Microplastic in surface waters of urban rivers: Concentration, sources, and associated bacterial assemblages. *Ecosphere*, 7: e01556.
- MILLER, M. E., M. HAMANN, and F. J. KROON. 2020. Bioaccumulation and biomagnification of microplastics in marine organisms: a review and meta-analysis of current data. *PLoS ONE*, 15: e0240792.
- NAPPER, I. E., and R. C. THOMPSON. 2020. Plastic debris in the marine environment: history and future challenges. *Global Challenges*, 4: 1900081.
- NUELLE, M. T., J. H. DEKIFF, D. REMY, and E. FRIES. 2014. A new analytical approach for monitoring microplastics in marine sediments. *Environmental Pollution*, 184: 161–169.
- OLIVEIRA, H. F. M. De, and L. M. S. AGUIAR. 2015. The response of bats (Mammalia: Chiroptera) to an incidental fire on a gallery forest at a Neotropical savanna. *Biota Neotropica*, 15: e0091.
- PALHETA, R. LEANDRA, L. GUSTAVO, S. LEANDRO, D. SILVA, and B. JENNIFER. 2020. The effect of urbanization on bats and communities of bat flies (Diptera: Nycteribiidae and Streblidae) in the Amazon, northern Brazil. *Acta Chiropterologica*, 22: 403–416.
- PASSOS, J. B., and M. PASSAMANI. 2003. *Artibeus lituratus* (Phyllostomidae): biologia e dispersão de sementes no Parque do Museu de Biologia Prof. Mello Leitão, Santa Teresa (ES). *Natureza On Line*, 1: 1–6.
- PAULY, J. L., S. J. STEGMEIER, H. A. ALLAART, R. T. CHENEY, P. J. ZHANG, A. G. MAYER, and R. J. STRECK. 1998. Inhaled cellulosic and plastic fibers found in human lung tissue. *Cancer Epidemiology Biomarkers and Prevention*, 7: 419–428.
- PEEL, M. C., B. L. FINLAYSON, and T. A. MCMAHON. 2007. Updated world map of the Köppen-Geiger climate classification. *Hydrology and Earth System Sciences*, 11: 1633–1644.
- PODLUTSKY, A. J., A. M. KHRITANKOV, N. D. OVODOV, and S. N. AUSTAD. 2005. A new field record for bat longevity. *Journals of Gerontology*, 60A: 1366–1368.

- R CORE TEAM. 2021. R: a language and environment for statistical computing, version 4.1 [Computer software]. R Foundation for Statistical Computing, Vienna. Retrieved from <https://cran.r-project.org>.
- RAGUSA, A., A. SVELATO, C. SANTACROCE, P. CATALANO, V. NOTARSTEFANO, O. CARNEVALI, F. PAPA, M. C. A. RONGIOLETTI, F. BAIOTTO, S. DRAGHI, *et al.* 2021. Plasticenta: first evidence of microplastics in human placenta. *Environment International*, 146: 106274.
- RAGUSA, A., V. NOTARSTEFANO, A. SVELATO, A. BELLONI, G. GIOACCHINI, C. BLONDEEL, E. ZUCHELLI, C. DE LUCA, S. D'AVINO, A. GULOTTA, *et al.* 2022. Raman microscopy detection and characterisation of microplastics in human breastmilk. *Polymers*, 14: 2700.
- REVEL, M., A. CHÂTEL, and C. MOUNEYRAC. 2018. Micro-(nano)plastics: a threat to human health? *Current Opinion in Environmental Science and Health*, 1: 17–23.
- RIBEIRO-BRASIL, D. R. G., L. S. BRASIL, G. K. O. VELOSO, T. P. de MATOS, E. S. de LIMA, and K. DIAS-SILVA. 2022. The impacts of plastics on aquatic insects. *Science of the Total Environment*, 813: 152436.
- RIBEIRO-BRASIL, D. R. G., N. R. TORRES, A. B. PICAÑO, D. S. SOUSA, V. S. RIBEIRO, L. S. BRASIL, and L. F. DE ASSIS MONTAG. 2020. Contamination of stream fish by plastic waste in the Brazilian Amazon. *Environmental Pollution*, 266: 115241.
- RILLIG, M. C., L. ZIERSCH, and S. HEMPEL. 2017. Microplastic transport in soil by earthworms. *Scientific Reports*, 7: 1362.
- RODRÍGUEZ-SAN PEDRO, A., J. L. ALLENDES, C. A. BELTRÁN, P. N. CHAPERON, M. M. SALDARRIAGA-CÓRDOBA, A. X. SILVA, and A. A. GREZ. 2020. Quantifying ecological and economic value of pest control services provided by bats in a vineyard landscape of central Chile. *Agriculture, Ecosystems and Environment*, 302: 107063.
- ROMAN, L., L. LOWENSTINE, L. M. PARSLEY, C. WILCOX, B. D. HARDESTY, K. GILARDI, and M. HINDELL. 2019. Is plastic ingestion in birds as toxic as we think? Insights from a plastic feeding experiment. *Science of the Total Environment*, 665: 660–667.
- SABOROWSKI, R., E. PAULISCHKIS, and L. GUTOW. 2019. How to get rid of ingested microplastic fibers? A straightforward approach of the Atlantic ditch shrimp *Palaemon varians*. *Environmental Pollution*, 254: 113068.
- SHRUTI, V. C., F. PÉREZ-GUEVARA, and G. KUTRALAM-MUNIASAMY. 2021. Wet wipes contribution to microfiber contamination under COVID-19 era: an important but overlooked problem. *Environmental Challenges*, 5: 100267.
- SRIDHARAN, S., M. KUMAR, L. SINGH, N. S. BOLAN, and M. SAHA. 2021. Microplastics as an emerging source of particulate air pollution: a critical review. *Journal of Hazardous Materials*, 418: 126245.
- STEENSGAARD, I., K. SYBERG, S. RIST, N. HARTMANN, A. BOLDRIN, and S. F. HANSEN. 2017. From macro- to microplastics — analysis of EU regulation along the life cycle of plastic bags. *Environmental Pollution*, 224: 289–299.
- SURAN, M. 2018. A planet too rich in fibre. *EMBO Reports*, 19: e46701.
- SURIPTO, B. 2021. Economic Contribution of Fruit Bats (Family Pteropodidae) Through Durian Fruit Production in the Agroecosystem in Java Island. *Advances in Social Science, Education and Humanities Research*, 528: 8–15.
- THOMPSON, R. C., S. H. SWAN, C. J. MOORE, and F. S. VOM SAAL. 2009. Our plastic age. *Philosophical Transactions of the Royal Society*, 364B: 1973–1976.
- TOKUNAGA, Y., H. OKOCHI, Y. TANI, Y. NIIDA, T. TACHIBANA, K. SAIGAWA, K. KATAYAMA, S. MORIGUCHI, T. KATO, and S. I. HAYAMA. 2023. Airborne microplastics detected in the lungs of wild birds in Japan. *Chemosphere*, 321: 138032.
- TRUONG, T. N. S., E. STRADY, T. C. KIEU-LE, Q. V. TRAN, T. M. T. LE, and Q. T. THUONG. 2021. Microplastic in atmospheric fallouts of a developing Southeast Asian megacity under tropical climate. *Chemosphere*, 272: 129874.
- VIEIRA, T. B., L. C. N. DA SILVA, L. M. de S. AGUIAR, M. OPREA, P. MENDES, and A. D. DITCHFIELD. 2021. Bat species composition associated with restinga lagoons from the Paulo César Vinha State Park, Espírito Santo, Brazil. *Papéis Avulsos de Zoologia*, 61: e20216132.
- WAGNER, M., C. SCHERER, D. ALVAREZ-MUÑOZ, N. BRENNHOLT, X. BOURRAIN, S. BUCHINGER, E. FRIES, C. GROBSOIS, J. KLASMEIER, T. MARTI, *et al.* 2014. Microplastics in freshwater ecosystems: what we know and what we need to know. *Environmental Sciences Europe*, 26: 1–9.
- WAITE, H. R., M. J. DONNELLY, and L. J. WALTERS. 2018. Quantity and types of microplastics in the organic tissues of the eastern oyster *Crassostrea virginica* and Atlantic mud crab *Panopeus herbstii* from a Florida estuary. *Marine Pollution Bulletin*, 129: 179–185.
- WANG, X., L. ZHU, K. LIU, and D. LI. 2022. Prevalence of microplastic fibers in the marginal sea water column off southeast China. *Science of the Total Environment*, 804: 150138.
- WEIER, S. M., V. M. G. LINDEN, A. HAMMER, I. GRASS, T. TSCHARNTKE, and P. J. TAYLOR. 2021. Bat guilds respond differently to habitat loss and fragmentation at different scales in macadamia orchards in South Africa. *Agriculture, Ecosystems & Environment*, 320: 107588.
- WEITZEL, S. L., J. M. FEURA, S. A. RUSH, R. B. IGLAY, and M. S. WOODREY. 2021. Availability and assessment of microplastic ingestion by marsh birds in Mississippi Gulf Coast tidal marshes. *Marine Pollution Bulletin*, 166: 112187.
- WILSON, D. R., B. J. GODLEY, G. L. HAGGAR, D. SANTILLO, and K. L. SHEEN. 2021. The influence of depositional environment on the abundance of microplastic pollution on beaches in the Bristol Channel, UK. *Marine Pollution Bulletin*, 164: 111997.
- WINDSOR, F. M., R. M. TILLEY, C. R. TYLER, and S. J. ORMEROD. 2019. Microplastic ingestion by riverine macroinvertebrates. *Science of the Total Environment*, 646: 68–74.
- WRIGHT, S. L., R. C. THOMPSON, and T. S. GALLOWAY. 2013. The physical impacts of microplastics on marine organisms: a review. *Environmental Pollution*, 178: 483–492.
- YIN, L., X. WEN, D. HUANG, C. DU, R. DENG, Z. ZHOU, J. TAO, R. LI, W. ZHOU, Z. WANG, and H. CHEN. 2021. Interactions between microplastics/nanoplastics and vascular plants. *Environmental Pollution*, 290: 117999.
- ZHANG, Q., E. G. XU, J. LI, Q. CHEN, L. MA, E. Y. ZENG, and H. SHI. 2020. A review of microplastics in table salt, drinking water, and air: direct human exposure. *Environmental Science and Technology*, 54: 3740–3751.
- ZHAO, S., L. ZHU, and D. LI. 2016. Microscopic anthropogenic litter in terrestrial birds from Shanghai, China: not only plastics but also natural fibers. *Science of the Total Environment*, 550: 1110–1115.
- ZICCARDI, L. M., A. EDGINGTON, K. HENTZ, K. J. KULACKI, and S. KANE DRISCOLL. 2016. Microplastics as vectors for bioaccumulation of hydrophobic organic chemicals in the

marine environment: a state-of-the-science review. *Environmental Toxicology and Chemistry*, 35: 1667–1676.

ZONG, X., J. ZHANG, J. ZHU, L. ZHANG, L. JIANG, Y. YIN, and H. GUO. 2021. Effects of polystyrene microplastic on uptake

and toxicity of copper and cadmium in hydroponic wheat seedlings (*Triticum aestivum* L.). *Ecotoxicology and Environmental Safety*, 217: 112217.

Received 27 May 2023, accepted 04 December 2023

Associate Editor: M. Corrie Schoeman