



Freshwater Mussels (Bivalvia: Unionida) of Vietnam: Diversity, Distribution, and Conservation Status

Authors: Do, Van Tu, Tuan, Le Quang, and Bogan, Arthur E.

Source: Freshwater Mollusk Biology and Conservation, 21(1) : 1-18

Published By: Freshwater Mollusk Conservation Society

URL: <https://doi.org/10.31931/fmbc.v21i1.2018.1-18>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

REGULAR ARTICLE

FRESHWATER MUSSELS (BIVALVIA: UNIONIDA) OF VIETNAM: DIVERSITY, DISTRIBUTION, AND CONSERVATION STATUS

Van Tu Do¹, Le Quang Tuan¹, and Arthur E. Bogan^{2*}

¹ Institute of Ecology and Biological Resources (IEBR), Vietnam Academy of Science and Technology, 18 Hoang Quoc Viet, Nghia Do, Cau Giay, Ha Noi, Vietnam, dovantu.iebr@gmail.com; penguin0343@gmail.com

² North Carolina Museum of Natural Sciences, 11 West Jones Street, Raleigh, NC 27601 USA

ABSTRACT

Vietnam has the second highest diversity of freshwater mussels (Unionida) in Asia after China. The purpose of this paper is to compile an up-to-date list of the modern unionid fauna of Vietnam and its current conservation status. Unfortunately, there has been relatively little research on this fauna in Vietnam. Fifty-nine species of Unionida have been recorded from Vietnam based on literature, museum records, and our fieldwork. Fifty were assessed in the International Union for Conservation of Nature (IUCN) Red List 2016 in the IUCN categories of Critically Endangered (four species, 6.8%), Endangered (seven species, 12%), Vulnerable (one species, 1.7%), Near Threatened (two species, 3.4%), Least Concern (23 species, 39%), Data Deficient (11 species, 18.6%), and Not Evaluated (11 species, 18.6%). Considering the impacts of pollution, timbering, agriculture, and damming of rivers, research on the diversity and conservation status of freshwater mussels is very urgently needed to propose specific conservation measures for these species in Vietnam. If all taxa listed as Data Deficient are found to be threatened, with around 42% of species threatened, this fauna would be one of the most threatened freshwater molluscan faunas in Asia.

KEY WORDS: Unionidae, Margaritiferidae, IUCN Red List, extinct, endangered, citizen science

INTRODUCTION

Freshwater bivalves are widely distributed in the freshwaters of the world and are considered one of the most imperiled animal groups (Bogan 1993, 2008; Lydeard et al. 2004; Strayer and Dudgeon 2010; Graf 2013; Haag and Williams 2014). Two major diversity hotspots of unionid bivalves are the southeastern United States (Neves et al. 1997; Bogan 2008; Haag 2012) and east and southeastern Asia (Bolotov et al. 2017; Zieritz et al. 2017). Our understanding of the distribution of unionids in Asia is limited and the coverage of information is uneven and mostly historical (Dudgeon et al. 2006; Zieritz et al. 2017).

Overviews of the freshwater bivalve fauna across Asia, including Vietnam, were provided by Haas (1910a, 1910b, 1910c–1920, 1923). Compendia covering the Unionida of the

world began with the *Synopsis* by Lea (1836, 1838, 1852, 1870) and were followed and expanded by Simpson (1900, 1914) and Modell (1942, 1949, 1964). Haas (1969a) wrote the last comprehensive survey of the Unionida, and followed it with a treatise (Haas 1969b) that covered both modern and fossil taxa to the generic level. Higher classification of the Unionida was overinflated by Starobogatov (1970). Brandt (1974) summarized the Thai and Southeast Asian unionids, but overlooked Haas (1969a, 1969b). Đặng et al. (1980) followed the classification proposed by Starobogatov (1970). The inflated taxonomy of the Unionidae erected by Starobogatov's classification was reduced by Graf (2007). A preliminary list of freshwater bivalve taxa from Vietnam from literature and museum data was included in the summary of the East and Southeast Asia freshwater bivalve fauna by Zieritz et al. (2017). Analyzing molecular data, the Unionidae has been confirmed as a monophyletic clade (Hoeh et al. 1998,

*Corresponding Author: arthur.bogan@naturalsciences.org

Table 1. Unionid taxa listed and/or figured by in the four volumes by Thach (2005, 2007, 2012, 2016b).

Species	2005	2007	2012	2016b
<i>Anodonta harlandi</i> Baird & Adams, 1867 ¹				Fig.
<i>Cristaria bialata</i> (Lea, 1829) ²	L/Fig.			
<i>Cristaria discoidea</i> ³		L/Fig.	Fig.	
<i>Cristaria truncata</i>			L/Fig.	
<i>Hyriopsis cumingii</i> ⁴			L/Fig.	
<i>Hyriopsis schlegeli</i> ⁵	L/Fig.			
<i>Hyriopsis</i> sp.			L/Fig.	
<i>Lamprotula leai</i>	L/Fig.		Fig.	
<i>Lanceolaria bilirata</i>				L
<i>Lanceolaria bogani</i>				L/Fig.
<i>Lanceolaria fruhstorferi</i>				Fig.
<i>Lanceolaria grayi</i>			L/Fig.	
<i>Lanceolaria grayana</i> (Lea, 1834)				Fig.
<i>Lanceolaria laevis</i>	L	Fig.		
<i>Lanceolaria yueyingae</i> He & Zhuang, 2013				Fig.
<i>Nodularia douglasiae</i>				L
<i>Oxynaia jourdyi</i>		L/Fig.		
<i>Oxynaia micheloti</i>	L/Fig.			
<i>Physunio inornatus</i>		L/Fig.		
<i>Physunio micropterus</i>		L/Fig.		
<i>Physunio modelli</i>		L/Fig.		
<i>Pilsbryconcha exilis</i>	L/Fig.		Fig.	
<i>Pilsbryconcha lemeslei</i>		L/Fig.		
<i>Pseudodon cambodjensis</i>		L/Fig.		
<i>Pseudodon contradens tumidula</i> (Lea, 1856) ⁶		Fig.		
<i>Pseudodon mouhoti</i> [sic] ⁷	L/Fig.			
<i>Pseudodon vondembuschianus chaperi</i> (Morgan, 1885)		L/Fig.		
<i>Sinanodonta elliptica</i>	Fig.			Fig.
<i>Sinanodonta hunganhi</i>				L/Fig.
<i>Sinanodonta woodiana</i>				Fig.
<i>Trapezoideus exolecens</i>		L/Fig.		
<i>Uniandra contradens tumidula</i> (Lea, 1856) ⁸	L			

L = listed; Fig.= figured.

¹*Anodonta harlandi* is a junior synonym of *Sinanodonta woodiana*.

²*Cristaria bialata* is a junior synonym of *C. plicata*.

³*Cristaria discoidea* is a junior synonym of *Pletholophus tenuis*.

⁴*Hyriopsis cumingii* moved to *Sinohyriopsis cumingii* (see Lopes-Lima et al. 2017).

⁵*Hyriopsis schlegeli* is misidentification of *Sinohyriopsis cumingii*.

⁶Misplaced species, belongs in *Contradens*.

⁷Incorrect spelling of *Pseudodon mouhotii*

⁸Correct genus is *Contradens*.

2001, 2002; Roe and Hoeh 2003; Graf and Cummings 2006; Breton et al. 2007, 2010; Doucet-Beaupré et al. 2010; Whelan et al. 2011; Pfeiffer and Graf 2015). The most recent phylogenetic classification within the modern Unionidae is that of Lopes-Lima et al. (2017) and Bolotov et al. (2017).

Early work on the Vietnamese molluscan fauna was typically performed by European malacologists describing freshwater bivalve species. These included the papers of Morelet (1865, 1866), Mabile (1887), Morlet (1886a, 1886b, 1891), Dautzenberg (1900), Bavay and Dautzenberg (1901), Martens (1902), Rochebrune (1904a, 1904b), Rolle (1904),

Dautzenberg and Fischer 1906a, 1906b, 1908), and Haas (1910a, 1910b, 1910c-1920, 1913, 1923). Isaac Lea, living in the United States, also described some unionids from Southeast Asia (see Scudder 1885). The only comprehensive treatment of the freshwater invertebrates, including freshwater bivalves for northern Vietnam, is that by Đặng et al. (1980).

Thach, a retired physician and avocational shell collector has published four volumes providing information and pictures of marine, terrestrial, and freshwater mollusks of Vietnam, including some Unionidae (Thach 2005, 2007, 2012, 2016b) (Table 1). A total of 21 species of freshwater mussels

from Vietnam were reported and figured in the 4 volumes by Thach. He also described two new species, *Lanceolaria bogani* Thach, 2016 (Thach 2016a) and *Sinanodonta hunganhi* Thach, 2016 (Thach 2016b). Thach divided the freshwater bivalves of Vietnam into Amblemidae and Unionidae, but did not comment on the Margaritiferidae. His report and figure of *Pilsbryconcha lemeslei* is the first record of this species from Vietnam (Thach 2007).

We recently have been surveying the freshwater mussel fauna of Vietnam (Bogan and Do 2011, 2013a, 2013b, 2014a, 2014b, 2016). The objective of the present paper is to develop an up-to-date list of the unionid fauna reported for Vietnam and their distribution (Table 2; see also Supplemental Species Range Maps). The modern unionid fauna of Vietnam is divided between two families, Margaritiferidae and Unionidae. Margaritiferidae is represented by a single species. The Unionidae fauna of Vietnam is represented by 28 genera and 58 species (Table 2).

METHODS

Study Area

Vietnam lies on the eastern side of the Indochina Peninsula encompassing 331,210 km² (Fig. 1). The northern part of the country is mountainous and contains the Red River basin, which drains to the east, emptying into the Gulf of Tonkin. The rivers draining to the west are tributaries of the Mekong River basin, draining into the Gulf of Thailand. Southern Vietnam is home to the Annamite Range, the Central or Western Highlands and the extensive Mekong River delta and its numerous distributaries. The major rivers of Vietnam are shown on Figure 1.

Historical literature was used to document species described from Vietnam and was compared with the list published by Đặng et al. (1980). Taxonomy of these species was challenging because Đặng et al. (1980) based their taxonomy on Starobogatov (1970), but see Graf's (2007) revision of Starobogatov's taxonomy. Taxonomy used in this paper is based on available literature, as presented by Lopes-Lima et al. (2017) and is the same as used by Zieritz et al. (2017). Additional information was collected from photographs of specimens from various museum collections available on the Internet, which were checked and identifications verified (Graf and Cummings 2017). Dates of publication were verified and resulted in minor changes (Bogan and Do 2011; Bogan 2015).

Our fieldwork was carried out at the following times: July 2010 in northern Vietnam; November 2012 across 12 northern provinces of Vietnam; March 2014 beginning in Hanoi, including the Central Highlands and the Mekong Delta area of southern Vietnam; and October–November 2016, concentrating on 7 northern provinces (Bogan and Do 2013a, 2013b, 2014a, 2014b, 2016). During each field trip, markets were visited in the early morning and people selling freshwater bivalves and gastropods were asked about where the animals

had been collected (Fig. 2). Other residents and fishermen were asked about local freshwater mussels. We searched streams and lakes crossed during our trips by wading and feeling for mussels or using hand dredges. Collection methods varied by location; some shell harvesters used handheld wire baskets on a rope to collect mussels and gastropods in northern Vietnam (Fig. 3). Long bamboo-handled rectangular wire frame nets were used to collect mussels from stream and lake bottoms in southern Vietnam (Fig. 4). One shell collector used a surface-supplied-air diving apparatus to dive and collect freshwater mussels.

The Vietnam Red Book was first developed in 1992, and it was a collaboration of the Institute of Ecology and Biological Resources (IEBR) and the International Union for Conservation of Nature (IUCN) with financial support from Sweden. The criteria used in the book were built on the IUCN Red List standards. The 1992 list was later revised and contained 13 species of freshwater mussels (Red Data Book of Vietnam 2000). This was expanded by the Institute of Science and Technology of Vietnam. A country-wide assessment of the conservation status of 416 species of animals in Vietnam resulted in the publication of the Vietnam Red Data Book (2007). This assessment included 11 species of freshwater mussels; 1 species was added and 3 species were dropped from the earlier lists: *Cristaria herculea*, *Lamprotula liedtkei*, and *Pilsbryconcha suilla* and was conducted using the IUCN conservation assessment protocol in place at the time. The Vietnam Red Data Book (2007) lists six Rare; four Vulnerable, and a single indeterminate mussel species (Table 2).

Conservation assessments of all species listed here were based upon the application of the criteria laid out in the IUCN Red List Categories and Criteria (2012) and explained in the IUCN Assessment Process (IUCN 2017). The categories of threatened status include Extinct, Extinct in the Wild, Critically Endangered, Endangered, Vulnerable, Near Threatened, Least Concern, Data Deficient, and Not Evaluated (see Table 2). Each category is based on five criteria: (1) reduction in population size, (2) geographic range, (3) population size estimated to number at listed level for each threatened status, (4) population size estimated to number less than a given level for the threatened status, and (5) quantitative analysis showing probability of extinction in the wild. The guidelines for the application of these criteria are carefully explained by the IUCN Standards and Petitions Subcommittee (2016). The standards and an explanation of required data is presented in tabular form based on the IUCN data entry format for assessing the conservation status of each species (IUCN 2013). The threatened species assessed by the IUCN from the Indo-Burma region were initially listed in Köhler et al. (2012). The conservation status of all the species listed in Table 2 and supporting information can be found on the IUCN Red List website (IUCN 2016).

This work has been registered with ZooBank and a copy has been archived at Zenodo.org.




Figure 1. Map of the major rivers of Vietnam.

RESULTS

Margaritiferidae Henderson, 1929

Recent molecular studies of the Margaritiferidae have considered it consists of a single genus, *Margaritifera* Schumacher, 1816 (Bolotov et al. 2016b; Araujo et al. 2017). Bolotov et al. (2016b) recognized three clades, which are treated as subgenera. *Margaritifera laosensis* is the only species belonging to the subgenus *Margaritanopsis* Haas, 1910. Historically, *M. laosensis* was known from Dien Bien Province in northwest Vietnam and in the adjacent part of Lao People's Democratic Republic (PDR). During recent fieldwork in West Central Vietnam (Bogan and Do 2014a), a local stated to have found similar shells but we did not procure any. Phuong (2011) provided an IUCN conservation assessment. The ecology and conservation of *M. laosensis* was detailed by Bolotov et al. (2014), who listed three known viable populations in the Lao PDR.

Unionidae Rafinesque, 1820

Aculamprotula nodulosa was reported from Cao Bang area of northern Vietnam by Đặng et al. (1980). Graf and Cummings (2007) continued to place this species in *Lamprotula*. However, Pfeiffer and Graf (2013) confirmed the split of *Lamprotula* and *Aculamprotula*. *Unio nodulosa* was placed in the genus *Aculamprotula* by He and Zhuang (2013) and Graf and Cummings (2017). Live specimens have been recently collected in northern Vietnam (Bogan and Do 2016).

Photographs presented by Graf and Cummings (2017) and a photograph provided by S. Schneider (University of Cambridge, Cambridge, UK, personal communication) are identified as *A. nodulosa*, but these shells are much rounder than the type figure of *A. nodulosa*.

Per Haas (1969a) *Chamberlainia hainesiana* [+ *Simpsonia demangei* Rochebrune, 1904 is listed as occurring in Thailand, Cambodia, and the Tonkin region of Vietnam. *Simpsonia demangei* is listed from "Rivière Claire entre Vietri et Tuyen-Quas, (Tonkin)" (Rochebrune, 1904). *Simpsonia demangei* was listed in the synonymy of *C. hainesiana* by Brandt (1974) and Graf and Cummings (2017). Such a disjunct distribution may suggest that the animal from Tonkin may be a separate and distinct species from *C. hainesiana*. New live specimens are needed to test the placement of this species described from Vietnam using molecular methods.

Uniandra Haas, 1912 was used as a senior synonym of *Conradens* Haas, 1913 by Brandt (1974). *Conradens* Haas, 1911 has priority over *Uniandra* Haas, 1912, based on the dates of publication of the sections of Haas (1910c–1920) (Bogan 2015).

Conradens conradens was listed by Brandt (1974) from Thailand, Lao PDR, Cambodia and southern Vietnam [+ *Uniandra conradens tumidula* (Lea, 1856)]. However, it was not reported by Đặng et al. (1980). This species has been collected from southern Vietnam (Bogan and Do 2014). Graf and Cummings (2017) figured lots identified as *Unio*

semidecoratus Morlet, 1889, Tonkin Muséum national d'Histoire naturelle, Paris (MNHN MP) 3861; *Unio dautzenbergi* Morlet, 1889, Tonkin; and *Physunio crossei* Deshayes, 1879, from Cochinchine MNHN MP 3774. All specimens were identified as *Conradens conradens* by Graf and Cummings (2017).

Considering that the type locality for *Conradens semmelincki* (Martens, 1891) is Borneo and the type locality for *Conradens fultoni* is Manson, Tonkin (Mau Son, Lang Son), *C. fultoni* is here considered a valid separate species. Graf and Cummings (2017) treat *C. fultoni* as a junior synonym of *C. semmelincki*.

Cristaria plicata is viewed as a valid, wide-ranging species (Lopes-Lima et al. 2017). Đặng et al. (1980) recognized four species of *Cristaria* in Vietnam: *Cristaria bellua* Morelet, 1866; *Cristaria bialata* Lea, 1829; *Cristaria herculea* Middendorff, 1847; and *Cristaria truncata* Đặng, 1980. Klishko et al. (2014, 2016) determined that *C. herculea* and *Cristaria tuberculata* Schumacher, 1817 are synonyms of *C. plicata*. *Cristaria bellua*, a misspelling of *Anodonta bellus*, and *Symphynota bialata* were listed by Brandt (1974) and He and Zhuang (2013) as synonyms of *C. plicata*.

Cristaria truncata is recognized as a valid species by Graf and Cummings (2017) but is poorly known and its relationship to *C. plicata* is unknown.

Cuneopsis demangei was described from Tonkin and recognized by Haas (1969a), Đặng et al. (1980), and Graf and Cummings (2017). This species is restricted to northern Vietnam but has not been collected in the last several decades.

Cuneopsis pisciculus (Heude, 1874) was illustrated by Graf and Cummings (2017) as University of Michigan Museum of Zoology (UMMZ) 110095, purportedly from Tonkin, Vietnam. The accompanying original label listed it from Ningpo and not from Tonkin. This is considered a spurious record and not part of the fauna of Vietnam.

Diaurora aurea was listed by Đặng et al. (1980), but has not been collected recently. Early specimens identified as *Diaurora* are housed in the Vietnam National University, Hanoi University of Science, Museum of Biology, Hanoi, but were not available for examination.

Ensidens sagittarius was reported by Brandt (1974) as a synonym of *Ensidens ingallsianus*, but Graf and Cummings (2017) treated *E. ingallsianus* as a separate species and illustrated specimens from Cochinchine. However, the specimens listed as *E. sagittarius* from Cochinchine all appear to be specimens of *E. ingallsianus*.

Gibbosula Simpson, 1900 was described containing only *Mya crassa* Wood, 1815. Haas (1969a, 1969b) included *Gibbosula* as a junior synonym of *Lamprotula* Simpson, 1900. He and Zhuang (2013) and Graf and Cummings (2017) recognized *Gibbosula* as a valid genus. Pfeiffer and Graf (2013) recognized two clades in *Lamprotula* but did not address the status of *Gibbosula* because the type species of *Lamprotula* is poorly understood. *Gibbosula* is retained here recognizing its uncertain status and placement.

Gibbosula crassa was listed as occurring in northern

Table 2. Complete list of freshwater mussel species found in Vietnam: 59 species, 28 genera, 2 families. The species conservation statuses are based on the International Union for Conservation of Nature (IUCN) Red Listing (2016) and the Red Data Book of Vietnam (1992, 2000) and Vietnam Red Data Book (2007).

Species	IUCN Red List	Vietnam Red Data Books	Distribution Area (Province) in Vietnam, Comments
<i>Aculamprotula nodulosa</i> (Wood, 1815)*	CR	R	Northeast (Bang Giang River in Cao Bang, Ky Cung River in Lang Son, Thuong River in Bac Giang, Tuyen Quang (IUCN listed as <i>Lamprotula nodulosa</i>)
<i>Chamberlainia hainesiana</i> (Lea, 1856)	DD	V	Northeast (Phu Tho, Lo River in Tuyen Quang)
<i>Contradens contradens</i> (Lea, 1838)	LC		Mekong River delta (An Giang, Long An) IUCN listed as <i>Uniandra contradens</i>
<i>Contradens fultoni</i> Haas, 1939 *	DD	R	Northeast (Mau Son in Lang Son) (IUCN Listed as <i>Uniandra semmelincki</i>)
<i>Cristaria plicata</i> (Leach, 1815)	DD	V	Northeast, Northwest, Red River delta
<i>Cristaria truncata</i> Đãng, 1980	EN		Northeast (Cao Bang, Bac Giang), Red River delta (Hai Duong, Ha Nam, Ninh Binh, Nam Dinh)
<i>Cuneopsis demangei</i> Haas, 1929	CR	K-indeterminate	Northeast (Da River in Phu Tho)
<i>Diuroira aureora</i> (Heude, 1883)	DD		Northeast (?)
<i>Ensidents ingallsianus</i> (Lea, 1852)	LC		Central Highlands, South Central Coast, Southeast, Mekong River delta
<i>Gibbousula crassa</i> (Wood, 1815)	CR	R	Northeast (Bang Giang River in Cao Bang, Ky Cung River in Lang Son) (IUCN listed as <i>Lamprotula crassa</i>)
<i>Harmandia somborensis</i> (Rochebrune, 1882)	DD		Southeast, Mekong River delta
<i>Hyriopsis gracilis</i> Haas, 1910	LC		Southeast, Mekong River delta
<i>Hyriopsis delaportei</i> (Crosse & Fischer, 1876)	LC		Mekong River delta (An Giang)
<i>Lamprotula bazini</i> (Heude, 1877)	DD		Northeast (Ha Noi)
<i>Lamprotula blaisei</i> (Dautzenberg & Fischer, 1905)*	VU	R	Northeast (Cao Bang, Lang Son, Bac Giang)
<i>Lamprotula contritus</i> (Heude, 1881)	EN		Northeast (Cao Bang, Bac Kan, Thai Thuyen)
<i>Lamprotula leai</i> (Gray in Griffith & Pidgeon, 1833)	LC	V	Northeast
<i>Lamprotula quadrangulosa</i> (Heude, 1881)	LC		Northeast (Bang Giang River) (IUCN treated as synonym of <i>L. leai</i> not listed)
<i>Lamprotula salaputium</i> (Martens, 1902)*	DD		Northeast (Tuyen Quang)
<i>Lanceolaria bilirata</i> (Martens, 1902)*	CR		Northeast (Cao Bang, Lang Son)
<i>Lanceolaria bogani</i> Thach, 2016*	NE		North Central Coast (Thua Thien Hue)
<i>Lanceolaria fruhstorferi</i> (Bavay & Dautzenberg, 1901)*	NE	R	Northeast (Cao Bang, Lang Son) (IUCN lists as synonym of <i>Lanceolaria grayana</i>)
<i>Lanceolaria gladiola</i> (Heude, 1877)	LC		Northeast, Northwest, Red River delta
<i>Lanceolaria grayi</i> (Gray in Griffith & Pidgeon, 1833)	DD		Northeast, Northwest, Red River delta
<i>Lanceolaria laevis</i> (Martens, 1902)*	NE		Northeast (Cong River in Thai Nguyen)
<i>Margaritifera laosensis</i> (Lea, 1863)	EN		Northwest (Dien Bien Phu)
<i>Nodularia dorri</i> (Wattebled, 1886)*	LC		Northeast, Northwest, Red River delta
<i>Nodularia douglasi</i> (Gray, 1833)	LC		Northeast, Northwest, Red River delta, North Central Coast
<i>Nodularia nuxpersica</i> (Dunker, 1848)	NE		Northeast (IUCN listed as <i>Unio douglasi</i>)
<i>Oxynta diespiter</i> (Mabille, 1887)*	EN		Northeast (Bac Kan, Thai Nguyen)
<i>Oxynta gladiator</i> (Ancey, 1881)*	DD		Northeast
<i>Oxynta jourdyi</i> (Morlet, 1886)*	NT		Northeast (Ha Noi, Vinh Phu, Phu Tho, Hoa Binh)
<i>Oxynta micheloti</i> (Morlet, 1886)*	EN		Northeast (Cao Bang, Phu Tho, Vinh Phuc)
<i>Physunio cambodiensis</i> (Lea, 1856)	LC		Central Highlands (Serepok River in Dak Lak), Mekong River delta (An Giang)

Table 2, continued.

Species	IUCN Red List	Vietnam Red Data Books	Distribution Area (Province) in Vietnam, Comments
<i>Physunio micropterus</i> (Morelet, 1866)	LC		Central Highlands (Dak Bla River tributary of Se San River in Kon Tum), Southeast (Dong Nai), Mekong River delta
<i>Physunio modelli</i> Brandt, 1974	LC		South Central Coast (Binh Thuan)
<i>Physunio superbus</i> (Lea, 1843)	LC		Mekong River delta (Can Tho)
<i>Pilsbryconcha compressa</i> (Martens, 1860)	LC		Southeast, Mekong River delta (IUCN listed as synonym of <i>P. exilis</i>)
<i>Pilsbryconcha exilis</i> (Lea, 1839)	LC		South Central Coast (Ninh Thuan), Southeast, Mekong River delta (Can Tho)
<i>Pilsbryconcha lemeslei</i> (Morelet, 1875)	LC		Central Highlands (Dak Bla River tributary of Se San River in Kon Tum, Khanh Hoa)
<i>Pletholophus</i> sp.*	NE		Northeast, Red River delta
<i>Pletholophus tenuis</i> (Gray in Griffith & Pidgeon, 1833)	LC		Throughout the country (IUCN listed as <i>Cristaria tenuis</i>)
<i>Protunio messageri</i> (Bavay & Dautzenberg, 1901)*	EN		Northeast (Bang Giang River in Cao Bang)
<i>Pseudobaphia</i> sp.	NE	R	Northeast (Bang Giang River in Cao Bang)
<i>Pseudodon cambodjensis</i> (Petit, 1865)	DD		Southeast, Mekong River delta
<i>Pseudodon inoscularis</i> (Gould, 1844)	LC		Southeast (Dong Nai), Mekong River delta
<i>Pseudodon mouhotii</i> (Lea, 1863)	LC		Southeast, Mekong River delta
<i>Pseudodon resupinatus</i> Martens, 1902*	EN		Northeast (Vinh Phuc, Lang Son)
<i>Pseudodon vondembuschianus</i> (Lea, 1840)	LC		Southeast, Mekong River delta
<i>Ptychorhynchus pfisteri</i> (Heude, 1874)	NT		Northeast (Vinh Phuc, Phu Tho), Northwest (Hoa Binh)
<i>Scabies crispata</i> (Gould, 1843)	LC		Northwest (Hoa Binh), Mekong River delta (An Giang)
<i>Sinanodonta hunganhi</i> Thach, 2016*	NE		North, Nghe An Province
<i>Sinanodonta jourdyi</i> (Mortlet, 1866)*	NE		Northeast, Northwest, Red River delta
<i>Sinanodonta lucida</i> (Heude, 1878)	NE		Northeast, Northwest, Red River delta
<i>Sinohyriopsis cumingii</i> (Lea, 1852)	LC	V	Northeast (Bac Giang, Ha Noi, Ha Nam, Nam Dinh, Ninh Binh) (IUCN listed as <i>Cristaria cumingii</i>)
<i>Solenia oleivora</i> (Heude, 1877)	NE		Northeast (Phan River in Vinh Phuc, Day River in Ha Noi and Ha Nam, Thuong River in Bac Giang)
<i>Trapezoideus exolescens</i> (Gould, 1843)	LC		Southeast, Mekong River delta
<i>Trapezoideus misellus</i> (Morelet, 1865)	DD		Northeast (Bang Giang River in Cao Bang, Vinh Phuc)
<i>Unionetta fabagina</i> (Deshayes in Deshayes & Julien, 1874)	LC		Central Highlands (Sa Thay River in Kon Tum)

CR = Critically Endangered; DD = Data Deficient; EN = Endangered; IUCN = International Union for Conservation of Nature; K = Indeterminate; LC = Least Concern; NE = Not Evaluated; NT = Near Threatened; R = Rare; VU = Vulnerable; V = Vulnerable.
*Endemic to Vietnam.



Figure 2. Pan with several unionid species for sale in a market, Ha Noi, Vietnam, Photograph by Arthur Bogan. November 18, 2014.

Vietnam by Đặng et al. (1980). Haas (1969a), Đặng et al. (1980), and He and Zhuang (2013) all placed *Lamprotula mansuyi* (Dautzenberg and Fischer, 1908), described from northern Vietnam in the synonymy of *Lamprotula crassa*. Live specimens of this species were recently collected from the Bang River, Cao Bang Province, Vietnam, for anatomical and molecular analyses (Bogan and Do 2016).

Harmandia somboriensis has been reported from the Mekong and Mun rivers, but not mentioned as occurring in Vietnam (Brandt 1974). Pfeiffer (personal communication) found it in shallow water around boulders. There are type specimens collected from “Cochinchine” preserved in the MNHN, Paris. This species collected from “Cochinchine” are assumed to be part of the Vietnamese fauna.

The genus *Hyriopsis* Conrad, 1853 has some taxonomic issues. *Limnoscapha* Lindholm, 1932 was proposed for a group of bivalve species occurring from the Miocene to Pliocene in what was, in 1978, the southern Soviet Union; these species became extinct at the end of the Pliocene (Gozhik 1978). This generic name was placed as a subgenus of the modern *Hyriopsis* and has been used as a modern subgenus (Modell 1950; Brandt 1974). Haas (1969b) and Graf and Cummings (2006) listed *Limnoscapha* as a synonym of *Hyriopsis*. *Limnoscapha* represents, in our opinion, an extinct fossil group not related to the modern *Hyriopsis* species of Asia.

Hyriopsis bialatus was listed from Malaysia, Thailand, Cambodia, southern Vietnam, and Tonkin (Brandt 1974) and recently confirmed from the Mekong Delta in Vietnam by Bogan and Do (2014b). However, *Hyriopsis bialatus* has been shown to be three separate species based on mitochondrial DNA sequence data with *H. bialatus* being described from the “in Songi flumine Malaccae” (Sungi River, Malacca, Malaysia) (Zieritz et al. 2016, 2017). Sungi is the Malay



Figure 3. Small clam rake used by local fishermen to collect freshwater mussels from the local rivers. Photograph by Van Tu Do. Ca Lo River, Soc Son District, Ha Noi, Vietnam. Photograph by Van Tu Do. November 23, 2012.

word for river, so the locality is unclear and Malaccae referred to the southern side of peninsular Malaysia. An available name for the species occurring in the Mekong River basin is *Hyriopsis gracilis* Haas, 1910 (Haas 1910b). We are using *H.*



Figure 4. Long-handled rake with basket from southern Vietnam. Photograph by Van Tu Do. April 4, 2014.

gracilis for what has been historically listed as *H. bialatus* only for the Mekong River basin populations.

Hyriopsis cumingii has been recognized by numerous authors including Haas (1969a, 1969b), Brandt (1974), and He and Zhuang (2013). Starobogatov (1970) described *Sinohyriopsis* with the type species *Unio cumingii* Lea, 1852 and simultaneously described *Nipponihyria* Starobogatov, 1970 type species *Hyriopsis schlegeli* Martens, 1861. Đặng et al. (1980) used the combination *Sinohyriopsis cumingii*. No use of the name *Nipponihyria* has been found in the literature since it was described. *Sinohyriopsis* is here given priority over *Nipponihyria* based on usage since both names were published in the same original work. *Hyriopsis cumingii* and *H. schlegeli* have been placed in *Sinohyriopsis* based on recent phylogenetic work that separated them from the type species of *Hyriopsis*, *Hyriopsis bialatus*, (Froufe et al. 2015; Lopes-Lima et al. 2017). *Sinohyriopsis cumingii* is used here.

Hyriopsis delaportei was reported for the first time in An Giang Province, Vietnam (Bogan and Do 2014b).

Lamprotula Simpson, 1900 was first divided by Wu (1998) into *Lamprotula* and *Aculamprotula* based on differences in shell shape, anatomy, and glochidial morphology. This distinction was confirmed with molecular sequence data by Zhou et al. (2007). Pfeiffer and Graf (2013) confirm this division but raise questions about the other generic names available for parts of this polyphyletic group.

Lamprotula bazini was reported from northern Vietnam by Đặng et al. (1980) and recognized as valid by He and Zhuang (2013) and Graf and Cummings (2017). This species has not been collected in several decades.

Lamprotula blaisei was listed from northern Vietnam by Đặng et al. (1980) and recognized as a valid by He and Zhuang (2013) and Graf and Cummings (2017). This species has not been collected in several decades.

Lamprotula contritus was reported from northern Vietnam by Đặng et al. (1980). He and Zhuang (2013) and Graf and Cummings (2017) consider this a synonym of *Lamprotula caveata* (Heude, 1877). This species has not been collected recently. Animals of this species from Vietnam do not appear to be the same as *L. caveata*, as they lack the characteristic depressions in the shell. We treat it here as a valid species pending further genetic analysis.

Lamprotula leai was reported from northern Vietnam by Đặng et al. (1980) and recognized as a valid species by He and Zhuang (2013) and Graf and Cummings (2017). This species is the most common species of this genus in northern Vietnam.

Lamprotula liedtkei (Rolle, 1904) was listed from northern Vietnam by Đặng et al. (1980) and recognized as a junior synonym of *Aculamprotula nodulosa* by He and Zhuang (2013) and Graf and Cummings (2017). However, the specimen of *Quadrula liedtkei* sp., with a Rolle manuscript name figured by Graf and Cummings (2017) is from Tonkin (United States National Museum, Smithsonian Institution [USNM] 187462) and is much rounder than the figure of the type of *Unio liedtkei* Rolle, 1904 and pictured by He and Zhuang (2013). This USNM specimen resembles the shell

shape of *Lamprotula similaris* (Simpson, 1900) from China (He and Zhuang 2013).

Lamprotula ponderosa (Dautzenberg and Fischer, 1905) was recognized as a variety separate from *Lamprotula leai*. However, it was listed as a junior synonym of *Lamprotula leai* by Haas (1969a). He and Zhuang (2013) and Graf and Cummings (2017) listed *Unio (Quadrula) leai* var. *ponderosa* Dautzenberg and Fischer, 1905 as a primary junior homonym of *Unio ponderosa* Rossmässler, 1842. It is also a junior homonym to *Unio ponderosa* Hanley, 1842 and *Unio ponderosa* Gray, 1825. The shell illustrated by Đặng et al. (1980) may be a separate species or variation of *L. leai*.

Lamprotula quadrangulosa was described from China and placed by Simpson (1914) as a junior synonym of *L. caveata* (Heude, 1877) also described from China. Đặng et al. (1980) viewed *L. quadrangulosa* as a valid species, separate from both *L. leai* and *L. caveata*. Haas (1969a), He and Zhuang (2013), and Graf and Cummings (2017) listed *L. quadrangulosa* as a synonym of *L. caveata*. *Lamprotula quadrangulosa* from Vietnam does not appear to be the same as *L. caveata*, lacking the characteristic depressions in the shell characteristic of *L. caveata*. We treat *L. quadrangulosa* as a valid species pending further genetic analysis.

Lamprotula salaputium described from Thuyen-Quan, Annam, was not figured by Martens. Martens (1902) noted he had a single specimen, making the specimen a holotype by monotypy. Graf and Cummings (2017) do not provide a picture of this species, but recognize it as a valid species. The mollusk collections in Berlin and Frankfurt museums have been queried and neither has this specimen. This species is an unknown species described from Vietnam.

Đặng et al. (1980) reported four species of *Lanceolaria* from northern Vietnam, *Lanceolaria bilirata*, *Lanceolaria fruhstorferi*, *Lanceolaria grayii* and *Lanceolaria laevis*. Graf and Cummings (2017) recognized three valid species from Vietnam: *L. bilirata*, *Lanceolaria gladiola*, and *L. grayii*. This genus needs a careful revision.

Lanceolaria bilirata was described from Tonkin, Vietnam, and is apparently restricted to northern Vietnam. Haas (1969a) placed this taxon as a subspecies of *Lanceolaria oxyrhyncha* (Martens, 1894) and Kondo (2008) listed it as a synonym of *Lanceolaria grayana* (Lea, 1834). The authors recognize this species.

Lanceolaria bogani was recently described from Thura Thien-Hue Province, Vietnam, and placed in the Unioninae by Thach (2016a).

Lanceolaria fruhstorferi was recognized as a valid species by Haas (1910c) but listed by He and Zhuang (2013) and Graf and Cummings (2017) as a junior synonym of *L. grayii*. This species was recognized as a separate valid species from Vietnam by Thach (2016a).

Lanceolaria gladiola was described from China but Haas (1911) and Graf and Cummings (2017) report it from Vietnam and figure specimens that appear to be *Lanceolaria grayii*. This is a doubtful record for Vietnam.

Lanceolaria grayii is the senior synonym for *Unio*

grayanus Lea, 1834 described from China (Bogan and Do 2011). Graf and Cummings (2017) include *L. fruhstorferi*, *L. laevis* and *Lanceolaria gracillimus* Rolle, 1904 as synonyms of *L. grayii*. All three were described from Tonkin. This group needs revision.

Lanceolaria laevis was originally described from Tonkin. It has been recognized and figured by Đặng et al. (1980) and Thach (2005, 2007). It was listed by He and Zhuang (2013) and Graf and Cummings (2017) as a junior synonym of *L. grayii*.

Nodularia Conrad, 1853 was erected with *Unio douglasiae* Griffith & Pidgeon, 1833 as the type species. Simpson (1900, 1914) and Starobogatov (1970) used *Nodularia* as a valid genus. However, *Nodularia* was considered a junior synonym of *Unio* Retzius, 1788 (Haas 1969a, 1969b). Đặng et al. (1980) included two taxa from northern Vietnam in *Nodularia*. Graf and Cummings (2007) recognized the genus and included four species, while Graf and Cummings (2017) included only three species. *Nodularidia* Cockerell, 1901, was an unnecessary replacement name for *Nodularia* when it was considered preoccupied. *Nodularidia* was used by He and Zhuang (2013) for a single species in China. *Nodularia* and *Unio* represent separate clades and may belong in separate tribes (Lopes-Lima et al. 2017). We recognize three taxa in *Nodularia* from Vietnam.

Nodularia dorri was reported by Đặng et al. (1980). Graf and Cummings (2017) also used this species and noted its distribution from the Gulf of Tonkin to the Mekong. It appears to be restricted to northern and central Vietnam.

Nodularia douglasiae crassidens Haas, 1910 (Haas 1910a) appeared in Simpson (1914) as a *Nodularia* (*Nodularia*) *douglasiae* var. *crassidens* and was recognized by Đặng et al. (1980). Haas (1969a) placed this taxon as a synonym of *Unio douglasiae douglasiae* but listed it as occurring in China and questionably in Cambodia. Graf and Cummings (2017) synonymize this subspecies under *Nodularia douglasiae*. *Nodularia douglasiae* is used here but it is recognized that *Nodularia* needs extensive revision.

Nodularia nuxpersica is a new addition to the fauna of Vietnam based on recent collections from markets in northern Vietnam. He and Zhuang (2013) figured and included *Unio nuxpersica* in the synonymy of *Scabies crispata* (Gould, 1843) and *Scabies chinensis* Liu et al., 1991 (He and Zhuang 2013). The Vietnam samples were shown to belong in *Nodularia* based on DNA sequence data (Lopes-Lima et al. 2017).

Oxynaia was described by Haas, but the date of publication has been confused. Haas (1969a) recorded it as 1911 but Haas (1969b) used 1913. Starobogatov (1970) used *Oxynaia* Haas, 1912. Graf and Cummings (2017) noted the generic name was associated with a figure in 1911 as a nomen nudum and only described in 1913 (Haas 1913) and subsequently redescribed in Haas (1913b). Based on the dates of publication for Haas (1910c–1920) reported by Bogan (2015), plates 14, 15, 16 in Haas (1911) are the first appearance of the generic name *Oxynaia* associated with four described species published in 1911. Thus, the generic name *Oxynaia* was available from the

date of publication of the plates. This is considered an indication for a genus named before 1931 under the International Commission on Zoological Nomenclature (ICZN 1999) Code Article 12.11.2.2. The generic name *Oxynaia* takes the date of publication as Haas, 1911 (see Bogan 2015). The type species of *Oxynaia* was designated as *Unio jourdyi* Morlet, 1886 by Haas (1913).

Oxynaia diespiter was described from Tonkin, and known from the single type specimen and conchologically is very close to *Oxynaia jourdyi*. Đặng et al. (1980) reported this species from northern Vietnam. Graf and Cummings (2017) illustrated a single specimen from Tonkin, MNHN_MO_2998, which may be the unmarked holotype.

Oxynaia gladiator was described from Tonkin and recognized as valid by Haas (1969a) and Graf and Cummings (2017). Haas (1969a) listed this species from Tonkin and Annam, Vietnam. However, it looks very much like *Oxynaia micheloti* (Morlet, 1886). Adding this to the fauna of Vietnam would bring the total *Oxynaia* species to four.

Oxynaia jourdyi is listed by Haas (1969a) as the type species for the genus; it was described from Bac Hat etang de la riviere Claire. Đặng et al. (1980) also listed this species.

Oxynaia micheloti was described from “Tonkin.” Haas (1969a) mentions this species is only known from the original description. Đặng et al. (1980) recognized the species. Based on the figures provided by Graf and Cummings (2017), *O. micheloti* does not have a very long shell and has a taller shell than many of the lots figured as that species. Some of the other figured lots appear to represent *Nodularia* species.

Physunio cambodiensis is listed from Cambodia, Lao PDR, and Thailand by Haas (1969a), Brandt (1974), and Graf and Cummings (2017). Davidson et al. (2006) collected this species from a site on the Serepok River in Dak Lak Province, and the Mekong River, An Giang and Long An Provinces, Vietnam. These are the first records for this species in Vietnam.

Physunio micropterus was reported by Brandt (1974) and Graf and Cummings (2017) from Cambodia and Thailand. Two specimens were reported from “Cochin China” or southern Vietnam (Graf and Cummings 2017). Davidson et al. (2006) collected this species from a site on the Dak Bla River tributary of the Se San River, Kon Tum Province, Vietnam. This species has been reported from Luy River, Binh Thuan province.

Physunio modelli was described from north central Thailand and reported from Thailand and Lao PDR (Brandt 1974). It has been reported from the Ham Tan district, Binh Thuan Province by Thach (2007:190, plate 61, fig. 1057).

Physunio superbus was described from New Holland in error and Brandt (1974) listed the distribution as possibly including southern Vietnam. Đặng et al. (1980) did not report this species. Graf and Cummings (2017) figure specimens from Cochinchine or the Mekong Delta area of southern Vietnam. There are specimens in the IEBR collections, Hanoi, with the only locality data recorded as Vietnam.

Pilsbryconcha compressa was recognized as a subspecies

of *Pilsbryconcha exilis* by Brandt (1974). Graf and Cummings (2017) listed *P. compressa* as a separate species. This species was reported from An Giang Province, Vietnam (Bogan and Do 2014b).

Pilsbryconcha exilis was reported from southern Vietnam (Brandt 1974). Graf and Cummings (2017) figured specimens from Vietnam.

Pilsbryconcha lemeslei was not mentioned by Đặng et al. (1980). Haas (1969a) listed it only from Thailand and Cambodia and Brandt (1974) did not mention this species from Vietnam. Thach (2007) mentions this species from Khanh Hoa, Vietnam, and figured a specimen that resembles specimens figured by Graf and Cummings (2017).

Pilsbryconcha suilla von Martens, 1902 was not originally figured and was placed in the genus *Pilsbryconcha* by Simpson (1914). Simpson remarked that Martens thought this species reminded him of *Pilsbryconcha*. Đặng et al. (1980) used the combination *Pilsbryconcha suilla* but did not figure this species. *Anodonta suilla* has been used as a junior synonym of *Sinanodonta woodiana* by Haas (1969a) and Graf and Cummings (2017). If it has a shell shape close to *Sinanodonta jourdyi* Morlet, it would become a synonym of *S. jourdyi* of Vietnam (Lopes-Lima, personal communication).

Cristaria discoidea (Lea, 1834) has been placed in *Cristaria* (*Pletholophus*) by Simpson (1900, 1914) and Haas (1969a). *Unio tenuis* Gray in Griffith and Pidgeon, has been dated as 1834 and listed as a junior synonym of *Unio discoidea* Lea, 1834 by Lea (1836, 1838, 1852, 1870), Simpson (1900, 1914) and Haas (1969a). Petit and Coan (2008) determined the date of publication of the figure of *Unio tenuis* Gray in Griffith and Pidgeon as 1833, and noted *Unio tenuis* has priority over the later name *Unio discoidea* Lea, 1834. This case does not meet the requirements of the Code 23.9 (ICZN 1999) for usage and the older name has date priority and must prevail. The correct name for *Cristaria discoidea* is *Cristaria tenuis* (Gray in Griffith and Pidgeon, 1833). *Cristaria tenuis* was used by He and Zhuang (2013) and Graf and Cummings (2017). Đặng et al. (1980) elevated *Pletholophus* to generic level and included three species: *Pletholophus swinhoi* (Adams, 1866); *Pletholophus inangulatus* (Haas, 1910a), and *Pletholophus discoideus* (Lea, 1834). All of them are considered synonyms under *Cristaria tenuis* (He and Zhuang 2013; Graf and Cummings 2017). Placement of *C. tenuis* in *Pletholophus* and separate from *Cristaria* was confirmed by Lopes-Lima et al. (2017).

DNA sequence analyses of *Pletholophus* samples from Vietnam revealed a second distinct species, here assumed to represent a new species, *Pletholophus* sp. (Lopes-Lima, personal communication).

Protunio Haas, 1912 was described by Haas but various dates have been listed for its publication. Haas (1969a, 1969b), Starobogatov (1970), and Graf and Cummings (2017) all listed 1913 (Haas 1913) as the date of publication of *Protunio*. Graf and Cummings (2017) cited Haas 1912 (plate 32) for *Protunio* but used 1913 as the date for the generic description (Haas 1913). Based on the dates of publication for Haas (1910c–

1920) reported by Bogan (2015), plate 32 is the first appearance of the generic name *Protunio* associated with a described species published in 1912 (Bogan 2015). Thus, the generic name *Protunio* was available from the date of publication of the plate and since *Protunio* was published in association with a single species, *Unio messengeri* Bavay and Dautzenberg, 1901 is the type species by monotypy. This is considered an indication for a genus named before 1931 under ICZN Code Article 12.112.2. The generic name *Protunio* takes the date of publication as Haas, 1912 (see Bogan 2015). *Protunio* was subsequently redescribed by Haas with the date on the signature of 1914 and the date on the cover of the *Lieferung* containing the four signatures of 1919 (Bogan 2015).

Protunio messengeri was recognized by Haas (1969a, 1969b) and Graf and Cummings (2017) as restricted to northern Vietnam. Đặng et al. (1980) reported this species from around Cao Bang and Lang Son. However, the specimen illustrated by Đặng et al. (1980) and Đặng and Hồ (in press) is not *Protunio messengeri* when compared with the original figures (Heude 1877) and figures provided by Graf and Cummings (2017). The specimen figured by Đặng et al. (1980) and Đặng and Hồ (in press) represents a different species. The figured shell shape is very similar to *Pseudobaphia biesiana* (Heude, 1877) figured by Haas (1910c–1920), He and Zhuang (2013), and Graf and Cummings (2017) but represents an undescribed species, *Pseudobaphia* sp. *Pseudobaphia* sp. is known from only three lots of specimens, one lot in IEBR, Hanoi, one lot in the North Carolina Museum of Natural Sciences, Raleigh, and a large lot in the Vietnam National University, Hanoi University of Science, Museum of Biology, Hanoi. This species has not been collected since 1971.

Pseudodon inoscularis is the type species of *Pseudodon* and has been reported from throughout Southeast Asia. Brandt (1974) recognized one species as a Rassenkreis or a ring species with a variety of subspecies. This group is in dire need of a taxonomic revision. Graf and Cummings (2017) claimed the range of *P. inoscularis* from South Vietnam but did not illustrate any specimens.

Pseudodon callifer (Martens, 1860) was listed by Brandt (1974) as subspecies of *P. inoscularis callifer* and reported from Thailand, Cambodia and southern Vietnam.

Pseudodon cambodjensis was not reported by Brandt (1974) or Đặng et al. (1980) from Vietnam, but Graf and Cummings (2017) list a specimen (Museum of Comparative Zoology, Harvard [MCZ] 37431) from Cochinchine and mapped it in southern Vietnam.

Pseudodon ellipticus Conrad, 1865 was listed by Graf and Cummings (2017) from Thailand, Cambodia, and southern Vietnam but they did not figure any specimens. It was described from Cambodia. Brandt (1974) used it as a subspecies of *Pseudodon vondembuschianus*.

Pseudodon inoscularis was recognized by Graf and Cummings (2017), who listed it from southern Vietnam. Haas (1969a) listed this species from Tenasserim, Myanmar [+]

Burma], only. Brandt (1974) treated the species as a ring species ranging from Myanmar, Thailand, Malaysia, Lao PDR, Cambodia, and southern Vietnam. This wide-ranging species appears to grade from one form to another across its range and should be carefully examined with molecular techniques.

Pseudodon mouhotii was recognized and listed from Vietnam by Brandt (1974) and Graf and Cummings (2017), but was not discussed by Đặng et al. (1980). *Pseudodon exilis* (Morelet, 1866) was considered by Brandt (1974) a junior synonym of *Pseudodon mouhotii*.

Pseudodon resupinatus was described from Than Moi, Tonkin, but was not originally figured. It is recognized as a valid species by Simpson (1914), Haas (1920), Đặng et al. (1980), and Graf and Cummings (2017). This species is endemic to northern Vietnam.

Pseudodon vondembuschianus, as used by Brandt (1974), contained three subspecies including *Pseudodon vondembuschianus ellipticus* Conrad, 1865, reported from southern Vietnam. Đặng et al. (1980) and Graf and Cummings (2017) do not list this species from Vietnam. It is found in southern Vietnam.

Ptychorhynchus pfisteri was reported by Đặng et al. (1980) from northern Vietnam. The species was not listed from Vietnam by Haas (1969a, 1969b), He and Zhuang (2013), or Graf and Cummings (2017). Recently specimens have been collected by the authors in Hanoi.

Scabies crispata was reported by Đặng et al. (1980) and Brandt (1974) from Vietnam. Graf and Cummings (2017) report this species from Cochinchine and Tonkin, Vietnam, but some of the specimens figured appear to be specimens of *Nodularia*. Recently, a field survey conducted by L. A. Prozorova and N. X. Quang (personal communication) recorded this species from Bung Binh Thien, An Phu, An Giang (southern Vietnam).

Sinanodonta hunganhi was described from around Vinh City, Nghệ An Province, in northern Vietnam and reported it living “along rivers” [sic] (Thach 2016b). Based on the published figure, this species appears to fit within the shell variation of *S. jourdyi*.

Sinanodonta jourdyi and *Sinanodonta elliptica* (Heude, 1878) were identified from northern Vietnam (Đặng et al. 1980). Specimens collected from Vietnam were identified as both species and were examined genetically. These specimens represent a single species, distinct from *Sinanodonta woodiana* of China (Lopes-Lima, personal communication). The name available for the Vietnamese species is *S. jourdyi*. Because *S. elliptica* was described from China, it is likely a synonym of *S. woodiana* and was not found in Vietnam (Lopes-Lima, personal communication).

Sinanodonta lucida was reported from North Vietnam (Đặng et al. 1980). Graf and Cummings (2017) treated *S. lucida* as a junior synonym of *S. woodiana*. Bolotov et al. (2016a) reported it as a separate, valid species but from China.

Sinanodonta woodiana has been assumed to be a wide-ranging and plastic species, with 103 synonyms listed for this

species (Graf and Cummings 2017). Haas (1969a) treated all three taxa reported from Vietnam as synonyms of *Anodonta* (*Anodonta*) *woodiana* (Lea, 1834). Đặng et al. (1980) recognized three species in the genus *Sinanodonta* in northern Vietnam, *S. jourdyi* (Morlet, 1886), *S. elliptica* (Heude, 1878), and *S. lucida* (Heude, 1878). He and Zhuang (2013) and Graf and Cummings (2017) listed all three taxa as synonyms of *S. woodiana*. Preliminary DNA sequence analyses have separated *S. jourdyi* from *S. woodiana* occurring in China. Because the type locality for *S. elliptica* is China, it is not recognized in Vietnam and is considered part of the greater *S. woodiana* complex (Lopez-Lima, personal communication). The complexity of *S. woodiana* has been documented for Asia and the species invading Europe (Bolotov et al. 2016a). Bolotov et al. (2016a) documented seven separate lineages within what has been named *Sinanodonta woodiana*. These analyses also separate *Sinanodonta lucida* as a separate lineage within the *S. woodiana* complex. This taxonomic puzzle will require further analyses to resolve this group.

Solenia oleivorus has been collected from the Phan River (Vinh Phuc Province), Day River (Ha Noi and Ha Nam provinces), and Thuong River (Bac Giang province) in Vietnam. This species was synonymized by He and Zhuang (2013) under *Solenia iridinea* (Heude, 1874) in China, but determined to be a separate and valid species by Ouyang et al. (2011). The authors recognize it as a valid species.

Trapezoideus was recognized as containing a single species with three subspecies by Brandt (1974). He remarked *Trapezoideus* might occur in southern Vietnam. Đặng et al. (1980) recognized *Trapezoideus misellus*. Graf and Cummings (2017) placed *T. misellus* as a junior synonym of *Trapezoideus excolescens* (Gould, 1843). Preliminary genetic analyses have suggested *Trapezoideus* is not a monotypic genus, but may harbor several cryptic species (Lopes-Lima, personal communication). Now, we have chosen to recognize the wide-spread species *T. excolescens* in southern Vietnam and *T. misellus* from northern Vietnam.

Unionetta fabagina was listed by Haas (1969a) from Cambodia and Lao PDR. Brandt (1974) reported *Unionetta fabagina* from the Mekong River, but not from Vietnam. A shell of *U. fabagina* has been collected from Sa Thay River, Kon Tun and Gia La provinces (Central Highlands), Vietnam.

DISCUSSION

Distribution of the Fauna

The unionid fauna of Vietnam can be divided along the major river basins draining the country (Fig. 1). The Red River basin is the major basin in the north, passing through Hanoi. There are representatives of the fauna in the far north of Vietnam that have relationships with the fauna of the Zhu River basin and the Yangtze River basin farther north in southern China (e.g., *Lamprotula* species). The western border areas of Vietnam, including Dien Bien Province and rivers of the Central Highland and southern Vietnam, are drained by

tributaries to the Mekong River basin in the north. The Mekong River fauna described by Brandt (1974) for Thailand, Lao PDR, and Cambodia extends into southern Vietnam in the distributaries in the Mekong Delta. The unionid fauna in the eastern rivers of the Central Highland is still poorly known.

Seventeen species are considered endemic to Vietnam and comprise 28.8% of the total species found in Vietnam (Table 2). Forty species were found in the north while 18 species were reported from the south and only one species (*Sinanodonta jourdyi*) is widely distributed throughout the country (Table 2; see also Supplemental Individual Species Maps).

Our recent surveys demonstrated that the northeastern area of Vietnam possess the highest diversity of freshwater mussels in Vietnam. Se San and Serepok, tributaries of the Mekong River, can be ranked as a second hotspot of diversity of this group. It is necessary to conduct more surveys in the northwest of Vietnam in Dien Bien Province and the Central Highlands where data are quite scarce. We have only spent part of a single field trip in southern Vietnam and need to spend more time in the Mekong Delta area of Vietnam.

Conservation Assessment

The IUCN hosted a training session in Phenom Penh, Cambodia, and a workshop in Vientiane, Lao PDR, to assess the conservation status of the freshwater fauna of the Indo-Burma area, including Vietnam (Allen et al. 2012). Freshwater bivalves and gastropods were a part of this program (Köhler et al. 2012). Our current conservation assessment began with the information from IUCN Red List (2016) and is supplemented with our recent field data and museum records. The IUCN conservation status of the freshwater bivalves of Vietnam is summarized in Table 2.

Based on the 59 species recorded in Vietnam (Table 2), those taxa assessed in the IUCN Red List (2016) included 4 species (6.8%) assessed as Critically Endangered, 7 species (12%) assessed as Endangered, 1 species (1.7%) assessed as Vulnerable, 2 species (3.4) assessed as Near Threatened, 23 species (39%) assessed as Least Concern, 11 species (18.6%) assessed as Data Deficient, and 11 species (18.6%) that were not evaluated.

Based on our survey results beginning in 2010 and using the IUCN Red List categories and criteria (IUCN 2012), we herein recommend that the conservation status of following four species should be changed and we will submit a revised species assessment to the IUCN:

(1) *Solenia oleivora* should be assessed as Vulnerable. It was found only in the Phan, Day, and Thuong rivers; population reduction was estimated about 50% over 10 yr; extent of occurrence was estimated as smaller than 20,000 km². This species is assessed only on the Vietnamese portion of its range. (2) *Lamprotula quadrangulosa* should be assessed as Vulnerable. It was found only in the Bang River; population reduction was estimated about 50% over 10 yr; extent of

occurrence was estimated as smaller than 20,000 km². This species is assessed only on the Vietnamese portion of its range. (3) *Trapezoideus misellus* should be assessed as Vulnerable. It was found only in the Bang River; population reduction was estimated about 50% over 10 yr; extent of occurrence was estimated as smaller than 20,000 km². This species assessed only on data from northern Vietnam.

(4) *Conradens fultoni* should be assessed at least as Endangered. It was found in only the Mau Son River, Lang Son Province, based on an early record from 1930; extent of occurrence was estimated as smaller than 5,000 km². This species is endemic to Vietnam.

We are particularly concerned about the status and continued survival of four species in the Bang River basin and tributaries of Li Chiang in northeastern Vietnam and China: *Aculamprotula nodulosa*, *Lamprotula contritus*, *Lamprotula quadrangulosa*, and *Gibbosula crassa*. These species have been very scarce in the past several decades. *Lamprotula blaisei*, *Conradens fultoni*, *Pseudobaphia* sp., and *Cuneopsis demangei* have not been seen or collected since the early 1970s. No shells or live specimens of these last four taxa were collected during our fieldwork, strongly suggesting that these taxa are extirpated from their former range in Vietnam. The first three species are endemic to Vietnam and they may already be extinct.

Impacts

Freshwater mussels are long-lived animals (10 to 200 yr) and have a unique life cycle with a parasitic larval stage on the gills or fins of fish, so negative impacts on the species may not be immediately apparent (Bogan 1993, 2006; Vaughn and Taylor 1999; Vaughn 2010; Haag 2012). Mussels provide a variety of ecosystem services that directly and indirectly impact the local human populations including biofiltration, food source for animals and humans, and resource materials (e.g., inlay materials) (Vaughn 2017). Negative impacts on freshwater mussels are numerous, including channel modification and habitat destruction from dredging; sedimentation; clear-cutting of watersheds; monoculture cropping; loss of riparian buffers along streams; pollution in many forms including fertilizers, pesticides, industrial effluents, and domestic sewage; mining; urbanization and damming of rivers; commercial exploitation; introduced species; expansion of nonnative parasite hosts; and the loss of native host fish (e.g., Dudgeon et al. 2006; Gillis 2012; Haag 2012; Vaughn 2012; Sousa et al. 2014; Zieritz et al. 2016). There is no research on the impacts of current agricultural and forestry practices on the unionid fauna of Vietnam; however, research in Malaysia (Zieritz et al. 2017) has indicated that such activities have had a negative effect on the distribution of freshwater mussels.

Many impacts on the freshwater environment were visible during our surveys including deforestation, road construction (Fig. 5), in-stream sand and gravel mining (Fig. 6), open-pit



Figure 5. Road construction and disposal of debris over the edge of the road and the impact on the local river; the river is the same color as the earth being dumped. Lang Son Province, Vietnam. Photograph by Arthur Bogan. November 11, 2012.

mining runoff (Fig. 7), harvesting for food (Fig. 2), domestic pollution, and construction of various dams and hydroelectric projects (Fig. 8). Streams have been locally modified by restriction of channels, diversion of water for rice fields, and terracing of flood plains for rice production. Most of the large trees have been cut from the mountains, affecting the rain runoff patterns and thus affecting water temperature and clarity and increasing runoff (see Naiman and Dudgeon 2011; Zieritz et al. 2016). These modifications are impacting freshwater mussels and gastropod populations. A few species are doing well in disturbed habitats and show up in the markets as food items, including *Sinanodonta* spp., *Cristaria plicata*, *Pletholophus tenuis*, and *Nodularia* spp. (V.T. Do, personal observations).

Current and planned dams in the Mekong River basin and their impacts have been summarized by Winemiller et al. (2016). Dams and reservoirs not only impact the mussel and



Figure 6. In-stream gravel mining, Dien Bien Province, Vietnam. Photograph by Arthur Bogan. November 19, 2012.



Figure 7. Open-pit mining, Cao Bang Province, Vietnam. Photograph by Arthur Bogan. November 13, 2012.

fish fauna in the footprint of the reservoir but also below the reservoir. Downstream of the dam (Fig. 8) water flow patterns, water temperature, and chemistry are changed and can dramatically impact mussels, their biology, and native host fish (Vaughn and Taylor 1999).

Mussels are used a food item and are intensively collected in some areas of Vietnam. A local mussel harvester commented that currently he is only able to collect about 100 kg per day where in the past he was collecting 500 to 800 kg per day to sell in local markets. This decline in harvest may be due to overharvesting or mussel population declines due to a combination of overharvesting, disturbance, and pollution. Consider the number of people across Vietnam collecting mussels each day and this impact on a yearly basis is staggering. Decline in the mussel fauna of Vietnam can be considered on a local scale as well as considering the national impact on the freshwater bivalve fauna. These impacts will also impact those species that range beyond Vietnam's borders.



Figure 8. Dam in central Vietnam with the riverbed downstream dewatered. Photograph by Arthur Bogan. November 18, 2014.

Conservation Recommendations

In addition to established national parks and conservation areas, beginning in 2008 and extending to 2020, Vietnam will establish 45 areas for protection of inland water bodies. Some parts of the Red and Da rivers (in northern Vietnam) will be included in protected areas. However, conservation activities for freshwater bivalves have never been mentioned. About 60% of the Vietnamese freshwater bivalve fauna is currently imperiled. Some freshwater bivalve species already appear to have disappeared from northern Vietnam and the future is bleak for endangered freshwater bivalve species.

New outreach materials need to be developed for distribution to local people, aquaculture agencies, primary and high schools, and universities which illustrate the Vietnamese freshwater bivalve fauna and their unique life cycle. Aquaculture programs can be encouraged to develop captive propagation programs that will identify the native host fish for species to assist in conservation. Public activities and education on the importance and role of freshwater bivalves have not been mentioned. Some of freshwater bivalve species already seem to have disappeared from northern Vietnam and there is reason to be pessimistic about the future of endangered freshwater bivalve species.

Knowledge of the taxonomy of marine mollusks is actively expanding due to activities of the amateur shell collectors who are regularly contributing to the description of new species (see Bouchet et al. 2016). Developing a public program not only to provide information on freshwater mollusks of Vietnam, but also a website to provide free information or assist with identifications, is needed. This would stimulate more public involvement in examining the local freshwater fauna. Shell clubs are common in the United States and Europe and there are already shell dealers and people interested in shells in Vietnam (see Thach 2005, 2007, 2012, 2016b). These people would be a great source of information and local knowledge about these animals and should be encouraged to participate. A template might be the Cornell FeederWatch project.

Future Research Needs

Our understanding of the phylogenetic relationships of the freshwater mussels of Vietnam is only beginning to be explored (e.g., Lopes-Lima et al. 2017). Using results of our expanding molecular analyses, we have recognized a small-sized species, *Nodularia nuxpersica*, new to Vietnam, and one undescribed new species, *Pletholophus* sp. By comparing photographs and museum collections, we have recognized a misidentified species as new, *Pseudobaphia* sp. If the results of the work in Malaysia are any indication, some of the currently recognized Vietnamese unionid species may be overturned as some species are being recognized as a species complex (e.g., *Hyriopsis bialatus*) (Zieritz et al. 2016). This effort will require continued collaboration with colleagues throughout Asia.

ACKNOWLEDGMENTS

We want to express our gratitude for the support of the Institute of Ecology and Biological Resources (IEBR), Vietnam Academy of Science and Technology (VAST), and North Carolina Museum of Natural Sciences for their support of this collaborative research. We also acknowledge the help of our assistants, Mr. Quoc Toan Phan, Nguyen Tong Cuong, Nguyen Dinh Tao, and our drivers, Mr. Nguyen Ngoc Quyen, who was our driver for three trips, and Mr. Dang Thanh Ha, who was our driver for the 2016 trip. We also appreciate the Nagao Natural Environment Foundation support for Do Van Tu's research project "The study of Unionoida biodiversity and conservation status in Bang and Ky Cung River basins in Northeast Vietnam." Ms. Jamie M. Smith and Mrs. Heather B. Leslie have been instrumental in databasing specimens deposited in the North Carolina Museum of Natural Sciences Mollusk Collection from Vietnam. Mrs. Cynthia M. Bogan and Ms. Jamie M. Smith have reviewed the manuscript for content and English composition. We thank our anonymous reviewers for their very helpful and thorough comments and suggestions; they have been very helpful.

LITERATURE CITED

- Allen, D. J., K. G. Smith, and W. R. T. Darwall (compilers). 2012. The Status and Distribution of Freshwater Biodiversity in Indo-Burma. IUCN, Cambridge, United Kingdom, and Gland, Switzerland. 158 pp.
- Araujo, R., S. Schneider, K. J. Roe, D. Erpenbeck, and A. Machordom. 2017. The origin and phylogeny of Margaritiferidae (Bivalvia, Unionoida): A synthesis of molecular and fossil data. *Zoologica Scripta* 46:289–307. doi: 10.1111/zsc.12217
- Bavay, A., and P. Dautzenberg. 1901. Description de deux Unio et d'un Corbicula nouveaux, provenant de l'Indo-Chine. *Journal de conchyliologie* 49:5–9, plate 1, figures 1–7.
- Bogan, A. E. 1993. Freshwater bivalve extinctions: Search for a cause. *American Zoologist* 33:599–609.
- Bogan, A. E. 2006. Conservation and extinction of the freshwater molluscan fauna of North America. Chapter 30. Pages 373–383 in C.F. Sturm, T. A. Pearce, and A. Valdes, editors. *The Mollusks: A Guide to Their Study, Collection, and Preservation*. American Malacological Society, Universal Publishers, Inc., Boca Raton, Florida.
- Bogan, A. E. 2008. Global diversity of freshwater mussels (Mollusca, Bivalvia) in freshwater. *Global diversity of freshwater animals*. *Hydrobiologia* 595:139–147.
- Bogan, A. E. 2015. Determining the date of publication for *Conradens* Haas and *Uniandra* Haas (Bivalvia; Unionidae). *The Nautilus* 129:175–178.
- Bogan, A. E., and V. T. Do. 2011. Research note: Clarification of the authorship and date of publication of three Asian species of Unionidae (Bivalvia): *The Nautilus* 125:171–172.
- Bogan, A. E., and V. T. Do. 2013a. Field research on the distribution of freshwater bivalves in northern Vietnam, November 2012. *Ellipsaria* 15:13–14.
- Bogan, A. E., and V. T. Do. 2013b. Conservation assessment of freshwater bivalves in northern Vietnam, 2012. *Tentacle* 21:10–20.
- Bogan, A. E., and V. T. Do. 2014a. Freshwater bivalve survey of Vietnam, Part II: Central Highlands and the Mekong Delta area. *Ellipsaria* 16:29–31.
- Bogan, A. E., and V. T. Do. 2014b. Two freshwater bivalve species new to the fauna of Vietnam (Mollusca: Bivalvia: Arcidae and Unionidae). *Tropical Natural History* 14:113–116.

- Bogan, A. E., and V. T. Do. 2016. Fieldtrip to northern Vietnam, 2016. *Ellipsaria* 18:27–29.
- Bolotov, I. N., Y. V. Bespalaya, M. Y. Gofarov, A. V. Kondakov, E. S. Konopleva, and I. V. Vikhrev. 2016a. Spreading of the Chinese pond mussel, *Sinanodonta woodiana*, across Wallacea: one or more lineages invade tropical islands and Europe. *Biochemical Systematics and Ecology* 67:58–64.
- Bolotov, I. N., I. V. Vikhrev, Y. Bespalaya, V. Artamonova, M. Gofarov, J. Kolosova, A. Kondakov, A. Makhrov, A. Frolov, S. Tumpeesuwan, A. Lyubas, T. Romanis, and K. Titova. 2014. Ecology and conservation of the endangered Indochinese freshwater pearl mussel, *Margaritifera laosensis* (Lea, 1863) in the Nam Pe and Nam Long rivers, Northern Laos. *Tropical Conservation Science* 7:706–719.
- Bolotov, I. N., I. V. Vikhrev, Y. V. Bespalaya, M. Y. Gofarov, A. V. Kondakov, E. S. Konopleva, N. N. Bolotov, and A. A. Lyubas. 2016b. Multi-locus fossil-calibrated phylogeny, biogeography and a subgeneric revision of the Margaritiferidae (Mollusca: Bivalvia: Unionoida). *Molecular Phylogenetics and Evolution* 103:104–121.
- Bolotov, I. N., I. V. Vikhrev, A. V. Kondakov, E. S. Konopleva, M. Y. Gofarov, O. V. Aksenova, and S. Tumpeesuwan. 2017. New taxa of freshwater mussels (Unionidae) from a species-rich but overlooked evolutionary hotspot in Southeast Asia. *Scientific Reports* 7:11573. doi:10.1038/s41598-017-11957-9
- Bouchet, P., S. Bary, V. Héros, and G. Marani. 2016. How many species of molluscs are there in the world oceans and who is going to describe them? Pages 9–24 in V. Héros, E. Strong, and P. Bouchet, editors. *Tropical Deep-Sea Benthos* 29. Muséum national d'Histoire naturelle, Paris.
- Brandt, R. A. M. 1974. The non-marine aquatic Mollusca of Thailand. *Archiv für Molluskenkunde* 105:1–423.
- Breton, S., H. Doucet-Beaupré, D. T. Stewart, W. R. Hoeh, and P. U. Blier. 2007. The unusual system of doubly uniparental inheritance of mtDNA: Isn't one enough? *Trends in Genetics* 23:465–474. doi: 10.1016/j.tig.2007.05.011
- Breton, S., D. T. Stewart, S. Shephardson, R. J. Trdan, A. E. Bogan, E. G. Chapman, A. J. Ruminas, H. Piontkivska, and W. R. Hoeh. 2010. Novel protein genes in animal mtDNA: A new sex determination system in freshwater mussels (Bivalvia: Unionoida)? *Molecular Biology and Evolution* 28:1645–1659. doi: 10.1093/molbev/msq345
- Đặng N. T., and T. H. Hồ (in press). *Freshwater Bivalves and Snails of Vietnam* (Mollusca: Gastropoda, Bivalvia). Publishing House for Science and Technology, Hanoi, Vietnam.
- Đặng, N. T., T. B. Thái, and V. M. Pham. 1980. Identification of freshwater invertebrates of North Vietnam. Sciences and Technology Publishing Co., Ha Noi, Vietnam. 573 pp. In Vietnamese: Định loại động vật không xương sống nước ngọt Bắc Việt Nam.
- Dautzenberg, P., 1900. Diagnose d'une nouvelle espèce d'Unio provenant de l'Indo-Chine. *Journal de conchyliologie* 48:429.
- Dautzenberg, P., and H. Fischer. 1906a. Contribution a la fauna malacologique de l'Indo-Chine. *Journal de conchyliologie*. 54:145–226, plates 5–7.
- Dautzenberg, P., and H. Fischer. 1906b. Liste des Mollusques récoltés par M. H. Mansuy en Indo-Chine et au Yunnan et description d'espèces nouvelles. *Journal de conchyliologie* 53:343–471, plates 8–10.
- Dautzenberg, P., and H. Fischer. 1908. Liste des Mollusques récoltés par M. Mansuy en Indo-Chine et description d'espèces nouvelles II. *Journal de conchyliologie* 56:169–217, plates 4–8.
- Davidson, S. P., T. Kunpradid, Y. Peerapornisai, T. M. L. Nguyen, B. Pathoumthong, C. Vongsambath, and A. D. Pham. 2006. Biomonitoring of the Lower Mekong and selected tributaries. MRC Technical Paper No. 13. Mekong River Commission, Vientiane, Lao PDR. 100 pp.
- Doucet-Beaupré, H., S. Breton, E. G. Chapman, P. U. Blier, A. E. Bogan, D. T. Stewart, and W. R. Hoeh. 2010. Mitochondrial phylogenomics of the Bivalvia (Mollusca): Searching for the origin and mitogenomic correlates of doubly uniparental inheritance of mtDNA. *BMC Evolutionary Biology* 10:50. doi:10.1186/1471-2148-10-50
- Dudgeon, D., A. H. Arthington, M. O. Gessner, Z.-I. Kawabata, D. J. Knowler, C. Lévêque, R. J. Naiman, A.-H. Prieur-Richard, D. Soto, M. L. J. Stiassny, and C. A. Sullivan. 2006. Freshwater biodiversity: Importance, threats, status and conservation challenges. *Biological Reviews* 81:163–182.
- Froufe, E., H. M. Gan, Y. P. Lee, J. Carneiro, S. Varandas, A. Teixeira, A. Zieritz, R. Sousa, and M. Lopes-Lima, 2015. The male and female complete mitochondrial genome sequences of the endangered freshwater mussel *Potomida littoralis* (Cuvier, 1798) (Bivalvia: Unionidae). *Mitochondrial DNA* 27:3571–3572. doi:10.3109/19401736.2015.1074223
- Gillis, P. L. 2012. Cumulative impacts of urban runoff and municipal wastewater effluents on wild freshwater mussels (*Lasmigona costata*). *Science of the Total Environment* 431:348–356.
- Gozhik, P. F. 1978. Fossil limnoscaphs in the southern Ukraine. *Malacological Review* 11:147–148.
- Graf, D. L. 2007. Palearctic freshwater mussel (Mollusca: Bivalvia: Unionoida) diversity and the Comparative Method as a species concept. *Proceedings of the Academy of Natural Sciences of Philadelphia* 156:71–88.
- Graf, D. L. 2013. Patterns of freshwater bivalve global diversity and the state of phylogenetic studies on the Unionoida, Sphaeriidae, and Cyrenidae. *American Malacology Bulletin* 31:135–153.
- Graf, D. L., and K. S. Cummings. 2006. Palaeoheterodont diversity (Mollusca: Trigonoida + Unionoida): What we know and what we wish we knew about freshwater mussel evolution. *Zoological Journal of the Linnean Society, London* 148:343–394.
- Graf, D. L., and K. S. Cummings. 2007. Review of the systematics and global diversity of freshwater mussel species (Bivalvia: Unionoida). *Journal of Molluscan Studies* 73:291–314.
- Graf, D. L., and K. S. Cummings. 2017. *Musselp. The Freshwater Mussels (Unionoida) of the World (and other less consequential bivalves)*. Available at <http://mussel-project.uwsp.edu/> (accessed January 3–May 10, 2017).
- Griffith, E., and E. Pidgeon, [1833]–1834, *The Mollusca and Radiata*. Vol. 12, in E. Griffith, editor [1824]–1835, *The Animal Kingdom Arranged in Conformity with its Organization*, by the Baron Cuvier, member of the Institute of France, &c. &c. with supplementary additions to each order, by Edward Griffith, F.L.S., A.S., corresponding member of the Academy of Natural Sciences of Philadelphia, &c. and others. Whittaker and Co., London.
- Haag, W. R. 2012. *North American Freshwater Mussels: Natural History, Ecology and Conservation*. Cambridge University Press, Cambridge, United Kingdom. 505 pp.
- Haag, W. R., and J. D. Williams. 2014. Biodiversity on the brink: An assessment of conservation strategies for North American freshwater mussels. *Hydrobiologia* 735:45–60.
- Haas, F. 1910a. New Unionidae from East Asia. *The Annals and Magazine of Natural History* (8th series) 6:496–499.
- Haas, F. 1910b. Neue Najaden. *Nachrichtsblatt der Deutschen Malakozoologischen Gesellschaft* 4:97–103.
- Haas, F. 1910c–1920. Die Unioniden. Neubearbeitung und Fortsetzung der Küsterschen und Clessinschen Monographien von *Unio* und *Anodonta*. Volume 9 (pt. 2, section 2), pages 1–344, plates 1–73, 12a; 15 text figures in H. C. Küster, editor. *Systematisches Conchylien-Cabinet von Martini und Chemnitz*. Bauer und Raspe, Nürnberg.
- Haas, F. 1913. Neue Najadengattungen. *Nachrichtsblatt der Deutschen*

- Malakozoologischen Gesellschaft 45:33–38. Date published January 27, 1913 *vide* Solem 1967:83
- Haas, F. 1923. Beiträge zu einer Monographie der asiatischen Unioniden. Sonderabdruck aus den Abhandlungen der Senkenbergischen Naturforschenden Gesellschaft 38:129–203, plates 15–16.
- Haas, F. 1969a. Superfamilia Unionacea. Das Tierreich (Berlin). Lieferung 88:i–x,1–663.
- Haas, F. 1969b. Superfamily Unionacea. Pages N411–N471 in R. C. Moore, editor. Treatise on Invertebrate Paleontology. Part N, Volume 1. Mollusca 6. Bivalvia. Geological Society of America and the University of Kansas, Lawrence, Kansas.
- He, J., and Z. Zhuang. 2013. The Freshwater Bivalves of China. ConchBooks, Harxheim, Germany. 198 pp.
- Heude, P. 1874. Diagnoses Molluscorum in fluminibus provinciae Nankingensis collectorum. Journal de conchyliologie 22:112–118.
- Heude, P. 1875–1885. Conchyliologie fluviatile de la province de Nanking et de la Chine centrale. Librairie F. Savy, Paris. 10 Fascicules, plates and unnumbered pages.
- Hoeh, W. R., M. B. Black, R. Gustafson, A. E. Bogan, R. A. Lutz, and R. C. Vrijenhoek. 1998. Testing alternative hypotheses of *Neotrigonia* (Bivalvia: Trigonioidea) phylogenetic relationships using cytochrome C oxidase subunit 1 DNA sequences. Malacologia 40:267–278.
- Hoeh, W. R., A. E. Bogan, K. S. Cummings, and S. I. Guttman. 2002. Evolutionary relationships among the higher taxa of freshwater mussels (Bivalvia: Unionoida): Inferences on phylogeny and character evolution from analyses of DNA sequence data. Malacological Review 31/32:123–141.
- Hoeh, W. R., A. E. Bogan, and W. H. Heard. 2001. A phylogenetic perspective on the evolution of morphological and reproductive characteristics in the Unionoida. Pages 257–280 in G. Bauer and K. Wächtler, editors. Ecology and Evolutionary Biology of Freshwater Mussels, Unionoida. Ecological Studies Vol. 145. Springer Verlag, Berlin.
- ICZN (International Commission on Zoological Nomenclature). 1999. International Code of Zoological Nomenclature. 4th ed. Incorporating Declaration 44, amendments of Article 74.7.3, with effect from December 31, 1999, and the Amendment on e-publication, amendments to Articles 8, 9, 10, 21 and 78, with effect from January 1, 2012. International Trust for Zoological Nomenclature, London. 306 pp. Available at <http://www.nhm.ac.uk/hosted-sites/iczn/code/> (accessed September 11, 2015).
- IUCN (International Union for Conservation of Nature and Natural Resources). 2012. IUCN Red List Categories and Criteria: Version 3.1. 2nd ed. IUCN, Gland, Switzerland and Cambridge, United Kingdom. 32 pp. Available at http://cmsdocs.s3.amazonaws.com/keydocuments/Categories_and_Criteria_en_web%2Bcover%2Bbckcover.pdf (accessed November 18, 2016).
- IUCN. 2013. Documentation standards and consistency checks for IUCN Red List assessments and species accounts. Version 2. Adopted by the IUCN Red List Committee and IUCN SSC Steering Committee. IUCN, Gland, Switzerland and Cambridge, United Kingdom. 65 pp. Available at http://cmsdocs.s3.amazonaws.com/keydocuments/RL_Standards_Consistency.pdf (accessed November 18, 2016).
- IUCN. 2016. Red List of threatened species. Edition 2016-2. Available at <http://www.iucnredlist.org/> (accessed June–November 2016).
- IUCN. 2017. IUCN Assessment Process for adding species to the IUCN Red List. Edition 2017-3. Available at <http://www.iucnredlist.org/technical-documents/assessment-process> (accessed December 21, 2017).
- IUCN Standards and Petitions Subcommittee. 2016. Guidelines for using the IUCN Red List Categories and Criteria. Version 12. Prepared by the Standards and Petitions subcommittee. International Union for Conservation of Nature and Natural Resources. Gland, Switzerland and Cambridge, United Kingdom. 101 pp. Available at <http://cmsdocs.s3.amazonaws.com/RedListGuidelines.pdf> (accessed November 19, 2016).
- Klishko, O., M. Manuel Lima-Lopez, E. Froufe, and A. E. Bogan. 2014. Are *Cristaria herculea* (Middendorff, 1847) and *Cristaria plicata* (Leach, 1815) (Bivalvia, Unionidae, Unioninae, Anodontini) separate species? Zookeys 438:1–15. doi:10.3897/zookeys.438.7493
- Klishko, O., M. Lopes-Lima, E. Froufe, A. E. Bogan, and V. Y. Abakumova. 2016. Systematics and distribution of *Cristaria plicata* (Bivalvia, Unionidae) from the Russian Far East. Zookeys 580:13–27. doi:10.3897/zookeys.580.7588
- Köhler, F., M. Seddon, A. E. Bogan, V. T. Do, P. Sri-Aroon, and D. Allen. 2012. The status and distribution of freshwater molluscs in the Indo-Burma region. Pages 66–88 in D. J. Allen, K. G. Smit, and W. R. T. Darwall, compilers. The Status and Distribution of Freshwater Biodiversity in Indo-Burma. IUCN, Cambridge, United Kingdom, and Gland, Switzerland.
- Kondo, T. 2008. Monograph of Unionoida in Japan (Mollusca: Bivalvia). Special Publication of the Malacological Society of Japan, Tokyo. No. 3, 69 pp.
- Lea, I. 1836. A Synopsis of the Family of Naiades. Cary, Lea and Blanchard, Philadelphia; John Miller, London. 59 pp., 1 plate, colored.
- Lea, I. 1838. A Synopsis of the Family of Naiades. 2nd ed., enlarged and improved. Carey, Lea and Blanchard, Philadelphia. 44 pp.
- Lea, I. 1852. A Synopsis of the Family of Naiades. 3rd ed., greatly enlarged and improved. Blanchard and Lea, Philadelphia. 88 pp.
- Lea, I. 1870. A Synopsis of the Family Unionidae. 4th ed., very greatly enlarged and improved. Henry C. Lea, Philadelphia. 184 pp.
- Lopes-Lima, M., E. Froufe, V. T. Do, M. Ghamizi, K. E. Mock, Ü. Kebapçı, O. Klishko, S. Kovitvadhi, U. Kovitvadh, O. S. Paulo, J. M. Pfeiffer III, M. Raley, N. Riccardi, H. Şereflişan, R. Sousa, A. Teixeira, S. Varandas, X.-P. Wu, D. T. Zanatta, A. Zieritz, and A. E. Bogan. 2017. Phylogeny of the most species-rich freshwater bivalve family (Bivalvia: Unionida: Unionidae): Defining modern subfamilies and tribes. Molecular Phylogenetics and Evolution 106:174–191. doi 10.1016/j.ympev.2016.08.021
- Lydeard, C., R. H. Cowie, W. F. Ponder, A. E. Bogan, P. Bouchet, S. A. Clark, K. S. Cummings, T. J. Frest, O. Gargominy, D. G. Herbert, R. Hershler, K. E. Perez, B. Roth, M. Seddon, E. E. Strong and F. G. Thompson. 2004. The global decline of nonmarine mollusks. BioScience 54:321–330.
- Mabille, J. 1887. Sur quelques Mollusques du Tonkin. Bulletins de la Société Malacologique de France 4:73–164, plates 1–4.
- Martens, E. V. 1902. Neue Unioniden aus Tonkin und Anam. Nachrichtenblatt der Deutschen Malakozoologischen Gesellschaft 34:130–135.
- Modell, H. 1942. Das natürliche system der Najaden. Archiv für Molluskenkunde 74:161–191.
- Modell, H. 1949. Das natürliche system der Najaden. 2. Archiv für Molluskenkunde 78:29–46.
- Modell, H. 1950. Tertiäre Najaden IV. Archiv für Molluskenkunde 79:25–36.
- Modell, H. 1964. Das natürliche system der Najaden. 3. Archiv für Molluskenkunde 93:71–126.
- Morelet, A. 1865. Rectifications et additions à la fauna malacologique de l'Indo-Chine. Journal de conchyliologie 13:19–23, 225–228.
- Morelet, A. 1866. Description d'espèces appartenant à la fauna malacologique de l'Indo-Chine. Journal de conchyliologie 14:62–64.
- Morlet, L. 1886a. Diagnoses Molluscorum novorum Tonkini. Journal de conchyliologie 34:75–78.
- Morlet, L. 1886b. Liste des Coquilles recueillies, au Tonkin, par M. Jourdy, chef d'escadron d'artillerie, et description d'espèces nouvelles. Journal de conchyliologie 34:257–295, plates 12–15.
- Morlet, L., 1891. Contributions à la faune malacologique de l'Indo-Chine. Journal de Conchyliologie 39:230–254, plates 5, 7.
- Naiman, R. J., and D. Dudgeon. 2011. Global alteration of freshwaters:

- Influences on human and environmental well-being. *Ecological Research* 26:865–873.
- Neves, R. J., A. E. Bogan, J. D. Williams, S. A. Ahlstedt, and P. D. Hartfield. 1998. Status of aquatic mollusks in the southeastern United States: A downward spiral of diversity. Pages 43–86 in G. W. Benz and D. E. Collins, editors. *Aquatic Fauna in Peril: The Southeastern Perspective*. Special Publication 1. Southeast Aquatic Research Institute, Lenz Design & Communications, Decatur, Georgia. 554 pp.
- Ouyang, J. X., X.-P. Wu, S. Ouyang, S. Li, and D. Zhao. 2011. Phylogenetic analysis of some Chinese freshwater Unionidae based on mitochondrial COI sequences. *Journal of Conchology* 40:543–548.
- Petit, R. E., and E. V. Coan. 2008. The molluscan taxa made available in the Griffith and Pidgeon (1833–1834) edition of Cuvier, with notes on the editions of Cuvier and on Wood's *Index Testaceologicus*. *Malacologia* 50:219–264.
- Pfeiffer, J. M. III, and D. L. Graf. 2013. Re-analysis confirms the polyphyly of *Lamprotula* Simpson, 1900 (Bivalvia: Unionida). *Journal of Molluscan Studies* 79:249–256.
- Pfeiffer, J. M. III, and D. L. Graf. 2015. Evolution of bilaterally asymmetrical larvae in freshwater mussels (Bivalvia: Unionida: Unionidae). *Zoological Journal of the Linnean Society* 175:307–318. doi: 10.1111/zoj.12282
- Phuong, T. D. N. 2011. *Margaritifera laosensis*. IUCN 2011. 2011 IUCN Red List of Threatened Species. Version 2011.2. Available at <http://www.iucnredlist.org> (accessed July 3, 2016).
- Red Data Book of Vietnam. 1992. Volume 1. Animals. Science and Technics Publishing House, Ha Noi. 396 pp. In Vietnamese.
- Red Data Book of Vietnam. 2000. Volume 1. Animals. Publishing House for Science and Technology, Ha Noi, 408 pp. In Vietnamese.
- Rochebrune, A.-T. 1904a. Monographie du genre *Harmandia*. *Bulletin du Muséum d'Histoire Naturelle* 10:138–141.
- Rochebrune, A.-T. 1904b. Essai monographique sur le Genre *Chamberlainia* Simpson. *Bulletin du Muséum d'Histoire Naturelle* 10:463–466.
- Roe, K. J., and W. R. Hoeh. 2003. Systematics of freshwater mussels (Bivalvia: Unionida). Pages 91–122 in C. Lydeard and D. R. Lindberg, editors. *Molecular Systematics and Phylogeography of Mollusks*. Smithsonian Series in Comparative Evolutionary Biology. Smithsonian Books, Washington, D.C.
- Rolle, H. 1904. Drei neue Najadeen aus Tonkin. *Nachrichtsblatt der Deutschen Malakozoologischen Gesellschaft* 36:25–26, plates 1–4.
- Scudder, N. P. 1885. The Published Writings of Isaac Lea, LL.D. *Biographies of American Naturalists II*. Bulletin of the U.S. National Museum, Number 23. Government Printing Office, Washington, DC. 278 pp.
- Simpson, C. T. 1900. Synopsis of the naiades, or pearly freshwater mussels. *Proceedings of the U.S. National Museum* 22:501–1044.
- Simpson, C. T. 1914. A Descriptive Catalogue of the Naiades, or Pearly Freshwater Mussels. Parts I–III. Bryant Walker, Detroit. 1,540 pp.
- Solem, A. 1967. New molluscan taxa and scientific writings of Fritz Haas. *Fieldiana: Zoology* 53:71–144.
- Sousa, R., A. Novais, R. Costa, and D. L. Strayer. 2014. Invasive bivalves in fresh waters: Impacts from individuals to ecosystems and possible control strategies. *Hydrobiologia* 735:233–251. doi 10.1007/s10750-012-149-1
- Starobogatov, Ya. I. 1970. Fauna Mollyuskov i Zoogeograficheskoe Raionirovanie Kontinental'nykh Vodoemov Zemnogo Shara [Mollusk Fauna and Zoogeographical Partitioning of Continental Water Reservoirs of the World]. Nauka, Leningrad. 372 pp., 39 figures, 12 tables. In Russian.
- Strayer, D. L., and D. Dudgeon. 2010. Freshwater biodiversity conservation: Recent progress and future challenges. *Journal of the North American Benthological Society* 29:344–358.
- Thach, N. N. 2005. *Shells of Vietnam*. ConchBooks, Hackenheim, Germany. 338 pp., 91 color plates.
- Thach, N. N. 2007. *Recently Collected Shells of Vietnam*. L'Informatore Piceno and N.N.T, Ancona, Italy. 384 pp., 118 color plates.
- Thach, N. N. 2012. *New Records of Molluscs from Vietnam*. 48HrBooks Company, Akron, Ohio. 276 pp., 151 color plates.
- Thach, N. N. 2016a. *Lanceolaria bogani* (Bivalvia: Unionida), a new species from Vietnam *Novapex* 17(1):9–11.
- Thach, N. N. 2016b. *Vietnamese New Mollusks, Seashells, Landsnails, Cephalopods with 59 New Species*. 48HrBooks Company, Akron, Ohio. 205 pp., 81 color plates.
- Vaughn, C. C. 2010. Biodiversity losses and ecosystem function in freshwaters: Emerging conclusions and research directions. *BioScience* 60:25–35.
- Vaughn, C. C. 2012. Life history traits and abundance can predict local colonization and extinction rates of freshwater mussels. *Freshwater Biology* 57:982–992.
- Vaughn, C. C. 2017. Ecosystem services provided by freshwater mussels. *Hydrobiologia*. doi 10.1007/s10750-017-3139-x
- Vaughn, C. C., and C. M. Taylor. 1999. Impoundments and the decline of freshwater mussels: A case study of an extinction gradient. *Conservation Biology* 13:912–920.
- Vietnam Red Data Book 2007. Part I. Animals. Publishing House for Science and Technology, Ha Noi. 515 pp. In Vietnamese.
- Whelan, N. V., A. J. Geneva, and D. L. Graf. 2011. Molecular phylogenetic analysis of tropical freshwater mussels (Mollusca: Bivalvia: Unionida) resolves the position of *Coelatura* and supports a monophyletic Unionida. *Molecular Phylogenetics and Evolution* 61:504–514.
- Winemiller, K. O., P. B. McIntyre, L. Castello, E. Fluet-Chouinard, T. Giarrizzo, S. Nam, I. G. Baird, W. Darwall, N. K. Lujan, I. Harrison, M. L. J. Stiassny, R. A. M. Silvano, D. B. Fitzgerald, F. M. Pelicice, A. A. Agostinho, L. C. Gomes, J. S. Albert, E. Baran, M. Petrere, Jr., C. Zarfl, M. Mulligan, J. P. Sullivan, C. C. Arantes, L. M. Sousa, A. A. Koning, D. J. Hoeninghaus, M. Sabaj, J. G. Lundberg, J. Armbruster, M. L. Thieme, P. Petry, J. Zuanon, G. Torrente Vilara, J. Snoeks, C. Ou, W. Rainboth, C. S. Pavanelli, A. Akama, A. van Soesbergen, and L. Sáenz. 2016. Balancing hydropower and biodiversity in the Amazon, Congo and Mekong. *Science* 351:128–129.
- Wu, X.-P. 1998. Studies on freshwater Mollusca in mid-lower reaches of the Yangtze (Chang Jiang) River. Doctoral Dissertation in Science. Institute of Hydrobiology, Chinese Academy of Sciences. Wuhan, Hubei Province, China. 197 pp.
- Zhou, C.-H., S. Ouyang, X.-P. Wu, and M. Li. 2007. Phylogeny of the genus *Lamprotula* (Unionida) in China based on mitochondrial DNA sequences of 16S rRNA and ND1 genes. *Acta Zoologica Sinica* 53:1024–1030.
- Zieritz, A., A. E. Bogan, E. Froufe, A. Klishko, T. Kondo, U. Kovitvadhi, S. Kovitvadhi, J. H. Lee, M. Lopes-Lima, J. M. Pfeiffer, R. Sousa, V. T. Do, I. Vikhrev, and D. T. Zanatta. 2017. Diversity, biogeography and conservation of freshwater mussels (Bivalvia: Unionida) in East and Southeast Asia. *Hydrobiologia*. doi 10.1007/s10750-017-3104-8
- Zieritz, A., M. Lopes-Lima, A. E. Bogan, R. Sousa, S. Walton, K. A. A. Rahim, J.-J. Wilson, P.-Y. Ng, E. Froufe, and S. McGowan. 2016. Factors driving changes in freshwater mussel (Bivalvia, Unionida) diversity and distribution in Peninsular Malaysia. *Science of the Total Environment*. 571:1069–1078. doi 10.1016/j.scitotenv.2016.07.098