

Notes on *Cyperus* sect. *Incurvi* (Cyperaceae) from the New World Tropics

Author: Tucker, Gordon C.

Source: *Willdenowia*, 44(2) : 253-261

Published By: Botanic Garden and Botanical Museum Berlin (BGBM)

URL: <https://doi.org/10.3372/wi.44.44207>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

GORDON C. TUCKER¹

Notes on *Cyperus* sect. *Incurvi* (*Cyperaceae*) from the New World Tropics

Abstract

Tucker G. C.: Notes on *Cyperus* sect. *Incurvi* (*Cyperaceae*) from the New World Tropics – Willdenowia 44: 253–261. 2014. – Version of record first published online on 8 July 2014 ahead of inclusion in August 2014 issue; ISSN 1868-6397; © 2014 BGBM Berlin-Dahlem.

DOI: <http://dx.doi.org/10.3372/wi.44.44207>

Based on study of herbarium specimens of *Cyperus* sect. *Incurvi* Kük., three species are described from the lowlands of N South America and Mesoamerica: *C. conservator-davidii* G. C. Tucker is a new species occurring in several countries and is morphologically similar to *C. simplex* Kunth; *C. lacustris* Schrad. ex Nees is an overlooked species that has been confused with *C. miliifolius* Poepp. & Kunth; *C. altsonii* Kük. and *C. hoppiifolius* Uittien pertain to the same species, for which the former is the earlier name. An identification key is provided for all species of *C. sect. Incurvi* occurring in the New World.

Additional key words: taxonomy, biogeography, identification key, South America, Central America, Mesoamerica, Amazonia, Guianas, sedges

Introduction

The genus *Cyperus* L. includes about 800 species, occurring worldwide in tropical and warm-temperate regions (Tucker 1994, 2001). The distribution of individual species ranges from nearly cosmopolitan (e.g. *C. odoratus* L. and *C. squarrosus* L.) to regional and narrow endemics (Kükenthal, 1935–1936). In the New World, areas of high diversity and endemism include Mexico, the Greater Antilles and E Brazil (Alves & al. 2007; Tucker 2007, 2008). While examining unidentified South American specimens of *Cyperaceae* in the herbaria of the Missouri Botanical Garden, Harvard University and the Royal Botanic Gardens, Kew, as well as loans from several other institutions, a new species and an overlooked species were noted. Additionally, two validly published names were determined to represent the same taxon. All cited specimens have been seen by the author unless indicated otherwise.

These species find their place in *Cyperus* [subg. *Pycnostachyus* C. B. Clarke] sect. *Incurvi* Kük. This pan-tropical section occurs in wet lowland forests and com-

prises about 30 species, ten of which are found in the New World Tropics (Kükenthal 1935–1936). The plants are perennials with simple or capitate inflorescences, with a wingless rachilla and floral scales that are saccate basally with a prolonged incurved apex. Most species have a rough, granular or papillose achene surface and several have pseudopetiolate leaves (Kükenthal 1935–1936). A dichotomous identification key is provided for all species of this section occurring in the New World.

The current paper represents a contribution toward a taxonomic monograph of the c. 225 species of *Cyperus* occurring in the New World Tropics. The taxonomic section of this paper consists of three parts: the description of a new species related to *C. simplex* Kunth and *C. grandisimplex* C. B. Clarke; the recognition of an overlooked, widespread species, related to *C. miliifolius* Poepp. & Kunth; and, the status of *C. hoppiifolius* Uittien as a synonym of *C. altsonii* Kük.

Provisional IUCN conservation assessments are made for the three species discussed herein, using IUCN (2012) criteria.

¹ Department of Biological Sciences, Eastern Illinois University, Charleston, Illinois 61920, U.S.A.; e-mail: gctucker@eiu.edu

Results and Discussion

Cyperus conservator-davidii G. C. Tucker, **sp. nov.** — Fig. 1.

Holotype: French Guiana, Monts d'Arawa, savane-roche central, 8 Jul 2002, *J. J. de Granville & al.* 15119 (US; isotypes: B, G, K, NY).

Diagnosis — Affinis *Cyperus simplex* Kunth sed radiis brevioribus, squamis angustis, acheniis oblongo-obovoid-eis et foliis bracteisque ventraliter papillois differt.

Description — *Herbs* perennial, rhizomatous. *Culms* triquetrous, 10–22 cm tall, (0.7–)1–1.5 mm thick, scabrelate on angles just below bracts. *Leaves* 4–8, V-shaped to inversely W-shaped, 10–15 cm long, 3–4 mm wide, abaxial surface papillose distally, adaxial surface and margin scabrelate. *Inflorescence bracts* 5 or 6, ascendant at 45°–60°, V-shaped, 1–15 cm long, 1.5–4 mm wide, abaxial surface papillose, adaxial surface and margin scabrelate; *secondary bracts* absent. *Rays* 5–8, 4–10(–12) cm long (longest 0.45–0.8 × as long as culm), scabrelate on angles; *secondary rays* absent. *Heads* digitate, 2–3 cm wide. *Spikelets* 1–5, lanceolate, compressed, 12–14 mm long, (2.5–)3–4 mm wide; *rachilla* persistent, greenish, wingless, successive scale scars 1.1–1.5 mm apart. *Floral scales* 12–16, deciduous, whitish, somewhat reddish or greenish tinged especially on margin apically, lanceolate, (2.2–)2.5–2.9 mm long, 1–1.5 mm wide, smooth, laterally weakly 2-nerved, in middle 1–3-nerved, apical portion c. 0.2 mm long, incurved parallel to rachilla. *Stamen* 1; *filament* 2–3 mm long; *anther* c. 1.2 mm long, connective apex c. 0.1 mm long. *Styles* 1–1.2 mm long; *stigmas* 3, (1.7–)2–3 mm long. *Achenes* white (sometimes brownish or reddish brown), oblong-obovoid, trigonous, 0.8–1.2 mm long, 0.7–0.8(–0.9) mm wide, faces slightly concave, papillate, base broadly rounded to cuneate, apex rounded to sub-truncate, apiculate.

Distribution — S Mexico; Central America: Costa Rica, Panama; South America: Bolivia, Brazil, Colombia, Ecuador, French Guiana, Guyana, Peru, Suriname, Venezuela.

Habitat and ecology — Tropical lowland forests; 100–950 m (however, a single collection from 2000 m).

Proposed IUCN status — Least Concern (LC). Widely distributed; apparently not threatened. This species is known from more than 25 collections, over a wide range, with numerous recent collections.

Specimens examined — BOLIVIA: PANDO: 80 km SW of Cobija, Limoero, 21 Jul 1989, *M. Prescott & J. M. Cardiel* 13 (K).

BRAZIL: ACRE: Jardim Zoobotânico of Federal Univ. of Acre, 15 Oct 1980, *Lowrie & al.* 535 (EIU, NY); Mpio. Buari, basin of Rio Purus, Rio Antimari, Floresta Estadu-

al do Antimari, 8 Mar 1997, *D. C. Daly & al.* 9298 (EIU, NY). — RORAIMA: Posto Mucajaí, margin of Rio Mucajaí, 18 Mar 1971, *G. Prance & al.* 11087 (AAU, NY, P). COLOMBIA: ANTIOQUÍA: Mpio. Dabeiba, 32 km SE de Mutatá, en la vía Mutatá-Dabeiba, 5 Aug 1987, *Callejas & al.* 5090 (EIU, NY); Mpio. Nariño, Vereda Puente Linda, 12 Sep 1994, *Fonnegra, Estudiantes de Taxonomía Plant. Vasc.* 5047B (MO with *C. simplex*); Mpio. Urrao, Parque Nac. Natural Las Orquídeas, Vereda Calles, 8 Dec 1993, *J. Pipoly & al.* 17928 (MO); *ibid.*, 26 Jan 2011, *Pedras-Peñalosa & al.* 1968 (GH, NY); Mpio. Frontino, road between Nutibara and La Blanquita, region of lower Murri, 7 Feb 1989, *J. M. MacDougal & al.* 3697 (MO). — META: 5 km arriba de la desembocadura de Caño Cabra en el Río Guayabero, 22 Feb 1969, *P. Pinto E. & C. Sastre* 972 (COL, P).

COSTA RICA: HEREDIA: Colonia Virgen del Socorro, 3 Feb 1984, *J. Gómez-Laurito* 9870 (F); 3 miles S of Cariblanco, 760 m, 28 May 1976, *T. B. Croat* 35838 (MO). ECUADOR: IMBABURA: Lita, 25 Apr 1949, *M. Acosta-Solis* 12253 (US). — NAPO: along Río Cuyabeno, c. 1 hour in canoe below Puerto Bolívar, 26 Jan 1984, *S. Laegaard* 51242 (AAU, MO). — SANTIAGO-ZAMORA: Patuca, 11 Jun 1947, *G. Harling* 1129 (MO).

FRENCH GUIANA: Monts d'Arawa, savane-roche central, 8 Jul 2002, *J. J. de Granville & al.* 15119 (B, G, K, NY, US) [type gathering]; Fleuve Approuague, Rivière Arataye, Saut Parare, 2 Sep 1977, *Sastre* 5876 (P).

GUYANA: UPPER TAKUTU-UPPER ESSEQUIBO: NW Kanuku Mtns., tributary of Nappi Creek, 14 Feb 1993, *Hoffmann & Artes* 3769 (US); Sand Creek-Rupununi River confluence, 29 May 1996, *Clarke & McPherson* 1809 (US); N Rupununi Savannas, 2.5 km S of Karanamb Ranch, 19 Feb 1992, *Hoffman & al.* 999 (EIU, US); S Pakaraima Mtns., escarpment to foot of Kopinang Falls, 2 Sep 1961, *B. Maguire & al.* 46077A (NY, US).

MEXICO: CHIAPAS: 3–5 km above Solosuchiapa along road to Tapilula, 26 July 1972, *Breedlove* 26442 (DS, DUKE, MO) [cited by Tucker (1994), as *C. simplex*]. — OAXACA: "Oaxaca," [no date], *Galeotti* 5855 (P); *ibid.*, 1845, *Buchinger s.n.* (P 01924897).

PANAMA: COLÓN: Brazos Brook Res., 22 Sep 1924, *Stevens* 702 (ILL); Canal Area, Madden Forest (Parque Nacional Soberanía), 23 July 1994, *Croat & Zhu* 77068 (MO).

PERU: AMAZONAS: Distrito Imaza, camino Putuim-Shimutaz, *R. Vasquez & al.* 21294 (MO). — LORETO: Iquitos, 28 Nov 1940, *E. Asplund* 14758 (P, S). — SAN MARTÍN: Mariscal Cáceres, Tocache Nuevo, [no date], *J. Schunke* V. 6508 (MO).

SURINAME: fluv. Corantijne, *Hulke* 88 (U); Zuid River, Kayser Airstrip, 26 Aug 1963, *H. S. Irwin & al.* 55249 (NY, US), and 22 Feb 1961, *K. U. Kramer & W. H. A. Hekking* 3000 (U); Nat. Res. Brownsberg, on border of plateau in line towards Weti Creek, 24 Oct 1918, *J. C. Lindeman & A. C. de Rooy* 691 (U).

VENEZUELA: ARAGUA: along ascent from Maracay to




Fig. 1. *Cyperus conservator-davidii* – A: habit; B: spikelet; C: achene. – Based on several cited specimens including the holotype. – Original artwork by Sarah Gillespie.

summit of Alto Choroni, 8 Feb 1973, T. B. Croat 21500 (MO). — MIRANDA: Cerros de Bachiller, near E end, 16–17 Mar 1978, J. A. Steyermark & G. Davidse 116310 (MO). — PORTUGUESA: Distrito Sucre en el limite con Estado Lara, Fila El Helechal, al norte de Chabasquén, 9 Feb 1984, B. Stergios & al. 6712 (MO). — TÁCHIRA: Distrito Córdoba, Cerro La Camiri, 6 Nov 1982, G. Davidse & A. C. González 21597 (MO).

Remarks — The type specimen, together with many collections of *Cyperus conservator-davidii*, were originally identified as *C. simplex*, from which the new species is quickly separated by its rays being shorter than the culms, its lanceolate floral scales and its ellipsoid achenes. Among *Cyperus* worldwide, *C. simplex* is one of only three species in which the rays (primary branches) of the inflorescence are longer than the culm. The other two species, *C. blysmoides* Hochst. of tropical Africa (Haines & Lye 1983) and *C. radians* Nees & Meyen of SE Asia (Dai & al. 2010), are members of *C.* subg. *Cyperus* and are not close relatives of *C. simplex*. In examining materials of *C. simplex* from throughout its range (Mexico to Brazil), I noted that numerous specimens have rays shorter than the culms. Other consistent differences were noted as well.

Cyperus conservator-davidii is also similar to *C. grandisimplex*, an uncommon species of E Brazil, in having rays shorter than the culm, but it can be distinguished by its lanceolate (rather than ovate-deltate) scales and papillate (rather than smooth) achenes. The presence of papillae on the leaves and bracts in *C. conservator-davidii* is a feature not otherwise known in *C.* subg. *Pycnostachyus*. In view of the large number of collections and the wide range over several countries, it is surprising that this species has gone unrecognized.

Eponymy — This new species is named for David A. Simpson (b. 1955), currently Keeper of the Herbarium at the Royal Botanic Gardens, Kew, and fellow specialist on the *Cyperaceae*.

Status of *Cyperus lacustris*

Cyperus miliifolius Poepp. & Kunth is a distinctive species of *Cyperaceae*, occurring in wet lowland and premontane forests from Costa Rica to Bolivia. In the most recent monograph of the genus, Kükenenthal (1935–1936) cited several collections from the lowlands of N South America. As I have had the opportunity to observe numerous specimens from the region, it became apparent that these lowland plants were readily separable from typical *C. miliifolius* (Fig. 2) and should be called *C. lacustris* Schrad. ex Nees.

The lectotype of *Cyperus lacustris* has leaves and bracts 4–7 mm wide, clearly within the range of the lowland plants rather than typical *C. miliifolius*. Under the synonymy of *C. miliifolius*, Kükenenthal also included “*C.*

tenuispicatus Boeck.” He did not elaborate on the dubious status of this name. Boeckeler’s name was, in fact, validly published and I believe it applies to this lowland relative of *C. miliifolius*. I have been unable to locate a type specimen, but its origin, French Guiana, as well as features of the protologue, such as acuminate leaf tips, support this interpretation. Below is provided a detailed description of *C. lacustris* based on numerous specimens cited below.

Cyperus lacustris Schrad. ex Nees in Martius, Fl. Brasil. 2(1): 31. 1842. — **Lectotype (designated here):** Brazil, Amazonas [Prov. Rio Negro], ripa fluv. Solimões et Rio Japurá (M not seen [photo F negative no. 18874!]). — Fig. 2. = *Cyperus tenuispicatus* Boeckeler in Linnaea 35: 525. 1868 — Type: “Guiana gallica. (Ex herb. Musei Paris.)”, [collector unknown], (B†; no corresponding collection located at P).

Description — *Herbs* perennial, 15–30 cm tall. *Rhizome* oblique. *Culms* triquetrous, 1–2 mm thick, smooth. *Leaves* 5–12, V-shaped to inversely W-shaped or plicate, 10–25 cm long, 4–8.5 mm wide, margin and veins scabrellate. *Inflorescence* bracts 3–6, horizontal to ascending at 45°–60°, V-shaped, inversely W-shaped, or plicate, 5–15 cm long, 3–7 mm wide, margin and veins scabrellate; *secondary bracts* absent. *Rays* 3–8, 2–6 cm long, sometimes absent and then inflorescence capitate; *secondary rays* sometimes present, 1–5 mm long. *Heads* digitate, 4–10 mm wide. *Spikelets* 1–3(–5), linear to slightly oblong-lanceolate, compressed ellipsoid in cross-section, 7–17 mm long, 2–3 mm wide; *rachilla* persistent, wingless, successive scale scars 1.2–1.5 mm apart. *Floral scales* 10–22(–50), deciduous, dusky whitish greenish brown, ovate to ovate-lanceolate, 1.75–2 mm long, 1.5–1.7 mm wide, laterally weakly 1–4-nerved, in middle nerveless or faintly 1–3-nerved, base saccate, apical portion 0.3–0.5 mm long, incurved parallel to rachilla. *Stamen* 1; *anther* 0.7–1 mm long. *Styles* 0.5–0.8 mm long; *stigmas* 3, 0.5–0.7 mm long. *Achenes* only partly covered by edge of scale at maturity, brown, ovoid, trigonous, (0.9–)1–1.2 mm long, (0.5–)0.65–0.7 mm wide, faces concave, surface papillose or granular, base stipitate, apex subacute, apiculate.

Distribution — N South America: N Brazil, Colombia, Ecuador, French Guiana, Guyana, Peru, Suriname, Venezuela.

Habitat and ecology — River banks, trailsides and moist shaded soils of lowland forests, from near sea level to 300 m.

Proposed IUCN status — Least Concern (LC). Widely distributed; apparently not threatened. This species is known from about 70 collections, over a wide range, with numerous recent collections.



Fig. 2. *Cyperus lacustris* – A: representative specimen, R. Spruce s.n. (E 153064) from Pará, Brazil. – *Cyperus miliifolius* – B (inset): detail of representative specimen showing leaves, bracts and inflorescence, L. Holm-Nielsen & al. 19880 (EIU) from Ecuador. – Scale bars: A & B = 2 cm.

Specimens examined — BRAZIL: AMAPÁ: Santo Antonio da Cachoeira, 30 Jul 1961, *Egler & Irwin* 46049 (MG, MO, NY); Rio Oiapoque, opposite Pedra Alice, 17 Aug 1960, *Irwin & al.* 47582 (MO, NY); Rio Araguari, between Camps 3 and 4 [01°55'N, 51°59'W], 6 Sep 1961, *Pires & al.* 50733 (MO, NY). — AMAZONAS: basin of Rio Demeni, vicinity of Tototobí, 27 Feb 1969, *G. T. Prance & al.* 10279 (K, M, MO, NY, P); Rio Cauabury, 3 Nov 1930, *E. G. Holt & E. R. Blake* 449 (F); Rio Purus S of Lago Prêto, 25 Jun 1971, *G. T. Prance & al.* 13652 (K, NY, P); Mpio. Itapiranga, Rio Uatumã, 21 Aug 1979, *C. A. Cid & al.* 634 (NY, US); Rio Branco, Jarú, *Kuhlmann* 361 (EIU, K, NY, RJ); Rio Curuquetê, São Paulo, 20 Jul 1971, *G. T. Prance & al.* 14437 (GH, K, MO, NY, P, US). — PARÁ: ad cataractas fl. Aripicuru, Dec 1849, *R. Spruce s.n.* (E 153064 [Fig. 2A]); banks of Rio Piria, N of km 90, road BR 22, "1965", *G. T. Prance & T. D. Pennington* 1751 (US); Mpio. Oriximiná, Rio Paru do Oeste, cachoeira Pancada, 5 Sep 1980, *C. A. Cid & al.* 2105 (US); Rio Plumina, 26 Nov 1928, *de Sampaio s.n.* (F, RDJ). — RONDÔNIA: margen do Rio Urupá, 8 Aug 1975, *M. R. Cordeiro* 470 (MO); Rio Demeni, vicinity of Tototobí, 27 Feb 1969, *G. T. Prance & al.* 10279 (MO, NY).

COLOMBIA: META: Río Guayabero, a 10 km de Caño Lozada, 17 Jan 1959, *E. Pinto & al.* 226 (COL, P). — VAUPES: Río Pacoa (tributary of Río Apaporis), 7–12 Feb 1952, *R. E. Schultes & I. Cabrera* 15249 (COL, K, P, US); margen del Río Cananarí, affluente del Apaporis, 29 Feb 1952, *L. E. Mora-O. & van der Hammen* APA-182 (COL). ECUADOR: NAPO: near Puerto Montúfar, 17 Feb 1980, *L. Holm-Nielsen & al.* 21240 (AAU); Río Tiputini, 10 km upstream from Río Napo, 24 Aug 1979, *L. Holm-Nielsen & al.* 19796 (AAU, MO).

FRENCH GUIANA: Cayenne, 1819, *Perollet s.n.* (P 01732973); Rivière Camopi, camp ORSTOM, 12 Feb 1968, *Oldeman & Sastre* 211 (P, U); Maroni River above mouth of Marouini River, 18 Aug 1987, *A. Weitzman* 212b (US); Bassin du Haut Marouini, camp no. 3 Akouba Booka goo Soula, 27 Aug 1987, *J. J. de Granville* 9788 (CAY, US); Bassin du Haut Marouini, Monpé Soula, 6 Sep 1987, *J. J. de Granville & al.* 10092 (NY, US); Roche Koutou, Bassin du Haut Marouini, 16 Aug 1987, *J. J. de Granville & al.* 9352 (CAY, NY, US); Grand Inini, aval de Bicade, *de Granville* C-18 (P, U); Saut Dalles, Bassin du Sinnamary, 20 Oct 1990, *M. Hoff* 6769 (CAY, MO, NY, P, U, US); Saut Takari-Tanté, Bassin du Sinnamary, 18 Nov 1989, *M. Hoff* 5909 (CAY, P, US); Grand Inini, en amont de l'embouchure de la Crique Saï, 27 Aug 1970, *Oldeman* 3587 (P, US); Haute Camopi-Mont Belvédère, 22 Nov 1984, *J. J. de Granville* 6949 (CAY, US); camp no. 1 Ouman fou Langa Soula, Bassin du Haut Marouini, 12 Aug 1987, *J. J. de Granville & al.* 9192 (CAY, NY, P, US); Saut Mapaou, Nov. 1955, *Hock s.n.* (P, U).

GUYANA: "British Guiana," [no date], *R. Schomburgk* 339 (US). — UPPER TAKUTU-UPPER ESSEQUIBO: basin of Kuyuwini River (Essequibo tributary), about 150 miles from mouth, 21 Nov 1937, *A. C. Smith* 2576 (NY, US);

S Rupununi Savanna, Takutu River, 16 Sep 1997, *M. J. Jansen-Jacobs & al.* 5522 (MO, U); near Aishalton Hospital, 19 Nov 1982, *A. L. Stoffers & al.* 479 (U, US); Kamoa River, 20 Sep 1989, *M. J. Jansen-Jacobs & al.* 1720 (K, U, US); Rupununi, Kuyuwini Landing, 7 Feb 1991, *M. J. Jansen-Jacobs* 2450 (MO, U, US); Kassakaityu River 15–20 km from juncture with Essequibo River, 22 Mar 1994, *T. W. Henkel & al.* 5289 (MO, US); Kanuku Mtns., Rupununi River, Puwib River, 13 Feb 1985, *M. J. Jansen-Jacobs & al.* 194 (MO, P, U, US); Kamoa River, 0–6 km W of camp, 9 Nov 1996, *D. Clarke* 3016 (MO, US); Upper Essequibo River, 1–4 km upstream from mouth of Kuyuwini River, 3 Oct 1993, *T. W. Henkel & al.* 3343 (MO, US). — POTARO-SIPARUNI: Kurupukari Falls, 11 Dec 1994, *P. Mutchnick & V. Funk* 635 (US); Pakutan Falls, 10 Nov 1995, *D. Clarke* 472 (US); Mazaruni River, sandy islands in the falls, 28 Oct 1952, *Forest Department* F3538 (U). — CUYUNI-MAZARUNI: Aurora, 9 Oct 1989, *L. J. Gillespie* 2229 (US).

PERU: LORETO: Maynas, Río Yaguasacu, below Brilla Nueva, 7 Nov 1977, *A. Gentry & J. Revilla* 20377 (MO). SURINAME: Toekoemoetoe-Kreek, fluv. Saramacca, 1 km S of Base Camp, 12 Oct 1959, *A. G. H. Daniëls & F. P. Jonker* 1318 (U, US); Saut Paloulou Icholi, bassin du haut-Maroni, 16 Sep 1994, *J. J. de Granville* 12513 (CAY, P, U, US); Avanavero vallen, 19 Mar 1920, *G. Stahel* 4615 (U); Lucie River, 2 km below affluence of Oost River, 22 Aug 1963, *H. S. Irwin & al.* 55198 (MO, NY); Lucie River, 2–10 km below Oost River, 10 Sep 1963, *H. S. Irwin & al.* 55561 (COL, NY, US); Zuid River, 2 km above confluence with Lucie River, 15 Sep 1963, *H. S. Irwin & al.* 55739 (F, NY, US); Coppenam River near Raleighfalls, 11 Sep 1933, *J. Lanjouw* 761 (U); Zuid R., nabij samevl. met Lucie R., 2 Jul 1963, *J. P. Schulz* 10015 (U); Suriname River below Kabelstation, 24 Jul 1953, *J. C. Lindeman* 4428 (U); Sipalwini savanna area on the Brazilian frontier, 29 Oct 1968, *F. H. F. Oldenburger & al.* 384 (U); Kabalebo, 20 km downstream of Kabalebo airstrip, 29 Dec 1964, *P. A. Florschütz & P. J. M. Maas* 2567 (U); distr. Brokopondo, inter Paranam et Afobaka, 27 Mar 1961, *Kramer & Hekking* 3181 (U); distr. Brokopondo, 2.4 km S of Gansee, 21 Apr 1964, *van Donselaar* 1231a (U); on the Tanjimama at Kamp VI, 19 Nov 1954, *A. M. W. Mennega* 472 (U); fluv. Suriname, 4 Oct 1901, *Went* 503 (U).

VENEZUELA: AMAZONAS: Río Casiquiare entre la boca del Siapa y el caño Momoni, 18 Feb–4 Mar 1986, *B. Stergios & G. Aymardo* 9184 (MO, NY, PORT not seen); Río Orinoco, Tamatama, 12–24 Jan 1930, *E. G. Holt & W. Gehrig* 265 (US). — APURE: Misión Río Mavaca, *B. Stergios & Yáñez* 15021 (MO, PORT not seen); *ibid.*, *B. Stergios & Yáñez* 15083 (MO, PORT not seen). — BOLÍVAR: Río Caura, arriba del Salto Para, 14 & 17 Jan 1977, *J. A. Steyermark & al.* 112919 (MO); Mpio. Foráneo Aripao, 1 km aguas abajo de Araguaña, 1 May 1988, *G. Aymard & L. Delgado* 6747 (MO); Río Casiquiare entre la boca del Siapa y el caño Momoni, 18 Feb–4 Mar 1986, *B. Ster-*

gios & *G. Aymard* 9184 (MO); Reserva Forestal Imataca, Río Cuyuni, Isla Anacoco, 22 May 1982, *B. Stergios & al.* 3957 (MO); *ibid.*, *B. Stergios & al.* 4979 (MO); bords de l'Orénoque [no date], *Chaffanjon* 512 (P). — TACHIRA: domingo del Tachira-Barinas, 7 km al E de El Pinal, 2 Oct 1967, *G. S. Bunting* 2400 (MO, VEN not seen).

Remarks — The small dark spikelets and narrow but obviously plicate leaves are good points of recognition for this species. It appears to be restricted to moist forests in N South America, especially the Guiana Shield region. With nearly 50 collections known, it is surprising it has gone essentially unrecognized. This species has long been included in *Cyperus miliifolius*, but can be readily distinguished in flower or fruit, as follows:

1. Culms 10–30 cm tall; leaves 4–8.5 mm wide; stamen 1; achenes stipitate *C. lacustris*
- Culms 36–60 cm tall; leaves 12–16(–22) mm wide; stamens 3; achenes sessile *C. miliifolius*

Status of *Cyperus hoppiiifolius*

In studies of the preceding two species, I noted that many specimens of both *Cyperus lacustris* and *C. conservator-davidii* had been determined as “*C. hoppiiifolius* Uittien”. This prompted the question of what the correct application of Uittien’s name might be. The type of *C. hoppiiifolius* is conserved at Utrecht (U). The plant has broad leaves and very large spikelets with up to 50 floral scales. It is a striking plant and not to be confused with *C. lacustris* or *C. conservator-davidii*, or with any neotropical *Cyperus*. Later, examining some specimens borrowed from Kew, I realized that the type of *C. altsonii* was the same distinctive species, with broad leaves and large spikelets (Fig. 3). The name *C. altsonii* has priority by less than a year. Evidently this is a rare species, known only from the type gatherings of these two names, and five fairly recent collections. A detailed description and synonymy follow.

The unusually wide spikelets are very distinctive among neotropical species. Denton (1978) noted that: “The spikelets and scales of the type collection [of *Cyperus altsonii*] are larger than those of the *Luzulae* Group, and the scales have many nerves rather than just three.” The species appears properly placed in *C. sect. Incurvi*. The distinctive broad pseudopetiolate leaves are a conspicuous feature of this striking, apparently rare, species. Only six species of *Cyperus* have such leaves; all the others are in tropical Africa, except *C. pseudopetiolatus* G. C. Tucker of Bahia, Brazil (Tucker 2007).

Cyperus altsonii Kük. in Bull. Misc. Inform. Kew 1932[7]: 322. 1932 [21 Sep 1932]. – Holotype: Guyana, Maicwac River at Kopinang River, 480 m, Apr 1926, *Altson* 492 (K; isotypes: BR, NY). – Fig. 3.

= *Cyperus hoppiiifolius* Uittien in Recueil Trav. Bot. Néerl. 30[2–4]: 186. 1933 [Jul 1933]. – Holotype: Suriname, Suriname River at Goddo, 26 Jan 1926, *Exped. Wilhelmina-Gebergte* 99 (U).

Description — Herbs perennial, glaucous. *Rhizome* ± erect, 1–2 cm long. *Culms* trigonous to triquetrous, 22–32 cm tall, 1.3–3.2 mm thick, scabrid on angles. *Leaves* 8–10, flat, 30–55 cm long (longer than culm), 10–14 mm wide, base narrowed to a conduplicate reddish pseudopetiole 7–15 cm long, margin and 3 main veins scabrellate, apex attenuate-acuminate. *Inflorescence bracts* 3 or 4, spreading, flat, 3–9 cm long, 3–12 mm wide, both surfaces and margin scabrellate, base narrowed but not pseudopetiolate; *secondary bracts* absent. *Rays* 5–7, compressed trigonous, 2–7.5 cm long, scabrellate (or smooth) on angles; *secondary rays* absent. *Heads* digitate, 20–60 mm wide. *Spikelets* 3–5(–10), linear-oblong, flattened, 22–34 mm long, 4–6 mm wide; *rachilla* persistent, wingless, c. 0.75 mm wide, c. 0.5 mm thick, successive scale scars 0.8–1.2 mm apart. *Floral scales* 45–65, deciduous, greenish white to light brown, ovate-lanceolate, 3.4–3.7 mm long, 1.4–2 mm wide, ± spinulose-scabrellate, laterally 4- or 5-nerved, in middle 3–5-nerved, narrow apical portion c. 1 mm long, straight to incurved. *Stamen* 1; *filament* 3.5–4.5 mm long, c. 0.2 mm wide; *anther* not observed. *Styles* 1.2–1.8 mm long; *stigmas* 3, 1–2 mm long. *Achenes* dark brown to black, obovoid, trigonous, 1–1.2 mm long, 0.75–0.9 mm wide, faces flat or slightly convex especially apically, surface coarsely papillose, base broadly cuneate, stipitate, apex obtuse, apiculate.

Distribution — Guianas Region of N South America: French Guiana, Guyana, Suriname.

Habitat and ecology — River banks, among moist rocks, trailsides and moist shaded soils of lowland forests, from 60–500 m.

Proposed IUCN status — Although the species occurs in all three countries of the Guiana region, apparently only seven collections exist; only five are recent, the most recent made nearly 20 years ago. The status of Endangered EN: B1a is suggested because the species is “known to exist at no more than five locations”.

Specimens examined — FRENCH GUIANA: Massif des Emerillons, au bord de la Haute Approuague, 18 Sep 1980, *J. J. de Granville* 3899 (P, U); Bassin du Haut Inipi, à 7 km WSW du Pic Coudreau, 29 Sep 1980, *J. J. de Granville* 4019 (P); Bassin du Haut-Marouini, Roche Koutou, pied du versant sud, 16 Aug 1987, *J. J. de Granville & al.* 9352 (P); Bassin du Sinnamary, Camp Eugène, 9 Feb 1995, *J. J. de Granville & G. Cremers* 12814 (K, P). GUYANA: Maicwac River at Kopinang River, Apr 1926, *Altson* 492 (K, NY).



Fig. 3. *Cyperus altsonii* – close-up of inflorescence, J. Florschütz & P. A. Florschütz 1414 (U) from Suriname.
– Scale bar = 2 cm.

SURINAME: Suriname River at Goddo, 26 Jan 1926, *Exp. ed. Wilhelmina-Gebergte* 99 (U [2 sheets]); Van Asch Van Wijcks Mountains, near Ebba Top base, 14 Feb 1951, J. Florschütz & P. A. Florschütz 1414 (U [Fig. 3]).

Key to species of *Cyperus* sect. *Incurvi* in the New World Tropics

1. Inflorescence capitate-contracted (without elongated primary rays; sometimes 1 short ray present); anther apex smooth 2

– Inflorescence expanded, with primary and sometimes secondary rays; spikelets solitary or digitate in groups of 2–6; anther apex setose 5

2. Spikelets dull white; inflorescence bracts lanceolate, 2–4 mm wide *C. dichromeniformis* Kunth
- Spikelets light to dark reddish or purplish brown; inflorescence bracts linear, 0.1–2 mm wide 3

3. Spikelets 2–4 mm long; scales obtuse; achenes narrowly obovoid; S Brazil (Minas Gerais, São Paulo, Santa Catarina) *C. inops* C. B. Clarke

– Spikelets 5–9 mm long; scales acute; achene lanceolate-ellipsoid or ovoid to obovoid; E Brazil (Bahia) to Guianas and Amazonia 4

4. Inflorescence bracts 1 or 2, strongly reflexed; achenes lanceolate-ellipsoid *C. consors* C. B. Clarke

– Inflorescence bracts 3–6, ascendent; achenes ovoid to obovoid *C. lacustris*

5. Leaves linear-lanceolate, 10–25 mm wide 6

– Leaves linear, (0.7–)2–6(–8.5) mm wide 7

6. Leaves 10–14 mm wide, with pseudopetiolate base; spikelets 4–6 mm wide *C. altsonii*

– Leaves without pseudopetiolate base; spikelets 2–3 mm wide *C. miliifolius*

7. Longest ray longer than culm; often proliferous . . .
 *C. simplex*
 – Longest ray shorter than the culm; rarely proliferous
 **8**
8. Scales ovate-lanceolate, ovate or suborbicular; stamens 3 per flower **9**
 – Scales ovate; stamen 1 per flower **11**
9. Culms 1.7–4 mm thick; successive scale scars c. 0.6 mm apart; anthers 0.4–0.5 mm long; achenes trigonous but dorsiventrally flattened
 *C. lundellii* O'Neill
 – Culms 0.5–1.2(–2) mm thick; successive scale scars 0.7–1.5 mm apart; anthers 0.9–2.2 mm long; achenes trigonous, not compressed **10**
10. Successive scale scars 0.7–0.9 mm apart; anthers c. 0.9 mm long; achenes ellipsoid, 0.4–0.5 mm wide *C. almensis* D. A. Simpson
 – Successive scale scars 1.1–1.5 mm apart; anthers 1.5–2.2 mm long; achenes obovoid, 0.7–0.8 mm wide
 *C. subcastaneus* D. A. Simpson
11. Rays compressed quadrate in cross-section . . . **12**
 – Rays trigonous in cross-section **13**
12. Culms smooth; anthers 0.6–0.8 mm long; achenes broadly obovoid *C. grandisimplex*
 – Culms scabrous on angles just below bracts; anthers c. 1.2 mm long; achenes ellipsoid
 *C. conservator-davidii*
13. Spikelets 2–3 mm wide; achenes ovoid to obovoid, 0.9–1.2 mm long; widespread: Amazonia and Guianas *C. lacustris*
 – Spikelets 2–2.5 mm wide; achenes ovoid-ellipsoid, 1.2–1.6 mm long; widespread: Paraguay, Bolivia, and SC Brazil (Goias) **14**
14. Spikelets oblong-ovate, 4–6 mm long, with 10–12 scales; scales laxly imbricate, deep brown, 3-nerved medially *C. anisitsii* Kük.
 – Spikelets linear, linear-oblong to slightly linear-lanceolate, 12–24 mm long, with 12–30 scales; scales densely imbricate, dull white, 9–11-nerved medially and laterally *C. pearcei* C. B. Clarke

Acknowledgements

I wish to thank the EIU Botany Alumni Fund for travel to K, NY and SP. The illustration was prepared by Sarah Gillespie. Kanchi Gandhi (GH) kindly confirmed the dates and places of publication of all names and synonyms. During the course of this study and my preceding studies of the Mexican and Central and South American species of *Cyperus*, some 30 000 herbarium specimens were borrowed from, received as gifts for determination from, or examined at, the herbaria listed below (codes follow Thiers 2008+). Thanks are extended to the curators of these institutions, without whose generosity and cooperation this study would not have been possible: AAU, AC, ARIZ, ASU, B, BD, BH, BKL, BM, BR, C,

CAS, CHAPA, CONN, CORD, CU, DAO, DAV, DS, DUKE, E, ECON, EIU, ENCB, F, FLAS, G, GH, HAL, IBUG, ILL, ILLS, IND, JE, K, LCU, LL, M, MA, MASS, MEXU, MG, MICH, MO, MSC, MT, MTMG, MU, ND, NHA, NY, NYS, P, PENN, PH, PMA, POM, PR, PRC, RDJ, RSA, S, SD, SIU, SMU, SP, TENN, TEX, TRT, UC, UCR, UEC, US, UTEP, VT, WIS, WRSL, WVA, YU and Z. In addition, two anonymous reviewers are thanked for their comments on an earlier draft of this paper.

References

- Alves M., Araújo A. C., Prata A. P., Vitta F. A., Hefler S. M., Trevisan R., Bragança Gil A., Martins S. & Thomas W. W. 2007: Diversidade de *Cyperaceae* no Brasil. – Pp. 286–290 in: Barbosa L. M. & Dos Santos Jr. N. A. (ed.), *A Botânica no Brasil: Pesquisa, Ensino e Políticas Públicas Ambientais*. – São Paulo: Sociedade Botânica do Brasil.
- Dai L. K., Tucker G. C. & Simpson D. A. 2010: *Cyperus* Linnaeus – Pp. 219–241 in: Wu Z. Y., Raven P. H. & Hong D. Y. (ed.), *Flora of China* **23**. – Beijing: Science Press; St. Louis: Missouri Botanical Garden Press.
- Denton M. F. 1978: A taxonomic treatment of the Luzulae group of *Cyperus*. – Contr. Univ. Michigan Herb. **11**: 197–271.
- Haines R. W. & Lye K. A. 1983: The sedges and rushes of East Africa. – Nairobi: East African Natural History Society.
- IUCN 2012: IUCN Red List categories and criteria: Version 3.1, ed. 2. – Gland & Cambridge: IUCN.
- Kükenthal G. 1935–1936: *Cyperaceae–Scirpoideae–Cypereae*. – In: Engler A. & Diels L. (ed.), *Das Pflanzenreich. Regni vegetabilis conspectus* **IV. 20 (Heft 101)**. – Leipzig: Wilhelm Engelmann.
- Thiers B. 2008+ [continuously updated]: Index herbariorum: a global directory of public herbaria and associated staff. – New York Botanical Garden: published at <http://sweetgum.nybg.org/ih/> [accessed 25 Jan 2014].
- Tucker G. C. 1994: Revision of the Mexican species of *Cyperus* L. (*Cyperaceae*). – *Syst. Bot. Monogr.* **43**: 1–213.
- Tucker G. C. 2001: *Cyperus* (*Cyperaceae*). – Pp. 740–757 in: Stevens W. D., Ulloa Ulloa C., Pool A., Montiel O. M., Arbeláez A. L. & Cutaia D. M. (ed.), *Flora de Nicaragua* (Tomo I). – Monogr. Syst. Bot. Missouri Bot. Gard. **85**.
- Tucker G. C. 2007: Systematics of *Cyperus* sect *Diffusi* Kunth (*Cyperaceae*) in the Neotropics. – Pp. 311–314 in: Barbosa L. M. & Dos Santos N. A. Jr. (ed.), *A botânica no Brasil: pesquisa, ensino e políticas públicas ambientais*. – São Paulo: Sociedade Botânica do Brasil.
- Tucker G. C. 2008: *Cyperus alvesii* (*Cyperaceae*), a new species from northeastern Brazil. – *Harvard Pap. Bot.* **13**: 237–240.