

Plumularia roxanae, a new epiphytic hydroid (Cnidaria: Hydrozoa: Plumulariidae) from the Indo-Pacific

Author: Galea, Horia R.

Source: Revue suisse de Zoologie, 127(2) : 367-376

Published By: Muséum d'histoire naturelle, Genève

URL: <https://doi.org/10.35929/RSZ.0026>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

***Plumularia roxanae*, a new epiphytic hydroid (Cnidaria: Hydrozoa: Plumulariidae) from the Indo-Pacific**

Horia R. Galea

Hydrozoan Research Laboratory, 405 Chemin Les Gatières, F-83170 Tourves, France. E-Mail: horia.galea@gmail.com

Abstract: A new epiphytic, small-sized species of *Plumularia* Lamarck, 1816, *P. roxanae* sp. nov., is described based on fertile material from Bali, Indonesia. Its deep hydrothecae are characteristically narrowed distally through a distinctive hypertrophy of the free portion of their adaxial wall. Its minute gonothecae, however, are comparable in shape and size to those of the well-known, circumtropical *P. floridana* Nutting, 1900.

Keywords: Taxonomy - new species.

INTRODUCTION

The highly speciose, circumglobal, ecologically diverse genus *Plumularia* Lamarck, 1816 comprises feather-shaped hydroids ranging in form from small, pinnate colonies with monosiphonic, unbranched stems to large structures composed of repeatedly branched, occasionally fascicled stems (Bouillon *et al.*, 2006). As noted by Schuchert (2013a, b), its taxonomy proves occasionally challenging, due to an important intraspecific variation (e.g. internode length, shape of hydro- and gonothecae) and a limited availability of diagnostic morphological characters. Although new species are still created based on infertile specimens (Watson, 2012), thus adding to the actual difficulties in the identification of numerous poorly-described nominal taxa erected by earlier authors (e.g. Mulder & Trebilcock 1911, 1914, 1915; Fraser 1938a, b, c, 1948), a small, epiphytic *Plumularia* with distinctive hydro- and gonothecae is described from material collected recently from Bali, Indonesia.

MATERIAL AND METHODS

Hydroid-covered algae freshly washed ashore were collected by hand during an ebb tide, and fixed immediately in 4% buffered formalin in seawater. The material was subsequently sorted in the laboratory and observations were made according to the methods described by Galea (2007, 2008). Holo- and paratype specimens are deposited in the invertebrate collection of the Muséum d'histoire naturelle de Geneva, Switzerland.

RESULTS

***Plumularia roxanae* sp. nov.**

Figs 1-2; Tables 1-2


Holotype: MHNG-INVE-137167; INDONESIA, Bali, Pererenan beach; -8.650831°, 115.119820°; 13 Jan. 2020; fertile colony on red alga.

Paratypes: MHNG-INVE-137168; INDONESIA, Bali, Pererenan beach; -8.650831°, 115.119820°; 13 Jan. 2020; fertile colony on red alga. – MHNG-INVE-137169; INDONESIA, Bali, Pererenan beach; -8.650831°, 115.119820°; 13 Jan. 2020; fertile colonies on 3 fragments of red algae.

Diagnosis: Small-sized, epiphytic *Plumularia* arising from stolon with perisarcal spurs, giving rise to monosiphonic stems divided homomerously by transverse nodes into rather short, collinear internodes, each with a subterminal cladial apophysis provided with an inconspicuous adaxial mamelon and a pair of axillar nematothecae; cladia heteromerously segmented by alternating straight and oblique nodes into short, ahydrothecate internodes occasionally provided with a nematotheca, and up to three, comparatively longer hydrothecate internodes accommodating a centrally-placed hydrotheca and its three associated nematothecae; hydrotheca deep, saccate, partly adnate, distal portion distinctly constricted through the hypertrophy of the free adaxial wall that adopts a triangular shape, considerably reducing the thecal lumen; abaxial wall slightly convex; gonothecae in two closely-set, parallel rows along the stem, borne on the cladial apophyses, minute, ovoid, thick-walled.

Manuscript accepted 22.07.2020

DOI: 10.35929/RSZ.0026






Fig. 2. *Plumularia roxanae* sp. nov. (A) Portion of ramified stolon, showing internal spurs. (B) Two stem internodes and proximal parts of their corresponding cladia. (C) Portion of stem with gonotheca. (D) Detail of stem internode. (E, F) Two hydrothecae seen laterally, note differences in the adaxial perisarc thickening. (G) Hydrotheca seen from above, note the position of lateral nematothecae. (H) Base of hydrotheca with mesial nematotheca. (I) Distal part of hydrotheca with lateral nematotheca. (J) Cladial ahydrothecate internode. (K) Gonotheca. Scale bars: 50 µm (D, H-K), 100 µm (E-G), 200 µm (B, C), 300 µm (A).

Fig. 1. *Plumularia roxanae* sp. nov. (A) Portion of cormoid. (B) Detail of a portion of stem with proximal parts of four successive cladia. (C) Detail of a stem internode, showing centrally an apophysis and the proximal part of the corresponding cladium. (D) Proximal most hydrotheca. (E) Distal most hydrotheca, note the absence of the distal part of its corresponding internode. (F) Portion of fertile stem with rows of gonothecae. (G) Gonotheca. (H) More detailed view of a gonotheca, showing apical aperture (arrowhead). Scale bars: 50 µm (H), 100 µm (C-E, G), 200 µm (B, F), 500 µm (A).

Etymology: It is my great pleasure to dedicate this new species to the memory of my late mother, Roxana S. Galea (née Florescu, 5 Oct. 1946 – 18 Jan. 1994), a Romanian biologist, who passed on to me her interest for natural sciences.

Description: Colonies epiphytic on algae, composed of numerous erect cormoids arising from creeping, branching, flattened stolonal tubes firmly attached to substrate; lumen with two lateral rows of perisarc spurs; a wide flange of thinner perisarc flanking outer lateral sides. Stems short, up to 5 mm high, with thick and firm perisarc, gradually thinning out distally; monosiphonic, divided by means of transverse nodes into a regular sequence of collinear, rather short internodes, length slightly increasing gradually towards tips of cormoids; each internode with two internal perisarc ridges (one above proximal node, the second below the distal node) and a subterminal cladial apophysis; apophyses alternate along the stem, the two rows either in one plane in some specimens or variably shifted on to the anterior side of the colony in others; apophyses well-developed, each with a subterminal, internal perisarc ridge, an inconspicuous mamelon (reduced to a mere pore) with small, circular aperture on adcauline side, and a pair of axillar nematothecae, one to each lateral side; distal node straight. Cladia given off at an angle of about 45° with the long axis of the stem; up to 1 mm long, divided heteromerously by an alternation of deeply-incised, straight and oblique nodes delimiting a- and hydrothecate internodes; first cladial internode ahydrothecate; all ahydrothecate internodes short, with proximal node straight and distal node oblique, with two internal perisarc ridges (one proximal, one distal) and, occasionally, with a nematotheca in middle of upper side; hydrothecate internodes, up to 3 per cladium, comparatively longer than their ahydrothecate counterparts, slightly sigmoid in overall shape, with proximal node oblique and distal node straight, a centrally-placed hydrotheca (together with its 3 associated nematothecae), occupying more than half the length of the internode, and two internal perisarc ridges near both ends. Hydrotheca deep, saccate, adnate for about 2/3 to its corresponding internode; abaxial wall thick, slightly convex proximally; free adaxial wall exceedingly hypertrophied, adopting a triangular shape, free angle upturned; adnate adaxial wall moderately thick, forming an L-shaped perisarc thickening at junction with the hydrothecal base, the latter pierced by circular foramen for the passage of the hydranth; hydrothecal aperture facing nearly upwards, rim entire, circular in apical view; mesial nematotheca well below the hydrothecal base, mounted on small prominence of the internode; lateral nematothecae borne on indistinct apophyses and hidden by the free portion of the adaxial wall when the hydrotheca is seen frontally, and typically inclined over this side of the thecal wall; hydranths

able to completely retract into their corresponding hydrothecae, with ca. 12 filiform tentacles in a whorl around a dome-shaped hypostome. All nematothecae of the colony movable, elongate-conical in shape and bithalamic; lower chamber comparatively taller than upper one, the latter with the wall slightly lowered on adaxial side. Gonothecae in two parallel, closely-set rows along the stem, arising singly from the anterior side of each cladial apophysis; likely male in the present material, with a homogenous, granular content; elongate-globular, thick-walled, with smooth outer surface, apically with a small funnel-shaped aperture in some specimens, delimited basally by transverse, thick diaphragm from a short, tubular pedicel.

Remarks: Upon a comparison with all shallow-water, small-sized (generally not surpassing 1 cm high), short-cladiate, nominal species of *Plumularia* Lamarck, 1816 from the tropical Indian and western Pacific oceans (Table 1), it appears that the material at hand is

Table 1. Measurements of *Plumularia roxanae* sp. nov., in μm .

Stem	
- internodes, length	220-285
- internodes, diameter at node	45-95
- cladial apophyses, length	40-45
Cladia	
- ahydrothecate internodes, length	65-95
- hydrothecate internodes, length	255-300
- diameter at transverse node	30-40
Hydrotheca	
- abaxial wall, length	180-195
- free adaxial wall, length	75-85
- adnate adaxial wall, length	110-120
- diameter at rim	100-115
Nematothecae	
- axillar, length	60-65
- axillar, diameter at rim	ca. 25
- of ahydrothecate internodes, length	ca. 40
- of ahydrothecate internodes, diameter at rim	ca. 25
- mesial, length	ca. 55
- mesial, diameter at rim	ca. 25
- lateral, length	55-60
- lateral, diameter at rim	ca. 35
Gonotheca	
- length	195-220
- maximum width	135-140
- pedicel, length	45-50
- pedicel, width	40-45

Table 2. Nominal species of *Plumularia* Lamarck, 1816 from the tropical Indo-west Pacific, and their distinctive features to *P. roxanae* sp. nov.

Nominal species	Distinctive features to <i>P. roxanae</i> sp. nov. Remarks. Habitat	Distribution and references
<i>P. amphibola</i> (Watson, 2011)	Nominal species indistinguishable morphologically from <i>P. crateriformis</i> Mulder & Trebilcock, 1911, from which it was separated only on the account of the presence of single hydrothecate internodes on cladia. Epiphytic	Australia (Watson, 2011)
<i>P. angusta</i> Stechow, 1923	Nominal species created by Stechow (1923: 226) for 3 varieties (a, b, d) of <i>P. setaceoides</i> Bale, 1882 dealt with by Mulder & Trebilcock (1911: pl. 3 figs 3-3b, pl. 3 fig. 6, pl. 2 fig. 9, respectively). Subsequently reported upon by Watson (1975: figs 30-31). No comprehensive description available, but morphology of both hydro- and gonothecae similar to that of the better-known <i>P. wilsoni</i> Bale, 1926 (see below), with which it could be conspecific, in that case having priority. Presumably epiphytic	Australia, Tasmania (Mulder & Trebilcock, 1911; Watson, 1975)
<i>P. australis</i> Kirchenpauer, 1876	Stem internodes moderately-long, geniculate; cladial apophyses with a mamelon and an axillar nematotheca; cladia composed of a proximal, short, athecate internode, followed by a hydrothecate internode; hydrotheca seated in a concavity of internode, slightly compressed laterally, abaxial wall distinctly cambered, adaxial wall projecting a strong, oblique intrathecal septum, rim weakly to noticeably sinuous; dioecious, 1-3 gonothecae on lower stem internodes, large, flask-shaped, body narrowing distally to an upwardly bent, cylindrical neck, with a wide, circular, thickened aperture with broadly everted rim. Epiphytic	Australia (Watson, 1990, 2011)
<i>P. caliculata</i> Bale, 1888	Stem internodes moderately-long, slightly geniculate, with internal septa at both ends, and 1 nematotheca; cladial apophyses with low mamelon and 1 axillar nematotheca; cladia heteromerous, internodes with septa at both ends, up to 5 hydrothecate internodes; hydrothecae cup-shaped, shallow, fully adnate, abaxial wall thickened and swollen basally; monoecious, gonothecae forming a continuous row on front of stem, attached to successive cladial apophyses just under mamelon, female globular with external marsupium, male elongated. Epiphytic	Australia, New Zealand, Japan (Vervoort & Watson, 2003)
<i>P. compressa</i> Bale, 1882	The hydro- and gonothecae of this nominal species appear indistinguishable from those of <i>P. australis</i> Kirchenpauer, 1876 (see above); compare Bale (1882: p. 15, figs 5-5a), Bale (1884: pl. 12 figs 9-10, pl. 19 figs 39-40) and Bale (1888: pl. 19 fig. 14) with Bale (1884: pl. 12 figs 7-8, pl. 19 figs 43-44) and Watson (1990: figs 177-182), respectively. If the synonymy is accepted, Kirchenpauer's species has priority. Epiphytic	Australia (Watson, 2005)
<i>P. corrugatissima</i> Mulder & Trebilcock, 1914	Nominal species inaccurately described. Stem internodes with 1 nematotheca; cladial apophyses with 2 nematothecae; cladia heteromerous, hydrothecate internodes distinctly swollen; hydrothecae small, cup-shaped, shallow, about as deep as wide, fully adnate, rim everted frontally, aperture somewhat oblique; gonothecae unknown. Presumably epiphytic	Australia (Mulder & Trebilcock, 1914; Bale, 1919)
<i>P. crateriformis</i> Mulder & Trebilcock, 1911	Stolon with nematothecae. Stems divided by strongly oblique nodes; internodes short, robust, with several thick complete internal septal ridges, and 1 nematotheca; cladial apophyses with 1 axillar nematotheca; ahydrothecate internodes with 1 nematotheca, with the exception of the proximal most; hydrothecae perched on bulges of upper side of hydrothecate internodes, alternating with 3 strong septal ridges; symmetrical, cup-shaped, walls expanding to margin, base rounded, aperture outrolled; 1-3 gonothecae in lower stem region, often of both sexes, urn-shaped, distally a dome-shaped cap, gonophore with row of refringent corpuscles. Epiphytic	Australia (Mulder & Trebilcock, 1911; Watson, 1990, 2005)
<i>P. crateroides</i> Mulder & Trebilcock, 1911	Nominal species inaccurately described. Stolon with nematothecae; stem internodes "with about three transverse wrinkles" and 1 nematotheca; cladial apophyses with 1 nematotheca; cladia heteromerous; hydrothecae cup-shaped, rather shallow, expanding upwards, partly adnate to the corresponding internode; gonothecae given off singly from base of stem, "somewhat flattened beneath, convex on top, obliquely truncated, aperture very large, margin everted". Presumably epiphytic	Australia (Mulder & Trebilcock, 1911)

Nominal species	Distinctive features to <i>P. roxanae</i> sp. nov. Remarks. Habitat	Distribution and references
<i>P. epibracteolosa</i> Watson, 1973	Stem divided by strongly oblique nodes; internodes with 3-4 internal ridges of perisarc, and 1 nematotheca; cladial apophyses with an axillar nematotheca besides the adaxial pore; cladial hydrothecate internodes with 4-5 internal ridges of perisarc; hydrothecae wide, shallow, cup-shaped; abaxial wall distinctly thickened; adaxial wall adnate to hydrocladium only near base, the remainder joined to the internode by a wedge of perisarc; margin sinuated, with a thickened outwardly rolled rim, the line of the margin curved down posteriorly to meet the internode; aperture truncated adaxially by a delicate transverse sheet of perisarc; a couple of large, piriform gonothecae arising from old, basal hydrocladial apophyses. Epiphytic	Australia, New Zealand (Watson, 1973; Vervoort & Watson, 2003)
<i>P. excavata</i> Mulder & Trebilcock, 1911	Nominal species inaccurately described. Stem "annulated", without nematothecae; cladia bearing single, concave hydrothecate internodes; hydrothecae cup-shaped, partly adnate, abaxial wall distinctly sigmoid (strongly concave proximally, convex distally); gonothecae unknown. Presumably epiphytic	Australia (Mulder & Trebilcock, 1911)
<i>P. filicaulis</i> Kirchenpauer, 1876	Colonies with sessile cladia and/or erect stems; stem divided by oblique nodes; internodes moderately-long, with 1 nematotheca; cladial apophyses with mamelon and 1 axillar nematotheca; cladia heteromerous, with up to 6 hydrothecate internodes; hydrothecae cup-shaped, basally adnate, abaxial wall slightly rounded, adaxial wall giving rise to an intrathecal septum, margin forming an angle of 45° with the internode, produced to form a peak on both the ad- and abaxial sides; gonothecae borne on hydrothiza, firmly applied to substratum, flattened, thick-walled, irregularly ovate in outline, aperture small, circular on upper surface of distal end, a number of minute pores on upper surface. Epiphytic	South Africa, Australia, Japan, Madagascar (Millard, 1975; Watson, 2005)
<i>P. flexuosa</i> Bale, 1894	Stem occasionally branched; internodes moderately-long, geniculate, with 1 nematotheca; cladial apophyses with 1-2 axillar nematothecae, as well as an additional nematotheca on the apophysis itself; hydrothecae deeply campanulate, expanding to aperture, abaxial wall weakly concave, adaxial wall almost straight, a small portion free from the internode; dioecious; gonothecae on lower stem internodes, elongate-ovoid, twice as long as wide, distal end tapering, truncate above and there provided with a rather small, somewhat oblique aperture, without submarginal teeth. Epiphytic and epizoic	Australia, New Zealand, Japan (Bale, 1894; Watson, 1973, 2000, 2005)
<i>P. floridana</i> Nutting, 1900	Stem internodes relatively long, slender, slightly geniculate, each with an internal perisarc ridge at each end, and 1 nematotheca; cladial apophyses with mamelon and 2 axillar nematothecae; cladia heteromerous, ahydrothecate internodes moderately-long and with 1 nematotheca, hydrothecate internodes with a centrally-placed hydrotheca; the latter fairly deep, cylindrical, half adnate; gonothecae small, globular, thin-walled, borne on cladial apophyses. Epiphytic	Circumglobal in warm waters (Ansín Agís <i>et al.</i> , 2001)
<i>P. fragilia</i> Watson, 2012	Stem internodes long, slender, with 2 nematothecae on opposite sides; cladial apophyses with a bun-shaped mamelon and 2 axillar nematothecae; cladia heteromerously segmented into long a- and hydrothecate internodes, the former with 1 nematotheca; hydrothecae seated about halfway along the internodes, small, deep bowl-shaped, abaxial wall deeply concave to straight, adaxial wall weakly concave and fully adnate to internode, margin oblique to internode axis, circular, rim everted; gonothecae unknown. Epizoic	Australia (Watson, 2012)
<i>P. goldsteini</i> Bale, 1882	Nominal species inadequately described, though easily recognizable through the peculiar shape of its hydrothecae. Cladia heteromerous, with up to 4 hydrothecate internodes; hydrotheca slipper-shaped, long axis strongly sigmoid, base distinctly swollen, aperture facing away from the internode; gonothecae unknown. Epiphytic	Australia (Bale, 1882, 1884 ; Watson, 1973)

Nominal species	Distinctive features to <i>P. roxanae</i> sp. nov. Remarks. Habitat	Distribution and references
<i>P. hyalina</i> Bale, 1882	Stem internodes of varied length, with or without internal septa, with 0-1 nematotheca; cladial apophysis with conical mamelon and 1 axillar nematotheca; hydrotheca deep bowl-shaped, abaxial wall slightly convex, adaxial wall concave and fully adnate to internode, margin a broad sinuous curve down to internode; one to several gonothecae in a row around lower stem, large, piriform, walls smooth to obscurely undulated, when mature flattened at top, and opening by a large, smooth-edged, circular aperture, gonophore with belt of refringent corpuscles. Epiphytic and epizoic	Australia (Mulder & Trebilcock, 1911; Trebilcock, 1928; Watson, 2011)
<i>P. lucerna</i> Mulder & Trebilcock, 1911	Very likely a stemless variety of <i>P. filicaulis</i> Kirchenpauer, 1876 (see above); compare Mulder & Trebilcock (1911: pl. 3 fig. 4) with Bale (1884: pl. 11 figs 6-7). Presumably epiphytic	Australia (Mulder & Trebilcock, 1911)
<i>P. meretricia</i> Watson, 1973	Stem internodes long, slender, occasionally with 1-2 nematothecae, often absent; cladial apophyses with prominent mamelon (with 1-2 apertures) adaxially and 1 axillar nematotheca, the latter often absent; hydrothecae campanulate, abaxial wall straight for most of its length, expanding below aperture; margin everted, slightly sinuated, curving down adaxially; gonothecae unknown. Epizoic	Australia (Watson, 1973, 2005)
<i>P. mooreana</i> Schuchert, 2013	Stem internodes relatively short, sturdy, each with 1 nematotheca; cladial apophyses with 1 axillar nematotheca; cladia heteromerous, with up to 4 relatively short hydrothecate internodes, each with a centrally-placed hydrotheca; the latter bowl-shaped, shallow, inclined downward at 45° with the internode; 1-2 gonothecae near stem bases, borne on laterally-set pedicels, urn-shaped, perisarc undulated, outer surface with scattered nematothecae. Inert substrates	French Polynesia, Fiji (Schuchert, 2013b)
<i>P. obesa</i> Blackburn, 1938	Stem internodes short, sturdy; cladial apophyses with mamelon and 1-2 nematothecae; cladia with single, short, concave hydrothecate internode, accommodating distally a hydrotheca; the latter laterally compressed, abaxial wall sigmoid, adaxial wall fully fused, concave, giving rise to a forwardly and upwardly-directed intrathecal ridge, margin sinuous, somewhat everted abaxially, distinctly lowered adaxially; gonothecae unknown. Epiphytic	Australia (Blackburn, 1938; Watson, 1990)
<i>P. obliqua</i> Johnston, 1847	Stem internodes moderately long, geniculate, with 1 nematotheca; cladial apophyses with conical mamelon and 1 axillar nematotheca; hydrothecate cladial internode with 2 internal septa behind the hydrotheca, the latter distal on internode, scoop-shaped, fully adnate, abaxial wall smoothly convex, thickened or not, margin cut obliquely back to internode, rim slightly everted, a perisarc thickening behind; gonothecae on lower stem internodes, large, urn-shaped, perisarc fairly thick, distally truncate, aperture closed by sheet of tissue; gonophore a medusoid. Epiphytic	Circumglobal in temperate to tropical waters (Watson, 2011)
<i>P. pulchella</i> Bale, 1882	Stem internodes robust, moderately long, with several transverse internal septa and 1 nematotheca; cladial apophyses with 2 axillar nematothecae; hydrothecate internodes short, concave, hydrotheca distally-placed, fully adnate, cup-shaped, slightly flaring, abaxial wall weakly concave, variably thickened; dioecious, gonothecae on lower stem internodes, at base of a cladial apophysis, large, globular, thick-walled, smooth to faintly rugose, aperture distal, set obliquely, aperture narrow, circular, with a row of large, internal submarginal cusps. Epiphytic	Australia (Watson, 1973, 2011)
<i>P. rotunda</i> Mulder & Trebilcock, 1911	Stem internodes moderately long, slightly geniculate, with 1 nematotheca; cladial apophyses with 1 axillar nematotheca; cladia heteromerously segmented, ahydrothecate internodes short and with 1 nematotheca, hydrothecate internodes up to 2, comparatively longer, each with a hydrotheca distally; hydrothecae elliptic, abaxial wall strongly arched, becoming thicker towards aperture where it is incurved, appearing as a submarginal septum, aperture small, set obliquely forwards, subrectangular, rim weakly lobate; gonothecae unknown. Epiphytic	Australia (Bale, 1919; Watson, 2015)
<i>P. spinulosa</i> Bale, 1882	Stem internodes moderately long and slender, with 1 nematotheca; cladial apophyses with mamelon and 1 axillar nematotheca; cladia with single, concave hydrothecate internode, ending in a distal spine of variable length, projecting beyond hydrotheca; the latter bonnet-shaped, fully adnate, laterally compressed, abaxial wall convex, a strong adaxial intrathecal septum projecting halfway across lumen, aperture sinuated adaxially; one to several gonothecae on lower cladial apophyses, large, elongated-conical, sometimes curved asymmetrically, with wide terminal aperture set perpendicularly to long axis. Epiphytic and epizoic	Australia, New Zealand, South Africa, Japan (Millard, 1975; Watson, 2011)

Nominal species	Distinctive features to <i>P. roxanae</i> sp. nov. Remarks. Habitat	Distribution and references
<i>P. strictocarpa</i> Pictet, 1893	Stolon with nematothecae; cauline internodes long, slender, slightly geniculate, with 1-2 nematothecae; cladial apophyses with mamelon and 1 axillar nematotheca; cladia heteromerous, ahydrothecate internodes relatively long, with internal septa at both ends and 1 nematotheca; hydrothecate internodes comparatively longer, with up to 3 internal septa; hydrothecae conical, fully adnate, margin rounded, tilted abaxially; gonothecae borne below lower cladial apophyses, large, partly or fully seated on substrate, barrel-shaped in frontal view, flattened dorso-ventrally, distinctly transversely ridged, aperture distal, circular. Epiphytic, epizoic, also growing on inert substrates	Circumglobal in warm waters (Ryland & Gibbons, 1991; Hirohito, 1974)
<i>P. togata</i> Watson, 1973	Cauline internodes moderately long and slender, with 1 nematotheca; cladial apophyses with conical mamelon and 1 axillar nematotheca; hydrotheca slipper-shaped, longitudinal axis S-shaped, abaxial wall considerably produced frontally, aperture narrowed and tilted adaxially; gonotheca borne beside apophysis of first stem internode, large, urn-shaped, distally truncate, walls smooth to weakly undulated, aperture closed by sheet of tissue. Epiphytic	Australia (Watson, 1973, 2011)
<i>P. tubacarpa</i> Watson, 2000	Cauline internodes long, each with 1-4 cladial apophyses (usually 2) and a variable number of nematothecae, 1 in axil of apophysis, 1 beside apophysis, 1 halfway between hydrocladia and sometimes 1 at side of internode just above node; cladia divided homomerously into quite long hydrothecate internodes; hydrothecae deep cup-shaped, adaxial wall entirely adnate, abaxial wall straight to family sinuous, concave just behind margin; gonothecae exceedingly long, tubular, rounded apically. Epizoic	Australia (Watson, 2000)
<i>P. warreni</i> Stechow, 1919	Stem internodes moderately long, with 1 frontal nematotheca; cladial apophyses with 1 axillar nematotheca; cladia heteromerous, ahydrothecate internodes short, with internal septa near ends, and 1 nematotheca, hydrothecate internode comparatively longer, with internal septa near ends and a centrally-placed hydrotheca; the latter inverted conical, completely adnate, margin slightly oblique; monoecious, gonothecae on cladial apophyses, elongate-ovoid, outline irregular, slightly flattened dorso-ventrally. Epiphytic	From South Africa to Australia (Millard, 1975)
<i>P. wilsoni</i> Bale, 1926	Stolon with internal pegs and nematothecae; stem internodes relatively short, with internal ridges near their ends, and 1 nematotheca on side opposite to cladial apophysis; the latter with 2 axillar nematothecae; cladia heteromerous, hydrothecate internodes relatively long, with a centrally-placed hydrotheca; abaxial wall sigmoid, thickened, adaxial wall free for 1/3 from corresponding internode, concave, aperture tilted abaxially; gonothecae borne on lower stem internodes, large, piriform, transversely ridged, distal end truncate. Presumably epiphytic, also epizoic	Australia, Tasmania, New Zealand (Ralph, 1961; Watson, 2015)

distinctive, mainly through the shape of its hydrothecae. Similar gonothecae are only found in *P. floridana* Nutting, 1900, a species of circumtropical occurrence, described under various names (Calder, 1997: 15).

The epiphytic hydroid erroneously assigned (Watson, 2011: 78) to *Monothecha hyalina* (Bale, 1882) by Vervoort & Watson (2003) resembles the present species, notably in the structure of the colony and the shape of hydrotheca, but shows several differences: 1) its stolon is devoid of perisarc spurs; 2) there is only one axillar nematotheca associated to the cladial apophyses of the stem; 3) the hydrothecae are devoid of the characteristic thickening of the free adaxial wall; 4) its (female) gonothecae are much bigger, ca. 1 mm high, as deduced from their fig. 90I, and they are piriform, with an apical, broad, circular aperture closed by an operculum. For these reasons, their record is not included in the synonymy of the present species, pending the discovery of colonies belonging to the opposite (female) sex.

Distribution: Known only from Bali, Indonesia, but presumably spread throughout the archipelago.

ACKNOWLEDGEMENTS

This study was made possible through a grant (App #40579/2019) from the PADI Foundation, Beverly Hills, CA, USA. I am grateful to Prof. Fran Ramil (University of Vigo, Spain) and Dr Henry H.C. Choong (Royal British Columbia Museum, Canada) for their thorough review of the manuscript.

REFERENCES

- Ansín Agís J., Ramil F., Vervoort, W. 2001. Atlantic Leptolida (Hydrozoa, Cnidaria) of the families Aglaopheniidae, Halopterididae, Kirchenpaueriidae and Plumulariidae collected during the CANCAP and Mauritania-II expeditions of the National Museum of Natural History, Leiden, the Netherlands. *Zoologische Verhandelingen, Leiden* 333: 1-268.
- Bale W.M. 1882. On the Hydroida of south-eastern Australia, with descriptions of supposed new species, and notes on the genus *Aglaophenia*. *Journal of the Microscopical Society of Victoria* 2(1): 15-48.
- Bale W.M. 1884. Catalogue of the Australian hydroid zoophytes. *Australian Museum, Sydney*, 198 pp.
- Bale W.M. 1888. On some new and rare Hydroida in the Australian Museum collection. *Proceedings of the Linnean Society of New South Wales* 2(3)2: 745-799.
- Bale W.M. 1894. Further notes on Australian hydroids, with descriptions of some new species. *Proceedings of the Royal Society of Victoria* 6: 93-117.
- Bale W.M. 1919. Further notes on Australian hydroids. IV. *Proceedings of the Royal Society of Victoria* 31(2): 327-362.
- Bale W.M. 1926. Further notes on Australian hydroids. V. *Proceedings of the Royal Society of Victoria* 38: 13-23.
- Blackburn M. 1938. Hydrozoa. In: The Sir Joseph Banks Islands. Reports of the expedition of the McCoy Society for Field Investigation and Research, 3. *Proceedings of the Royal Society of Victoria* 50(2): 312-328.
- Bouillon J., Gravili C., Pagès F., Gili J.M., Boero F. 2006. An introduction to Hydrozoa. *Mémoires du Muséum national d'Histoire naturelle, Paris* 194: 1-591.
- Calder D.R. 1997. Shallow-water hydroids of Bermuda: superfamily Plumularioidea. *Royal Ontario Museum Life Sciences Contributions* 161: 1-84.
- Fraser C.M. 1938a. Hydroids of the 1934 Allan Hancock Pacific Expedition. *Allan Hancock Pacific Expeditions* 4(1): 1-105.
- Fraser C.M. 1938b. Hydroids of the 1936 and 1937 Allan Hancock Pacific Expeditions. *Allan Hancock Pacific Expeditions* 4(2): 107-127.
- Fraser C.M. 1938c. Hydroids of the 1932, 1933, 1935, and 1938 Allan Hancock Pacific Expeditions. *Allan Hancock Pacific Expeditions* 4(3): 129-153.
- Fraser C.M. 1948. Hydroids of the Allan Hancock Pacific Expeditions since March, 1938. *Allan Hancock Pacific Expeditions* 4(5): 179-343.
- Galea H.R. 2007. Hydroids and hydromedusae (Cnidaria: Hydrozoa) from the fjords region of southern Chile. *Zootaxa* 1597: 1-116.
- Galea H.R. 2008. On a collection of shallow-water hydroids (Cnidaria: Hydrozoa) from Guadeloupe and Les Saintes, French Lesser Antilles. *Zootaxa* 1878: 1-54.
- Hirohito, Emperor of Japan. 1974. Some hydrozoans of the Bonin Islands. *Publications of the Biological Laboratory, Imperial Household, Tokyo* 11: 1-55.
- Johnston G. 1847. A history of the British zoophytes. *John van Voorst, London*, XVI + 488 pp.
- Kirchenpauer G.H. 1876. Ueber die Hydroidenfamilie Plumularidae, einzelne Gruppen derselben und ihre Fruchtbehälter. II. *Plumularia* und *Nemertesia*. *Abhandlungen aus dem Gebiete der Naturwissenschaften, Hamburg* 6(2): 1-59.
- Lamarck J.B. 1816. Histoire naturelle des animaux sans vertèbres. *Verdière, Paris*, 568 pp.
- Millard N.A.H. 1975. Monograph on the Hydroida of southern Africa. *Annals of the South African Museum* 68: 1-513.
- Mulder J.F., Trebilcock R.E. 1911. Notes on Victorian Hydroida, with description of new species. (Continued). *Geelong Naturalist* 4(4): 115-124.
- Mulder J.F., Trebilcock R.E. 1914. Victorian Hydroida. With description of new species. Part IV. *Geelong Naturalist* 6(2): 38-47.
- Mulder J.F., Trebilcock R.E. 1915. Victorian Hydroida. With description of new species. Part V. *Geelong Naturalist* 6(3): 51-59.
- Nutting C.C. 1900. American hydroids. Part I. The Plumulariidae. *Special Bulletin of the United States National Museum* 4(1): 1-285.
- Pictet C. 1893. Etude sur les Hydrides de la Baie d'Amboine. *Revue suisse de Zoologie* 1: 1-64.
- Ralph P.M. 1961. New Zealand thecate hydroids. Part IV. Family Plumulariidae. *Transactions of the Royal Society of New Zealand* 1(3): 19-74.
- Ryland J.S., Gibbons M.J. 1991. Intertidal and shallow water hydroids from Fiji. II. Plumulariidae and Aglaopheniidae. *Memoirs of the Queensland Museum* 30(3): 525-560.
- Schuchert P. 2013a. The status of *Plumularia lagenifera* Allman, 1885 (Cnidaria, Hydrozoa) and related species. *Zootaxa* 3613(2): 101-124.

- Schuchert P. 2013b. *Plumularia mooreana*, a new marine hydroid from French Polynesia (Hydrozoa, Cnidaria). *Revue suisse de Zoologie* 120(3): 405-413.
- Stechow E. 1919. Zur Kenntnis des Hydroidenfauna des Mittelmeeres, Amerikas und anderer Gebiete, nebst Angaben über einige Kirchenpauer'sche Typen von Plumulariden. *Zoologische Jahrbücher* 42(1): 1-172.
- Stechow E. 1923. Zur Kenntnis des Hydroidenfauna des Mittelmeeres, Amerikas und anderer Gebiete. II. Teil. *Zoologische Jahrbücher* 47(1): 29-270.
- Trebilcock R.E. 1928. Notes on New Zealand Hydroids. *Proceedings of the Royal Society of Victoria*. New series. 41: 1-33, plates 1-7.
- Vervoort W., Watson J.E. 2003. The marine fauna of New Zealand: Leptothecata (Cnidaria: Hydrozoa) (thecate hydroids). *NIWA Biodiversity Memoir* 119: 1-538.
- Watson J.E. 1973. Pearson Island Expedition 1969. – 9. Hydroids. *Transactions of the Royal Society of South Australia* 97(3): 153-200.
- Watson J.E. 1975. Hydroids of Bruny Island, southern Tasmania. *Transactions of the Royal Society of South Australia* 99(4): 157-176.
- Watson J.E. 1990. Studies on Australian hydroids. The genus *Eudendrium* and the fauna of the seagrass *Amphibolis*. *Deakin University, Melbourne*, 288 pp.
- Watson J.E. 2000. Hydroids (Hydrozoa: Leptothecata) from the Beagle Gulf and Darwin Harbour, northern Australia. *The Beagle, Records of the Museum and Art Galleries of the Northern Territory* 16: 1-82.
- Watson J.E. 2005. Hydroids of the Archipelago of the Recherche and Esperance, Western Australia: annotated list, redescription of species and description of new species [pp. 495-611]. In: Wells F.E., Walker D.I., Kendrick G. (eds), *The Marine Flora and fauna of Esperance, Western Australia. Western Australian Museum, Perth*, 727 pp.
- Watson J.E. 2011. Review of the genus *Monothea* (Hydrozoa: Leptolida) from Australia with description of a new species and a note on *Monothecella* Stechow, 1923. *Memoirs of Museum Victoria* 68: 71-91.
- Watson J.E. 2012. Some hydroids (Hydrozoa: Hydroidolina) from Dampier, western Australia: annotated list with description of two new species. *Memoirs of Museum Victoria* 69: 355-363.
- Watson J.E. 2015. A new species of *Halopteris* (Hydrozoa: Leptothecata) and redescription of *Plumularia rotunda* from Victoria, Australia. *Memoirs of the Museum Victoria* 73: 41-46.