

Citrus (Rutaceae) SNP Markers Based on Competitive Allele-Specific PCR; Transferability Across the Aurantioideae Subfamily

Authors: Garcia-Lor, Andres, Ancillo, Gema, Navarro, Luis, and Ollitrault, Patrick

Source: Applications in Plant Sciences, 1(4)

Published By: Botanical Society of America

URL: <https://doi.org/10.3732/apps.1200406>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

CITRUS (RUTACEAE) SNP MARKERS BASED ON COMPETITIVE ALLELE-SPECIFIC PCR; TRANSFERABILITY ACROSS THE AURANTIOIDEAE SUBFAMILY¹

ANDRES GARCIA-LOR², GEMA ANCILLO², LUIS NAVARRO^{2,4}, AND PATRICK OLLITRAULT^{2,3,4}

²Centro de Protección Vegetal y Biotecnología, Instituto Valenciano de Investigaciones Agrarias (IVIA), Apartado Oficial, 46113 Moncada (Valencia), Spain; and ³CIRAD, UMR AGAP, F-34398, Montpellier, France

- **Premise of the study:** Single nucleotide polymorphism (SNP) markers based on Competitive Allele-Specific PCR (KASPar) were developed from sequences of three *Citrus* species. Their transferability was tested in 63 *Citrus* genotypes and 19 relative genera of the subfamily Aurantioideae to estimate the potential of SNP markers, selected from a limited intrageneric discovery panel, for ongoing broader diversity analysis at the intra- and intergeneric levels and systematic germplasm bank characterization.
- **Methods and Results:** Forty-two SNP markers were developed using KASPar technology. Forty-one were successfully genotyped in all of the *Citrus* germplasm, where intra- and interspecific polymorphisms were observed. The transferability and diversity decreased with increasing taxonomic distance.
- **Conclusions:** SNP markers based on the KASPar method developed from sequence data of a limited intrageneric discovery panel provide a valuable molecular resource for genetic diversity analysis of germplasm within a genus and should be useful for germplasm fingerprinting at a much broader diversity level.

Key words: Competitive Allele-Specific PCR; genetic diversity; Rutaceae; single-nucleotide polymorphisms (SNPs).

Single nucleotide polymorphisms (SNPs) are the most frequent type of DNA sequence polymorphism. Their abundance and uniform distribution in genomes make them very powerful genetic markers. Several SNP genotyping methods have been developed. For low-to-medium throughput genotyping, the KBioscience Competitive Allele-Specific PCR genotyping system (KASPar; KBioscience Ltd., Hoddesdon, United Kingdom) appears to be an interesting approach (Cuppen, 2007) that has been successfully applied in animals and plants (Nijman et al., 2008; Bauer et al., 2009; Cortes et al., 2011). For genetic diversity studies with SNP markers, it is very important to determine the representativeness of the discovery panel (Albrechtsen et al., 2010). Ascertainment bias of the SNP markers affects the evaluation of genetic parameters, as was observed for the *Citrus* L. genus using SNP markers mined in a single Clementine cultivar (Ollitrault et al., 2012). Recently, Garcia-Lor et al. (2013) sequenced 27 amplified nuclear gene fragments for 45 genotypes of *Citrus*, which resulted in the identification of 1097 SNPs. Taking advantage of these previously obtained SNP data, the objective of this work was to implement a set of polymorphic SNP markers for systematic germplasm bank characterization within the *Citrus* genus and to investigate their transferability across the Aurantioideae [Engler] subfamily. More generally,

the objective was to estimate the usefulness of SNP markers developed using KASPar technology, which were selected from a limited intrageneric discovery panel, for broader diversity analysis at the intra- and intergeneric levels.

METHODS AND RESULTS

The 42 SNP markers used in this study were selected from SNPs identified by Garcia-Lor et al. (2013) in 27 nuclear genes. Most cultivated citrus (except for *C. aurantifolia* (Christm.) Swingle) arose from interspecific hybridization of three ancestral taxa: *C. medica* L., *C. reticulata* Blanco, and *C. maxima* (Burm.) Merr. (Nicolosi et al., 2000; Barkley et al., 2006; Garcia-Lor et al., 2012). Therefore, we selected SNPs between and within these three taxa (based on seven *C. reticulata*, five *C. maxima*, and five *C. medica* accessions). Primers were defined by KBioscience (<http://www.kbioscience.co.uk/>) from each SNP-locus flanking sequence (Appendix S1). Two allele-specific oligonucleotides and one common oligonucleotide were defined for each locus (Table 1). The KASPar system uses two Förster resonance energy transfer (FRET) cassettes, where fluorometric dye is conjugated to the primer but quenched via resonance energy transfer. In this system, sample DNA is amplified in a thermal cycler using allele-specific primers, leading to the separation of fluorometric dye and quencher when the FRET cassette primer is hybridized with DNA (Cuppen, 2007). Normalized signals of each SNP allele (*x* and *y*) were provided by KBioscience. Automatic allele calls provided by KlusterCaller software were visually checked with two-dimensional plot representations using SNPViewer software (KBioscience Ltd.).

Eighty-four accessions (Appendix 1) were genotyped for the 42 SNP markers. The sample set included representatives of the two tribes of the Aurantioideae (Clausenae and Citreae). In Clausenae, the subtribe Clauseniae was represented by four genotypes (three genera). Within the Citreae, three subtribes were represented: Triphasilineae (one genus was included), Balsamocitrinae (represented by six genera), and Citrinae (11 genera represented). For the Citrinae, we adopted the subdivision of this tribe into three groups (as proposed by Swingle and Reece, 1967), namely the primitive citrus fruit group (four accessions of four genera), the near citrus fruit group (three accessions of two

¹Manuscript received 3 August 2012; revision accepted 26 September 2012.

This work was supported by a grant (Prometeo/2008/121) from the Generalitat Valenciana, Spain, and by a grant (AGL2011-26490) from the Ministry of Economy and Innovation, Fondo Europeo de Desarrollo Regional (FEDER).

⁴Author for correspondence: patrick.ollitrault@cirad.fr, lnavarro@ivia.es

TABLE 1. Characteristics of 41 SNP primers used for genotyping of the Aurantioidae subfamily.

ID ^a	Gene	SNP-specific primers ^b	Common primer ^c	AlleleX	AlleleY	GenBank accession ^d no.
EMA-M30	Malic enzyme (EMA)	AlleleX: GCCTATTCAATAAATTTAGATGTCAGAAA AlleleY: CCTATTCATATAAATTTAGATGTCAGGAA	GTTTAGCCCGCAGCTTTCTTTCTCTTT	T	C	JX630064
ACO-P353	Aconitase (ACO)	AlleleX: ATGCTGCAGAGAAAACCCAGTAAATG AlleleY: CAATGCTGCAGAGAAAACCCAGTAAATA	TCTCTGTTTTGAAGCTAAATCCCAGCTCAA	C	T	JX630065
ACO-C601	Aconitase (ACO)	AlleleX: ATAAAGGCTTATGAAGAAGTTTCAACTC AlleleY: CATAAAGGCTTATGAAGAAGTTTCAACTT	CTGAAGCTAAATTTGCAGACATGGAACATT	G	A	JX630065
F3'H-P30	Flavonoid 3'-hydroxylase (F3'H)	AlleleX: CCCACTTGGCTACGACGCT AlleleY: CCACCTGGCTACGACGCC	CTCGGACCAATAATCAGCAAAAGACCAT	T	C	JX630066
F3'H-M309	Flavonoid 3'-hydroxylase (F3'H)	AlleleX: ACGTCATGAGTCTACACCATA AlleleY: CGTCATGAGCTTACCCACATG	GACCAAAGGACAGAAATCTAATGAGTTTA	T	C	JX630066
F3'H-C341	Flavonoid 3'-hydroxylase (F3'H)	AlleleX: GAGCTCATGAGCTAGCTGGAAT AlleleY: GAGCTCATGAGCTAGCTGGATA	GCAAATCGAGGGTATAAATTCACCAATGTT	T	A	JX630066
PEPC-M316	Phosphoenolpyruvate carboxylase (PEPC)	AlleleX: TAAAGAGCAATGAATTTCTTCAAACTAA AlleleY: AAAGAGCAATGAATTTCTTCAAACTAG	GTGCAATTAAGAACTGAGAAGGCAATAGAA	T	C	JX630067
PEPC-C328	Phosphoenolpyruvate carboxylase (PEPC)	AlleleX: TAAAGTGACTTAAAGAGCAATGAATTC AlleleY: CTTAAAGGCTGACTTAAAGAGCAATGAATTT	GAAGCATAGAATATTCCTAYTAGGTTTGAA	G	A	JX630067
SOS1-M50	Salt overly sensitive 1 (SOS1)	AlleleX: GGTTTAGTACTGAGTAAGTACTTGG AlleleY: AAATGGTTTTAGTACTGAGTAAGTACTTGT	GGACTTTTTCAGGTTTTTCACGGCCGCAAT	G	A	JX630068
CCC1-M85	Cation chloride cotransporter (CCC1)	AlleleX: CATTTGGTTATGAGGTATCCAGAG AlleleY: AACATTTGGTTATGAGGTATCCAGAA	CAGTAAGGTTTTTCACGGCCGCAAT	G	A	JX630069
CCC1-P727	Cation chloride cotransporter (CCC1)	AlleleX: ATCAACCACCAGCTTACTGGTAT AlleleY: CAACCACCAGCTTACTGGTAC	GGCACATTTCTACTAACAATAATCCATGTA	T	C	JX630069
TRPA-M593	Vacuolar citrate/H ⁺ symporter (TRPA)	AlleleX: AACGTGGCAGCAGAGTGAIG AlleleY: AACGTGGCAGCAGAGTGAIC	TCCCAGTGGCCACTGGCAGTCAAT	C	G	JX630070
INVA-M437	Acid invertase (INVA)	AlleleX: GTTCAGCAGATCTCTCGCTGGAA AlleleY: CAGCAGATCTCTCGCTGGAG	ACAGGGAGTCCAAATGTTGGAGTTTA	T	C	JX630071
INVA-P855	Acid invertase (INVA)	AlleleX: GGCACGTCAATAGAACTCTCACAAT AlleleY: GACGTCAATAGAACTCTCACAAC	CCTGCAATATACATACACAATGTTCCAAA	T	C	JX630071
MDH-MP69	Malate dehydrogenase (MDH)	AlleleX: AGGCCACTGAACTCACAAGTAT AlleleY: GGCCACTGAACTCACAAGTATG	CTGGTGTGAGGTTCAACTCCAAGAA	A	C	JX630072
MDH-M519	Malate dehydrogenase (MDH)	AlleleX: CAGCTCAACCAAGTCTTTACTATA AlleleY: AGCCTCAACCAAGTCTTTACTATG	GATGACCTCTTCAACATCAACGCCAAA	T	C	JX630072
ATMR-C372	MRP-like ABC transporter (ATMR)	AlleleX: GAATCATTTTATGATGGAATCGACATTTCCG AlleleY: AGAATCATTTTATGATGGAATCGACATTTCA	ACCTTAGGTCATGAAGCCCAACAA	G	A	JX630073
ATMR-M728	MRP-like ABC transporter (ATMR)	AlleleX: GTTTGATTTAAGGAGTCATATGATCTTTTTT AlleleY: TGATTTAATGGAGTCATATGATCTTTTTG	AAAGTTCACATTTTGGCAATGTTTAGCTTT	T	G	JX630073
CHS-P57	Chalcone synthase (CHS)	AlleleX: CAAGTATGGTAGTTTCAGAAAGTGGTT AlleleY: CAAGTATGGTAGTTTCAGAAAGTGGTT	AAAAACAACCTTGGAAAGCCCGGTTTTT	T	A	JX630074
CHS-M183	Chalcone synthase (CHS)	AlleleX: GTTGGAGCTGACCCATCTCTG AlleleY: GTTGGAGCTGACCCATCTCTC	GTTAAGTTCATGAAGGAGAAAGACTCTTT	G	C	JX630074
CHI-M598	Chalcone isomerase (CHI)	AlleleX: CGTCACTTTCACGGCTCCG AlleleY: CGTCACTTTCACGGCTCC	TGGACTTTTGTGATCTCTGGAGGTT	C	G	JX630075
PKF-C64	Phosphofructokinase (PKF)	AlleleX: ACTCCCTCTCCCTCTGTTCTC AlleleY: CACTCCCTCTCCCTCTGTTCTA	GGCCATCGACGATTTTGAAGGGGTT	C	A	JX630076
PKF-M186	Phosphofructokinase (PKF)	AlleleX: CGTCCGTAAATACAGATTCGAAGT AlleleY: CGTCCGTAAATACAGATTCGAAGC	CCGAACAGATTTTGGAAACAATTTCCGAAT	T	C	JX630076
NADK2-M285	NADH kinase (NADK2)	AlleleX: CATCTTCTTTGGTGATACAGAAGAA AlleleY: ATCTTCTTCTTTGGTGATACAGAAGAG	AACTCATTTCTAGATCTGTGATGAGGAGGTT	T	C	JX630077
DFR-M240	Dihydroflavono 4-reductase (DFR)	AlleleX: CCGAAGGGAACCTTTGATGAAG AlleleY: CCGAAGGGAACCTTTGATGAAG	GAAAAACTCCAGTCCAGCCCTCGAAT	G	C	JX630078
LAPX-M258	Ascorbate peroxidase (LAPX)	AlleleX: GAATTGACCATGGTTGTGTTTATTTTC AlleleY: GAATTGACCATGGTTGTGTTTATTTTG	GGCAACAACCTCCAGCCCAACTCAA	C	G	JX630079

TABLE 1. Continued.

ID ^a	Gene	SNP-specific primers ^b	Common primer ^c	AlleleX	AlleleY	GenBank accession ^d no.
PSY-M30	Phytoene synthase (PSY)	AlleleX: GTCCATTTGATATGCTTGCTGG AlleleY: GTCCATTTGATATGCTTGCTGG	CGACAGGAAATTTGGTTACTGTATCTGAT	G	C	JX630080
PSY-C461	Phytoene synthase (PSY)	AlleleX: CGCAGCCCTAFTAAACTCTTTGTCA AlleleY: CGCAGCCCTAFTAAACTCTTTGTCT	AAGTTCTGCATGCTACCCCTCTCAATAATT	T	A	JX630080
AOC-M290	Ascorbate oxydase (AOC)	AlleleX: AAAGGGTGCATCTGAGCCCAAAG AlleleY: AAAGGGTGCATCTGAGCCCAA	CTGCGTTGAAAACATAATGGTACTGTACTTT	C	T	JX630081
AOC-C593	Ascorbate oxydase (AOC)	AlleleX: GCCATACCCATGGAATTCGGCT AlleleY: GCCATACCCATGGAATTCGGCA	GGGTTAACTGGAGGCTCCATTT	T	A	JX630081
DXS-C545	1-deoxyxylulose 5-phosphate synthase (DXS)	AlleleX: ACCAAATGCATGAACGGCTTTC AlleleY: ACCAAATGCATGAACGGCTTTCG	GGGCTTGCAGGATTCGCCAAA	G	C	JX630082
DXS-M618	1-deoxyxylulose 5-phosphate synthase (DXS)	AlleleX: GGTCTTGGTATGTACTTCG AlleleY: CTGCTGGTCTTGGTATGTACTTCA	CCTACAAATTTCTCTAGATTGATGAAAAGGAA	G	A	JX630082
FLS-P129	Flavonol synthase (FLS)	AlleleX: GGCTTCCCGCATGGAACGTG AlleleY: GGCTTCCCGCATGGAACGTG	CGATCTCGACGACCCCGCTTCAA	T	C	JX630083
FLS-M400	Flavonol synthase (FLS)	AlleleX: CCGTCTTCTAFTCACTACCGCTTT AlleleY: CGTCTTCTAFTCACTACCGCTTC	TTCAACCGGTAAGAAGGAGGCTTGT	T	C	JX630083
LCY2-M379	Lycopene β-cyclase 2 (LCY2)	AlleleX: TGTGATGAGTTTGAAGACATAGACTTA AlleleY: GTTGATGAGTTTGAAGACATAGACTTA	CGGCCAAGTTTTGTCCAAAACAGTCTA	G	A	JX566716
LCYB-M480	Lycopene β-cyclase (LCYB)	AlleleX: GAATAACCTTAATAACTTTAGCTTGGTGG AlleleY: GAATAACCTTAATAACTTTAGCTTGGTGA	GCTGCAAAATGCATAACCAATGGTGTTA	C	T	JX630084
LCYB-P736	Lycopene β-cyclase (LCYB)	AlleleX: GATTGCGATCTGAACAACAATTCGG AlleleY: CGCATCTGAACAACAATTCGG	GAAAAGTAGGAAATTTGCTATTTGCCTCTT	G	C	JX630084
HYB-M62	β-Carotene hydroxylase (HYB)	AlleleX: AAAACAACAATACCGTGAAAGAGTTGAT AlleleY: AACAAAACAATACCGTGAAAGAGTTGAG	GGCTTCTTTAATGGCAAAAACCCGAAAGAAA	A	C	AF315289
HYB-C433	β-Carotene hydroxylase (HYB)	AlleleX: GAGCAAAATGCCCCAACATTTTCAGT AlleleY: AGAGCAAAATGCCCCAACATTTTCAGT	GTACAGGGTGGAGAGGTGCCTT	G	A	JX630087
TSC-C80	Trehalose-6-phosphate synthase (TSC)	AlleleX: TCTTGACCCTTGGAAAATGTTCTTT AlleleY: CTTTGACCCTTGGAAAATGTTCTTT	GCCTCTTTTGCACAAACAACAGGCTCAT	T	G	JX630084
NCED3-M535	9-cis-epoxy hydroxy carotenoid dioxygenase 3 (NCED3)	AlleleX: GACACCTTGTCTTGCATAAATCACA AlleleY: ACACCTTGTCTTGCATAAATCACC	CAAGTGGTGTTCAAAGTTGAATGAGATGAT	T	G	JX630086

^aID = SNP locus name.

^bAllele X and Y forward primers.

^cReverse primer.

^dGenBank accession numbers for the genomic fragment gene sequences of *C. resini* (corresponding sequences with identification of each SNP marker are also given in Appendix S1).

TABLE 2. Results of initial primer screening in different *Citrus* species and subtribes of the Aurantioideae subfamily.

Marker	<i>C. reticulata</i> (N = 12)		<i>C. maxima</i> (N = 11)		<i>C. medica</i> (N = 6)		<i>Citrus</i> (N = 32)		True citrus* (N = 16)		Balsamocitrinae (N = 6)		Near Citrus (N = 3)		Primitive Citrus (N = 4)		Triphasitinae (N = 1)		Clauseniae (N = 4)		Aurantioideae (N = 84)			
	A	H ₀	A	H _e	A	H ₀	A	H _e	A	MD	A	MD	A	H ₀	A	MD	A	MD	A	MD	A	H ₀	H _e	
EMA-M30	2	0.73	1	0.00	1	0.00	2	3.13	1	0.00	1	66.67	0	100.00	1	0.00	1	0.00	1	75.00	2	0.29	0.26	
ACO-P353	1	0.00	2	0.55	1	0.00	2	0.00	1	0.00	2	33.33	1	0.00	2	0.00	1	0.00	1	25.00	2	0.16	0.27	
ACO-C601	1	0.00	1	0.00	1	0.00	2	0.00	1	0.00	1	33.33	1	0.00	1	0.00	1	0.00	1	25.00	2	0.06	0.19	
F3'H-P30	1	0.00	2	0.10	1	0.00	2	6.25	1	0.00	1	16.67	1	0.00	0	100.00	1	0.00	1	0.00	2	0.10	0.24	
F3'H-M309	2	0.33	0.30	1	0.00	2	0.00	2	0.00	1	0.00	16.67	1	0.00	1	0.00	1	0.00	1	0.00	2	0.07	0.09	
F3'H-C341	1	0.00	1	0.00	2	0.17	0.37	2	0.00	1	0.00	0.00	1	0.00	1	0.00	1	0.00	1	0.00	2	0.04	0.10	
PEPC-M316	1	0.00	1	0.00	1	0.00	2	3.13	1	0.00	1	16.67	1	0.00	1	25.00	1	0.00	1	50.00	2	0.11	0.29	
PEPC-C328	1	0.00	1	0.00	1	0.00	2	0.00	1	0.00	1	16.67	1	0.00	1	0.00	1	0.00	1	75.00	2	0.08	0.18	
SOS1-M50	1	0.00	1	0.00	1	0.00	2	0.00	2	0.00	1	0.00	1	0.00	1	0.00	1	0.00	1	75.00	2	0.12	0.35	
CCCI-M85	2	0.67	0.37	1	0.00	1	0.00	2	0.00	2	0.00	0.00	1	0.00	1	0.00	1	0.00	1	50.00	2	0.24	0.31	
CCCI-P727	2	0.58	0.37	1	0.00	1	0.00	2	0.00	1	0.00	0.00	1	0.00	1	0.00	1	0.00	1	25.00	2	0.16	0.34	
TRPA-M593	2	0.58	0.33	1	0.00	1	0.00	2	0.00	1	0.00	0.00	1	0.00	1	0.00	1	0.00	1	25.00	2	0.20	0.37	
INVA-M437	1	0.00	1	0.00	1	0.00	2	0.00	1	0.00	1	50.00	1	0.00	1	0.00	1	0.00	1	50.00	2	0.13	0.29	
INVA-P855	1	0.00	1	0.00	2	0.33	0.24	2	0.00	2	0.00	16.67	2	0.00	2	0.00	2	1.00	2	0.00	2	0.22	0.37	
MDH-MP69	1	0.00	1	0.00	1	0.00	2	0.00	2	6.25	1	33.33	1	0.00	1	0.00	1	0.00	1	50.00	2	0.18	0.37	
MDH-M519	2	0.42	0.33	1	0.00	1	0.00	2	0.00	1	0.00	16.67	1	0.00	1	25.00	1	0.00	1	50.00	2	0.18	0.24	
ATMR-C372	1	0.00	1	0.00	2	0.50	0.37	2	0.00	1	0.00	16.67	1	0.00	1	0.00	1	0.00	1	25.00	2	0.09	0.12	
ATMR-M728	1	0.00	1	0.00	1	0.00	2	3.13	2	6.25	1	83.33	1	33.33	1	0.00	1	0.00	0	100.00	2	0.15	0.37	
CHS-P57	1	0.00	1	0.00	1	0.00	2	0.00	1	0.00	2	0.00	1	0.00	1	0.00	1	0.00	1	0.00	2	0.08	0.26	
CHS-M183	1	0.00	1	0.00	1	0.00	2	0.00	1	0.00	1	66.67	1	0.00	1	0.00	1	0.00	1	25.00	2	0.06	0.31	
CHI-M598	2	0.09	0.08	1	0.00	2	0.17	0.14	2	3.13	2	33.33	1	0.00	2	0.00	1	0.00	1	50.00	2	0.22	0.37	
PKF-C64	1	0.00	1	0.00	1	0.00	2	3.13	1	0.00	1	0.00	1	0.00	1	0.00	1	0.00	1	50.00	2	0.05	0.14	
PKF-M186	1	0.00	1	0.00	1	0.00	2	0.00	1	0.00	0	100.00	1	33.33	1	75.00	0	100.00	1	25.00	2	0.13	0.37	
NADK2-M285	1	0.00	1	0.00	2	0.33	0.24	2	0.00	2	6.25	1	50.00	1	33.33	1	25.00	0	100.00	1	25.00	2	0.20	0.34
DFR-M240	1	0.00	1	0.00	1	0.00	2	3.13	2	0.00	1	83.33	2	33.33	2	0.00	0	100.00	1	25.00	2	0.17	0.37	
LAPX-M238	2	0.50	0.35	1	0.00	1	0.00	2	0.00	1	0.00	0.00	1	0.00	1	0.00	1	0.00	1	75.00	2	0.22	0.26	
PSY-M30	2	0.67	0.35	1	0.00	1	0.00	2	0.00	1	0.00	0.00	1	0.00	1	0.00	1	0.00	1	25.00	2	0.27	0.37	
PSY-C461	1	0.00	1	0.00	1	0.00	2	0.00	1	0.00	1	33.33	1	0.00	1	0.00	1	0.00	1	25.00	2	0.06	0.17	
AOC-M290	2	0.45	0.29	1	0.00	1	0.00	2	6.25	1	16.67	2	0.00	2	0.00	1	0.00	1	25.00	2	0.28	0.30		
AOC-C593	1	0.00	1	0.00	1	0.00	2	0.00	1	0.00	2	16.67	2	0.00	2	0.00	1	0.00	1	25.00	2	0.06	0.17	
DXS-C545	1	0.00	1	0.00	2	0.33	0.35	2	0.00	1	0.00	16.67	1	0.00	1	0.00	1	0.00	1	25.00	2	0.07	0.13	
DXS-M618	2	0.50	0.30	1	0.00	1	0.00	2	0.00	2	0.00	16.67	2	0.00	1	0.00	1	0.00	1	50.00	2	0.21	0.27	
FLS-P129	1	0.00	1	0.00	2	0.27	0.34	1	0.00	2	0.00	0.00	1	0.00	1	0.00	1	0.00	1	0.00	2	0.18	0.26	
FLS-M400	2	0.50	0.30	1	0.00	1	0.00	2	0.00	1	0.00	16.67	1	0.00	1	0.00	1	0.00	1	0.00	2	0.18	0.27	
LCY2-M379	2	0.67	0.35	1	0.00	1	0.00	2	0.00	1	0.00	0.00	1	0.00	1	0.00	1	0.00	1	0.00	2	0.27	0.26	
LCYB-M480	2	0.33	0.24	1	0.00	1	0.00	2	0.00	2	0.00	0.00	1	0.00	1	0.00	1	0.00	1	25.00	2	0.23	0.34	
LCYB-P736	1	0.00	1	0.00	2	0.18	0.37	1	0.00	2	0.00	66.67	1	0.00	1	0.00	1	0.00	1	25.00	2	0.06	0.16	
HYB-M62	2	0.42	0.33	1	0.00	1	0.00	2	3.13	2	0.00	0.00	1	0.00	1	25.00	1	0.00	1	75.00	2	0.27	0.37	
HYB-C433	1	0.00	1	0.00	1	0.00	2	0.00	1	0.00	2	0.00	1	0.00	1	0.00	1	0.00	1	0.00	2	0.10	0.19	
TSC-C80	1	0.00	1	0.00	1	0.00	2	3.13	1	0.00	1	0.00	1	0.00	1	0.00	1	0.00	1	0.00	2	0.07	0.16	
NCED3-M535	1	0.00	1	0.00	1	0.00	2	0.00	1	0.00	1	16.67	1	0.00	1	0.00	1	0.00	1	25.00	2	0.13	0.29	
Mean	1.37	0.18	0.11	1.10	0.03	0.03	1.15	0.04	0.04	2	0.91	1.32	0.61	1.22	6.50	1.12	6.71	0.93	9.78	1.07	26.83	2	0.15	0.26

Note: A = number of alleles; H_e = expected heterozygosity; H₀ = observed heterozygosity; MD = missing data (%); N = sample size.
 * True citrus excluding the *Citrus* genus.




Fig. 1. Neighbor-joining analysis based on simple matching dissimilarities from 41 SNP loci for 50 accessions belonging to the genus *Citrus*, including secondary species and hybrids. Numbers near nodes are bootstrap values based on 1000 resamplings (only values >50% are indicated).

genera), and the “true citrus fruit trees” group (48 accessions of six genera). High-molecular-weight genomic DNA was extracted from leaf samples using a DNeasy Plant Mini Kit (QIAGEN, Madrid, Spain) according to the manufacturer’s instructions.

From the 42 SNP primers tested, only one did not produce polymorphisms. To check the accuracy of the allele call for the 41 other markers, we compared the KASPar genotyping data with Sanger sequencing data available for 35 accessions of the “true citrus fruit trees” (Garcia-Lor et al., 2013). The conformity level was 95.41%, while 2.99% did not agree and 1.60% were missing data.

The allele number and the percentage of missing data are presented for each taxon (Table 2). The expected (H_e) and observed heterozygosity (H_o) were evaluated for *C. reticulata*, *C. maxima*, *C. medica*, the *Citrus* genus, and the “true citrus fruit trees” excluding the *Citrus* genus. Data analysis was conducted with PowerMarker version 3.25 (Liu and Muse, 2005) and DARwin (Perrier and Jacquemoud-Collet, 2006) software.

The missing data rate was very low in *Citrus* (0.9%) and, generally, in the “true citrus fruit trees” group (0.6%, excluding the *Citrus* genus). The missing data rate increased to 6.5% and 6.7% in the close citrus and primitive citrus groups of the Citrinae subtribe, respectively, reaching a level of 9.8% and 22.4% for the two other subtribes of the Citreae tribe, the Triphasilineae and the Balsamocitrinae, respectively. Missing data reached 26.8% in the Clauseninae tribe. These results indicate an increasing loss of transferability with increasing taxonomic distance. As expected due to the discovery panel, the *Citrus* genus was the most polymorphic (an average of two alleles per locus; $H_e = 0.30$; $H_o = 0.23$), followed by the “true citrus fruit trees” group excluding the *Citrus* genus (alleles per locus [A] = 1.32; $H_e = 0.09$; $H_o = 0.02$). Diversity within and between the other taxa decreased considerably (data not shown). However, despite this important loss of polymorphism, all citrus relatives were differentiated when missing amplification was considered to represent null alleles, providing molecular fingerprinting for traceability in germplasm bank management.

Among the *Citrus* ancestral taxa, *C. reticulata* was the most polymorphic ($A = 1.37$; $H_e = 0.11$), followed by *C. medica* ($A = 1.15$; $H_e = 0.04$), and *C. maxima* ($A = 1.10$; $H_e = 0.03$). Considering as subpopulations the three species used in the discovery panel, the F_{ST} value was very high (0.842). The high level of differentiation between *C. reticulata*, *C. maxima*, and *C. medica* for this SNP panel was well illustrated by neighbor-joining analysis (Fig. 1). The relative position of the accessions of secondary species (*C. aurantium* L., *C. aurantifolia*, *C. limon* (L.) Osbeck, *C. paradisi* Macfad., and *C. sinensis* (L.) Osbeck) and hybrids (Clementine, tangor, and tangelo) agrees with previous molecular studies (Nicolosi et al., 2000; Ollitrault et al., 2012; Garcia-Lor et al., 2012). Therefore, these markers should be useful as phylogenetic tracers of DNA fragments in secondary cultivated citrus species.

CONCLUSIONS

Forty-one SNP markers were successfully developed from SNP loci mined by Sanger sequencing in a discovery panel including 17 genotypes of the three main cultivated *Citrus* ancestral taxa. The genotyping data displayed high conformity with previous sequencing data. Genotyping was highly successful within the *Citrus* genus, and the genetic organization displayed by this SNP marker panel was in agreement with previous studies. The frequency of missing data was higher for the citrus relatives and increased with taxonomic distances within the Aurantioideae subfamily, suggesting incomplete transferability. The polymorphism revealed within the relatives of the “true citrus fruit trees” group remained relatively high but decreased

strongly when considering the other citrus relatives. However, all citrus relative genotypes were differentiated. The markers that were developed appeared to be useful for phylogenetic studies within the “true citrus fruit trees” group. Therefore, SNP markers based on the KASPar method developed from sequence data of a limited intragenomic discovery panel provide a valuable molecular resource for genetic diversity analysis of germplasm within a genus and should be useful for germplasm fingerprinting at a much broader diversity level.

LITERATURE CITED

- ALBRECHTSEN, A., F. C. NIELSEN, AND R. NIELSEN. 2010. Ascertainment biases in SNP chips affect measures of population divergence. *Molecular Biology and Evolution* 27: 2534–2547.
- BARKLEY, N. A., M. L. ROOSE, R. R. KRUEGER, AND C. T. FEDERICI. 2006. Assessing genetic diversity and population structure in a citrus germplasm collection utilizing simple sequence repeat markers (SSRs). *Theoretical and Applied Genetics* 112: 1519–1531.
- BAUER, F., C. C. ELBERS, R. A. H. ADAN, R. J. F. LOOS, N. C. ONLAND-MORET, D. E. GROBBEE, J. VAN VLIET-OSTAPCHOUK, ET AL. 2009. Obesity genes identified in genome-wide association studies are associated with adiposity measures and potentially with nutrient-specific food preference. *American Journal of Clinical Nutrition* 90: 951–959.
- CORTES, A. J., M. C. CHAVARRO, AND M. W. BLAIR. 2011. SNP marker diversity in common bean (*Phaseolus vulgaris* L.). *Theoretical and Applied Genetics* 123: 827–845.
- CUPPEN, E. 2007. Genotyping by allele-specific amplification (KASPar). *Cold Spring Harbor Protocols* 9: 172–173.
- GARCIA-LOR, A., F. LURO, L. NAVARRO, AND P. OLLITRAULT. 2012. Comparative use of InDel and SSR markers in deciphering the interspecific structure of cultivated citrus genetic diversity: A perspective for genetic association studies. *Molecular Genetics and Genomics* 287: 77–94.
- GARCIA-LOR, A., F. CURK, R. MORILLON, H. SNOUSSI, G. ANCILLO, F. LURO, L. NAVARRO, AND P. OLLITRAULT. 2013. A nuclear phylogenetic analysis: SNPs, indels and SSRs deliver new insights into the relationships in the “true citrus fruit trees” group (Citrinae, Rutaceae) and the origin of cultivated species. *Annals of Botany* 111: 1–19.
- LIU, K., AND S. V. MUSE. 2005. PowerMarker: An integrated analysis environment for genetic marker analysis. *Bioinformatics* 21: 2189–2192.
- NICOLOSI, E., Z. N. DENG, A. GENTILE, S. LA MALFA, G. CONTINELLA, AND E. TRIBULATO. 2000. Citrus phylogeny and genetic origin of important species as investigated by molecular markers. *Theoretical and Applied Genetics* 100: 1155–1166.
- NUMAN, I. J., S. KUIPERS, M. VERHEUL, V. GURYEV, AND E. CUPPEN. 2008. A genome-wide SNP panel for mapping and association studies in the rat. *BMC Genomics* 9: 95.
- OLLITRAULT, P., J. TEROL, A. GARCIA-LOR, A. BÉRARD, A. CHAUVEAU, Y. FROELICHER, C. BELZILE, ET AL. 2012. SNP mining in *C. clementina* BAC end sequences; transferability in the *Citrus* genus (Rutaceae), phylogenetic inferences and perspectives for genetic mapping. *BMC Genomics* 13: 13.
- PERRIER, X., AND J. P. JACQUEMOUD-COLLET. 2006. DARwin software: Dissimilarity Analysis and Representation for Windows. Website <http://darwin.cirad.fr/darwin> [accessed 1 March 2013].
- SWINGLE, W., AND P. REECE. 1967. The botany of *Citrus* and its wild relatives. In W. Reuther, H. J. Webber, and L. D. Batchelor [eds.], *The citrus industry*, vol. 1, 190–430. University of California, Berkeley, California, USA.

APPENDIX 1. Accessions analyzed in this study. Information presented: species name, Latin name or common name, accession number, ex-situ germplasm bank. IVIA = Carretera Moncada, Naquera, Km 4.4, Apartado Oficial, 46113 Moncada (Valencia), Spain; INRA/CIRAD = Station INRA, 20230 San Giuliano, France.

1. Citraea

Balsamocitrinae: *Aegle marmelos* (L.) Corrêa: 345, IVIA; *Aeglopsis chevalieri* Swingle: 308, IVIA; *Afraegle paniculata* (Schum. & Thonn.) Engl.: 273, IVIA; *Balsamocitrus dawei* Stapf: 372, IVIA; *Feroniella oblata* Swingle: 585, IVIA; *Swinglea glutinosa* (Blanco) Merr.: 292, IVIA.

Citrinae

True citrus fruit:

Citrus: *C. maxima* (Burm.) Merr.: Azimboa, 420, IVIA; Chandler, 207, IVIA; Da xanh, 589, IVIA; Deep red, 277, IVIA; Flores, 673, INRA/CIRAD; Gil, 321, IVIA; Nam roi, 590, IVIA; Pink, 275, IVIA; Sans Pepins, 710, INRA/CIRAD; Tahiti, 727, INRA/CIRAD; Timor, 707, INRA/CIRAD. *C. medica* L.: Arizona, 169, IVIA; Buddha hand, 202, IVIA; Corsican, 567, IVIA; Diamante, 560, IVIA; Humpang, 722, INRA/CIRAD; Poncire Commun, 701, INRA/CIRAD. *C. reticulata* Blanco: Bombay, 518, INRA/CIRAD; Dancy, 434, IVIA; De soe, 713, INRA/CIRAD; Imperial, 576, IVIA; Fuzhu, 571, IVIA; Ladu, 595, INRA/CIRAD; Ladu ordinaire, 590, INRA/CIRAD; Ponkan, 482, IVIA; Swatow, 175, INRA/CIRAD; Szinkom, 597, INRA/CIRAD; Vohangisany ambodiampoly, 437, SRA; Willow leaf, 154, IVIA. **Papeda:** *C. hystrix* DC.: Combava, 178, IVIA; *C. ichangensis* Swingle: Papeda Ichang, 358, IVIA; *C. micrantha* Wester: Micrantha, IVIA.

Secondary species: *C. aurantifolia* (Christm.) Swingle: Alemow, 288, IVIA; Calabria, 254, IVIA; Mexican, 164, IVIA. *C. aurantium* L.: Bouquet de fleurs, 139, IVIA; Cajel, 108, IVIA; Seville, 117, IVIA. *C. limon* (L.) Osbeck: Eureka frost, 297, IVIA; Rough lemon, 333, IVIA; Volkamer lemon, 432, IVIA; *C. paradisi* Macfad.: Duncan, 274, IVIA; Marsh, 176, IVIA; Rio red, 289, IVIA. *C. sinensis* (L.) Osbeck: Lane late, 198, IVIA; Sanguinelli, 34, IVIA; Valencia late, 363, IVIA.

Hybrids: Clementine, Clemenules, 22, IVIA; Tangelo, Orlando, 101, IVIA; Tangor, King, 477, IVIA.

Clymenia: *C. polyandra* (Tanaka) Swingle: 584, IVIA.

Eremocitrus: *E. glauca* (Lindl.) Swingle: 346, IVIA.

Fortunella: *F. crassifolia* Swingle: 280, IVIA; *F. hindsii* Swingle: 281, IVIA; *F. japonica* (Thunb.) Swingle: 381, IVIA; *F. margarita* (Lour.) Swingle: 38, IVIA; *Fortunella* sp.: 98, IVIA.

Microcitrus: *M. australasica* Swingle: 150, IVIA; *M. australis* Swingle: 313, IVIA; *M. australis* × *M. australasica*: 378, IVIA; Australian Wild Lime, 314, IVIA; New Guinea Wild Lime, 315, IVIA.

Poncirus trifoliata (L.) Raf.: Flying Dragon, 537, IVIA; Pomeroy, 374, IVIA; Rich 75, 236, IVIA; Rubidoux, 217, IVIA.

Near citrus fruit: *Atalantia ceylanica* (Arn.) Oliv.: 172, IVIA; *Atalantia citroides* Pierre ex Guillaumin, 284, IVIA; *Citropsis gillettiana* Swingle & M. Kellern.: 517, IVIA.

Primitive citrus fruit: *Hesperethusa crenulata* (Roxb.) M. Roem.: 580, IVIA; *Pleiospermium* sp., 380, IVIA; *Severinia buxifolia* (Poir.) Ten.: 147, IVIA; *Severinia disticha* (Blanco) Swingle: 418, IVIA.

Triphasilineae: *Triphasia trifolia* (Burm. f.) P. Wilson: 182, IVIA.

2. Clauseneae

Clauseniae: *Clausena excavata* Burm. f.: 311, IVIA; *Clausena lansium* (Lour.) Skeels: 343, IVIA; *Glycosmis pentaphylla* (Retz.) DC.: 148, IVIA; *Murraya koenigii* (L.) Spreng.: 377, IVIA.