



## **Development of Nuclear Microsatellites for the Arcto-Tertiary Tree *Zelkova carpinifolia* (Ulmaceae) Using 454 Pyrosequencing**

Authors: Maharramova, Elmira H., Muller, Ludo A., Korotkova, Nadja, and Borsch, Thomas

Source: Applications in Plant Sciences, 2(3)

Published By: Botanical Society of America

URL: <https://doi.org/10.3732/apps.1300072>

---

BioOne Complete ([complete.BioOne.org](http://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## DEVELOPMENT OF NUCLEAR MICROSATELLITES FOR THE ARCTO-TERTIARY TREE *ZELKOVA CARPINIFOLIA* (ULMACEAE) USING 454 PYROSEQUENCING<sup>1</sup>

ELMIRA H. MAHARRAMOVA<sup>2,3,5</sup>, LUDO A. MULLER<sup>4</sup>, NADJA KOROTKOVA<sup>2,4</sup>,  
AND THOMAS BORSCH<sup>2,4</sup>

<sup>2</sup>Botanischer Garten und Botanisches Museum Berlin-Dahlem, Freie Universität Berlin, Königin Luise-Straße 6–8, 14195 Berlin, Germany; <sup>3</sup>Institute of Botany, Azerbaijan National Academy of Sciences, 40 Badamdar Highway, Baku AZ1073, Azerbaijan; and <sup>4</sup>Institut für Biologie-Botanik, Dahlem Centre of Plant Sciences, Freie Universität Berlin, Altensteinstraße 6, 14195 Berlin, Germany

- *Premise of the study:* The current study aimed at developing nuclear microsatellite markers for the relict tree species *Zelkova carpinifolia*, which is threatened in its natural range in the South Caucasus.
- *Methods and Results:* Pyrosequencing of an enriched microsatellite library on the Roche FLX platform using the 454 Titanium kit produced 86,058 sequence reads, most of which contained short tandem repeats. Eighty microsatellite loci identified using the software package QDD version 1 were selected and tested for proper PCR amplification. Of these, 13 allowed proper amplification and were shown to be polymorphic among a sample of 25 *Z. carpinifolia* specimens from various geographic origins.
- *Conclusions:* The set of microsatellite markers will be suitable for the assessment of genetic diversity in *Z. carpinifolia*. They will allow for an examination of phylogeographic patterns as well as of population structure and gene flow within this species.

**Key words:** conservation genetics; microsatellites; phylogeography; pyrosequencing; Ulmaceae; *Zelkova carpinifolia*.

*Zelkova carpinifolia* (Pall.) K. Koch. (Ulmaceae) is a representative species of the Arcto-Tertiary relict flora (Wang et al., 2001; Milne and Abbot, 2002; Kozłowski et al., 2012). It occurs naturally in Hyrcan and Colchic forests in Iran, Azerbaijan, Georgia, and Turkey (Browicz and Zielinski, 1982; Denk and Grimm, 2005; Kvavadze and Connor, 2005; Akhiani et al., 2010). Its valuable timber has led to uncontrolled logging, which has resulted in a general population decline. Therefore, *Z. carpinifolia* was listed under the International Union for Conservation of Nature (IUCN) category “Near Threatened” (Güner and Zielinski, 1998). To plan conservation strategies for *Z. carpinifolia*, investigation of its genetic diversity is of great importance (Kozłowski et al., 2012). The phylogeography of the species needs to be understood as there may be specific refugia within the range of the species that harbor relatively high genetic diversities and regionally adapted genotypes may occur.

The first studies of genetic diversity in *Zelkova* Spach employed inter-simple sequence repeat (ISSR) and amplified fragment

length polymorphism (AFLP) markers (Fineschi et al., 2004). Primers targeting microsatellite loci in *Z. serrata* (Thunb.) Makino (Fukatsu et al., 2005) did not amplify well in *Z. carpinifolia* and revealed only limited polymorphism (unpublished data). The current study therefore aimed to develop polymorphic microsatellite markers for *Z. carpinifolia* using 454 pyrosequencing.

### METHODS AND RESULTS

**Plant material and DNA extraction**—*Zelkova carpinifolia* samples were collected from six populations (10–20 individuals per population) in Azerbaijan and Iran (Appendix 1). Genomic DNA was extracted from silica gel-dried leaves using either the NucleoSpin Plant II kit (Macherey Nagel, Düren, Germany) following the manufacturer’s protocol or a three-fraction cetyltrimethylammonium bromide (CTAB) method (Borsch et al., 2003).

**Microsatellite marker development**—Construction of the enriched microsatellite libraries was carried out according to the protocol of Glenn and Schable (2005) with modifications. Genomic DNA was digested in 25- $\mu$ L reactions containing 1 $\times$  NEBuffer 3.1, 0.1 mg/mL bovine serum albumin (BSA), 0.05 M NaCl, 20 U *EcoRI*, 10 U *MseI* restriction enzymes (New England Biolabs, Frankfurt, Germany), and 2.4  $\mu$ g genomic DNA during a 2-h incubation at 37°C. Digested DNA was run on a 2% agarose gel, and 200–400-bp fragments were extracted using the Gel/PCR DNA Fragment Extraction Kit (Avegene, Zollikofen, Switzerland) following the manufacturer’s protocol. To prepare double-stranded adapters, forward (*EcoRI*: 5'-CTCGTAGACTGCGTACC, *MseI*: 5'-GACGATGAGTCCTGAG) and reverse (*EcoRI*: 5'-AATGGTACGCAGTCTACGAG, *MseI*: 5'-TACTCAGGACTCATCGTC) adapters (Metabion, Martinsried, Germany) were mixed in equimolar amounts (10  $\mu$ M), incubated at 95°C for 5 min, and allowed to slowly cool down to room temperature. Adapters were ligated to the restriction fragments in 40- $\mu$ L reactions containing 0.8  $\mu$ M of each of the double-stranded *EcoRI* and *MseI* adapters, 1 $\times$  T4 DNA

<sup>1</sup>Manuscript received 4 September 2013; revision accepted 17 November 2013.

Pyrosequencing was performed at the Berlin Center for Genomics in Biodiversity Research (BeGenDiv). We especially wish to thank S. Mbedi for laboratory assistance and M. Jafari for providing *Zelkova* samples from Iran. The study was carried out as part of the project “Developing tools for conserving the plant diversity of the Transcaucasus” funded by VolkswagenStiftung. This is publication number 005 of BeGenDiv.

<sup>5</sup>Author for correspondence: e.maharramova@bgbm.org

doi:10.3732/apps.1300072

*Applications in Plant Sciences* 2014 2(3): 1300072; <http://www.bioone.org/loi/apps> © 2014 Maharramova et al. Published by the Botanical Society of America. This work is licensed under a Creative Commons Attribution License (CC-BY-NC-SA).

TABLE 1. Characteristics of 13 nuclear microsatellite loci for *Zelkova carpinifolia*.

Locus	Primer sequences (5'–3')	Fluorescent label	Repeat motif	$T_a$ (°C)	Allele size range (bp)	GenBank/ENA accession no.
ZMS_1	F: AAGGTTTCATCCCATGAGCAC* R: GTCCAGTCCGGCATCTTTT	6-FAM	(CTT) <sub>7</sub>	60	238–259	KF408341
ZMS_2	F: CAAGCTGCCAATACTCCACA* R: GGTTTGGTTTCTGCTTGGAA	VIC	(AGA) <sub>7</sub>	60	181–213	KF408342
ZMS_3	F: CCCAACAGCACTTGAAGCTA* R: TTTGTGTAGTCTTTGGCCCC	6-FAM	(CA) <sub>8</sub>	60	136–138	KF408343
ZMS_4	F: ACCGTTGGAACAAAACACTCG* R: AGGGCAGTAGAGCTTGATCG	NED	(AG) <sub>9</sub>	52	105–131	KF408344
ZMS_5	F: TTTTGTTCGTTGTTGCATGTC* R: AGGCTTAGGTTGGGTGGTTT	PET	(GAA) <sub>6</sub>	60	99–114	KF408345
ZMS_6	F: TTTGGTGCCTTAGCATGTGA* R: TCTTTTGTTTGCGATAAGTCG	6-FAM	(CT) <sub>8</sub>	58	101–120	KF408346
ZMS_7	F: TCTCTGTTATGATCACGCTCCA R: GAGCTAAACCCCATTTCAAATATA*	NED	(AC) <sub>9</sub>	60	120–128	KF408347
ZMS_8	F: ACGAACAAGATCCGACAGAGG* R: ATTGGATGGCACACGTAATTC	PET	(TC) <sub>12</sub>	60	188–204	KF408348
ZMS_9	F: CAGGTGATTGGAATTGGAAAA* R: CCTCACAGAGTACGAGGGATG	6-FAM	(AC) <sub>7</sub>	62	171–196	KF408350
ZMS_10	F: CTGCGAGGGCAGTAGGTAAG* R: TCCTAGATCCTTTGATATCTTTCTCA	VIC	(GA) <sub>8</sub>	62	175–189	KF408351
ZMS_11	F: AGTAGGCTTGGAAAGCAAATGA R: TTGGAACCTGCAATCGCTTT*	VIC	(GT) <sub>9</sub>	58	96–115	KF408352
ZMS_12	F: CTTCCAATCATTCGAGAGC* R: TGACATCTCAAAGGTTGCCA	PET	(CA) <sub>7</sub>	58	100–104	KF408353
ZMS_13	F: AACAAAGAACTTTCATCCCC* R: GCATTTGGTTTTACCTAAAGTTACAG	NED	(TG) <sub>9</sub>	58	110–116	KF408354
ZMS_14	F: GCGGTTGAGAGGAGAAAGAA R: ACCAAACCCATCAACCATTG*	6-FAM	(AGG) <sub>10</sub>	57	105 <sup>†</sup>	HG737351
ZMS_15	F: GCTCCTCTTCAGCCAAACAC* R: CTTTCATCCTCGTTTGCATCA	6-FAM	(AGG) <sub>12</sub>	60	128 <sup>†</sup>	HG737352
ZMS_16	F: GCCGGTACTGATTCTTCAA* R: GGACCATCTCACCAACATCC	6-FAM	(AAG) <sub>5</sub>	60	233 <sup>†</sup>	HG737353
ZMS_17	F: AGCCCGTGTCTTTGAATATG R: GGATAAGGCTATTTTGGCCC*	6-FAM	(GA) <sub>8</sub>	55	167 <sup>†</sup>	HG737354
ZMS_18	F: GCCACGTAACACAGATAAGGG* R: GCATGTTGGTGCTTTGAGAA	6-FAM	(TG) <sub>7</sub>	58	103 <sup>†</sup>	HG737355

Note: ENA = European Nucleotide Archive;  $T_a$  = optimal annealing temperature.

\*Labeled primer.

<sup>†</sup>Expected size.

ligase buffer, and 600 U T4 DNA ligase (New England Biolabs). Reactions were performed at 16°C for 12 h and successful ligation was tested by PCR amplification in 50- $\mu$ L reactions containing 4  $\mu$ L of the adapter-ligated DNA, 1 $\times$  *Taq* DNA polymerase buffer, 1 $\times$  PCR enhancer solution, 150  $\mu$ M of each dNTP, 2 U *Taq* DNA polymerase (PeqLab, Erlangen, Germany), 35 ng/ $\mu$ L BSA, 0.5  $\mu$ M *Eco*RI primer (5'-CTCGTAGACTGCGTACCAATTC), and 0.5  $\mu$ M *Mse*I primer (5'-GACGATGAGTCTCTGAGTAA) (Metabion) using the following temperature profile: 95°C for 2 min; 20 cycles of 95°C for 20 s, 60°C for 20 s, 72°C for 1.5 min; and 72°C for 10 min. Two mixtures of 3'-biotinylated oligonucleotides [1: (AG)<sub>12</sub>, (TG)<sub>12</sub>, (AAG)<sub>8</sub>, (AAT)<sub>12</sub>, (ACT)<sub>12</sub> and 2: (ACAG)<sub>6</sub>, (ACCT)<sub>6</sub>, (ACTC)<sub>6</sub>, (ACTG)<sub>6</sub>] were used to generate two separate libraries of adapter-ligated genomic DNA enriched for repetitive motifs. Hybridization of biotinylated oligonucleotides and adapter-ligated genomic DNA, followed by the capturing of hybridized DNA on Dynabeads (Life Technologies, Darmstadt, Germany) and subsequent washing and resuspension, followed the protocol of Glenn and Schable (2005). Recovery PCR was performed with 2  $\mu$ L of enriched DNA in the same reaction mixture and the temperature profile as described above, but with 25 cycles. The enrichment procedure was repeated and the second recovery PCR was carried out with an *Eco*RI oligonucleotide primer extended with a Roche 454 pyrosequencing adapter (5'-GCCTC-CCTCGGCCATCAG-3') and a specimen-specific barcode sequence and an *Mse*I oligonucleotide primer extended with a Roche 454 B sequencing adapter (5'-GCCTTGCCAGCCCGCTCAG-3'; Appendix 2). Two enrichment libraries were prepared for each of six *Z. carpinifolia* specimens (population ZE053\_IR, Appendix 1). Enrichment libraries were purified using the Amplicon Library Preparation Protocol (Roche, Branford, Connecticut, USA) according to the manufacturer's instructions. Lib-L Kits and the GS FLX Titanium Sequencing Kit XLR70 (Roche) were used for the library sequencing on the Roche GS FLX

System. The resulting DNA sequences were analyzed using the R package SeqinR (Charif and Lobry, 2007). The 454 pyrosequencing yielded 86,058 total reads ranging from 40 to 577 bp with an average length of 138 bp.

**Microsatellite screening**—Screening for microsatellite loci and primer design was performed using QDD software version 1 (Megléc et al., 2010), which identified 136 microsatellite loci containing di-, tri-, and tetranucleotide repeat motifs. A total of 80 microsatellite loci with 5–45 repeats, including three loci with tetranucleotide repeats, 17 loci with trinucleotide repeats, and 60 loci with dinucleotide repeats, were selected randomly, and oligonucleotide primer combinations with a GC content of 35–60% and melting temperature ( $T_m$ ) ranging between 57°C and 60°C were tested for proper PCR amplification on genomic DNA extracted from three specimens (Table 1). A gradient PCR was carried out for each primer pair in a 25- $\mu$ L reaction volume containing 20–40 ng DNA, 0.4  $\mu$ M each of forward and reverse primer (Eurofins MWG Operon, Ebersberg, Germany), 1 $\times$  *Taq*Buffer S, 1.5 mM MgCl<sub>2</sub>, 250  $\mu$ M of each dNTP, and 0.75 U Hot *Taq* polymerase (PeqLab). The temperature profile was as follows: 96°C for 2 min, 54  $\pm$  8°C for 30 s, 72°C for 30 s; followed by 34 cycles of 95°C for 30 s, 54  $\pm$  8°C for 30 s, 72°C for 30 s; and 72°C for 15 min. PCR performance was assessed by gel electrophoresis. Of the initial 80 loci tested, 40 primer pairs failed to amplify a product. The reactions with the other 40 primer pairs were repeated in 50- $\mu$ L reaction volumes. PCR products were run on 1.5% agarose gel and extracted using the NucleoSpin Gel and PCR Clean-up Kit (Macherey-Nagel) following the manufacturer's protocol. PCR products were sequenced at Macrogen (Seoul, Korea) using standard Sanger sequencing to confirm locus-specificity of the oligonucleotide primers. Only the 20 primer pairs that specifically amplified the targeted microsatellite loci were retained for further analyses. Polymorphism

TABLE 2. Results of initial primer screening of specimens from five populations of *Zelkova carpinifolia*.

Locus	ZE001_AZ (N = 5)			ZE007_AZ (N = 5)			ZE013_AZ (N = 5)			ZE017_AZ (N = 5)			ZE021_AZ (N = 5)			Total (N = 25)			
	A	H <sub>o</sub>	H <sub>e</sub>	A	H <sub>o</sub>	H <sub>e</sub>	A	H <sub>o</sub>	H <sub>e</sub>	A	H <sub>o</sub>	H <sub>e</sub>	A	H <sub>o</sub>	H <sub>e</sub>	A	H <sub>o</sub>	H <sub>e</sub>	PIC
ZMS_1	2	0.60	0.4667	4	1.00	0.7778	2	0.80	0.5333	4	1.00	0.7111	4	0.60	0.6444	5	0.80	0.6294	0.5536
ZMS_2	3	1.00	0.7111	4	1.00	0.7333	4	0.60	0.5333	4	0.80	0.7333	3	0.80	0.6444	8	0.84	0.7053	0.6439
ZMS_3	2	1.00	0.5556	2	0.80	0.5333	2	0.80	0.5333	2	0.20	0.5556	2	0.20	0.4667	2	0.60	0.4971	0.3685
ZMS_4	3	0.80	0.6000	4	0.80	0.8000	5	1.00	0.8000	5	1.00	0.8444	5	0.40	0.8000	8	0.76	0.8188	0.7811
ZMS_5	3	0.40	0.3778	2	0.40	0.3556	1	0	0	1	0	0	3	0.60	0.5111	5	0.28	0.2596	0.2455
ZMS_6	2	0	0.3556	2	0	0.3556	3	0	0.6222	5	0.40	0.8667	5	0.20	0.8222	8	0.12	0.8049	0.7599
ZMS_7	3	1.00	0.6444	4	1.00	0.7778	4	0.60	0.6444	2	0.80	0.5333	3	0.80	0.6000	4	0.84	0.7094	0.6413
ZMS_8	5	1.00	0.8222	4	1.00	0.7333	4	1.00	0.7333	3	0.80	0.6000	4	0.20	0.6444	7	0.80	0.7351	0.7351
ZMS_9	3	0.40	0.6222	3	0	0.7111	4	0.60	0.7333	3	0.60	0.5111	3	0.20	0.5111	5	0.36	0.7127	0.6506
ZMS_10	2	0.20	0.5556	4	0.60	0.7333	3	0.40	0.6000	2	0.20	0.4667	4	0.80	0.7778	5	0.44	0.6588	0.6065
ZMS_11	1	0	0	3	0.40	0.7111	2	0	0.3556	5	0.40	0.8222	2	0	0.3556	6	0.16	0.5037	0.4526
ZMS_12	2	0.60	0.4667	2	0.40	0.3556	2	0.20	0.2000	2	0.20	0.2000	1	0	0	3	0.28	0.2539	0.2316
ZMS_13	3	0.60	0.6444	3	1.00	0.6444	2	0.80	0.5333	2	0.20	0.2000	2	0.20	0.2000	4	0.56	0.5102	0.4431

Note: A = number of alleles; H<sub>e</sub> = expected heterozygosity (Nei's unbiased gene diversity); H<sub>o</sub> = observed heterozygosity; N = sample size; PIC = polymorphism information content.

assessment used genomic DNA extracted from 25 *Z. carpinifolia* specimens (five specimens per population; Appendix 1) and fluorescently labeled forward or reverse primers (Applied Biosystems, Warrington, United Kingdom), and the same reaction conditions as in the previous step. Two primer pairs did not amplify a product in all individuals and were excluded from further analysis. The rest of the PCR products were sent to Macrogen for fragment analysis.

**Microsatellite marker data analysis**—Electropherograms from the fragment analysis were analyzed using GeneMarker software version 1.95 (SoftGenetics, State College, Pennsylvania, USA). Analyzed amplification products with unscorable peaks were not considered to be useful in the studies of variability due to the possible misinterpretation of data. Thirteen primer pairs produced polymorphic and heterozygous loci (accession numbers KF408341–KF408348, KF408350–KF408354). The Excel Microsatellite Toolkit (Park, 2001) was used for the calculation of the number of alleles, observed (H<sub>o</sub>) and expected (H<sub>e</sub>) heterozygosities, and polymorphism information content (PIC) values. Allele sizes ranged from two to eight alleles per locus across 25 individuals from five different populations, with H<sub>o</sub> and H<sub>e</sub> varying between 0.12 and 0.84, and 0.25 and 0.81, respectively (Table 2).

### CONCLUSIONS

In the current study, we used 454 pyrosequencing to obtain sequence reads for microsatellite primer development in *Z. carpinifolia*. After thorough evaluation, we present 13 primer pairs for amplifying polymorphic microsatellite loci in *Z. carpinifolia*. These 13 loci are proposed to be used for the microsatellite screening of naturally distributed *Z. carpinifolia* populations, but they may also cross-amplify in closely related species. Analysis of the obtained allele frequencies and knowledge of the individual genotypes will help us to understand the population structure and relationships between and within different populations of *Zelkova* species.

### LITERATURE CITED

AKHANI, H., M. DJAMALI, A. GHORBANALIZADEH, AND E. RAMEZANI. 2010. Plant biodiversity of Hyrcanian relict forests, N Iran: An overview of the flora, vegetation, palaeoecology and conservation. *Pakistan Journal of Botany* 42: 231–258.

BORSCH, T., K. W. HILU, D. QUANDT, V. WILDE, C. NEINHUIS, AND W. BARTHOLOTT. 2003. Noncoding plastid *trnT-trnF* sequences reveal a well resolved phylogeny of basal angiosperms. *Journal of Evolutionary Biology* 16: 558–576.

BROWICZ, K., AND J. ZIELINSKI. 1982. Chronology of trees and shrubs in south-west Asia and adjacent regions. Polish Academy of Sciences, Institute of Dendrology, Polish Scientific Publishers, Warsaw, Poland.

CHARIF, D., AND J. LOBRY. 2007. SeqinR 1.0-2: A contributed package to the R project for statistical computing devoted to biological sequences retrieval and analysis. In U. Bastolla, M. Porto, H. E. Roman, and M. Vendruscolo [eds.], Structural approaches to sequence evolution, 207–232. Springer, Berlin, Germany.

DENK, T., AND G. W. GRIMM. 2005. Phylogeny and biogeography of *Zelkova* (Ulmaceae sensu stricto) as inferred from leaf morphology, ITS sequence data and the fossil record. *Botanical Journal of the Linnean Society* 147: 129–157.

FINESCHI, S., S. COZZOLINO, M. MIGLIACCIO, AND G. G. VENDRAMIN. 2004. Genetic variation of relic tree species: The case of Mediterranean *Zelkova abelicea* (Lam.) Boissier and *Z. sicula* Di Pasquale, Garfi and Quézel (Ulmaceae). *Forest Ecology and Management* 197: 273–278.

FUKATSU, E., K. ISODA, T. HIRAO, M. TAKAHASHI, AND A. WATANABE. 2005. Development and characterization of simple sequence repeat DNA markers for *Zelkova serrata*. *Molecular Ecology Notes* 5: 378–380.

GLENN, T. C., AND N. A. SCHABLE. 2005. Isolating microsatellite DNA loci. In E. A. Zimmer and E. H. Roalson [eds.], *Methods in enzymology*, 202–222. Academic Press, San Diego, California, USA.

GÜNER, A., AND J. ZIELINSKI. 1998. *Zelkova carpinifolia*. In IUCN 2013–IUCN Red List of Threatened Species. Version 2013.1. Website <http://www.iucnredlist.org> [accessed 22 August 2013].

- KOZŁOWSKI, G., D. GIBBS, F. HUAN, D. FREY, AND J. GRATZFELD. 2012. Conservation of threatened relict trees through living ex situ collections: Lessons from the global survey of the genus *Zelkova* (Ulmaceae). *Biodiversity and Conservation* 21: 671–685.
- KVAVADZE, E. V., AND S. E. CONNOR. 2005. *Zelkova carpinifolia* (Pallas) K. Koch in Holocene sediments of Georgia: An indicator of climatic optima. *Review of Palaeobotany and Palynology* 133: 69–89.
- MEGLÉCZ, E., C. COSTEDOAT, V. DUBUT, A. GILLES, T. MALAUSA, N. PECH, AND J. F. MARTIN. 2010. QDD: A user-friendly program to select microsatellite markers and design primers from large sequencing projects. *Bioinformatics (Oxford, England)* 26: 403–404.
- MILNE, R. I., AND R. J. ABBOT. 2002. The origin and evolution of tertiary relict floras. *Advances in Botanical Research* 38: 281–314.
- PARK, S. D. E. 2001. Trypanotolerance in west African cattle and the population genetic effects of selection. Ph.D. thesis, University of Dublin, Dublin, Ireland.
- WANG, Y.-F., D. K. FERGUSON, R. ZETTER, T. DENK, AND G. GARFI. 2001. Leaf architecture and epidermal characters in *Zelkova*, Ulmaceae. *Botanical Journal of the Linnean Society* 136: 255–265.

APPENDIX 1. Locations and voucher information for populations of *Zelkova carpinifolia* used in this study. Vouchers are deposited at the herbarium of the Botanischer Garten und Botanisches Museum Berlin-Dahlem (B), Berlin, Germany.

Population	Locality	Geographic coordinates	Collection no.
ZE001_AZ	Zuvand Mtn., Lerik, Azerbaijan	38°43'33"N, 48°25'53"E	EM01
ZE007_AZ	Azfilial, Lankaran, Azerbaijan	38°40.938'N, 48°47.222'E	EM13
ZE013_AZ	Tezekend, Lankaran, Azerbaijan	38°39.091'N, 48°48.284'E	EM20
ZE017_AZ	Zungulesh, Astara, Azerbaijan	38°27.175'N, 48°47.575'E	EM28
ZE021_AZ	Khanbulan, Lankaran, Azerbaijan	38°39.809'N, 48°46.649'E	EM33
ZE053_IR*	Gorgan, Golestan, Iran	36°43'35"N, 54°34'49"E	

\* Used for construction of the genomic library.

APPENDIX 2. *EcoRI*-A and *MseI*-B primers used for PCR recovery of the enriched DNA library.

Primer name	Primer sequence (5'–3') <sup>a</sup>
EcoRI_A001	CCA TCT CAT CCC TGC GTG TCT CCG ACT CAG <b>ACG AGT GCG</b> TCT CGT AGA CTG CGT ACC AAT TC
EcoRI_A002	CCA TCT CAT CCC TGC GTG TCT CCG ACT CAG <b>ACG CTC GAC</b> ACT CGT AGA CTG CGT ACC AAT TC
EcoRI_A003	CCA TCT CAT CCC TGC GTG TCT CCG ACT CAG <b>AGA CGC ACT</b> CCT CGT AGA CTG CGT ACC AAT TC
EcoRI_A004	CCA TCT CAT CCC TGC GTG TCT CCG ACT CAG <b>AGC ACT GTA</b> GCT CGT AGA CTG CGT ACC AAT TC
EcoRI_A005	CCA TCT CAT CCC TGC GTG TCT CCG ACT CAG <b>ATC AGA CAC</b> GCT CGT AGA CTG CGT ACC AAT TC
EcoRI_A006	CCA TCT CAT CCC TGC GTG TCT CCG ACT CAG <b>ATA TCG CGA</b> GCT CGT AGA CTG CGT ACC AAT TC
MseI_B	CCT ATC CCC TGT GTG CCT TGG CAG TCT CAG GAC GAT GAG TCC TGA GTA A

<sup>a</sup> Specimen-specific barcodes (tags) are shown in bold.