



## Highly Polymorphic Microsatellite Markers in *Pulsatilla vulgaris* (Ranunculaceae) Using Next-Generation Sequencing

Authors: DiLeo, Michelle F., Graf, René, Holderegger, Rolf, Rico, Yessica, and Wagner, Helene H.

Source: Applications in Plant Sciences, 3(7)

Published By: Botanical Society of America

URL: <https://doi.org/10.3732/apps.1500031>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## HIGHLY POLYMORPHIC MICROSATELLITE MARKERS IN *PULSATILLA VULGARIS* (RANUNCULACEAE) USING NEXT-GENERATION SEQUENCING<sup>1</sup>

MICHELLE F. DiLEO<sup>2,5</sup>, RENÉ GRAF<sup>3</sup>, ROLF HOLDEREGGER<sup>3</sup>, YESSICA RICO<sup>4</sup>,  
AND HELENE H. WAGNER<sup>2</sup>

<sup>2</sup>Department of Biology, University of Toronto Mississauga, 3359 Mississauga Road, Mississauga, Ontario L5L 1C6, Canada;

<sup>3</sup>WSL Swiss Federal Research Institute, Zürcherstrasse 111, CH-8903 Birmensdorf, Switzerland; and <sup>4</sup>Department of Natural History, Royal Ontario Museum, 100 Queens Park, Toronto, Ontario M5S 2C6, Canada

- *Premise of the study:* We developed novel microsatellite markers for the perennial plant *Pulsatilla vulgaris* (Ranunculaceae) to investigate the effects of fragmentation on gene flow in this imperiled species.
- *Methods and Results:* We identified microsatellites and developed primers based on 454 shotgun sequences. We identified 14 markers that were polymorphic and produced clean bands. Of these, eight could be analyzed as diploids. Genotyping of 97 individuals across two populations revealed these markers to be highly polymorphic with seven to 17 alleles per locus and observed heterozygosity from 0.41 to 0.83.
- *Conclusions:* The markers are highly informative and will be used to test if the reintroduction of shepherding in southern Germany improves genetic connectivity among fragmented populations of *P. vulgaris*. The combination of diploid and tetraploid markers presented here will be useful in resolving the polyploidization history of this and related species.

**Key words:** 454 sequencing; microsatellites; *Pulsatilla vulgaris*; Ranunculaceae; simple sequence repeat (SSR) markers.

*Pulsatilla vulgaris* Mill. (Ranunculaceae) is an early-flowering perennial herb of conservation concern and a flagship and specialist species of calcareous grasslands across central Europe, ranging from France in the south to Sweden at its northern limit (Wells and Barling, 1971; Pfeifer et al., 2002). Over the last century, *P. vulgaris* has witnessed rapid decline and local extirpation across its range, and is listed as “near threatened” by the International Union for Conservation of Nature (IUCN, 2014). Its decline is linked to the abandonment of traditional grazing practices, which has resulted in the severe loss and fragmentation of calcareous grasslands following afforestation (Butaye et al., 2005), and increased above-ground competition from coarse grasses (Walker and Pinches, 2011). Knowledge of the landscape-scale distribution of genetic variation is required to create effective management plans for fragmented populations, and evaluations of genetic diversity and inbreeding will allow the identification of populations that are at highest risk of extirpation. This, however, requires genetic markers with high resolution such as microsatellites. No such markers are yet available for *P. vulgaris*, and any potential genetic analyses are

complicated as this species is suspected to be an allotetraploid ( $2n = 32$ ; Böcher, 1934). We therefore de novo developed microsatellite markers for this calcareous plant species, specifically selecting loci that can be analyzed as diploid or double-diploids in downstream genetic analyses.

### METHODS AND RESULTS

We extracted genomic DNA from homogenized leaf tissue from two populations in the Franconian Alb, Germany (A25: 48°57'38.7"N, 10°56'32.2"E; A03: 48°59'51.7"N, 11°3'35.4"E), using the QIAGEN DNeasy Plant Mini Kit (QIAGEN, Mississauga, Ontario, Canada) following the manufacturer's protocol. A voucher from the study area is deposited at the Herbarium at the University of Toronto Mississauga (TRTE; voucher no.: BR-07052015; locality: Laubenthal, Landkreis Weissenburg, Germany; 48°58'48.7130"N, 11°03'14.2254"E). DNA from five individuals was mixed, and a 4- $\mu$ L sample was sent to LGC Genomics (Berlin, Germany) for 454 shotgun sequencing on 1/8 of a plate using GS FLX Titanium (Roche, 454 Life Sciences, Branford, Connecticut, USA). In total, 92,833 reads were returned and assembled into 4088 contigs. We used MSATCOMMANDER (version 1.0.8; Faircloth, 2008) to identify di-, tri-, and tetranucleotide microsatellites to develop primers in regions flanking the identified microsatellites. A total of 457 microsatellites were identified, and 76 suitable primer pairs could be designed. We amplified 75 of these pairs by PCR in an initial screen of 10 individuals, and of these, 18 showed clear peaks and 14 of these were polymorphic (Table 1). Eight markers showed evidence of disomic inheritance (e.g., had a maximum of two alleles per individual) and were retained for further analysis (Table 2).

We amplified these eight microsatellites in two multiplex reactions in 97 individuals with the QIAGEN Multiplex PCR Kit (QIAGEN) in 10- $\mu$ L final reaction volumes with 0.2  $\mu$ M of each primer, 4.6  $\mu$ L of Multiplex Mix, 1.2  $\mu$ L of DNase-free water, and 5–30 ng of template DNA. Thermocycling conditions for Multiplex A (Table 1) strictly followed the manufacturer's protocol. Multiplex B was amplified using a touchdown procedure with an initial denaturation at 95°C for 15 min; 10 cycles of denaturation at 95°C for 60 s, annealing at

<sup>1</sup>Manuscript received 23 March 2015; revision accepted 30 April 2015.

The authors thank Nimesh Patel, Amaneet Lochab, and Qasim Muhammad for assistance with DNA extraction and genotyping. We also thank Juergen Boehmer and Mr. Dadrich for help in the field. Funding was provided by the Natural Sciences and Engineering Research Council of Canada (NSERC CGS-D and Michael Smith Foreign Study Supplement to M.F.D., and Discovery Grant to H.H.W.) and the Government of Central Franconia, Bavaria, Germany.

<sup>5</sup>Author for correspondence: michelle.dileo@mail.utoronto.ca

doi:10.3732/apps.1500031

*Applications in Plant Sciences* 2015 3(7): 1500031; <http://www.bioone.org/loi/apps> © 2015 DiLeo et al. Published by the Botanical Society of America.

This work is licensed under a Creative Commons Attribution License (CC-BY-NC-SA).

TABLE 1. Characteristics of 18 de novo–developed microsatellite markers for *Pulsatilla vulgaris*.

Locus	Primer sequences (5'–3')	Repeat motif	T <sub>a</sub> (°C)	Multiplex	Fluorescent dye <sup>a</sup>	Allele size range (bp)	A	Marker ploidy	GenBank accession no.
PV2	F: GTTTGCGATGATCATGAAGTGC R: AGAACTTCCACGAACAAGGC	(AAAT) <sub>6</sub>	55	A	HEX	410–426	4	diploid	KP885677
PV7	F: ACCGCCAACATAGCAAACAC R: ACCCACCACAACCTTGAGAGG	(AG) <sub>10</sub>	TD 65–55	B	6-FAM	326–362	8	diploid	KP885678
PV9	F: GAACTAACCTGCTTCCGTGC R: GCAAGCCAAAGTCCACTCTG	(AG) <sub>11</sub>	55	single	HEX	283–309	12	tetraploid	KP885679
PV16	F: TTGTTGGGTGCGTTGAGAAG R: ATCACTTGTAGCCTCCGCTC	(AC) <sub>12</sub>	55	single	6-FAM	330–362	11	tetraploid	KP885680
PV18	F: CCATGAACCCCTTGTACAGAG R: AGGACCCAAAGATCTCAAGCC	(AC) <sub>10</sub>	55	single	6-FAM	415–451	11	tetraploid	KP885681
PV27	F: AACCCCTTGCACCAACTTG R: AATCTTATCTGGCGGGGAGG	(AG) <sub>8</sub>	55	A	6-FAM	392–454	5	diploid	KP885682
PV32	F: CATGCCTTTGTACCCTGCTG R: ACGACCTTTGTTTGACCGTC	(AAG) <sub>9</sub>	55	single	6-FAM	356–422	8	tetraploid	KP885683
PV33	F: AGCCTTGGTGTATTTGGGG R: GCTCACTTGGACCAACTCCC	(AC) <sub>9</sub>	TD 65–55	B	6-FAM	442–452	8	diploid	KP885684
PV44	F: GTATGTGTGTGCCAAGGGTC R: TGCTTAAAGTAGCATGCCG	(ATC) <sub>10</sub>	55	single	HEX	429	1	—	KR109213
PV48	F: CGGGCTGTAAGTATGCTTC R: GCAAGTGAGCCAGTCCATC	(AG) <sub>10</sub>	55	single	HEX	235	1	—	KR109214
PV50	F: GATGGTGTGAGGGTTTGGG R: TGCCACCTACTTTCCACACC	(AAT) <sub>12</sub>	55	single	HEX	406–466	10	tetraploid	KP885685
PV52	F: TTGGTCAAATGGTCGCAACG R: GGTGCTCAAGATTATCGGGC	(AG) <sub>10</sub>	55	A	6-FAM	262–280	7	diploid	KP885686
PV54	F: TACTCGCGACTGACAAGCTC R: TTGTGGGAGTGGAAAGAAC	(AT) <sub>10</sub>	55	single	6-FAM	204	1	—	KR109215
PV56	F: CGTCACGTGTGGGCTAAATC R: TCCTTTGTACTCTCCAACGG	(AC) <sub>11</sub>	TD 65–55	B	6-FAM	236–270	9	tetraploid	KP885687
PV57	F: GTGCAAATTAACACTGCAG R: TGCTCGAAACCATAAGTCTGC	(AC) <sub>10</sub>	55	single	6-FAM	419–429	5	tetraploid	KP885688
PV59	F: TGGACAACGTACCTTACACAG R: AGCTTCCACACCTCAACTGG	(AG) <sub>10</sub>	55	single	HEX	171	1	—	KR109216
PV64	F: AGCAATCTCTGTCCGGCTAG R: GTGACTGCAGATGTGGTGG	(AC) <sub>13</sub>	TD 65–55	B	HEX	368–404	7	diploid	KP885689
PV65	F: ACGGACGCAAATCTTCTGAC R: GAGAACGACGCCATGACAG	(AG) <sub>10</sub>	55	A	6-FAM	a: 114–138 b: 150–174	6 6	diploid diploid	KP885690

Note: A = total number of alleles based on initial screen of 10 individuals; T<sub>a</sub> = annealing temperature.

<sup>a</sup>Fluorescent dye on forward primers used for fragment analysis.

65–55°C for 60 s, and extension at 72°C for 90 s; 25 cycles of 95°C for 60 s, 55°C for 60 s, and 72°C for 90 s; and a final extension at 72°C for 10 min. PCR products were diluted 20× and 1–2 μL of product was loaded to an ABI 3730xl Capillary Sequencer (Applied Biosystems, Burlington, Ontario, Canada) using GeneScan 500 LIZ (Life Technologies, Burlington, Ontario, Canada) size standard for fragment analysis at the Centre for Applied Genomics (The Hospital

for Sick Children, Toronto, Canada). Genotyping was performed in GeneMarker 2.4.0 (SoftGenetics, State College, Pennsylvania, USA).

All diploid markers except for PV65 amplified a maximum of two alleles per individual. PV65 amplified a maximum of four alleles per individual in two sets of size-separated fragments (PV65a and PV65b, Table 1). Loci showed high levels of polymorphism with seven to 17 alleles per locus, and observed heterozygosity from 0.41 to 0.83 per locus (Table 2). Exact tests performed in GENEPOP 4.2 (Rousset, 2008) revealed three loci with significant departures from Hardy–Weinberg equilibrium (HWE), but departures were not consistent across populations, except for locus PV64. It is possible that selfing in one or both populations, or the presence of population substructure (i.e., the Wahlund effect; Wahlund, 1928), led to this homozygote excess. Test for departures from linkage equilibrium expectations in GENEPOP showed that all pairs of loci were unlinked (results not shown).

We used open-pollinated progeny arrays to confirm that the identified “diploid” loci followed a disomic pattern of inheritance by amplifying the eight loci in 10 seeds collected from each of 15 mother plants. For PV65, all offspring genotypes had maternal contributions from both of the size-separated amplicons, suggesting that PV65a and PV65b represent independent genomic complements with disomic inheritance. For the remaining loci, disomic inheritance was confirmed by the observation of a maximum of two alleles for all offspring, with each offspring sharing an allele with the maternal genotype.

TABLE 2. Results from HWE exact tests for diploid markers developed for *Pulsatilla vulgaris* in two populations.<sup>a</sup>

Locus	A	A03 (n = 41)			A25 (n = 56)		
		H <sub>o</sub>	H <sub>e</sub>	P <sup>b</sup>	H <sub>o</sub>	H <sub>e</sub>	P <sup>b</sup>
PV2	8	0.67	0.75	0.87	0.69	0.75	1.00
PV7	16	0.59	0.64	0.87	0.76	0.86	0.43
PV27	17	0.83	0.87	0.86	0.79	0.86	0.88
PV33	7	0.59	0.58	0.32	0.71	0.70	0.66
PV52	13	0.75	0.78	0.86	0.74	0.84	0.30
PV56	13	0.55	0.69	0.24	0.65	0.87	<0.001
PV64	11	0.41	0.77	<0.001	0.52	0.79	0.003
PV65a	13	0.41	0.79	<0.001	0.64	0.64	0.62
PV65b	13	0.68	0.67	0.54	0.75	0.77	0.62

Note: A = total number of alleles; H<sub>e</sub> = expected heterozygosity; H<sub>o</sub> = observed heterozygosity; n = number of individuals sampled.

<sup>a</sup>Geographic coordinates for the populations are A03: 48°59'51.7"N, 11°3'35.4"E; A25: 48°57'38.7"N, 10°56'32.2"E.

<sup>b</sup>P values are corrected for multiple testing.

## CONCLUSIONS

These new microsatellite markers are highly informative and will be used to quantify gene flow across fragmented

populations of *P. vulgaris* in southern Germany. We will test whether the reintroduction of shepherding is a suitable conservation measure to improve genetic connectivity among populations of this species. The combination of diploid and tetraploid markers presented here will be useful in clarifying the polyploidization history of this and related species.

#### LITERATURE CITED

- BÖCHER, T. W. 1934. Beiträge zur Zytologie der Gattung Anemone. *Botanisk Tidsskrift* 42: 183–206.
- BUTAYE, J., D. ADRIAENS, AND O. HONNAY. 2005. Conservation and restoration of calcareous grasslands: A concise review of the effects of fragmentation and management on plant species. *Biotechnology, Agronomy, Society and Environment* 9: 111–118.
- FAIRCLOTH, B. C. 2008. MSATCOMMANDER: Detection of microsatellite repeat arrays and automated, locus-specific primer design. *Molecular Ecology Resources* 8: 92–94.
- IUCN. 2014. The IUCN Red List of Threatened Species, Version 2014.2 [online]. Website [www.iucnredlist.org](http://www.iucnredlist.org) [accessed 15 December 2014].
- PFEIFER, E., R. HOLDEREGGER, D. MATTHIES, AND R. RUTISHAUSER. 2002. Investigation on the population biology of a flagship species of dry meadows: *Pulsatilla vulgaris* Mill. in north-eastern Switzerland. *Botanica Helvetica* 112: 153–172.
- ROUSSET, F. 2008. GENEPOP'007: A complete reimplementation of the GENEPOP software for Windows and Linux. *Molecular Ecology Resources* 8: 103–106.
- WAHLUND, S. 1928. Zusammensetzung von Populationen und Korrelationserscheinungen vom Standpunkt der Vererbungslehre aus betrachtet. *Hereditas* 11: 65–106.
- WALKER, K. J., AND C. E. PINCHES. 2011. Reduced grazing and the decline of *Pulsatilla vulgaris* Mill. (Ranunculaceae) in England, UK. *Biological Conservation* 144: 3098–3105.
- WELLS, T. C. E., AND D. M. BARLING. 1971. *Pulsatilla vulgaris* Mill. (*Anemone pulsatilla* L.). *Journal of Ecology* 59: 275–292.