



Microsatellite Markers for the *Pilosella alpicola* Group (Hieraciinae, Asteraceae) and Their Cross-Amplification in Other Hieraciinae Genera

Authors: Vít, Petr, Šingliarová, Barbora, Zozomová-Lihová, Judita, Marhold, Karol, and Krak, Karol

Source: Applications in Plant Sciences, 3(8)

Published By: Botanical Society of America

URL: <https://doi.org/10.3732/apps.1500048>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

MICROSATELLITE MARKERS FOR THE *PILOSELLA ALPICOLA* GROUP (HIERACIINAE, ASTERACEAE) AND THEIR CROSS-AMPLIFICATION IN OTHER HIERACIINAE GENERA¹

PETR VÍT^{2,3}, BARBORA ŠINGLIAROVÁ⁴, JUDITA ZOZOMOVÁ-LIHOVÁ⁴, KAROL MARHOLD^{4,5},
AND KAROL KRAK^{2,3,6}

²Institute of Botany, Academy of Sciences of the Czech Republic, Zámek 1, CZ-252 43 Průhonice, Czech Republic; ³Faculty of Environmental Sciences, Czech University of Life Sciences Prague, Kamýcká 129, Prague CZ-165 21, Czech Republic;

⁴Institute of Botany, Slovak Academy of Sciences, Dúbravská cesta 9, SK-845 23 Bratislava, Slovak Republic; and ⁵Department of Botany, Faculty of Science, Charles University, Benátská 2, CZ-128 01 Prague, Czech Republic

- *Premise of the study:* Microsatellite markers were developed for the *Pilosella alpicola* group (Asteraceae), comprising four closely related species distributed in subalpine areas of Europe. These species are believed to have diverged recently, but display contrasting cytogeographic patterns and variation in breeding systems, representing a promising model system for studying plant speciation, adaptation, and recent polyploidization.
- *Methods and Results:* We developed 17 microsatellite markers for the *P. alpicola* group using 454 sequencing. Sixteen markers were polymorphic, with the number of alleles per locus ranging from seven to 16 and observed and expected heterozygosity ranging from 0.45 to 0.84 and 0.72 to 0.92, respectively. Ten and five loci amplified in the related species, *P. echioides* and *P. officinarum*, respectively, but only two in *Andryala* and one in *Hieracium* s. str.
- *Conclusions:* The developed microsatellite markers have high potential to become useful tools to study microevolutionary processes in the *P. alpicola* group and related *Pilosella* species.

Key words: hybridization; *Pilosella alpicola*; polyploidy; population genetics; simple sequence repeat (SSR) markers.

The *Pilosella alpicola* group (Asteraceae) comprises four vicariant species of perennial herbs distributed disjunctly in subalpine areas of Europe (the Alps, Carpathians, and Balkan mountains). In this study, we followed the generic circumscriptions outlined by Bräutigam and Greuter (2007) in our treatment of the *P. alpicola* group. Our recent studies (Šingliarová et al., 2011a, b) showed that although the taxa of this group are very closely related and diverged relatively recently, they are morphologically well defined and display contrasting cytogeographic patterns and variation in breeding systems. Two species (*P. ullepitschii* (Błocki) Szlach from the Carpathians and *P. serbica* (F. W. Schultz & Sch. Bip.) Szlach from Serbia) are exclusively diploid and outcrossing. The Balkan taxon *P. rhodopea* (Griseb.) Szlach represents a unique diploid-polyploid complex with up to five cytotypes (diploids to hexaploids) occurring mostly in mixed-ploidy populations. All

P. rhodopea cytotypes are outcrossing, although polyploids exhibit severely reduced fertility (Šingliarová et al., 2011b). The alpine *P. alpicola* F. W. Schultz & Sch. Bip. (s. str.) has two allopatric cytotypes (tetraploids and pentaploids) with presumably polytopic allopolyploid origins and agamospermic mode of reproduction (Šingliarová et al., 2011a, b). All species of the *P. alpicola* group are rather rare, and their populations are often small and endangered by human activities. Recently, the conservation status of all taxa of the group was evaluated and their inclusion in relevant national Red Lists was proposed (Šingliarová et al., 2013). Interesting distributional patterns and a variety of evolutionary mechanisms playing a role in the differentiation of the *P. alpicola* group suggest it as a promising model system for studying plant speciation, adaptation, and recent polyploidization.

Microsatellite markers have already been developed for the related *P. officinarum* F. W. Schultz & Sch. Bip. (Zini and Komjanc, 2007) and the closely related genus *Hieracium* L. (s. str.) (Jönsson et al., 2010). The cross-amplification of the *P. officinarum* markers in other *Pilosella* Hill species and *Hieracium* s. str. gave unsatisfactory results (P. Trávníček, personal communication; Jönsson et al., 2010). Low cross-amplification success reported so far for the developed markers in *Hieracium* s. str. and *Pilosella* led us to deduce their poor transferability also to the *P. alpicola* group; therefore, a new set of markers is developed here. In addition, other *Pilosella*, *Hieracium* s. str., and *Andryala* L. taxa were used here to evaluate the potential cross-amplification of the developed

¹Manuscript received 28 April 2015; revision accepted 31 May 2015.

The authors thank S. Pířová for technical assistance and J. Chrtek and J. Zahradníček for plants for the cross-amplification experiments. The study was financially supported by the Research and Development Support Agency (grant no. APVV-0320-10 to K.M.), by the long-term research development project no. RVO 67985939 (to P.V. and K.K.), and by the project ITMS6240120014 (Centre of Excellence for Biodiversity and Land-Use Conservation), supported by the Research and Development Operational Program funded by the European Regional Development Fund (ERDF).

⁶Author for correspondence: karol.krak@ibot.cas.cz

doi:10.3732/apps.1500048

markers in other phylogenetic lineages within the subtribe Hieraciinae (Fehrer et al., 2007; Krak and Mráz, 2008).

METHODS AND RESULTS

Total genomic DNA was extracted from fresh or silica gel-dried leaf material using the DNeasy Plant Mini Kit (QIAGEN, Hilden, Germany) according to the manufacturer's instructions. DNA from one diploid individual of *P. rhodopea* was used for 454 sequencing. Genomic library preparation and 1/4 run on the Roche GS FLX platform was performed at GATC Biotech (Konstanz, Germany). MSATCOMMANDER version 0.8.2 (Faircloth, 2008) was used to identify reads containing di-, tri-, tetra-, penta-, and hexanucleotide repeats and to design primers. Primers were successfully designed to 179 microsatellite-containing reads.

Sixty primer pairs were selected and tested for amplification in three diploid individuals, each representing a different species of the studied group. Each reaction contained 1× PCR buffer with KCl (Fermentas, St. Leon, Germany), 2.5 mM MgCl₂, 0.2 mM of each dNTP, 0.2 μM of both primers, 0.5 units of *Taq* DNA polymerase (Fermentas), and 20 ng of DNA. The cycling conditions were: 3 min at 95°C; followed by 35 cycles of 95°C for 30 s, locus-specific annealing temperature (see Table 1) for 30 s, and 72°C for 1 min; and a final extension at 72°C for 10 min. PCR products were checked on 2.5% agarose gels. Twenty-three primer pairs that amplified in the three tested species of the *P. alpicola* group were further applied to the complete set of 31 diploid individuals, originating from five distinct and geographically well-separated natural populations (Appendix 1). In this step, the amplification protocol was modified

as described in Schuelke (2000) to facilitate fluorescent labeling of the PCR products by 6-FAM (Applied Biosystems, Foster City, California, USA). One microliter of each PCR product was mixed with 0.1 μL of GeneScan 500 LIZ internal size standard (Applied Biosystems) and 12 μL of Hi-Di formamide (Applied Biosystems) and electrophoresed using ABI PRISM 3130 Genetic Analyzer (Applied Biosystems). Allele calling was performed using GeneMarker version 2.4.0 (SoftGenetics, State College, Pennsylvania, USA). Six primer pairs amplified three or four alleles in the majority of samples, indicating possible duplications of these loci; these were therefore discarded. One locus, Palpi21, was invariable. Two other loci, Palpi25 and Palpi35, showed multiple peaks in four samples belonging to populations COZ, DOD, and MEN (see Appendix 1 for population localities). For these particular samples, the allele lengths were scored as missing data, and the markers were retained for genetic diversity estimates. For the summary information on the 16 polymorphic loci see Table 1.

The genetic diversity of these 16 loci was then estimated. Number of alleles per locus, expected and observed heterozygosity, and polymorphic information content were calculated based on allele frequencies for each population as well as for the complete data set using CERVUS version 3.0.3 (Field Genetics, London, United Kingdom). All markers showed a relatively high degree of polymorphism (Table 2). The number of detected alleles ranged from seven for Palpi 1 to 16 for Palpi18, Palpi34, and Palpi37 (average 11.4). The observed heterozygosity ranged from 0.45 at Palpi32 to 0.84 at Palpi49 (average 0.64). The values of expected heterozygosity varied from 0.72 at Palpi1 to 0.92 at Palpi34 (average 0.85). The values of the polymorphic information content varied from 0.67 at Palpi1 to 0.89 at Palpi34 (average 0.81). The level of variability within the tested populations was comparable for the populations of

TABLE 1. Characteristics of 17 newly developed microsatellite markers for the *Pilosella alpicola* group. Locus Palpi21 is monomorphic; the 16 other loci are polymorphic.

Locus	Primer sequences (5'–3')	Repeat motif	Allele size range (bp)	T _a (°C)	Cross-species amplification ^a	GenBank accession no.
Palpi1	F: TGCATTCTTCGGTCCCAC R: TGAAGAACATCGACATAATGCAC	(AT) ₁₃	232–247	55	—	KJ415260
Palpi5	F: TGCATTCTTCGGTCCCAC R: TGAAGAACATCGACATAATGCAC	(AC) ₁₄	216–230	55	Po, Pe	KJ415261
Palpi8	F: AGGCAATATACTATGTGTTGACCTG R: TCAGTGGCCCTCTATTATACCC	(AC) ₁₄	458–474	57	Po, Pe	KJ415262
Palpi18	F: GCACGCATCTTACCAGACG R: TCGGTTAATCCTCTGGCGG	(ATT) ₂₃	182–257	53	—	KJ415263
Palpi19	F: TGTAATTCGGCTGCAAAATCG R: TCACGAGTGGTTTCCTTACTC	(AAT) ₁₉	253–292	55	—	KJ415264
Palpi20	F: TGTTTGATGGCGATTACGAG R: GCCCAGTTCCTTTGAACCC	(AAT) ₂₀	331–383	53	—	KJ415265
Palpi21 ^b	F: GCCACAGCTTACGAGCATC R: TTGCGGTGAAGACGGTG	(CT) ₁₂	220	55	NA	KR862872
Palpi25	F: TCCCGCTGGTTCGAAAGAC R: GCCCTCAAGGCCTCAACC	(GT) ₁₂	142–176	55	Po, Pe	KJ415266
Palpi26	F: CCCATCTGGTTAAAGCGCC R: TCGAAGTTGTTACGAGTTGCAG	(AC) ₁₂	192–230	53	—	KJ415267
Palpi29	F: CTGCCATTGTTGGTGGCTC R: AGGGATAGCACTTGCTGGG	(AT) ₁₂	377–396	55	Po, Pe, Hu, Hs, Al, Ai	KJ415268
Palpi32	F: CTGTGAGGTACATACGTGGC R: TTTGTGGTTTGAGTTTACCCG	(ATT) ₁₃	377–413	55	Pe	KJ415269
Palpi33	F: TCACTCATGAAATCTGTAACATCCG R: CCATAAAGGCAACTTTAGTCCC	(AAT) ₁₁	224–273	56	Pe	KJ415270
Palpi34	F: TGGTTAACGGTGGGCGATC R: CTAGCAGGTGGCCTTGACC	(AAC) ₁₂	406–436	52	Pe	KJ415271
Palpi35	F: AAGTCGACACCTTTAGTCCC R: GCTGCTAATCCGTGGTCAG	(ATT) ₁₇	243–274	54	—	KJ415272
Palpi37	F: TCCTCTAAGTGGTCTTTTGC R: ATGCCTCACGAGACGTTCC	(CTT) ₁₁	451–505	52	Po, Pe	KJ415273
Palpi45	F: GGCAGAAGGCTAAGAAGTGC R: GGAAGAATTTGATGGGTTGAC	(AC) ₁₁	277–319	55	Pe	KJ415274
Palpi49	F: GGGCTGCCTATTCCTGATG R: GGGTGTATGACGACGAGTTTG	(AG) ₁₀	240–265	55	Pe, Ai	KJ415275

Note: T_a = annealing temperature.

^aLoci that were successfully cross-amplified in other Hieraciinae species: *Pilosella officinarum* (Po), *P. echioides* (Pe), *Hieracium umbellatum* (Hu), *H. stelligerum* (Hs), *Andryala laevis* (Al), and *A. integrifolia* (Ai); NA = information not available.

^bMonomorphic locus.

TABLE 2. Genetic diversity parameters estimated for 16 polymorphic microsatellite loci for each analyzed population and the complete data set of the *Pilosella alpicola* group.

Locus	<i>P. rhodopea</i> (population COZ; n = 6)			<i>P. rhodopea</i> (population DOD; n = 5)			<i>P. ulleipschii</i> (population MEN; n = 7)			<i>P. ulleipschii</i> (population SMARE; n = 7)			<i>P. serbica</i> (population KOP; n = 6)			Overall (n = 31)				
	A	H _o	H _e	A	H _o	H _e	A	H _o	H _e	A	H _o	H _e	A	H _o	H _e	A	H _o	H _e	PIC	
	Palpi1	2	1	0.545	4	0.8	0.78	3	0.86	0.58	0.45	2	1	0.54	3	0.33	0.59	7	0.81	0.72
Palpi5	4	0.67	0.697	6	1	0.89	3	0.71	0.6	0.47	2	0	0.26	4	0.6	0.53	9	0.57	0.77	0.73
Palpi8	4	0.67	0.682	4	0.4	0.71	6	0.86	0.87	0.78	3	0.57	0.6	6	1	0.8	9	0.71	0.85	0.82
Palpi18	4	0.67	0.803	6	0.75	0.93	6	0.27	0.84	0.74	2	0	0.3	8	0.83	0.89	16	0.48	0.9	0.87
Palpi19	7	0.83	0.924	3	0	0.71	9	1	0.9	0.82	3	0.71	0.54	5	0.67	0.8	15	0.7	0.9	0.88
Palpi20	3	0.33	0.439	4	1	0.78	6	0.86	0.86	0.77	3	0.14	0.47	3	0.6	0.71	11	0.57	0.83	0.79
Palpi25	3	0.6	0.711	2	0.33	0.33	3	0.5	0.44	0.36	3	0.57	0.56	4	0.67	0.65	9	0.56	0.85	0.81
Palpi26	7	1	0.894	5	0.8	0.84	3	0.71	0.6	0.47	2	0	0.3	7	0.6	0.91	8	0.62	0.81	0.77
Palpi29	4	0.83	0.712	4	0.6	0.73	6	0.86	0.79	0.7	4	0.57	0.69	4	1	0.71	9	0.77	0.83	0.79
Palpi32	4	0.33	0.636	3	0.2	0.51	4	0.86	0.81	0.72	2	0.14	0.14	5	0.67	0.79	11	0.45	0.89	0.86
Palpi33	6	1	0.848	3	1	0.71	5	0.71	0.74	0.65	1	0	0	2	0.33	0.3	12	0.58	0.87	0.84
Palpi34	6	0.67	0.894	5	0.8	0.76	6	0.86	0.82	0.73	4	0.57	0.71	4	0.8	0.79	16	0.73	0.92	0.89
Palpi35	5	0.6	0.822	4	0.67	0.87	5	0.5	0.74	0.64	4	0.27	0.71	3	0.67	0.62	10	0.52	0.9	0.87
Palpi37	5	0.83	0.758	6	1	0.89	6	0.43	0.88	0.79	4	0.43	0.69	6	0.67	0.76	16	0.65	0.88	0.86
Palpi45	4	0.67	0.561	2	1	0.56	4	0.83	0.77	0.65	3	0.14	0.28	4	0.83	0.77	13	0.67	0.81	0.77
Palpi49	9	0.83	0.939	4	0.8	0.8	7	0.57	0.89	0.8	2	1	0.54	4	1	0.74	10	0.84	0.84	0.8
Average	4.18	0.72	0.74	4.06	0.7	0.74	5.3	0.71	0.76	0.67	2.8	0.38	0.46	4.5	0.7	0.71	11.4	0.64	0.85	0.81

Note: A = number of alleles per locus; H_e = expected heterozygosity; H_o = observed heterozygosity; n = number of individuals; PIC = polymorphic information content.

P. rhodopea, *P. serbica*, and one population of *P. ulleipschii* (population MEN). The other population of *P. ulleipschii* (SMARE) showed lower values for all estimated parameters (see Table 2 for details).

To evaluate the potential utility of the newly developed markers in other phylogenetic lineages of the Asteraceae subtribe Hieraciinae (Fehrer et al., 2007; Krak and Mráz, 2008), cross-amplification experiments were performed in one individual of *P. echioides* F. W. Schultz & Sch. Bip., *P. officinarum*, *Hieracium umbellatum* L., *H. stelligerum* Froel., *Andryala laevisomentosa* (Nyar. ex Sennikov) Greuter, and *A. integrifolia* L. The results of cross-amplification tests are summarized in Table 1. Higher cross-amplification success was reached within the genus *Pilosella*, where 10 markers could be cross-amplified in *P. echioides* and five in *P. officinarum*. Only one marker (Palpi29) was cross-amplified in all tested taxa. No other marker gave a positive result in the three *Hieracium* s. str. species and *A. laevisomentosa*. In *A. integrifolia*, cross-amplification of Palpi49 was positive as well.

CONCLUSIONS

In this study, we have developed 17 novel microsatellite markers for the *P. alpicola* group. Although only a relatively small sample set was used to evaluate their variability, 16 markers showed high levels of polymorphism. The transferability of these markers to two additional *Pilosella* species and to the closely related genera *Andryala* and *Hieracium* s. str. is restricted. Five markers were cross-amplified in *P. officinarum* and 10 in *P. echioides*, suggesting high species specificity of these markers. The cross-amplification in the other Hieraciinae genera was even less successful. Only one marker amplified in all four species and an additional one in *A. integrifolia*.

LITERATURE CITED

BRÄUTIGAM, S., AND W. GREUTER. 2007. A new treatment of *Pilosella* for the Euro-Mediterranean flora. *Willdenowia* 37: 123–137.

FAIRCLOTH, B. C. 2008. MSATCOMMANDER: Detection of microsatellite repeat arrays and automated, locus-specific primer design. *Molecular Ecology Resources* 8: 92–94.

FEHRER, J., B. GEMEINHOLZER, J. CHRTEK JR., AND S. BRÄUTIGAM. 2007. Incongruent plastid and nuclear DNA phylogenies reveal ancient intergeneric hybridization in *Pilosella* hawkweeds (*Hieracium*, Cichorieae, Asteraceae). *Molecular Phylogenetics and Evolution* 42: 347–361.

JÖNSSON, J., M. WELLENREUTHER, AND T. TYLER. 2010. Ten polymorphic microsatellite markers for *Hieracium* s.str. (Asteraceae). *Conservation Genetics Resources* 2: 295–300.

KRAK, K., AND P. MRÁZ. 2008. Trichomes in the tribe Lactuceae (Asteraceae): Taxonomic implications. *Biologia* 63: 616–630.

SCHUELKE, M. 2000. An economic method for the fluorescent labeling of PCR fragments. *Nature Biotechnology* 18: 233–234.

ŠINGLIAROVÁ, B., J. CHRTEK, I. PLAČKOVÁ, AND P. MRÁZ. 2011a. Allozyme variation in diploid, polyploid and mixed-ploidy populations of the *Pilosella alpicola* group (Asteraceae): Relation to morphology, origin of polyploids and breeding system. *Folia Geobotanica* 46: 387–410.

ŠINGLIAROVÁ, B., I. HODÁLOVÁ, AND P. MRÁZ. 2011b. Biosystematic study of the diploid-polyploid *Pilosella alpicola* group with variation in breeding system: Patterns and processes. *Taxon* 60: 450–470.

ŠINGLIAROVÁ, B., R. ŠUVADA, AND P. MRÁZ. 2013. Allopatric distribution, ecology and conservation status of the *Pilosella alpicola* group (Asteraceae). *Nordic Journal of Botany* 31: 122–128.

ZINI, E., AND M. KOMJANC. 2007. Identification of microsatellite markers in *Hieracium pilosella* L. *Conservation Genetics* 9: 487–489.

APPENDIX 1. Locality and voucher information of the *Pilosella alpicola* group and representatives of other lineages within the subtribe Hieraciinae used for the current study.

Species	Population code	Voucher specimen accession no.	Collection locality ^a	Geographic coordinates	<i>n</i>
<i>Pilosella rhodopea</i> (Griseb.) Szeląg	COZ	BS_COZ_6, 10, 12, 13, 19, 20 ^b	RO; Cozia Mtns., Mt. Cozia, 1592 m a.s.l.	45°19'04"N, 24°20'17"E	6
<i>P. serbica</i> (F. W. Schultz & Sch. Bip.) Szeląg	DOD	BS_DOD_1, 6–8, 10 ^b	BG; Rila Mtns., Mt. Dodov vrah, 2540 m a.s.l.	42°09'59"N, 23°20'23"E	5
	KOP	BS_KOP_1, 2, 4–7 ^b	RS; Kopaonik Mtns., Mt. Suvo Rudište, 1917 m a.s.l.	43°16'28"N, 20°48'55"E	6
<i>P. ullepitschii</i> (Błocki) Szeląg	MEN	BS_MEN_12, 14, 18, 19, 27, 29, 30 ^b	SK; Vysoké Tatry Mtns., Mengusovská dolina valley, 1800–1875 m a.s.l.	49°09'57"N, 20°03'40"E	7
	SMARE	BS_SMARE_11–15, 17, 18 ^b	RO; Nemira Mtns., Mt. Sandru Mare, 1590–1640 m a.s.l.	46°11'57"N, 26°20'21"E	7
<i>P. officinarum</i> F. W. Schultz & Sch. Bip.	1484	PRA_PV_1484 ^c	CZ; Prague, Bohnice–Zámky, 205 m a.s.l.	50°08'26"N, 14°23'56"E	1 ^d
<i>P. echioides</i> F. W. Schultz & Sch. Bip.	TR608	PRA_PV_TR608 ^c	CZ; Beroun, Trubínský vrch hill, 325 m a.s.l.	49°56'37"N, 13°59'47"E	1 ^d
<i>Hieracium umbellatum</i> L.	1021	PRA_PV_1021 ^c	PL; Województwo pomorskie, Baltic coast, 5 m a.s.l.	54°20'00"N, 19°02'27"E	1 ^d
<i>H. stelligerum</i> Froel.	1233	PRA_KK_1233 ^c	FR; Vallon Pont d'Arc, along the road D 390, 500 m a.s.l.	44°24'25"N, 04°24'10"E	1 ^d
<i>H. intybaceum</i> All.	2509a	PRA_KK_2509a ^c	AT; Tirol, Tuxer Alpen, 3.3 km NE of the village of Navis, 1878 m a.s.l.	47°09'22"N, 11°33'48"E	1 ^d
<i>Andryala integrifolia</i> L.	AZ3	PRA_JZ_AZ3 ^c	DZ; Alger, town district of Kouba, 90 m a.s.l.	36°43'00"N, 03°05'00"E	1 ^d
<i>A. laevitomentosa</i> (Nyár. ex Sennikov) Greuter	D5	PRA_JZ_D5 ^c	RO; Bukovina, Mt. Pietrosul Broștenilor, 1780 m a.s.l.	47°22'31"N, 25°32'29"E	1 ^d

Note: *n* = number of individuals.

^aAbbreviations for countries: AT = Austria, BG = Bulgaria, CZ = Czech Republic, DZ = Algeria, FR = France, PL = Poland, RO = Romania, SK = Slovakia, RS = Serbia.

^bVoucher specimens deposited in the herbarium of the Institute of Botany, Slovak Academy of Sciences (SAV), Bratislava, Slovakia.

^cVoucher specimens deposited in the herbarium of the Institute of Botany, Czech Academy of Sciences (PRA), Prague, Czech Republic.

^dSamples used to evaluate the potential cross-amplification of the markers within Hieraciinae.