



Characterization of Microsatellite Loci in the Lichen-Forming Fungus *Cetraria aculeata* (Parmeliaceae, Ascomycota)

Authors: Lutsak, Tetiana, Fernández-Mendoza, Fernando, Greshake, Bastian, Grande, Francesco Dal, Ebersberger, Ingo, et al.

Source: Applications in Plant Sciences, 4(9)

Published By: Botanical Society of America

URL: <https://doi.org/10.3732/apps.1600047>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

CHARACTERIZATION OF MICROSATELLITE LOCI IN THE LICHEN-FORMING FUNGUS *CETRARIA ACULEATA* (PARMELIACEAE, ASCOMYCOTA)¹

TETIANA LUTSAK^{2,7}, FERNANDO FERNÁNDEZ-MENDOZA³, BASTIAN GRESHAKE⁴,
FRANCESCO DAL GRANDE⁵, INGO EBERSBERGER⁴, SIEGLINDE OTT⁶,
AND CHRISTIAN PRINTZEN²

²Senckenberg Research Institute and Natural History Museum, D-60325 Frankfurt am Main, Germany; ³Institute of Plant Sciences, Karl-Franzens-Universität, Graz, A-8010 Graz, Austria; ⁴Department of Applied Bioinformatics, Goethe-Universität Frankfurt am Main, D-60438 Frankfurt am Main, Germany; ⁵Senckenberg Biodiversity and Climate Research Centre (BiK-F), D-60325 Frankfurt am Main, Germany; and ⁶Institute of Botany, Heinrich-Heine-Universität Düsseldorf, D-40225 Düsseldorf, Germany

- *Premise of the study:* Polymorphic microsatellite markers were developed for the lichen species *Cetraria aculeata* (Parmeliaceae) to study fine-scale population diversity and phylogeographic structure.
- *Methods and Results:* Using Illumina HiSeq and MiSeq, 15 fungus-specific microsatellite markers were developed and tested on 81 specimens from four populations from Spain. The number of alleles ranged from four to 13 alleles per locus with a mean of 7.9, and average gene diversities varied from 0.40 to 0.73 over four populations. The amplification rates of 10 markers (CA01–CA10) in populations of *C. aculeata* exceeded 85%. The markers also amplified across a range of closely related species, except for locus CA05, which did not amplify in *C. australiensis* and *C. “panamericana,”* and locus CA10 which did not amplify in *C. australiensis*.
- *Conclusions:* The identified microsatellite markers will be used to study the genetic diversity and phylogeographic structure in populations of *C. aculeata* in western Eurasia.

Key words: *Cetraria aculeata*; lichen; microsatellites; Parmeliaceae.

In lichens, microsatellite markers have been reported for a number of species and were successfully applied to assess levels of genetic diversity and dispersal patterns (e.g., Walser et al., 2005; Jones et al., 2015). In this paper, we develop microsatellite primers for the lichen species *Cetraria aculeata* (Schreb.) Fr. (Parmeliaceae), a dark brown, fruticose soil lichen characterized by an extremely wide distributional range and ecological niche, which makes it a good model organism to study the impact of environmental factors on population genetic structure and diversity. The species originated in the Northern Hemisphere and dispersed through South America into the Antarctic during the Pleistocene (Fernández-Mendoza and Printzen, 2013), establishing one of its centers of diversity in the Mediterranean region (Fernández-Mendoza et al., 2011; Printzen et al., 2013). Mediterranean haplotypes of *C. aculeata* mycobionts are similar to those from the Afro-alpine mountain ranges (Lutsak et al., 2015), while central and northern European haplotypes differ

more strongly (Fernández-Mendoza et al., 2011). The fine-scale structure of both symbionts is not sufficiently resolved by DNA sequence data. Therefore, microsatellite markers are of high importance to study the phylogeographic structure and genetic diversity of *C. aculeata* across western Eurasia.

METHODS AND RESULTS

Total genomic DNA was extracted from one thallus of *C. aculeata* (isolate 3041, Ukrainian Nature Steppe Reserve, FR-0261072) in four single reactions with the DNeasy 96 Plant Kit (QIAGEN, Hilden, Germany). The samples were frozen in liquid nitrogen, ground, and incubated overnight at 37°C in a mixture of enzymes: 400 µL of Glusulase (PerkinElmer, Boston, Massachusetts, USA) and 500 µL of 0.1 mg mL⁻¹ Zymolyase 20T (Seikagaku Biobusiness, Tokyo, Japan) in 500 µL of phosphate buffer (pH 7.4). The DNA extracts were pooled, and libraries for the Illumina HiSeq and MiSeq sequencing platforms were created. A total of 25,727,973 paired-end reads with an average length of 100 bp and 15,120,929 paired-end reads with an average length of 250 bp were obtained from the HiSeq and MiSeq runs, respectively. Illumina adapters and primers were trimmed and quality filtering was performed using Trimmomatic software version 0.27 (Bolger et al., 2014). The lichen metagenome was assembled using Velvet software version 1.2.10 (Zerbino and Birney, 2008). Using the BLAST tool v2.2.25+, algal and bacterial contigs were filtered out. The latest available National Center for Biotechnology Information (NCBI) nucleotide database (NT Database, 24 May 2013) was used for aligning the assembled contigs. The alignment file was processed by using MEGAN5 version 4.70.4 software under default settings (Huson et al., 2007). Plots generated by MEGAN were used for assessing the bacterial, fungal, and algal contigs, which served as a reference for mapping the raw reads in the second round of genome assembly. The inferred contigs were screened for tri-, tetra-, penta-, and hexanucleotide repeats with at

¹Manuscript received 13 April 2016; revision accepted 3 June 2016.

This study received financial support from the Adolf Messer-Stiftung, the Marga und Kurt Möllgaard-Stiftung, and the research funding program Landes-Offensive zur Entwicklung wissenschaftlich-ökonomischer Exzellenz (LOEWE) of Hesse's Ministry of Higher Education, Research and the Arts. The work of T.L. is supported by a grant from the German Academic Exchange Service (DAAD).

⁷Author for correspondence: tetiana.lutsak@senckenberg.de

doi:10.3732/apps.1600047

Applications in Plant Sciences 2016 4(9): 1600047; <http://www.bioone.org/loi/apps> © 2016 Lutsak et al. Published by the Botanical Society of America. This work is licensed under a Creative Commons Attribution License (CC-BY-NC-SA).

TABLE 1. Overview of the microsatellite loci developed for the mycobiont of *Cetraria aculeata*.

| Locus | Primer sequences (5'–3') | Repeat motif ^a | Allele size range (bp) | T _a (°C) | Fluorescent dye | PCR multiplex | Amplification rate (%) | GenBank accession no. |
|-------|---|---------------------------|------------------------|---------------------|-----------------|---------------|------------------------|-----------------------|
| CA01 | F: GTATGGTGGTGTACGGGTT R: CGGTGGTGAGAAGTGAGAGG | (GAAG) _n | 214–238 | 55 | FAM | 1 | 100 | KU361362 |
| CA02 | F: TGATGTCATCGAAGCCCTGG R: ATATCCGTTGCCGTGGTGGT | (TTGA) _n | 247–267 | 53 | NED | 1 | 100 | KU361368 |
| CA03 | F: TAGTAGCGGGCAGTCGAAAC R: TGCTTGAGCTTGTCTTCTCGA | (CTCTG) _n | 297–362 | 53 | VIC | 1 | 100 | KU361361 |
| CA04 | F: GAAACTGAGGGGAAGTGCCA R: GATGGCTGGTCCCAATGACA | (CTCCAC) _n | 318–344 | 54 | PET | 1 | 100 | KU361364 |
| CA05 | F: AACCCCAACCCACAGAATC R: GCGAACGAGGATGATTGTGC | (CACG) _n | 123–135 | 54 | FAM | 2 | 100 | KU361363 |
| CA06 | F: AGAACGGCAGGAAGAAGAGC R: CCCTGGTCGGCTCCCAATATC | (TTGA) _n | 201–269 | 55 | NED | 2 | 100 | KU361367 |
| CA07 | F: AATGAAACACGTGGCTGTGG R: CAGACCGGGTGTGTCAATCA | (GTAG) _n | 205–253 | 53 | VIC | 3 | 100 | KU361365 |
| CA08 | F: AGTGCGGTGAATGTACGAG R: CGGGCCGCTCCATTTGTATA | (AATGA) _n | 221–276 | 54 | FAM | 3 | 100 | KU361370 |
| CA09 | F: ATGAGGCACGCACAGAATGA R: CACTCCACTCGTTTACCC | (CCGAGT) _n | 312–380 | 54 | PET | 3 | 100 | KU361369 |
| CA10 | F: TGACTAAGGAGGATGTGCG R: TTGACTGCTTCCGAGGAGAA | (TCAA) _n | 318–350 | 53 | FAM | 3 | 85.1 | KU361366 |
| CA11 | F: CCTTTTACACGGTCGCTTC R: CCCCTCCTTCTAACTCCGAC | (ATAC) _n | 364–424 | 55 | PET | 2 | 76.7 | KU764488 |
| CA12 | F: CGTCTCCGTGTACCATAGCC R: CAGCAGCGTTATCAGCAAGC | (TTGCTG) _n | 184–284 | 55 | PET | 1 | 60 | KU764489 |
| CA13 | F: GCTCACCTCTCAGCAGATC R: TGGCTGCGTTCTTTACAGG | (GAAGAG) _n | 265–443 | 55 | VIC | 2 | 76.7 | KU764490 |
| CA14 | F: GCAACGTGCATGGAACGTT R: TCTCGTTCCGGCAGTTGAGAC | (GTAAA) _n | 219–264 | 53 | PET | 2 | 45 | KU764491 |
| CA15 | F: CGCTTGTGATATCGTCCGGA R: ACATCATCCGCAGCTTCCAA | (CTCTTG) _n | 188–318 | 53 | FAM | 2 | 70 | KU764492 |

Note: T_a = annealing temperature.

^an stands for numerous repeats.

least five repeats using the MicroSAteellite identification tool (MISA; Thiel et al., 2003; <http://pgrc.ipk-gatersleben.de/misa/misa.html>). We selected 40 loci with perfect (uninterrupted) microsatellites and a repeat size of four to six base pairs for testing. Primers were designed using the Primer3 plugin in Geneious R7.1.9 (created by BioMatters, <http://www.geneious.com/>).

On a subset of eight specimens of *C. aculeata*, 24 primer pairs amplified successfully and showed significant variability. Single PCR reactions were performed using illustra PuReTaq Ready-To-Go PCR Beads (GE Healthcare, Little Chalfont, Buckinghamshire, England) according to the manufacturer's protocol with the following conditions: denaturation for 5 min at 95°C; followed by 10 cycles of 1 min at 94°C, 1 min at 65–57°C (touchdown of –1° per cycle), and 1 min at 72°C; followed by 20 cycles of 1 min at 94°C and 1 min at 54, 53, or

52°C (depending on primer pair); and a final extension of 10 min at 72°C. The 24 primer pairs were also tested in PCR reactions with three axenic cultures of *Trebouxia jamesii* (Hildreth & Ahmadjian) Gärtner, a photobiont of *C. aculeata*, isolated from specimens from Antarctica, Spain, and Germany. Eight primer pairs produced PCR products with at least one of the cultures and were discarded. The remaining 16 primer pairs were assumed to be fungal specific. Forward primers were labeled with fluorescent dyes and used in multiplex PCRs with the Type-It Multiplex Kit (QIAGEN). PCR reactions were performed in a total volume of 25 µL, which contained 12.5 µL of Type-It Multiplex PCR Master Mix, 2.5 µL of primer mix, 5 µL of RNA-free water, and 5 µL of sample DNA. The PCR conditions were set to: denaturation for 5 min at 95°C; followed by five cycles of 30 s at 95°C, 90 s at 57°C, and 30 s at 72°C; then 15 cycles of

TABLE 2. Characteristics of microsatellite loci CA01–CA10 in populations of *Cetraria aculeata* from Spain.^a

| Locus | Total | | Spa1 (n = 10) | | Spa2 (n = 23) | | Spa3 (n = 24) | | Spa4 (n = 24) | |
|-------|-------|----------------|---------------|----------------|---------------|----------------|---------------|----------------|---------------|----------------|
| | A | H _e | A | H _e | A | H _e | A | H _e | A | H _e |
| CA01 | 6 | 0.423 | 3 | 0.511 | 3 | 0.466 | 3 | 0.409 | 3 | 0.304 |
| CA02 | 5 | 0.404 | 2 | 0.356 | 4 | 0.704 | 3 | 0.475 | 2 | 0.083 |
| CA03 | 10 | 0.713 | 5 | 0.667 | 4 | 0.668 | 8 | 0.812 | 4 | 0.707 |
| CA04 | 5 | 0.501 | 3 | 0.711 | 3 | 0.372 | 3 | 0.359 | 3 | 0.562 |
| CA05 | 4 | 0.524 | 3 | 0.622 | 3 | 0.502 | 3 | 0.627 | 2 | 0.344 |
| CA06 | 13 | 0.726 | 5 | 0.822 | 6 | 0.771 | 8 | 0.804 | 2 | 0.507 |
| CA07 | 9 | 0.564 | 4 | 0.733 | 6 | 0.613 | 5 | 0.486 | 4 | 0.424 |
| CA08 | 10 | 0.679 | 4 | 0.778 | 8 | 0.834 | 6 | 0.746 | 3 | 0.359 |
| CA09 | 9 | 0.674 | 3 | 0.644 | 5 | 0.708 | 4 | 0.634 | 4 | 0.710 |
| CA10 | 8 | 0.522 | 4 | 0.711 | 4 | 0.575 | 1 | 0.000 | 6 | 0.801 |
| Mean | 7.9 | 0.573 | 3.6 | 0.656 | 4.6 | 0.621 | 4.4 | 0.535 | 3.3 | 0.480 |

Note: A = number of alleles; H_e = Nei's unbiased gene diversity; n = total number of samples per population analyzed.

^aPopulation locality and voucher information are provided in Appendix 1.

TABLE 3. Cross-amplification of *Cetraria aculeata* microsatellite markers with related species of the genus *Cetraria*.

| Species | No. of alleles observed | | | | | | | | | | | | | | |
|--------------------------|-------------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| | CA01 | CA02 | CA03 | CA04 | CA05 | CA06 | CA07 | CA08 | CA09 | CA10 | CA11 | CA12 | CA13 | CA14 | CA15 |
| <i>C. australiensis</i> | 1 | 1 | 1 | 1 | 0 | 1 | 2 | 1 | 2 | 0 | 0 | 0 | 0 | 0 | 0 |
| <i>C. crespoae</i> | 2 | 1 | 1 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 0 | 0 | 2 | 0 | 0 |
| <i>C. muricata</i> | 1 | 2 | 1 | 2 | 2 | 1 | 2 | 2 | 2 | 2 | 1 | 1 | 1 | 0 | 0 |
| <i>C. odontella</i> | 2 | 1 | 1 | 2 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 0 | 0 | 0 | 0 |
| <i>C. "panamericana"</i> | 1 | 1 | 1 | 2 | 0 | 1 | 1 | 1 | 1 | 1 | 0 | 0 | 1 | 0 | 0 |

Note: 0 = locus did not amplify; 1 = locus amplified in a single specimen; 2 = locus amplified in two specimens tested.

30 s at 95°C, 90 s at 55°C, and 30 s at 72°C; then 10 cycles of 30 s at 95°C, 90 s at 54°C, and 30 s at 72°C; and 10 cycles of 30 s at 95°C, 90 s at 53°C, and 30 s at 72°C; with a final extension of 30 min at 60°C. One more primer pair was eliminated, as it yielded products of inconsistent size.

A data set of 81 specimens of *C. aculeata* was used to test the amplification rates and the variability of selected microsatellite loci. The data set consisted of four populations of 10 to 24 individuals collected in Spain (Appendix 1). Sequences and primers were submitted to GenBank (Table 1). Alleles were sized using GeneMarker version 1.90 (SoftGenetics, State College, Pennsylvania, USA). The variability of the microsatellite loci was measured by counting the number of alleles and calculating Nei's unbiased gene diversity using GenAlEx 6.5 (Peakall and Smouse, 2012).

The microsatellite markers CA01–CA09 showed an amplification rate of 100%. CA10 amplified in 85% of the samples (Table 1). Tetranucleotide microsatellites (six out of 10) were the most common microsatellite motif. The microsatellite loci showed between four (CA05) and 13 (CA06) alleles per locus, with a mean of 7.9 and average gene diversities between 0.40 and 0.73 across four populations (Table 2). The other five markers (CA11–CA15) were not considered for the statistical analysis due to lower amplification rates. The primers developed were also tested for cross-species amplification with specimens of *C. muricata* (Ach.) Eckfeldt, *C. crespoae* (Barreno & Vázquez) Kärnefelt, *C. odontella* (Ach.) Ach., *C. australiensis* W. A. Weber ex Kärnefelt, and the still undescribed *C. "panamericana"* (listed in Appendix 1). The polymorphic markers CA01–CA10 successfully amplified in all the species, except for locus CA05 in *C. australiensis* and *C. "panamericana,"* and locus CA10 in *C. australiensis* (Table 3). Linkage disequilibrium in pairwise combinations of the loci within populations and across the total sample was tested using GENEPOP 4.2 (Raymond and Rousset, 1995). Holm-Bonferroni sequential correction of *P* values (Holm, 1979) using the implementation by Gaetano (2013) and a significance threshold of 0.05 resulted in nine significant deviations from linkage equilibrium out of 45 pairwise comparisons across all populations. The deviations involved all loci except CA06 and CA10. Only five out of 171 pairwise comparisons within populations showed significant deviation from linkage equilibrium. These deviations are more likely explained by the predominantly clonal reproduction of *C. aculeata* or population structure than by physical linkage of microsatellite loci.

CONCLUSIONS

The microsatellite markers developed here will facilitate the studies of genetic diversity and population structure of *C. aculeata* along geographic and bioclimatic gradients. These studies will increase our understanding of dispersal patterns and local adaptation in lichens. The fact that primers amplify across a broader range of species may also contribute to a better understanding of species boundaries and speciation within the genus.

LITERATURE CITED

- BOLGER, A. M., M. LOHSE, AND B. USADEL. 2014. Trimmomatic: A flexible trimmer for Illumina sequence data. *Bioinformatics (Oxford, England)* 30: 2114–2120.
- FERNÁNDEZ-MENDOZA, F., S. DOMASCHKE, M. A. GARCÍA, P. JORDAN, M. P. MARTÍN, AND C. PRINTZEN. 2011. Population structure of mycobionts and photobionts of the widespread lichen *Cetraria aculeata*. *Molecular Ecology* 20: 1208–1232.
- FERNÁNDEZ-MENDOZA, F., AND C. PRINTZEN. 2013. Pleistocene expansion of the bipolar lichen *Cetraria aculeata* into the Southern hemisphere. *Molecular Ecology* 22: 1961–1983.
- GAETANO, J. 2013. Holm-Bonferroni sequential correction: An Excel calculator (1.2). Website https://www.researchgate.net/publication/242331583_Holm-Bonferroni_Sequential_Correction_An_EXCEL_Calculator_-_Ver._1.2 [accessed 25 August 2016].
- HOLM, S. 1979. A simple sequentially rejective multiple test procedure. *Scandinavian Journal of Statistics* 6: 65–70.
- HUSON, D., A. AUCH, J. QI, AND S. SCHUSTER. 2007. MEGAN analysis of metagenome data. *Genome Research* 17: 377–386.
- JONES, T. C., I. D. HOGG, R. J. WILKINS, AND T. G. GREEN. 2015. Microsatellite analyses of the Antarctic endemic lichen *Buellia frigida* Darb. (Physciaceae) suggest limited dispersal and the presence of glacial refugia in the Ross Sea region. *Polar Biology* 38: 941–949.
- LUTSAK, T., F. FERNÁNDEZ-MENDOZA, P. KIRIKA, M. WONDAFRASH, AND C. PRINTZEN. 2015. Mycobiont-photobiont interactions of the lichen *Cetraria aculeata* in high alpine regions of East Africa and South America. *Symbiosis* 68: 25–37.
- PEAKALL, R., AND P. SMOUSE. 2012. GenAlEx 6.5: Genetic analysis in Excel. Population genetic software for teaching and research. *Bioinformatics (Oxford, England)* 28: 2537–2539.
- PRINTZEN, C., S. DOMASCHKE, F. FERNÁNDEZ-MENDOZA, AND S. PÉREZ-ORTEGA. 2013. Biogeography and ecology of *Cetraria aculeata*, a widely distributed lichen with a bipolar distribution. *MycKeys* 6: 33–53.
- RAYMOND, M., AND F. ROUSSET. 1995. GENEPOP (version 1.2): Population genetics software for exact tests and ecumenicism. *Journal of Heredity* 86: 248–249.
- THIEL, T., W. MICHALEK, R. K. VARSHNEY, AND A. GRANER. 2003. Exploiting EST databases for the development and characterization of gene-derived SSR-markers in barley (*Hordeum vulgare* L.). *Theoretical and Applied Genetics* 10: 411–422.
- WALSER, J. C., R. HOLDEREGGER, F. GUGERLI, S. E. HOEBEE, AND C. SCHEIDEGGER. 2005. Microsatellites reveal regional population differentiation and isolation in *Lobaria pulmonaria*, an epiphytic lichen. *Molecular Ecology* 14: 457–467.
- ZERBINO, D. R., AND E. BIRNEY. 2008. Velvet: Algorithms for de novo short read assembly using de Bruijn graphs. *Genome Research* 18: 821–829.

APPENDIX 1. Localities and collection data for *Cetraria* species used in this study.

| Species | n | Geographic coordinates | Locality data | Collector and collection year | Voucher no. ^a |
|---|----|---------------------------------|---|-------------------------------------|---------------------------|
| <i>C. aculeata</i> (Schreb.) Fr. | 10 | 40°31'26.3568"N, 3°27'45.8706"W | Spain (Spa1): Provincia de Madrid, Canencia | Fernández-Mendoza, 2014 | FR-0261059 |
| <i>C. aculeata</i> | 23 | 43°6'41.58"N, 5°33'8.9316"W | Spain (Spa2): Provincia de Asturias, Gamoniteiro | Fernández-Mendoza, 2014 | FR-0261060 |
| <i>C. aculeata</i> | 24 | 41°2'5.172"N, 3°10'20.5674"W | Spain (Spa3): Provincia de Jaén, Despeñaperros | Fernández-Mendoza, 2011 | FR-0261062 |
| <i>C. aculeata</i> | 24 | 39°47'47.4612"N, 1°16'26.3316"W | Spain (Spa4): Provincia de Cuenca, Casillas de Ranera | Fernández-Mendoza, 2014 | FR-0261063 |
| <i>C. australiensis</i> W. A. Weber ex Kärnefelt | 2 | 36°27'26.5278"S, 148°16'4.962"E | Australia: New South Wales, Mount Kosciuszko | Fernández-Mendoza & de Miquel, 2009 | FR-0261079, FR-0261080 |
| <i>C. crespocae</i> (Barreno & Vázquez) Kärnefelt | 2 | 39°31'58.0008"N, 4°57'7.9986"W | Spain: Toledo, National Park de Cabañeros | Pérez-Ortega, 2010 | FR-0261081, FR-0261082 |
| <i>C. muricata</i> (Ach.) Eckfeldt | 1 | 49°17'00"N, 87°53'00.66"E | Russian Federation: Altai Republic, Altai Mts. | Lustyk & Kočí, 2011 | hb. P. Resl |
| <i>C. muricata</i> | 1 | 56°35'56.997"N, 4°44'50.5206"W | United Kingdom: Scotland, Rannoch Moor | Fernández-Mendoza, 2010 | FR-0261084 |
| <i>C. odontella</i> (Ach.) Ach. | 1 | 68°32'59.9994"N, 27°22'30"E | Finland: Lake Inari | Westberg | LD1188229 |
| <i>C. odontella</i> | 1 | 44°57'56.7504"N, 6°36'19.1478"E | France: Granon | Fernández-Mendoza, 2013 | FR-0261085 |
| <i>C. "panamericana"</i> | 2 | 51°10'0.0006"N, 117°24'0"W | Canada: British Columbia, Glacier National Park | Goward, 2005 | FR-0261086, FR-0261087 |

Note: n = number of individuals.

^a Vouchers are deposited in Herbarium Senckenbergianum Frankfurt (FR), Lund University Herbarium (LD), and the private herbarium of Philipp Resl.