



Development and Evaluation of Microsatellite Markers for the Critically Endangered Birch *Betula chichibuensis* (Betulaceae)

Authors: Igarashi, Yuji, Aihara, Hiroki, Handa, Yoshihiro, Katsumata, Hiroshi, Fujii, Masanori, et al.

Source: Applications in Plant Sciences, 5(5)

Published By: Botanical Society of America

URL: <https://doi.org/10.3732/apps.1700016>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

DEVELOPMENT AND EVALUATION OF MICROSATELLITE MARKERS FOR THE CRITICALLY ENDANGERED BIRCH *BETULA* *CHICHIBUENSIS* (BETULACEAE)¹

YUJI IGARASHI², HIROKI AIHARA³, YOSHIHIRO HANDA^{3,5}, HIROSHI KATSUMATA^{3,5}, MASANORI FUJII⁴,
KOICHIRO NAKANO^{3,5}, AND TOSHIHIDE HIRAO^{2,6}

²The University of Tokyo Chichibu Forest, Graduate School of Agricultural and Life Sciences, The University of Tokyo, 1-1-49 Hinoda-machi, Chichibu, Saitama 368-0034, Japan; ³FASMAC Co. Ltd., 5-1-3 Midorigaoka, Atsugi, Kanagawa 243-0041, Japan; and ⁴Institute for Environmental Sciences, 1-7 Inenome, Obuchi, Rokkasho, Kamikita, Aomori 039-3212, Japan

- *Premise of the study:* Microsatellite markers were developed and characterized for the critically endangered birch *Betula chichibuensis* (Betulaceae) to investigate the genetic structure of this species for conservation purposes.
- *Methods and Results:* Sixteen microsatellite markers with di-, tri-, and tetranucleotide repeat motifs were developed and optimized using MiSeq paired-end sequencing. Of these, 14 were polymorphic, with two to five alleles per locus, in 47 individuals from two newly discovered populations of *B. chichibuensis* in Japan. Observed and unbiased expected heterozygosities per locus ranged from 0.000 to 0.617 and from 0.000 to 0.629, respectively. These markers were tested for cross-species amplification in *B. maximowicziana*, *B. platyphylla* var. *japonica*, and *B. schmidtii*.
- *Conclusions:* This set of microsatellite markers, the first developed for *B. chichibuensis*, will help elucidate spatial patterns of gene flow and levels of inbreeding in this species to aid its conservation.

Key words: *Betula chichibuensis*; Betulaceae; conservation genetics; critically endangered species; microsatellites; MiSeq sequencing.

The genus *Betula* L. (Betulaceae) comprises approximately 60 tree species distributed in boreal and cool-temperate zones of the Northern Hemisphere (Furlow, 1990). Individuals of *B. chichibuensis* H. Hara (subgenus *Aspera*) are small trees endemic to Japan (Ashburner and McAllister, 2013). Partly because its habitat is limited to limestone outcrops, this species is narrowly confined to the Chichibu (McAllister, 1993; Igarashi and Yoshida, 2013) and Kitakami (Nagato and Shimai, 2007) mountains in central and northeastern Honshu, respectively. Because only a few small populations have been recorded in these locations, *B. chichibuensis* is listed as critically endangered on the IUCN Red List (Shaw et al., 2014). The small population sizes and restricted distribution of *B. chichibuensis* make this species susceptible to diseases and natural disasters, and seriously impede gene flow (Ministry of the Environment, 2015). Analysis of *B. chichibuensis* genetic structure and maintenance of its

genetic diversity are therefore essential for both in situ and ex situ conservation.

In this study, we developed microsatellite markers for *B. chichibuensis* to investigate the current genetic status of the remaining populations. We also examined the transferability of these developed markers to three other *Betula* species: *B. maximowicziana* Regel (subgenus *Acuminata*), *B. platyphylla* Sukaczew var. *japonica* (Miq.) H. Hara (subgenus *Betula*), and *B. schmidtii* Regel (subgenus *Aspera*).

METHODS AND RESULTS

Plant material and DNA extraction—Plant materials of *B. chichibuensis* were collected from two newly discovered populations growing on limestone outcrops on western Futago Mountain (WF) and along the Oku-Chichibu Forest Road (OC) in the Chichibu Mountains of Japan (Appendix 1). Shoots of 23 and 24 individuals were collected from WF and OC, respectively. Genomic DNA was extracted from freeze-dried leaves and winter buds using a DNeasy Plant Mini Kit (QIAGEN, Hilden, Germany). The concentration of genomic DNA was determined with a Qubit 2.0 Fluorometer (Life Technologies, Carlsbad, California, USA) and by gel electrophoresis.

Microsatellite marker development—A total of 400 ng of genomic DNA from an individual OC sample was sheared with NEBNext dsDNA Fragmentase (New England Biolabs, Ipswich, Massachusetts, USA). A paired-end library for MiSeq sequencing (Illumina, San Diego, California, USA) was generated using a NEBNext Ultra DNA Library Prep Kit for Illumina (New England Biolabs). A single 301-bp paired-end sequencing run yielded 20,746,148 reads (DNA Data Bank of Japan [DDBJ] Sequence Read Archive accession no.:

¹Manuscript received 19 February 2017; revision accepted 30 March 2017.

The authors thank Toyotaro Iwata for helping with fieldwork. This study was supported in part by the 25th Pro Natura Fund (2014) of the Pro Natura Foundation of Japan and a joint project between the University of Tokyo Chichibu Forest and the Suntory Natural Water Sanctuary.

⁵Current address: Bioengineering Laboratory Co. Ltd., 3068 Tenko 7th Building, Sakai, Atsugi, Kanagawa 243-0022, Japan

⁶Author for correspondence: hirao@uf.a.u-tokyo.ac.jp

doi:10.3732/apps.1700016

Applications in Plant Sciences 2017 5(5): 1700016; <http://www.bioone.org/loi/apps> © 2017 Igarashi et al. Published by the Botanical Society of America. This is an open access article distributed under the terms of the Creative Commons Attribution License (CC-BY-NC-SA 4.0), which permits unrestricted noncommercial use and redistribution provided that the original author and source are credited and the new work is distributed under the same license as the original.

DRA005642). Raw reads with quality scores less than 20 and lengths shorter than 20 bp were filtered using Sickle version 1.33 (Joshi and Fass, 2011). De novo assembly using Velvet version 1.2.10 (Zerbino and Birney, 2008) produced 204,911 contigs, where parameters were set as *k*-mer as 91, auto coverage cut-off, and minimum contig length of 300. The data sets were collated and filtered in QDD version 3.1 (Megléc et al., 2014) to generate sequences containing microsatellites and to design PCR primers. A total of 125 perfect microsatellite loci consisting of di-, tri-, tetra-, penta-, and hexanucleotide repeat motifs were identified according to the following parameters: GC content of 40–60%, a melting temperature of 57°C to 63°C, and a maximum difference of 2°C between forward and reverse primers.

Microsatellite marker screening—For initial screening, of the 125 loci identified, 56 were selected based on repeat number and fragment size. For these loci, PCR amplification and polymorphism were tested using 10 samples. Individual primer pairs were assayed in 10-μL reaction mixtures containing 4 ng of genomic DNA, 0.05 μM of M13–21-tagged (5'-TGTAACGACGCGCCAGT-3') forward primer, 0.2 μM of reverse primer, 0.2 μM of universal primer labeled with 6-FAM fluorescent dye (Applied Biosystems, Foster City, California, USA), 0.2 μL of PrimeSTAR GXL DNA polymerase (TaKaRa Bio Inc., Tokyo, Japan), 2 μL of 5× PrimeSTAR GXL Buffer, and 0.2 mM of dNTP mixture. Thermal cycling conditions consisted of 98°C for 5 min; followed by 38 cycles of 98°C for 20 s, touchdown annealing (65°C for four cycles, 63°C for four cycles, 60°C for 20 cycles, and 53°C for 10 cycles) for 20 s, and 68°C for 40 s; and a final step of 68°C for 2 min. Based on the test results, 16 primer pairs were selected (Table 1).

Polymorphism of the 16 markers was examined in 47 samples from the two distinct WF and OC populations (Appendix 1). Following a modified version of the

efficient genotyping method described by Blacket et al. (2012), locus-specific forward primers were tagged with 5' sequences (Table 1), while universal primers were labeled with different fluorescent dyes (6-FAM, VIC, NED, or PET; Applied Biosystems). Two sets of 8-plex PCR amplifications were performed in 10-μL reaction mixtures containing 10 ng of genomic DNA, 0.05 μM of forward primer, 0.2 μM of reverse primer, 0.2 μM of fluorescently labeled primer, 0.5 μL of PrimeSTAR GXL DNA polymerase (TaKaRa Bio Inc.), 2 μL of 5× PrimeSTAR GXL Buffer, and 0.3 mM of dNTP mixture. To obtain high-quality amplification product, a modification of the touchdown PCR procedure of Korbie and Mattick (2008) was carried out using the following cycling conditions: 98°C for 5 min; followed by 50 cycles of 98°C for 30 s, touchdown annealing (63°C to 57°C [decreasing 1°C every two cycles] for 14 cycles, 56°C for 15 cycles, 53°C to 51°C [decreasing 1°C every two cycles] for six cycles, and 50°C for 15 cycles) for 90 s, and 68°C for 40 s; and a final step of 68°C for 15 min. Finally, 1 μL of PCR product was mixed with 0.5 μL of GeneScan 600 LIZ Size Standard (Applied Biosystems) and 8.5 μL Hi-Di formamide (Applied Biosystems) and sequenced on an ABI 3730xl DNA Analyzer (Applied Biosystems). Genotypes were scored by analyzing fragment sizes using Peak Scanner version 2.0 (Applied Biosystems).

Microsatellite marker evaluation—Descriptive statistics were computed for the assayed markers using CERVUS version 3.0.7 (Kalinowski et al., 2007). Of the 16 loci tested, 14 were polymorphic, with two to five alleles per locus detected across 47 individuals from the WF and OC populations (Table 2). The mean number of alleles per locus was 2.438, with mean observed and unbiased expected heterozygosities per locus of 0.327 (0.000–0.617) and 0.350 (0.000–0.629), respectively. The mean number of alleles per locus was 2.250 in the WF population and 2.313 in the OC population. For the WF population, mean observed and unbiased expected

TABLE 1. Characteristics of 16 microsatellite markers developed for *Betula chichibuensis*.^a

Locus	Primer sequences (5'–3')	Repeat motif	Allele size range (bp)	Fluorescent dye ^b (Multiplex set no.)	GenBank accession no.
Bcc3	F: CTTGTCCCTCATCAGCCTTG R: GCCATTGGTGTGTCATATCTT	(CT) ₁₂	281–283	PET2 (1)	LC214895
Bcc4	F: CCTGGTAACAAGATCATTGCAT R: GGGTTCCTCATCAAACCCCTGA	(CT) ₁₂	243–245	NED1 (1)	LC214896
Bcc7	F: GGAATAGCCCATTCGACCTT R: GAGATCAGCGATTTCCAGT	(AG) ₁₁	277–283	FAM2 (1)	LC214897
Bcc10	F: GACCCGGCACAAACTTAATC R: GGGAAAGGATCGTGGAGAAAT	(AG) ₁₁	230–234	VIC1 (1)	LC214898
Bcc11	F: CATGCCTTGGAACTGGAAG R: CAATGTCAGAACCTGCTCCT	(CT) ₁₁	254–256	PET1 (1)	LC214899
Bcc13	F: GGGACCCTTGAAGATGACA R: TGAATGCCGTCCTTAGCTCT	(TC) ₁₁	234	PET1 (2)	LC214900
Bcc16	F: CAACCAGGAGGGCGTATTAG R: GATCAGATCCGCAATGCTAT	(AG) ₁₁	260–262	VIC2 (2)	LC214901
Bcc18	F: GGACTCCCATCTGGTAGGAT R: GCCACTCCCTAATCATCAACTT	(AT) ₁₁	260–276	NED2 (2)	LC214902
Bcc22	F: GTGCAAACCTGACTCTGGCG R: ATATGCATGTCGAGCCTGT	(CAA) ₈	302–305	PET2 (2)	LC214903
Bcc25	F: ACCATTGTTGTGGAGTCCG R: AAAGCGCGTAGAGCTCGTTT	(AGA) ₈	156–160	FAM1 (2)	LC214904
Bcc27	F: CGTACACCCAAACTGACCCT R: CTTTCGCCATTGAGTTCTCC	(CCA) ₇	203–215	NED1 (2)	LC214905
Bcc30	F: GCGATAGCCGTTCTTCCATA R: CGGCTTAATGCCATAGAAGG	(AAG) ₇	221–224	FAM1 (1)	LC214906
Bcc34	F: GGGTGAAGCAGCTAGAGAC R: CCAACTTGGTGCCATTAGTGT	(AAG) ₇	200–205	VIC1 (2)	LC214907
Bcc38	F: ACGTGTGGCAAATTCATTG R: TTGCTGCATGAACCTCTTTC	(CTTT) ₆	241	FAM2 (2)	LC214908
Bcc46	F: GCACCGTTAAGTGATCAAA R: ACATACGCCTTGAATCGCA	(GAAA) ₅	271–283	VIC2 (1)	LC214909
Bcc50	F: TACAGGTTGGTTGCCAAAT R: TCCCGTATTAAGGACCATCTG	(TTAT) ₅	300–308	NED2 (1)	LC214910

^a Touchdown annealing temperatures (63°C to 57°C [–1°C every two cycles] for 14 cycles, 56°C for 15 cycles, 53°C to 51°C [–1°C every two cycles] for six cycles, and 50°C for 15 cycles) were used.

^b Sequences of fluorescent labels were as follows: FAM1 = 5'-TGTAACGACGCGCCAGT-3' (M13–21), VIC1 = 5'-CGCATTCTCATTGCATAC-3' (CMV-Fw), NED1 = 5'-ATGCTAGTTATTGCTCAG-3' (pBAD-F), PET1 = 5'-CAGTAATCAGCTATGACG-3' (M13-P5), FAM2 = 5'-TGTAACGACGACGATCGT-3' (modified M13–21), VIC2 = 5'-CGCATTCTCAACTCTATG-3' (modified CMV-Fw), NED2 = 5'-ATGCTAGTTATCTGCAGT-3' (modified pBAD-F), PET2 = 5'-CAGTAATCAGGATCTGAC-3' (modified M13-P5).

TABLE 2. Genetic variation of 16 microsatellite loci in two natural populations of *Betula chichibuensis* in central Honshu, Japan.^a

Locus	Western Futago Mountain (n = 23)				Oku-Chichibu Forest Road (n = 24)				Total (n = 47)			
	A	H _o ^b	H _e	Null	A	H _o ^b	H _e	Null	A	H _o ^b	H _e	Null
Bcc3	2	0.478	0.414	-0.082	2	0.250	0.284	0.053	2	0.362	0.351	-0.021
Bcc4	2	0.391	0.322	-0.107	2	0.458	0.510	0.043	2	0.426	0.454	0.027
Bcc7	3	0.130	0.127	-0.024	3	0.542	0.513	-0.056	3	0.340	0.356	0.002
Bcc10	3	0.391	0.492	0.101	3	0.542	0.536	-0.020	3	0.468	0.563	0.092
Bcc11	1	0.000	0.000	—	2	0.250	0.223	-0.062	2	0.128	0.121	-0.024
Bcc13	1	0.000	0.000	—	1	0.000	0.000	—	1	0.000	0.000	—
Bcc16	2	0.043	0.043	-0.004	1	0.000	0.000	—	2	0.021	0.021	-0.001
Bcc18	2	0.304	0.264	-0.079	2	0.500	0.383	-0.142	2	0.404	0.326	-0.111
Bcc22	3	0.348	0.414	0.066	3	0.167	0.159	-0.035	3	0.255	0.298	0.059
Bcc25	3	0.304	0.425	0.167	3	0.708	0.681	-0.031	3	0.511	0.613	0.100
Bcc27	4	0.565	0.641	0.052	4	0.667	0.593	-0.067	5	0.617	0.628	0.006
Bcc30	2	0.391	0.414	0.018	2	0.417	0.496	0.077	2	0.404	0.461	0.060
Bcc34	2	0.043	0.043	-0.004	3	0.500	0.465	-0.074	3	0.277	0.284	-0.009
Bcc38	1	0.000	0.000	—	1	0.000	0.000	—	1	0.000	0.000	—
Bcc46	2	0.522	0.487	-0.046	2	0.417	0.422	-0.004	2	0.468	0.500	0.027
Bcc50	3	0.652	0.633	-0.033	3	0.458	0.606	0.133	3	0.553	0.629	0.058
Average	2.250	0.285	0.295	0.002	2.313	0.367	0.367	-0.014	2.438	0.327	0.350	0.019

Note: A = number of alleles; H_e = unbiased expected heterozygosity; H_o = observed heterozygosity; n = number of individuals sampled; Null = null allele frequency estimate.

^aVoucher and locality information are provided in Appendix 1.

^bNo significant deviation from Hardy–Weinberg equilibrium was detected (P < 0.05).

heterozygosities per locus were 0.285 (0.000–0.652) and 0.295 (0.000–0.641), respectively; for the OC population, the corresponding values were 0.367 (0.000–0.708) and 0.367 (0.000–0.681). GENEPOP version 4.2 (Rousset, 2008) was used to test for deviations from Hardy–Weinberg equilibrium. No significant deviations (P < 0.05) were observed at any of the loci in either population. Null allele frequency estimates were nearly zero or negative except for Bcc10 and Bcc25 in the WF population and Bcc50 in the OC population. Cross-amplifications were carried out to test marker transferability to closely related species (Appendix 1). Polymorphic variation was detected at six loci in *B. maximowicziana* and *B. platyphylla* var. *japonica* and at seven loci in *B. schmidtii* (Table 3). These results are consistent with the close phylogenetic relationship of *B. chichibuensis* and *B. schmidtii* (Wang et al., 2016).

TABLE 3. Cross-amplification of 16 microsatellite loci in three species closely related to *Betula chichibuensis*.^a

Locus	<i>B. maximowicziana</i> (n = 5)			<i>B. platyphylla</i> var. <i>japonica</i> (n = 5)			<i>B. schmidtii</i> (n = 5)		
	A	H _o	H _e	A	H _o	H _e	A	H _o	H _e
Bcc3	2	0.400	0.356	2	0.200	0.200	2	0.200	0.200
Bcc4	—	—	—	—	—	—	—	—	—
Bcc7	1	0.000	0.000	1	0.000	0.000	2	0.400	0.356
Bcc10	1	0.000	0.000	2	0.200	0.200	—	—	—
Bcc11	2	0.400	0.356	3	0.400	0.600	—	—	—
Bcc13	2	0.200	0.200	2	0.200	0.200	1	0.000	0.000
Bcc16	—	—	—	—	—	—	—	—	—
Bcc18	1	0.000	0.000	4	0.600	0.711	—	—	—
Bcc22	1	0.000	0.000	—	—	—	—	—	—
Bcc25	1	0.000	0.000	—	—	—	1	0.000	0.000
Bcc27	—	—	—	—	—	—	2	0.000	0.533
Bcc30	1	0.000	0.000	1	0.000	0.000	2	0.400	0.356
Bcc34	—	—	—	1	0.000	0.000	2	0.400	0.533
Bcc38	5	1.000	0.844	1	0.000	0.000	2	0.400	0.356
Bcc46	2	0.200	0.556	3	0.400	0.600	1	0.000	0.000
Bcc50	2	0.000	0.356	1	0.000	0.000	2	0.200	0.200
Average	1.313	0.138	0.167	1.313	0.125	0.157	1.063	0.125	0.158

Note: — = failed or nonspecific amplification (i.e., three or more polymorphic bands detected); A = number of alleles; H_e = unbiased expected heterozygosity; H_o = observed heterozygosity; n = number of individuals sampled.

^aTesting for Hardy–Weinberg equilibrium and estimation of null allele frequency were not performed because of small sample sizes.

CONCLUSIONS

We developed 16 microsatellite markers for the critically endangered birch *B. chichibuensis* using MiSeq paired-end sequencing. These markers will facilitate understanding of spatial patterns of gene flow and levels of inbreeding, information essential for the conservation of the small isolated populations of this species. Some of the markers were successfully transferred to closely related *Betula* species.

LITERATURE CITED

- ASHBURNER, K., AND H. A. McALLISTER. 2013. The genus *Betula*: A taxonomic revision of birches. Royal Botanic Gardens, Kew, Richmond, United Kingdom.
- BLACKETT, M. J., C. ROBIN, R. T. GOOD, S. F. LEE, AND A. D. MILLER. 2012. Universal primers for fluorescent labelling of PCR fragments—An efficient and cost-effective approach to genotyping by fluorescence. *Molecular Ecology Resources* 12: 456–463.
- FURLOW, J. J. 1990. The genera of Betulaceae in the southeastern United States. *Journal of the Arnold Arboretum* 71: 1–67.
- IGARASHI, Y., AND Y. YOSHIDA. 2013. List of vascular plants of the University of Tokyo Chichibu Forest. *University of Tokyo Forests* 54: 107–155 (in Japanese).
- JOSHI, N. A., AND J. N. FASS. 2011. Sickle: A sliding-window, adaptive, quality-based trimming tool for FastQ files, version 1.33. Website <https://github.com/najoshi/sickle> [accessed 30 January 2017].
- KALINOWSKI, S. T., M. L. TAPER, AND T. C. MARSHALL. 2007. Revising how the computer program CERVUS accommodates genotyping error increases success in paternity assignment. *Molecular Ecology* 16: 1099–1106.
- KORBIE, D. J., AND J. S. MATTICK. 2008. Touchdown PCR for increased specificity and sensitivity in PCR amplification. *Nature Protocols* 3: 1452–1456.
- McALLISTER, H. A. 1993. Cytology and the conservation of rare birches. In D. R. Hunt [ed.], *Betula*: Proceedings of the IDS *Betula* symposium, 61–66. International Dendrology Society, Morpeth, Northern Ireland.
- MEGLÉCZ, E., N. PECH, A. GILLES, V. DUBUT, P. HINGAMP, A. TRILLES, R. GRENIER, AND J.-F. MARTIN. 2014. QDD version 3.1: A user-friendly computer program for microsatellite selection and primer design revisited: Experimental validation of variables determining genotyping success rate. *Molecular Ecology Resources* 14: 1302–1313.

- MINISTRY OF THE ENVIRONMENT. 2015. Red data book 2014: Threatened wildlife of Japan, vol. 8: Vascular plants. Gyosei, Tokyo, Japan (in Japanese).
- NAGATO, K., AND S. SHIMAI. 2007. On the *Betula chichibuensis* community developing in the limestone of Kataiwa area in Tono City, Iwate Prefecture, Northeastern Japan. *Bulletin of Daito Bunka University* 45: 1–16 (in Japanese).
- ROUSSET, F. 2008. GENEPOP'007: A complete re-implementation of the GENEPOP software for Windows and Linux. *Molecular Ecology Resources* 8: 103–106.
- SHAW, K., S. ROY, AND B. WILSON. 2014. *Betula chichibuensis*. The IUCN Red List of Threatened Species 2014: e.T194282A2309490. Website <http://dx.doi.org/10.2305/IUCN.UK.2014-3.RLTS.T194282A2309490.en> [accessed 30 January 2017].
- WANG, N., H. A. MCALLISTER, P. R. BARTLETT, AND R. J. A. BUGGS. 2016. Molecular phylogeny and genome size evolution of the genus *Betula* (Betulaceae). *Annals of Botany* 117: 1023–1035.
- ZERBINO, D. R., AND E. BIRNEY. 2008. Velvet: Algorithms for de novo short read assembly using de Bruijn graphs. *Genome Research* 18: 821–829.

APPENDIX 1. Voucher information for species used in the development and evaluation of microsatellite markers for *Betula chichibuensis*.

Taxon	Population	Location	Geographic coordinates ^a	N	Voucher no. ^b
<i>B. chichibuensis</i> H. Hara	Western Futago Mountain (WF)	Ogano, Saitama, Japan	36°04'N, 138°51'E	23	UTCFCB 00001–00015, 00017–00024
	Oku-Chichibu Forest Rd. (OC)	Chichibu, Saitama, Japan	35°57'N, 138°44'E	24	UTCFCB 00073–00096
<i>B. maximowicziana</i> Regel	Tochimoto	Chichibu, Saitama, Japan	35°57'N, 138°49'E	5	UTCFFT 00039.1–00039.5
<i>B. platyphylla</i> Sukaczew var. <i>japonica</i> (Miq.) H. Hara	Tochimoto	Chichibu, Saitama, Japan	35°56'N, 138°49'E	5	UTCFFT 00040.1–00040.5
<i>B. schmidtii</i> Regel	Ochigawa	Chichibu, Saitama, Japan	35°54'N, 138°59'E	5	UTCFFT 00041.1–00041.5

Note: N = number of samples.

^aFor conservation reasons, low-resolution geographic coordinates are given.

^bAll vouchers are stored at the Herbarium of the University of Tokyo Chichibu Forest (UTCF), The University of Tokyo, Saitama, Japan.