



Isolation and Characterization of Microsatellite Loci from *Arthropodium cirratum* (Asparagaceae)

Authors: Bulgarella, Mariana, Biggs, Patrick J., Lange, Peter J. de, and Shepherd, Lara D.

Source: Applications in Plant Sciences, 5(8)

Published By: Botanical Society of America

URL: <https://doi.org/10.3732/apps.1700041>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

ISOLATION AND CHARACTERIZATION OF MICROSATELLITE LOCI FROM *ARTHROPODIUM CIRRATUM* (ASPARAGACEAE)¹

MARIANA BULGARELLA², PATRICK J. BIGGS³, PETER J. DE LANGE⁴, AND LARA D. SHEPHERD^{2,5}

²Museum of New Zealand Te Papa Tongarewa, P.O. Box 467, Wellington, New Zealand; ³Institute of Fundamental Sciences, Massey University, Palmerston North, New Zealand; and ⁴Science and Capability Group, Terrestrial Ecosystems, Department of Conservation, Newton, Auckland, New Zealand

- **Premise of the study:** Microsatellite markers were developed for *Arthropodium cirratum* (Asparagaceae) to study population genetic structure and translocation of this species. These markers were tested for cross-amplification in two other *Arthropodium* species.
- **Methods and Results:** Sixteen microsatellite markers were developed from a genomic library and tested in three populations of *A. cirratum*. The loci exhibited one to five alleles per locus, with private alleles present in each of the populations. Cross-amplification tests in the two other New Zealand *Arthropodium* species revealed that many of the loci amplify and demonstrate polymorphism in *A. bifurcatum*.
- **Conclusions:** These markers will be useful for determining genetic structure in *A. cirratum* and for determining the origins of translocated populations of this species.

Key words: *Arthropodium*; Asparagaceae; nuclear microsatellites; rengarenga; translocation.

Arthropodium R. Br. (Asparagaceae) is a genus of nine species (Heenan et al., 2004) of perennial, lily-like herbs found in Australia, New Zealand, New Caledonia, New Guinea, and Madagascar. Within New Zealand, there are three endemic species: *A. candidum* Raoul (small renga lily), *A. cirratum* (G. Forst.) R. Br., and *A. bifurcatum* Heenan, A. D. Mitch. & de Lange (the latter two species both have the common names rengarenga and New Zealand rock lily).

Arthropodium cirratum was cultivated as a food source for Māori and translocated beyond its natural range (Shepherd et al., 2016). A recent phylogeographic study of *A. cirratum* using chloroplast sequences revealed a very high level of structuring, with many populations fixed for unique chloroplast haplotypes (Shepherd et al., 2016). Microsatellite markers will be useful for testing whether the nuclear genome demonstrates the high genetic structuring found for the chloroplast genome (Shepherd et al., 2016). They will also aid in examining the origins of the populations that derive from translocation by Māori and testing proposed hybridization between *A. cirratum* and *A. bifurcatum* (Heenan et al., 2004).

¹Manuscript received 19 April 2017; revision accepted 22 May 2017.

This study was funded by a Rutherford Discovery Fellowship from the Royal Society of New Zealand (contract no. RDF-MNZ1201) to L.D.S. The authors thank all iwi and hapu who granted us permission to collect from within their rohe. Samples were collected under Department of Conservation permits WA-23814-FLO, BOP-23814-FLO, TT-23661-FLO, NO-233360-FLO and Otari Wilton's Bush permit 145.

⁵Author for correspondence: lara.shepherd@tepapa.govt.nz

doi:10.3732/apps.1700041

METHODS AND RESULTS

We extracted DNA from leaf tissue of four *A. cirratum* individuals (Appendix 1), each from different populations, using a modified cetyltrimethylammonium bromide (CTAB) method (steps 1, 3–7 from table 1 in Shepherd and McLay, 2011). The extracted DNA was pooled and amplified using a REPLI-g kit (QIAGEN, Hilden, Germany) following the manufacturer's protocol to generate sufficient template for library construction. An Illumina paired-end genomic library was constructed using the TruSeq Nano DNA Library Prep Kit (Illumina, San Diego, California, USA) following the manufacturer's instructions. The library was sequenced in a single lane using the Illumina MiSeq platform to generate 2 × 250-bp reads at the Massey Genome Service (Massey University, Palmerston North, New Zealand).

We assembled the resulting 10,955,497 paired sequence reads using MEGAHIT (Li et al., 2015), as this software required sequence reads of the same length. A set of four assembly parameters was tried with the reads, and the resulting contigs were merged to make a set of longest unique contigs. This resulted in 1.589 Gb of assembled sequence, comprising 2,618,361 contigs, with a maximum length of 18,007 bp, average GC content of 34.74%, and an N50 of 1513 bp when analyzed using QUAST (Gurevich et al., 2013). The SSR_pipeline (Miller et al., 2013) was used to detect di- and tetranucleotide repeats on this contig set with a minimum of 250-bp flanking sequence on each side to allow for PCR primer design. We used WebSat (Martins et al., 2009) to develop primers for 33 loci, which had at least eight tetra- or 15 dinucleotide repeat units. An M13 tag (TGTAACAACGACGGCCAGT) was added to the 5' end of the forward primer of each locus. These primer pairs were tested on five samples, which included three samples of *A. cirratum* and one sample each of *A. candidum* and *A. bifurcatum*. Each locus was initially amplified individually in 10-μL PCR reactions that contained 1 μL of diluted template DNA, 0.02 μM forward primer, 0.8 μM reverse primer, 0.8 μM M13 primer (labeled with FAM, NED, PET, or HEX), 1× MyTaq mix (Bioline, London, United Kingdom), and 0.1 M betaine. PCR thermocycling conditions were an initial denaturation of 94°C for 5 min; 30 cycles of 94°C for 30 s, 55°C for 45 s, and 72°C for 45 s; followed by eight cycles of 94°C for 30 s, 53°C for 45 s, and 72°C for 45 s; and a final extension at 72°C for 15 min.

Of the 33 primer pairs tested, 16 amplified in at least two species and were polymorphic. These 16 loci were subsequently screened using 63 samples from three populations of *A. cirratum* and additional samples of *A. bifurcatum* and

Applications in Plant Sciences 2017 5(8): 1700041; <http://www.bioone.org/loi/apps> © 2017 Bulgarella et al. Published by the Botanical Society of America. This is an open access article distributed under the terms of the Creative Commons Attribution License (CC-BY-NC-SA 4.0), which permits unrestricted noncommercial use and redistribution provided that the original author and source are credited and the new work is distributed under the same license as the original.

TABLE 1. Primer sequences and thermal cycling conditions for 16 microsatellite loci developed for *Arthropodium cirratum*.

Locus	Primer sequences (5′–3′)	Repeat motif	Allele size range (bp)	T _a (°C)	Fluorescent dye (Pooling group)	GenBank accession no.
ArtCir1	F: AAAACACAGCAGACAAACACA R: ATTGTA CTCCGCTCATTGTTC	(CCTC) ₇	351–359	52	FAM (3)	KY907147
ArtCir4	F: CAGTTTCGCTAAAGGACGGAG R: TAATTGGACCTCTCTCATCGGT	(TATT) ₇	207–219	55	FAM (3)	KY907148
ArtCir7	F: AATTGCCTTCAACGTCTTTAGC R: CGAATACGAACCCCATATTGAC	(AATA) ₇	200–215	55	FAM (1)	KY907149
ArtCir9	F: GCCGAAGCTGACAAATGAAA R: CCCACATAATCAAAACCTCCAT	(TCTT) ₇	255–267	55	FAM (2)	KY907150
ArtCir12	F: CCTACCTGCATCTTGACCTTGT R: GTTGAGAGAATGACACTTGGGC	(TTTG) ₈	360–372	55	NED (3)	KY907151
ArtCir13	F: TTCGATAGAGAGTGGTGACGAG R: AAATCAATCCCCTCCGTTAGAT	(TATT) ₇	260–272	55	HEX (1)	KY907152
ArtCir18	F: CTTGTAAAGTCAAGCTCATCGGT R: ACCGGACATCCAACAATTAGAA	(TAAA) ₁₁	318–336	55	HEX (1)	KY907153
ArtCir22	F: ACATCTTTTCATACACGGGCTT R: CTCTCAAGGATCACAAGGAACC	(ATAA) ₉	382–405	52	PET (1)	KY907154
ArtCir23	F: CGAAAACGACTAACGTGAAGAA R: TATGTGTTGGTTGAAGGAGAGC	(ATGT) ₇	344–360	55	FAM (2)	KY907155
ArtCir26	F: GGGCCACTCATATTTTCAATTTTC R: GTAGGTGCTATCCTCCCTTCCT	(CATA) ₇	380–401	52	NED (2)	KY907156
ArtCir32	F: CCGTACTTCTCTCTGTTTGTGT R: ACCCAACCCTCATTTTATCTCC	(TAAA) ₇	367–385	56	NED (1)	KY907157
ArtCir38	F: AGCTATGCCCTCTTTTAGTCA R: ACCAAGATTGCTCCATCAAAGT	(ATTT) ₇	274–289	55	HEX (2)	KY907158
ArtCir43	F: TAAAGAGGAAATGGGTAGGT R: TCTTCTACAACAACCCGAGAA	(TA) ₁₈	388–409	55	PET (2)	KY907159
ArtCir48	F: TTCGCAAAGGATATTAGGTGTG R: TACGAGAACAAGGGAGGGATTA	(AT) ₂₀	305–409	55	PET (2)	KY907160
ArtCir50	F: GGCTAATTTTAAATGTGCTTGGC R: ATGGATGAGAGAGAAAGGACCA	(AT) ₁₈	358–403	55	NED (3)	KY907161
ArtCir59	F: CTATCTCACCATATCGCTGC R: TCGTTTCAAGACAGAAGGCAT	(AT) ₁₈	290–305	55	PET (1)	KY907162

Note: T_a = annealing temperature.

A. candidum (Appendix 1). For this trial, some loci were coamplified in the same PCR reaction (ArtCir13 with ArtCir18, ArtCir9 with ArtCir23, and ArtCir43 with ArtCir48). For these combined PCR reactions, 1 μL of diluted template DNA was combined with 0.02 μM each forward primer, 0.8 μM each reverse primer, 1.2 μM M13 primer (labeled with FAM, NED, PET, or HEX), 1× MyTaq mix (Bioline), and 0.075 M betaine. The PCR annealing temperatures are reported in Table 1. Genotyping was performed on an ABI 3130xl Genetic

Analyzer (Applied Biosystems, Foster City, California, USA) at the Massey Genome Service. Alleles were sized using the internal size standard GeneScan 500 LIZ (Applied Biosystems) and scored using Geneious version 10.0.2 (Biomatters Ltd., Auckland, New Zealand).

The number of alleles and observed and expected heterozygosities for the three *A. cirratum* populations were determined using GenAEX 6.5 (Peakall and Smouse, 2012). Observed and expected heterozygosities ranged from 0.000

TABLE 2. Genetic diversity measures for three populations of *Arthropodium cirratum*.^a

Locus	Maunganui Bluff (N = 22)			Matapouri Bay (N = 20)			Hick's Bay (N = 21)			Total (N = 63)
	A	H _o	H _e	A	H _o	H _e	A	H _o	H _e	A _T
ArtCir1	1	0.000	0.000	2	0.100	0.261	1	0.000	0.000	2
ArtCir4	1	0.000	0.000	2	0.100	0.455*	2	0.048	0.046	3
ArtCir7	1	0.000	0.000	3	0.100	0.261	1	0.000	0.000	4
ArtCir9	1	0.000	0.000	1	0.000	0.000	1	0.000	0.000	1
ArtCir12	1	0.000	0.000	1	0.000	0.000	1	0.000	0.000	2
ArtCir13	1	0.000	0.000	1	0.000	0.000	2	0.286	0.444	3
ArtCir18	2	1.000	0.500*	2	0.800	0.480*	2	1.000	0.500*	4
ArtCir22	1	0.000	0.000	2	0.000	0.495*	1	0.000	0.000	2
ArtCir23	1	0.000	0.000	2	0.100	0.320*	2	0.143	0.278	4
ArtCir26	1	0.000	0.000	2	0.100	0.180	2	0.000	0.172	5
ArtCir32	1	0.000	0.000	1	0.000	0.000	2	0.095	0.091	3
ArtCir38	1	0.000	0.000	2	0.850	0.489*	3	0.952	0.544*	5
ArtCir43	1	0.000	0.000	1	0.000	0.000	1	0.000	0.000	2
ArtCir48	1	0.000	0.000	2	0.100	0.095	1	0.000	0.000	3
ArtCir50	1	0.000	0.000	1	0.000	0.000	2	0.810	0.495	2
ArtCir59	2	0.045	0.044	2	0.150	0.139	1	0.000	0.000	3

Note: A = number of alleles; A_T = total number of alleles; H_e = expected heterozygosity; H_o = observed heterozygosity; N = number of samples.

^aLocality and voucher information are provided in Appendix 1.

*Significant departure from Hardy–Weinberg equilibrium (HWE) at P < 0.05 following sequential Bonferroni correction.

TABLE 3. Cross-amplification of 16 *Arthropodium cirratum* microsatellites in *A. bifurcatum* and *A. candidum*, showing fragment sizes of each allele.

Locus	<i>A. bifurcatum</i> (<i>N</i> = 6)	<i>A. candidum</i> (<i>N</i> = 3)	<i>A. cirratum</i> (<i>N</i> = 63)	<i>A_T</i> (<i>N</i> = 72)
ArtCir1	351, 355, 359	363	351, 359	4
ArtCir4	207, 215, 219	219	208, 211, 219	5
ArtCir7	200, 208, 212	189	200, 212, 213, 215	6
ArtCir9	267, 256	267	255	3
ArtCir12	360, 363, 372	368	363, 368	4
ArtCir13	268	—	260, 268, 272	3
ArtCir18	336	—	318, 334, 336, 325	4
ArtCir22	386, 389, 402, 405	—	382, 386	5
ArtCir23	344, 355	—	345, 348, 357, 360	6
ArtCir26	380, 388, 390, 401	—	383, 387, 393, 397, 399	9
ArtCir32	375, 379	—	367, 369, 385	5
ArtCir38	281, 276	—	274, 281, 282, 285, 289	6
ArtCir43	388, 394, 403	391	408, 409	6
ArtCir48	313	—	303, 313, 315	3
ArtCir50	358, 377, 385	403	358, 403	4
ArtCir59	290	290	290, 297, 305	3

Note: *A_T* = total number of alleles; *N* = number of samples.

to 1.000 and 0.044 to 0.544, respectively (Table 2). Alleles per locus ranged from one to five in *A. cirratum* (mean = 3). All three of the *A. cirratum* populations exhibited private alleles, and 14 of the loci had private alleles in at least one of the three populations. Tests of pairwise linkage disequilibrium were performed using GENEPOP 4.2 (Rousset, 2008). No significant linkage disequilibrium was detected among paired loci comparisons after sequential Bonferroni correction (Holm, 1979). Deviation from Hardy–Weinberg equilibrium was tested for each locus with GenAEx 6.5. Following sequential Bonferroni correction, significant deviation from Hardy–Weinberg equilibrium was observed for five loci (Table 2). This is unsurprising for a species with delayed autonomous self-pollination (Zhou et al., 2012). ArtCir18 showed fixed heterozygote genotypes for all the screened individuals in the Maunganui Bluff and Hick’s Bay populations, but each population was fixed for different alleles.

All 16 loci amplified in the closely related species *A. bifurcatum*, and 12 of these were polymorphic (Table 3). Eight loci amplified in the more distantly related *A. candidum*, but none of the three samples screened were polymorphic at these loci.

CONCLUSIONS

We developed 16 variable microsatellite markers for *A. cirratum* using Illumina MiSeq data. Although most of the markers had a low number of alleles, many showed fixed allelic

differences between the populations examined. These markers will be useful for characterizing genetic diversity and structure in *A. cirratum* and for examining the translocation of this species.

LITERATURE CITED

- GUREVICH, A., V. SAVELIEV, N. VYAHHI, AND G. TESLER. 2013. QUAST: Quality assessment tool for genome assemblies. *Bioinformatics (Oxford, England)* 29: 1072–1075.
- HEENAN, P. B., A. D. MITCHELL, AND P. J. DE LANGE. 2004. *Arthropodium bifurcatum* (Asparagaceae), a new species from northern New Zealand. *New Zealand Journal of Botany* 42: 233–246.
- HOLM, S. 1979. A simple sequential rejective method procedure. *Scandinavian Journal of Statistics* 6: 65–70.
- LI, D., C. M. LIU, R. LUO, K. SADAKANE, AND T. W. LAM. 2015. MEGAHIT: An ultra-fast single-node solution for large and complex metagenomics assembly via succinct de Bruijn graph. *Bioinformatics (Oxford, England)* 31: 1674–1676.
- MARTINS, W. S., D. C. S. LUCAS, K. F. S. NEVES, AND D. J. BERTIOLI. 2009. WebSat—A Web software for microsatellite marker development. *Bioinformatics* 3: 282–283.
- MILLER, M. P., B. J. KNAUS, T. D. MULLINS, AND S. M. HAIG. 2013. SSR_pipeline—Computer software for the identification of microsatellite sequences from paired-end Illumina high-throughput DNA sequence data (ver. 1.1, February 2014). U.S. Geological Survey Data Series 778. U.S. Geological Survey, Reston, Virginia, USA. Website <https://pubs.er.usgs.gov/publication/ds778> [accessed 17 July 2017].
- PEAKALL, R., AND P. E. SMOUSE. 2012. GenAEx 6.5: Genetic analysis in Excel. Population genetic software for teaching and research—An update. *Bioinformatics (Oxford, England)* 28: 2537–2539.
- ROUSSET, F. 2008. GENEPOP’007: A complete re-implementation of the GENEPOP software for Windows and Linux. *Molecular Ecology Resources* 8: 103–106.
- SHEPHERD, L. D., AND T. G. B. MCLAY. 2011. Two micro-scale protocols for the isolation of DNA from polysaccharide-rich plant tissue. *Journal of Plant Research* 124: 311–314.
- SHEPHERD, L. D., P. J. DE LANGE, S. COX, P. A. MCLENACHAN, N. R. ROSKRUGE, AND P. J. LOCKHART. 2016. Evidence of a strong domestication bottleneck in the recently cultivated New Zealand endemic root crop, *Arthropodium cirratum* (Asparagaceae). *PLoS ONE* 11: e0152455.
- ZHOU, W., D.-Z. LI, AND H. WANG. 2012. Coexistence of delayed autonomous self-pollination and deceptive pollination in *Arthropodium cirratum* (Asparagaceae). *Plant Diversity and Resources* 34: 187–191.

APPENDIX 1. Location and voucher information for *Arthropodium* species used in this study.

Species	<i>n</i>	Voucher no. ^a	Location	Geographic coordinates
<i>Arthropodium cirratum</i> (G. Forst.) R. Br.	1	WELT SP103437 ^b	In cultivation, ex. Surville Cliffs, Northland, NZ	−34.3956, 173.0124
	1	WELT SP104032 ^b	Waikawa, East Cape, NZ	−37.6783, 177.7483
	1	AK 311376 ^b	Haparapara, East Cape, NZ	−37.7929, 177.6679
	1	CHR 473343 ^b	Papanui Point, Waikato, NZ	−37.8898, 174.7636
	21	AK 311414	Hick's Bay, East Cape, NZ	−37.5683, 178.2866
	22	AK 308946	Maunganui Bluff, Northland, NZ	−35.7783, 173.5703
	20	WELT SP103515	Matapouri Bay, Northland, NZ	−35.5623, 174.5094
<i>A. bifurcatum</i> Heenan, A. D. Mitch. & de Lange	1	WELT SP103440	In cultivation, ex. Hen Island, Northland, NZ	−35.8917, 174.7274
	2	WELT SP103512	In cultivation, ex. Poor Knights Islands, Northland, NZ	−35.4688, 174.7365
	1	WELT SP103511	In cultivation, ex. Surville Cliffs, Northland, NZ	−34.3956, 173.0124
	1	AK 309832	Surville Cliffs, Northland, NZ	−34.3956, 173.0124
<i>A. candidum</i> Raoul	1	WELT SP103534	Great Island, Three Kings, Northland, NZ	−34.1575, 172.1387
	2	WELT SP103527	Golden Bay, NW Nelson, NZ	−40.8873, 172.8122
	1	—	Lake Wakatipu, Otago, NZ	−40.0424, 168.63704

Note: *n* = number of sampled individuals; NZ = New Zealand.

^aVouchers are deposited in the herbaria of Auckland Museum (AK), Auckland, New Zealand; Landcare Research (CHR), Lincoln, New Zealand; or the Museum of New Zealand (WELT), Wellington, New Zealand. One representative voucher sample was collected per population.

^bSamples used for initial library construction.