

A Termite from the Late Oligocene of Northern Ethiopia

Authors: Engel, Michael S., Pan, Aaron D., and Jacobs, Bonnie F.

Source: Acta Palaeontologica Polonica, 58(2) : 331-334

Published By: Institute of Paleobiology, Polish Academy of Sciences

URL: <https://doi.org/10.4202/app.2011.0198>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



A termite from the Late Oligocene of northern Ethiopia

MICHAEL S. ENGEL, AARON D. PAN, and BONNIE F. JACOBS

Termites of the family Stolotermitidae are a relict lineage of primitive Isoptera. The fossil record of Stolotermitidae is exceptionally poor, with only two Miocene (Neogene) species documented to date. Herein, a new genus and species of Paleogene termites is described and figured from the Late Oligocene (28–27 Ma, Early Chattian) of northwestern Ethiopia (Amhara Region, Chilga Woreda). *Chilgatermes diamatensis* gen. et sp. nov., is most similar to genera of the Stolotermitidae, Archotermopsidae, and Termopsidae but can be distinguished on the basis of forewing venational details. The genus is tentatively placed in the Stolotermitidae: Porotermitinae. *Chilgatermes diamatensis* is the first fossil termite from Ethiopia and, indeed, the first from the entire African continent.

Introduction

The family Stolotermitidae comprise ten living species in Australia, sub-Saharan Africa, and the Neotropics (Emerson 1942; Krishna et al. 2013). The family has historically been included in the Termopsidae s.l., a heterogeneous lineage of primitive Isoptera widely known as the “dampwood termites”. Recently, Termopsidae has been recognized as paraphyletic and its constituents segregated into three separate families—Termopsidae for the fossil genus *Termopsis* Heer, 1849, Stolotermitidae for the former subfamilies Porotermitinae (*Porotermes* Hagen, 1858, three extant species) and Stolotermitinae (*Stolotermes* Hagen, 1858, seven extant species), and Archotermopsidae for the genera *Archotermopsis* Desneux, 1904, *Zootermopsis* Emerson, 1933, *Hodotermopsis* Holmgren, 1911, *Parotermes* Scudder, 1883, and likely also *Gyatermes* Engel and Gross, 2009 (Engel et al. 2009). This arrangement has garnered support from morphological, paleontological, behavioral, biochemical, and DNA sequence data sources (Engel et al. 2009; Ware et al. 2010; Lacey et al. 2011).

Only two fossil species were documented to date, both of *Stolotermes* and both from the Miocene (Neogene). The first species attributed to the genus was *Stolotermes amanoi* Fujiyama, 1983, described from a shed wing recovered from the Late Miocene Anadozawa Formation of Japan (Fujiyama 1983). The most recent record came from a similarly shed forewing described as *Stolotermes kupe* Kaulfuss, Harris, and Lee, 2010 from the Early Miocene (Otaian Stage) of New Zealand (Kaulfuss et al. 2010). Aside from these two specimens, no other records for the family have been documented. *Tanytermes* Engel, Grimaldi, and Krishna, 2007 and *Dharmatermes* Engel, Grimal-

di, and Krishna, 2007 from the mid-Cretaceous of Myanmar have wing venations superficially similar to Stolotermitidae and were compared with *Stolotermes* and *Porotermes* when first discovered (Engel et al. 2007b). However, in all cladistic treatments *Tanytermes* and *Dharmatermes* group elsewhere, albeit often near Stolotermitidae—e.g., related to the Kalotermitidae + Neoisoptera clade (= Icoisoptera) (Engel et al. 2009) or as progressive sisters to Neoisoptera (Ware et al. 2010) or Kalotermitidae alone (Lo and Eggleton 2010).

Herein is described an enigmatic fossil species, again based on a pair of shed wings, from the Late Oligocene (Paleogene) of northeastern Africa. The new fossil is loosely attributable to the Stolotermitidae although it is plesiomorphic in many details to the two subfamilies, Stolotermitinae and Porotermitinae, and may represent a stem-group to Stolotermitidae or Porotermitinae (less likely). While wings alone often do not provide a sufficiently rich source of character data for definitive placement of a particular fossil, this species is noteworthy as it is the first fossil record of a termite, of any age, from Africa thereby representing a significant new record for the paleontology of Isoptera.

Institutional abbreviations.—CH, Chilga collections, National Museum of Ethiopia, Addis Ababa, Ethiopia.

Geological setting

Fossiliferous sediments are located on the Ethiopian Plateau, approximately 60 km west of Gondar. The regional geology consists of massive (approximately 2000 m thick) Oligocene trap basalts with interspersed tuffs, lignites, and fluvial volcanoclastic and clastic sediments exposed along streams and gully cuts (Hoffmann et al. 1997; Kappelman et al. 2003; Jacobs et al. 2005). The study site is located in a nearly 100 m thick sedimentary section of strata that overlies a 32.4 ± 1.6 Ma whole rock K/Ar dated basalt next to the Guang River (Kappelman et al. 2003). Compression fossils, including the specimen discussed herein, were excavated from a 22 to 36 cm thick, greenish-gray to yellow-green massive mudstone layer within the section. The layer represents an over-bank (or pond) deposit derived from a weathered ash (Pan 2007). The fossil termite was found with a relatively diverse autochthonous or parautochthonous tropical moist forest paleoflora, the Guang River flora, composed of compressions of leaves and reproductive structures, logs, in situ stumps, and pollen (Jacobs et al. 2005; Pan 2007, 2010; Pan and Jacobs 2009; Pan et al. 2010). All of these fossils are dated to 27.23 ± 0.1 Ma by a $^{206}\text{Pb}/^{238}\text{U}$ analysis of zircon crystals extracted from an ash layer stratigraphically correlated with the mudstone (Pan 2010).




Fig. 1. Termite *Chilgatermes diamatensis* gen. et sp. nov. (CH 52-70), Chilga Woreda, Ethiopia, Early Chattian (Oligocene). Photomicrographs of part (A) and counterpart (B). Photomicrographs by MSE.

Systematic paleontology

Morphological terminology follows that of Engel et al. (2007a, b, 2011a), and Grimaldi et al. (2008), while the format for the description is that of Engel et al. (2007b), Engel and Gross (2009), and Engel et al. (2011b). The classification adopted herein is that of Engel et al. (2009).

Family Stolotermitidae Holmgren, 1910

Subfamily Porotermitinae? Emerson, 1942

Genus *Chilgatermes* nov.

Type species: *Chilgatermes diamatensis* gen. et sp. nov.; see below.

Etymology: A combination of Chilga, from the name of the geographic region, and *termes*, “wood-borer”, common stem for isopteran genera (gender masculine).

Diagnosis.—As for the type species; see below.

Chilgatermes diamatensis sp. nov.

Figs. 1–2.

Etymology: The specific epithet is taken from the ancient kingdom of D’mt (a.k.a. Diamat), which ruled portions of modern Eritrea and northern Ethiopia from ca. 700–400 BCE, and which encompassed in its southernmost regions the Chilga deposits.

Holotype: Alate wings, overlapping with only forewing venation discernible; CH 52-70 (a and b), part and counterpart; deposited in the Chilga collections, National Museum of Ethiopia, Addis Ababa, Ethiopia.

Type locality: Sublocality 2 (CH 52) of Guang River flora, northwestern Ethiopia (Amhara Region, Chilga Woreda, approximately 60 km west of Gondar (Jacobs et al. 2005).

Type horizon: Late Oligocene (Early Chattian, 28–27 Ma).

Diagnosis.—Alate forewing: Membrane reticulate; all veins originating inside scale; basal suture gently convex, not straight or oblique (straight and oblique in modern Stolotermitidae); radial field relatively wide, not as narrow as in extant Stolotermitidae, of approximately same width for most of wing length, widening to encompass wing apex in ex-




Fig. 2. Reconstruction of forewing venation of termite *Chilgatermes diamatensis* gen. et sp. nov. (CH 52-70) from Chilga Woreda, Ethiopia, Early Chattian (Oligocene), membrane reticulations omitted. Drawing by Ismael A. Hinojosa-Díaz. Abbreviations: CuA, anterior cubital vein; M, medial vein; R, radial vein; Rs, radial sector vein.

tre apical portion of wing; R (radial vein) simple, extending to about one-third wing length, apparently with a single, short veinlet extending to costal margin in basal third; Rs (radial sector vein) strong, with several (8), largely-simple branches extending to costal margin; M (medial vein) strong, running closer to CuA (anterior cubital vein) than to Rs in proximal third of wing, extensively developed, with numerous (7), mostly dichotomously-branching primary branches extending to posterior wing margin, apicalmost branch terminating just posterior to wing apex, basalmost branch terminating on posterior margin near wing midlength; CuA extensively developed in proximal half of wing, with numerous (at least 10), largely-simple branches.

Description.—As in the diagnosis, with the following minor additions: Total length (as preserved) from basal suture to apex 17.4 mm; maximum width 5.9 mm. Sc (subcostal vein) not evident; apex of Rs stem branching dichotomously just posterior to wing apex, apicalmost superior branch forking near its base, posteriormost fork terminating at wing apex; penultimate branch of CuA forking near posterior wing margin.

Stratigraphic and geographic range.—Type locality only.

Discussion and conclusions

Chilgatermes differs from extant Stolotermitidae, including *Poro-termes*, in the broader radial field; the much more extensively-developed medial vein, with more than seven primary branches; the gently convex basal suture; and the more densely-branched CuA. In the absence of the wing scale and structures of the body it is impossible to give a more precise phylogenetic association of the fossil. However, it is likely that its biology was similar to other stolotermitids, living in somewhat temperate regions and nesting in dead wood, often stumps and root systems but logs (i.e., not standing erect and with roots still embedded in the soil) are also prone. Nests are built entirely within the timber, with colonies established by reproductives entering gaps or cracks on surfaces exposed above the soil level. However, nests are extended downward by the workers, typically into the root systems, thereby developing a connection with the soil, necessary for the maintenance of sufficient moisture levels within the colony. The nests themselves consist of irregular, flattened chambers which either border

one another directly or are connected by narrow tunnels. Given the abundance of plant material from Chilga it is entirely possible that entire nests may be recovered from the deposits.

Hopefully continued exploration in Chilga will reveal not only more complete material of *C. diamatensis* but additional termite and other insect species. The Cenozoic record of insects in Africa is confined to a small handful of localities, with about 120 specimens recorded from the most “prolific” of these (e.g., Schlüter 2003). Given the diversity of insects today and in the past (Grimaldi and Engel 2005), this is a paltry record for such an ecologically significant and diversity dominant lineage of terrestrial animals. By the Cenozoic insects had already been around for at least 345 million years, diversified into all of their major lineages, and nearly all of these had been dominant in their respective ecosystems for eons (Grimaldi and Engel 2005). By the time of the Late Oligocene the termites had diversified and arisen in abundance, becoming one of the most significant recyclers of carbon (Engel et al. 2009), with diverse faunas documented in the preceding and following epochs of Chilga (e.g., Engel et al. 2007a, 2009, 2011a; Wappler and Engel 2006; Engel and Gross 2009; Krishna and Grimaldi 2009). The ecosystem of Chilga in the Late Oligocene was certainly ideal habitat for termites of several families. Given the usual development of large numbers of workers and soldiers, particularly in the Rhinotermitidae and Termitidae, as well as swarms of reproductives at certain times of the year, the potential for isopteran material is considerable and wings and other termite fragments should be sought actively in the region.

Acknowledgements.—The senior author is grateful to the staff of the Fort Worth Museum of Science and History for making his visit possible, and to Southern Methodist University for permitting and assisting with work in their facilities. We would like to thank the Authority for Research and Conservation of Cultural Heritage and the Ministry of Culture and Tourism, Ethiopia for permission to conduct continuing research on the Ethiopian Plateau, the National Museum of Ethiopia for their assistance and support in Addis Ababa, and the Gondar ARCC and Chilga Ministry of Culture and Sport Affairs for permission to conduct field research and logistical support. We are indebted to Ismael A. Hinojosa-Díaz (University of Kansas, Lawrence, USA) for providing the reconstruction of the forewing venation, to Kumar Krishna (Ameri-

can Museum of Natural History, New York, USA) for insightful comments on the characters and placement of *Chilgatermes*, and to Kumar Krishna and Uwe Kaulfuss (University of Otago, Dunedin, New Zealand) for critical reviews of the manuscript. This is a contribution of the Division of Entomology, University of Kansas Natural History Museum. Support for work in Ethiopia was provided by the National Science Foundation (EAR-0001259, EAR-0240251, and EAR-0617306), the National Geographic Society, and the Dallas Paleontological Society.

References

- Emerson, A.E. 1942. The relations of a relict South African termite (Isoptera, Hodotermitidae, *Stolotermes*). *American Museum Novitates* 1187: 1–12.
- Engel, M.S. and Gross, M. 2009. A giant termite from the Late Miocene of Styria, Austria (Isoptera). *Naturwissenschaften* 96: 289–295.
- Engel, M.S., Grimaldi, D.A., and Krishna, K. 2007a. A synopsis of Baltic amber termites (Isoptera). *Stuttgarter Beiträge zur Naturkunde, Serie B (Geologie und Paläontologie)* 372: 1–20.
- Engel, M.S., Grimaldi, D.A., and Krishna, K. 2007b. Primitive termites from the Early Cretaceous of Asia (Isoptera). *Stuttgarter Beiträge zur Naturkunde, Serie B (Geologie und Paläontologie)* 371: 1–32.
- Engel, M.S., Grimaldi, D.A., and Krishna, K. 2009. Termites (Isoptera): Their phylogeny, classification, and rise to ecological dominance. *American Museum Novitates* 3650: 1–27.
- Engel, M.S., Grimaldi, D.A., Nascimbene, P.C., and Singh, H. 2011a. The termites of Early Eocene Cambay amber, with the earliest record of the Termitidae (Isoptera). *ZooKeys* 148: 105–123.
- Engel, M.S., Nel, A., Azar, D., Soriano, C., Tafforeau, P., Néraudeau, D., Colin, J.-P., and Perrichot, V. 2011b. New, primitive termites (Isoptera) from Early Cretaceous ambers of France and Lebanon. *Palaeodiversity* 4: 29–39.
- Fujiyama, I. 1983. Neogene termites from northeastern districts of Japan, with references to the occurrence of fossil insects in the districts. *Memoirs of the National Science Museum, Tokyo* 16: 83–99.
- Grimaldi, D. and Engel, M.S. 2005. *Evolution of the Insects*. xv + 755 pp. Cambridge University Press, Cambridge.
- Grimaldi, D.A., Engel, M.S., and Krishna, K. 2008. The species of Isoptera (Insecta) from the Early Cretaceous Crato Formation: A revision. *American Museum Novitates* 3626: 1–30.
- Hoffman, C., Courtillot, V., Feraud, G., Rochette, P., Yirgus, G., Ketefos, E., and Pik, R. 1997. Timing of the Ethiopian flood basalt event and implications for plume birth and global change. *Nature* 389: 838–841.
- Holmgren, N. 1910. Das System der Termiten. *Zoologischer Anzeiger* 35: 284–286.
- Jacobs, B., Tabor, N., Feseha, M., Pan, A., Kappelman, J., Rasmussen, T., Sanders W., Wiemann, M., Crabaugh, J., and Garcia Massini, J.L. 2005. Oligocene terrestrial strata of northwestern Ethiopia: A preliminary report on paleoenvironments and paleontology. *Palaeontologia Electronica* 8 (1): 1–19 [25A].
- Kappelman, J., Rasmussen, D.T., Sanders, W.J., Feseha, M., Bown, T.M., Copeland, P., Crabaugh, J., Fleagle, J.G., Glantz, M., Gordon, A., Jacobs, B.F., Maga, M., Muldoon, K., Pan, A., Pyne, L., Richmond, B., Ryan, T.J., Seiffert, E.R., Sen, S., Todd, L., Wiemann, M.C., and Winkler, A. 2003. New Oligocene mammals from Ethiopia and the pattern and timing of faunal exchange between Afro-Arabia and Eurasia. *Nature* 426: 549–552.
- Kaulfuss, U., Harris, A.C., and Lee, D.E. 2010. A new fossil termite (Isoptera, Stolotermitidae, *Stolotermes*) from the Early Miocene of Otago, New Zealand. *Acta Geologica Sinica* 84: 705–709.
- Krishna, K. and Grimaldi, D. 2009. Diverse Rhinotermitidae and Termitidae (Isoptera) in Dominican amber. *American Museum Novitates* 3640: 1–48.
- Krishna, K., Grimaldi, D.A., Krishna, V., and Engel, M.S. 2013. Treatise on the Isoptera of the world. *Bulletin of the American Museum of Natural History* 377: 1–2704.
- Lacey, M.J., Sémon, E., Krasulová, J., Sillam-Dussès, D., Robert, A., Cornette, R., Hoskovec, M., Žáček, P., Valterová, I., and Bordereau, C. 2011. Chemical communication in termites: *syn*-4,6-dimethylundecan-1-ol as trail following pheromone, *syn*-4,6-dimethylundecanal and (*5E*)-2,6,10-trimethylundeca-5,9-dienal as the respective male and female sex pheromones in *Hodotermopsis sjoestedti* (Isoptera, Archotermopsidae). *Journal of Insect Physiology* 57: 1585–1591.
- Lo, N. and Eggleton, P. 2010. Termite phylogenetics and co-cladogenesis with symbionts. In: D.E. Bignell, Y. Roisin, and N. Lo (eds.), *Biology of Termites: A Modern Synthesis*, 27–50. Springer Verlag, Berlin.
- Pan, A.D. 2007. *The Late Oligocene (28–27 Ma) Guang River Flora from the Northwestern Plateau of Ethiopia*. 235 pp.. Unpublished Doctoral Dissertation, Southern Methodist University, Dallas.
- Pan, A.D. 2010. Rutaceae leaf fossils from the Late Oligocene (27.23 Ma) Guang River flora of northwestern Ethiopia. *Review of Palaeobotany and Palynology* 159: 188–194.
- Pan, A.D. and Jacobs, B.F. 2009. The earliest record of the genus *Cola* (Malvaceae *sensu lato*: Sterculioideae) from the Late Oligocene (28–27 Ma) of Ethiopia and leaf characteristics within the genus. *Plant Systematics and Evolution* 283: 247–262.
- Pan, A.D., Jacobs, B.F., and Herendeen, P.S. 2010. Detarieae *sensu lato* (Fabaceae) from the Late Oligocene (27.23 Ma) Guang River flora of north-western Ethiopia. *Botanical Journal of the Linnean Society* 163: 44–54.
- Schlüter, T. 2003. Fossil insects in Gondwana-localities and palaeodiversity trends. *Acta zoologica cracoviensia* 46 (supplement): 345–371.
- Wappler, T. and Engel, M.S. 2006. A new record of *Mastotermes* from the Eocene of Germany (Isoptera: Mastotermitidae). *Journal of Paleontology* 80: 380–385.
- Ware, J.L., Grimaldi, D.A., and Engel, M.S. 2010. The effects of fossil placement and calibration on divergence times and rates: An example from the termites (Insecta: Isoptera). *Arthropod Structure and Development* 39: 204–219.

Michael S. Engel [msengel@ku.edu], Division of Entomology (Paleoentomology), Natural History Museum, and Department of Ecology & Evolutionary Biology, University of Kansas, 1501 Crestline Drive – Suite 140, Lawrence, Kansas 66045, USA; and Division of Invertebrate Zoology, American Museum of Natural History, Central Park West at 79th, New York, New York 10024-5192, USA;

Aaron D. Pan [apan@dhdc.org], Don Harrington Discovery Center, 1200 Streit Drive, Amarillo, Texas 79106-1759, USA, and Botanical Research Institute of Texas, 1700 N. University Drive, Fort Worth, Texas 76107-3400, USA;

Bonnie F. Jacobs [bjacobs@smu.edu], Roy M. Huffington Department of Earth Sciences, Southern Methodist University, PO Box 750395, Dallas, Texas 75275-0395, USA.

Received 16 December 2012, accepted 12 April 2012, available online 20 April 2012.

Copyright © 2013 M.S. Engel et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.