

## **Non-Exotic Invasion of Great-Tailed Grackles *Quiscalus mexicanus* in a Tropical Dry Forest Reserve**

Authors: MacGregor-Fors, Ian, Vázquez, Leopoldo, Vega-Rivera, Jorge H., and Schondube, Jorge E.

Source: *Ardea*, 97(3) : 367-369

Published By: Netherlands Ornithologists' Union

URL: <https://doi.org/10.5253/078.097.0312>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## Non-exotic invasion of Great-tailed Grackles *Quiscalus mexicanus* in a tropical dry forest reserve

Ian MacGregor-Fors<sup>1</sup>, Leopoldo Vázquez<sup>2</sup>, Jorge H. Vega-Rivera<sup>3</sup> & Jorge E. Schondube<sup>1,\*</sup>

MacGregor-Fors I., Vázquez L., Vega-Rivera J.H. & Schondube J.E. 2009. Non-exotic invasion of Great-tailed Grackles *Quiscalus mexicanus* in a tropical dry forest reserve. *Ardea* 97(3): 367–369.

We report two recordings of Great-tailed Grackles *Quiscalus mexicanus* in the Chamela Biosphere Reserve, which is an important conservation area for pristine tropical dry forests in West Mexico. These recordings are important for three main reasons: (1) they reveal the recent encroachment of the reserve by human-disturbed areas; (2) the recorded grackles fought with West Mexican Chachalacas *Ortalis poliocephala*, and won, underlining their possible negative effects towards other native bird species; and (3) if Great-tailed Grackles successfully invade the conservation area, it could turn into an unmanageable scenario with possible negative consequences within an important area for bird conservation in the dry tropics.

Key words: dispersal, Great-tailed Grackle, *Quiscalus mexicanus*, tropical dry forest, Chamela, Mexico

<sup>1</sup>Centro de Investigaciones en Ecosistemas, Universidad Nacional Autónoma de México, Campus Morelia, Apartado Postal 27-3, Morelia, Michoacán, 58190, México; <sup>2</sup>Instituto de Biología, Universidad Nacional Autónoma de México, Ciudad Universitaria Delegación Coyoacán, México, D.F. 04510, México; <sup>3</sup>Estación de Biología Chamela, Instituto de Biología, Universidad Nacional Autónoma de México, Apartado Postal 21, San Patricio, Jalisco, 22630, México;

\*corresponding author (chon@oikos.unam.mx)



Invasive species play a significant role in global change, representing a threat to biodiversity (Vitousek *et al.* 1997). Although a great proportion of invasive bird species are exotic (e.g. House Sparrow *Passer domesticus* in the Americas), some native bird species are also able to invade new areas by dispersing through matrixes of human-disturbed habitats (e.g. Canyon Towhee *Pipilo fuscus*, Great-tailed Grackle *Quiscalus mexicanus*). Successful native invasive bird species could have dramatic effects on local avifaunas, a process tended to be ignored because the invaders are native to the country where their invasions occur.

The Great-tailed Grackle is a large-sized icterid presently distributed from southern Canada to Peru (Stiles & Skutch 1989, Howell & Webb 1995, Sibley 2001). This grackle has broadened its distribution in the last 150 years, expanding from southern Mexico, to

central-northern Mexico, most of the USA, and part of Canada. This expansion is closely related to landscape matrixes generated by agricultural and urban activities (Wehtje 2003). In Mexico, the Great-tailed Grackle is highly successful in urban areas, filling a niche similar to what Rock Pigeons *Columba livia* and European Starlings *Sturnus vulgaris* occupy in Europe (Christensen 2000).

### Results and discussion

Although the Great-tailed Grackle is well distributed along human-disturbed landscapes in Mexico, it does not enter natural habitats. We were surprised to record this grackle within the tropical dry forest of the Chamela-Cuixmala Biosphere Reserve. Our first sighting-recording was in February 2008, when a male Great-tailed Grackle was recorded within the build-up area of

the Chamela biological station (19°29'52"N, 105°02'39"W, 90 m a.s.l.). Our second observation was conducted in early May 2009, when we sight-recorded four males and one female Great-tailed Grackles in the edge between the forest and the biological station, competing for food-leftovers (vegetables and fruit) besides the station's kitchen, where other wildlife species feed daily (e.g. West Mexican Chachalaca *Ortalis poliocephala*, San Blas Jay *Cyanocorax sanblasianus*, Yellow-winged Cacique *Cacicus melanicterus*, Back-streaked Oriole *Icterus pustulatus*, Collared Peccary *Tayassu tajacu*, White-nosed Coati *Nasua narica*).

The nearest areas from where the grackles could have dispersed to the biological station are a trash dumping site and a crop-field area, both located ~2.7 km away. Although several urban settlements, where the grackle is present and abundant, are quite close to the biological station (Careyes: 6.7 km; San Mateo: 9.4 km; Juan Gil: 11.4; Zapata/Villa: 14.9 km), this species has never been recorded in the station grounds since the reserve was established in 1971 (Hutto *et al.* 1985, Arizmendi *et al.* 1990, Ornelas *et al.* 1993). Furthermore, during intensive mist netting and point-counts from 1999 to 2001, and from 2004 to 2008, we did not capture or observe grackles within the Biosphere Reserve (Vega-Rivera, Schondube & MacGregor-Fors, unpubl. data).

We consider these recordings important for three reasons. First, finding the Great-tailed Grackle in the build-up area of the Chamela Biological Station suggests that human disturbance encroaches upon the reserve (Vega-Rivera *et al.* 2004). This could explain why the grackles were able to find a small patch of suitable habitat completely surrounded by pristine forest. Indeed, an analysis of regional landcover change using remote sensing shows a decrease in forest cover and an increase in fragmentation around the biosphere reserve in the last seven years (Sánchez-Azofeifa *et al.* 2009). Secondly, we recorded two male grackles fighting successfully for food with West Mexican Chachalacas, a species which is almost three times heavier. This pinpoints their possible effect towards local neotropical bird species. Finally, although we do not know the impact of Great-tailed Grackles on the bird community of the preserve, if this species settles in the area we expect a negative effect on the local avifauna. This is worrisome because the Chamela-Cuixmala biosphere reserve is a crucial area for bird conservation in the dry tropics (Arizmendi & Márquez-Valdelamar 2000, Arriaga *et al.* 2000, Stattersfield *et al.* 1998).

While we are starting to understand the effects that invasive bird species have on native avifaunas

(Blackburn *et al.* 2009, MacGregor Fors *et al.* 2009), we know even less of the consequences that the invasion of new habitats by a native non-local species can have on bird communities. Usually this problem does not get public attention because it represents a conflict of native local vs. native non-local species. Due to the negative effects that human-commensal native non-local invasive bird species could have on native local communities, we suggest that Great-tailed Grackles should be removed from the area to avoid an unmanageable invasion scenario. However, given the present context of fragmentation and habitat modification by human activities that favour a few generalist human-commensal bird species, we can wonder whether the removal of Great-tailed Grackles from the Chamela Biological Station is going to be helpful. In other words, will we be able to safeguard specific native local bird communities in the light of the changing world? And at what cost?

We thank the Estación de Biología Chamela (IBUNAM). We also thank two anonymous reviewers for their useful comments. Field observations were carried out as part of the "Curso teórico-práctico de ecología poblacional de aves" funded by the Posgrado en Ciencias Biológicas of the Universidad Nacional Autónoma de México. IM-F (175447) and LV (23046), being part of the Posgrado en Ciencias Biológicas of the Universidad Nacional Autónoma de México, received a Ph.D. and Master's scholarship from CONACYT, respectively.

## References

- Arizmendi M.C., Berlanga H., Márquez-Valdelamar L., Navarrijo L. & Ornelas, F. 1990. Avifauna de la región de Chamela, Jalisco. Instituto de Biología-UNAM, México.
- Arriaga L., Espinoza J.M., Aguilar C., Martínez E., Gómez L. & Loa E. 2000. Regiones terrestres prioritarias de México. CONABIO, México.
- Blackburn T.M., Lockwood J.L. & Cassey O.P. 2009. Avian invasions. The ecology and evolution of exotic birds. Oxford University Press, Oxford.
- Christensen A.F. 2000. The fifteenth- and twentieth-century colonization of the Basin of Mexico by the Great-tailed Grackle (*Quiscalus mexicanus*). Global. Ecol. Biogeogr. 9: 415–420
- Howell S.N.G. & Webb S. 1995. A guide to the birds of Mexico and Northern Central America. Oxford University Press, Oxford.
- Hutto R.L., Hendrics P. & Pletschet S. 1985. Un censo invernal de las aves de la Estación de Biología Chamela, Jalisco, Mexico. Ann. Inst. Biol. UNAM 56: 945–954.
- MacGregor-Fors I., Morales-Pérez L., Quesada J. & Schondube J.E. 2009. Relationship between the presence of House Sparrows (*Passer domesticus*) and Neotropical bird community structure and diversity. Biol. Invasions 11: doi:10.1007/s10530-009-9432-5.

- Ornelas J.F., Arizmendi M.C., Marquez-Valdelamar L., Navarrijo M.L. & Berlanga H.A. 1993. Variability profiles for line transect bird censuses in a tropical dry forest in Mexico. *Condor* 95: 422–441.
- Sánchez-Azofeifa G.A., Quesada M., Cuevas-Reyes P., Castillo A. & Sánchez-Montoya G. 2009. Land cover and conservation in the area of influence of the Chamela-Cuixmala Biosphere Reserve, Mexico. *Forest. Ecol. Manag.* 258: 907–912.
- Sibley D.A. 2001. *The Sibley guide to birds*. Alfred A. Knopf, New York.
- Stattersfield A.J., Crosby M.J., Long A.J. & Wege D.C. 1998. *Endemic bird areas of the World: priorities for biodiversity conservation*. Birdlife Conservation Series No. 7. Birdlife International, Cambridge.
- Stiles F. & Skutch A. 1989. *A guide to the birds of Costa Rica*. Comstock Publishing Associates, Ithaca.
- Vega-Rivera J.H., Alvarado F., Valdivia T. & Rappole J. 2004. First report of cowbird parasitism on the Black-capped Gnatcatcher in western Mexico. *Southwest. Nat.* 49: 277–278.
- Vitousek P.M., D'Antonio C.M., Loope L.L., Rejmánek M. & Westbrooks R. 1997. Introduced species: a significant component of human-caused global change. *New. Zeal. J. Ecol.* 21: 1–16
- Wehtje W. 2003. The range expansion of the great-tailed grackle (*Quiscalus mexicanus* Gmelin) in North America since 1880. *J. Biogeogr.* 30: 1593–1607.

### Samenvatting

In 2008 en 2009 werden in het droog- tropische bosreservaat Chamela-Cuixmala in het westen van Mexico twee waarnemingen van Langstaarttroepialen *Quiscalus mexicanus* gedaan. Deze soort komt van het zuiden van Canada tot in Peru voor. Het was een schok deze commensaal van mensen in een ongerept tropisch bos aan te treffen. Dat te meer daar een conflict om voedsel met de driemaal zo zware West-Mexicaanse Chachalaca *Ortalis poliocephala* door de Langstaarttroepiaal werd gewonnen. Bij de problematiek rond invasieve soorten wordt er gewoonlijk van uitgegaan dat de kolonisten exoten zijn. In dit geval, echter, gaat het om een inheemse soort die een kwetsbaar en ongerept habitat binnendringt, daartoe in staat gesteld door negatieve ontwikkelingen buiten het park (habitatversnippering en verstedelijking). De vraag is: wat te doen? Verwijdering van de binnendringers zal op den duur, bij voortschrijdende menselijke invloed rond het reservaat, steeds moeilijker en kostbaarder worden. Een niet ingrijpen zou echter grote consequenties kunnen hebben voor de kwetsbare lokale avifauna. (RGB)

*Corresponding editor: Rob G. Bijlsma*

*Received 31 July 2009; accepted 22 September 2009*

# ARDEA

TIJDSCHRIFT DER NEDERLANDSE ORNITHOLOGISCHE UNIE (NOU)

*ARDEA* is the scientific journal of the Netherlands Ornithologists' Union (NOU), published bi-annually in spring and autumn. Next to the regular issues, special issues are produced frequently. The NOU was founded in 1901 as a non-profit ornithological society, composed of persons interested in field ornithology, ecology and biology of birds. All members of the NOU receive *ARDEA* and *LIMOSA* and are invited to attend scientific meetings held two or three times per year.

## NETHERLANDS ORNITHOLOGISTS' UNION (NOU)

**Chairman** – J.M. Tinbergen, Animal Ecology Group, University of Groningen, P.O. Box 14, 9750 AA Haren, The Netherlands

**Secretary** – P.J. van den Hout, Royal Netherlands Institute for Sea Research (NIOZ), P.O. Box 59, 1790 AB Den Burg, Texel, The Netherlands (hout@nioz.nl)

**Treasurer** – E.C. Smith, Ir. van Stuivenbergweg 4, 6644 AB Ewijk, The Netherlands (ekko.diny@planet.nl)

**Further board members** – E. Boerma, G.J. Gerritsen, J. Komdeur, J. Ouwehand, G.L. Ouweneel, J.J. de Vries

**Membership NOU** – The 2010 membership fee for persons with a postal address in The Netherlands is €42 (or €25 for persons <25 years old at the end of the year). Family members (€9 per year) do not receive journals. Foreign membership amounts to €54 (Europe), or €65 (rest of the world). Payments to ING-bank account 285522 in the name of Nederlandse Ornithologische Unie, Sloetmarke 41, 8016 CJ Zwolle, The Netherlands (BIC: INGBNL2A and IBAN: NL36INGB0000285522). Payment by creditcard is possible.

Correspondence concerning membership, payment alternatives and change of address should be sent to: Erwin de Visser, Sloetmarke 41, 8016 CJ Zwolle, The Netherlands (nou ledenadmin@gmail.com).

**Research grants** – The NOU supports ornithological research and scientific publications through its Huib Kluijver Fund and the 'Stichting Vogeltekstation'. Applications for grants can be addressed to the NOU Secretary. Donations to either fund are welcomed by the NOU treasurer.

**Internet** – [www.nou.nu](http://www.nou.nu)

## ARDEA

**Editors of ARDEA** – Rob G. Bijlsma, Wapse (Editor in chief); Christiaan Both, Groningen; Niels J. Dingemanse, Groningen; Dik Heg, Bern; Ken Kraaijeveld, Leiden; Jouke Prop, Ezinge (Technical editor); Julia Stahl, Oldenburg; B. Irene Tieleman, Groningen; Yvonne I. Verkuil, Groningen

**Dissertation reviews** – Popko Wiersma, Groningen

**Editorial address** – Jouke Prop, Allersmaweg 56, 9891 TD Ezinge, The Netherlands (ardea.nou@planet.nl)

**Graphics** – Dick Visser, Haren

**Artwork** – Jos Zwarts, Bunnik

**Internet** – [www.ARDEAJournal.nl](http://www.ARDEAJournal.nl)

**Subscription ARDEA** – Separate subscription to *ARDEA* is possible. The 2010 subscription rates are €36 (The Netherlands), €42 (Europe), and €50 (rest of the world). Institutional subscription rates are €53, €69, and €78, respectively). Papers that were published more than five years ago can be freely downloaded as pdf by anyone through *ARDEA*'s website. More recent papers are available only to members of the NOU and subscribers of *ARDEA*-online. Receiving a hard-copy with additional access to *ARDEA*-online costs €55 (The Netherlands and Europe), €70 (rest of the world), or €110 (institutions). Subscriptions to *ARDEA*-online (without receiving a hard copy) cost €40 (individuals worldwide), or €85 (institutions). Payments to ING-bank account 125347, in the name of Nederlandse Ornithologische Unie, Ir. van Stuivenbergweg 4, 6644 AB Ewijk, The Netherlands (BIC: INGBNL2A and IBAN: NL16INGB0000125347). Correspondence concerning subscription, change of address, and orders for back volumes to: Ekko Smith, Ir. van Stuivenbergweg 4, 6644 AB Ewijk, The Netherlands (ekko.diny@planet.nl).

**Exchange of publications** – All periodicals sent in exchange for *ARDEA* should be addressed to: Library of the Netherlands Ornithologists' Union (NOU), c/o Tineke Prins, Institute of Systematics and Population Biology, Zoological Museum, P.O. Box 94766, 1090 GT Amsterdam, The Netherlands.

**Books for review** – should be addressed to: *ARDEA* Secretariat, c/o J. Prop, Animal Ecology Group, Biological Centre, P.O. Box 14, 9750 AA Haren, The Netherlands. After review, the books will be deposited in the NOU Library in Haren.

**NOU Library (journals)** – Mauritskade 57, Amsterdam, Mo–Fr 10:00–16:00 (to check in advance by telephone + 31 20 525 6614).

**NOU Library (books)** – Library Biological Centre, Kerklaan 30, Haren (G.), Mo–Thu 09:00–17:00 (to check at [www.rug.nl/bibliotheek/locaties/bibfwn/index](http://www.rug.nl/bibliotheek/locaties/bibfwn/index)).

© Nederlandse Ornithologische Unie (NOU), 2009

Layout by Dick Visser, Haren, The Netherlands

Printed by Van Denderen, Groningen, The Netherlands, October 2009

Downloaded From: <https://bioone.org/journals/Ardea> on 27 May 2024

Terms of Use: <https://bioone.org/terms-of-use>