

Diet of Wintering Long-Eared Owl *Asio otus* in Zabol, Southeastern Iran

Authors: Khaleghizadeh, Abolghasem, Arbabi, Tayebbeh, Noori, Gholamreza, Javidkar, Mohammad, and Shahriari, Alireza

Source: *Ardea*, 97(4) : 631-633

Published By: Netherlands Ornithologists' Union

URL: <https://doi.org/10.5253/078.097.0432>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Diet of wintering Long-eared Owl *Asio otus* in Zabol, southeastern Iran

Abolghasem Khaleghizadeh^{1,*}, Tayebah Arbabi², Gholamreza Noori³,
Mohammad Javidkar² & Alireza Shahriari²



Khaleghizadeh A., Arbabi T., Noori G., Javidkar M. & Shahriari A. 2009. Diet of wintering Long-eared Owl *Asio otus* in Zabol, southeastern Iran. In: Johnson D.H., Van Nieuwenhuysen D. & Duncan J.R. (eds) Proc. Fourth World Owl Conf. Oct–Nov 2007, Groningen, The Netherlands. *Ardea* 97(4): 631–633.

We examined 250 pellets of Long-eared Owls *Asio otus* in 2006 and 2007 from winter roost sites in southeastern Iran. In contrast to the diet of wintering owls (e.g. mainly small mammals) reported elsewhere in the world, the diet of wintering Long-eared Owls at Zabol was predominantly larger rodents (c. 150 g). Specifically, big-size rodents, including the Indian Gerbil *Tatera indica*, and a Bandikoot Rat *Nesokia indica* made up 72.9% of the total biomass in the diet of the owls at Zabol. In addition to small mammal and bird species found in Long-eared Owl pellets from other regions, we found *Meriones* spp. and *Gerbillus* spp. (both are types of Gerbils) that, thus far, had not been reported in the diet of this owl.

Key words: Long-eared Owl, *Asio otus*, Iran, pellet, diet, rodentia, *Tatera indica*

¹Ornithology Laboratory, Agricultural Zoology Res. Dep., Iranian Research Institute of Plant Protection, Tehran, Iran; ²Department of Environment, University of Zabol, Zabol, Iran; ³Faculty of Geography, University of Seistan & Baluchestan, Zahedan, Iran;

*corresponding author (akhaleghizadeh@gmail.com)

INTRODUCTION

There is extensive literature on the diet of Long-eared Owls from the Western Palearctic (Birrer 2009), but no data on this species from Iran. Thus far in Iran, diet analysis on owls has been limited to the Little Owl *Athene noctua* (Obuch & Kristin 2004) and Barn Owl *Tyto alba* (Khaleghizadeh & Rokni 2008). In Iran, the Long-eared Owl *Asio otus* is resident in the regions of the Alburz and Zagross Mountains and is a winter migrant in other parts of the country (Mansoori 2001, D.A. Scott pers. comm.). It is a winter migrant to Zabol, southeastern Iran (Noori *et al.* 2007). We examined pellets of the Long-eared Owl, collected from the outskirts of the town of Zabol, southeastern Iran, to offer the first report on the diet of this owl for the region.

METHODS

Study area

Long-eared Owl pellets were collected at the Emam Ali (Jazinak) Forest Park (51 ha; 30°53'N, 61°33'E), c. 15 km southeast of Zabol, southeastern Iran. This area is characterized by a flat and dry region with a seasonal river. Twenty-three plant species belonging to 12 families such as Poaceae, Asteraceae, Chenopodiaceae were identified in the patchy cover of the Forest Park. *Atriplex lentiformis* was the most frequently encountered plant, while *Populus euphratica* had the lowest frequency. This vegetation is typical of the surrounding Sistan Plain. The Zabol area (elevation 489 m a.s.l.) is found within the Sistan Plain in Sistan and Baluchestan Province, and borders Afghanistan and Pakistan. The

Sistan area is a dry region of about 15 200 km² with average annual temperature of 22.7°C; the warmest month is July and the coldest January. The annual precipitation is 59.6 mm with the maximum precipitation observed in March (14.6 mm). Under such conditions, life depends on the inland delta of the Helmand River originating from the mountains of the southern Hindu Kush (Afghanistan) and the associated wetland, Hamoon. The water cover in the Hamoon is extensive but shallow (less than 3 m at the highest water levels). Strong winds (>9 m/s) in this very dry region makes evaporation more than 3 m annually. Thus, this water source is very sensitive to climatic fluctuations and modifications of water inflow by humans. Agricultural lands encompass 25 861 ha, with wheat the dominant crop (about 50% of the area); barley, alfalfa, watermelon, melon and grapes are also grown. Other green areas consist of rangelands with about 474 595 ha, natural forests with about 11 000 ha and afforested areas with 1633 ha. Zabol's population is estimated about 130 642 people (Noori *et al.* 2007b).

Data collection

Long-eared Owl pellets were collected at a winter roost site in *Tamarix stricta* in the Emam Ali Forest Park. Owls were seen on several occasions and about 40 owls were counted at this roost on 28 December 2006 (Noori *et al.* 2007a). Pellets were collected under owl roosts located in *Tamarix* vegetation. About 250 pellets were collected on 26 January 2006, 12 February and 18 March 2007. Intact pellets were air dried and weighed, and their length and diameter recorded. Each pellet was soaked in 95% alcohol and teased apart using a pair of forceps and a needle. Bone remains and

skulls of Rodentia in each pellet were placed in separate containers. The mammal remains were identified following Etemad (1978). Frequency (numbers of each prey) and biomass of the prey were calculated based on Ziaie (2008).

RESULTS AND DISCUSSION

The colour of pellets was dark grey to black, average dry weight was 2.5 g. On average, pellets were 37.5 mm in length and 21.0 mm in diameter. From 250 pellets, 258 prey items were found (average 1.03 prey items per pellet). Rodentia were present in 99% of the pellets. Of the 258 prey identified, the composition was as follows (Table 1): rodents (not identified to species) 47.7%, *Gerbillus* spp. 26.7%, and Indian Gerbil *Tatera indica* 17.8%. Other identified mammals included the Bandikoot Rat *Nesokia indica*, jerbil/jird *Meriones* spp., mouse *Mus* spp. and shrews Soricidae, each with < 1% of the total number of prey taken. Fifteen (5.8% by frequency) of the prey found in the pellets were birds. Importantly, big-size rodents (c. 150 g) including *Tatera indica* and *Nesokia indica* made up 72.9% of the total biomass in the diet of the owls. The respective percent biomass for all prey is shown in Table 1.

Overall, the prey taken by Long-eared Owls wintering in our study area was consistent with that taken by wintering owls in many other locations, i.e. majority of prey were mammals, with a few birds also taken. However, the majority of prey here were big-size rodents, specifically *Tatera indica*. Neither *Meriones* spp. nor *Gerbillus* spp. (types of jirds/gerbils) had previously been reported in the diet of this owl (Cramp

Table 1. Diet of the Long-eared Owl at Zabol, Sistan and Baluchestan Province, Iran. In total, 258 prey items were identified.

Prey species		Frequency	Frequency (%)	Average weight (g)	Biomass (%)
<i>Tatera indica</i>	Indian Gerbil	45	17.4	163.0	33.1
<i>Nesokia indica</i>	Bandikoot Rat	1	0.4	155.0	0.7
<i>Meriones</i> spp.	jird	1	0.4	150.0	0.7
<i>Gerbillus</i> spp.	gerbil	69	26.7	25.0	7.8
<i>Mus</i> spp.	mouse	2	0.8	21.0	0.2
Rodentia	small-size rodents	20	7.8	30.0	2.7
	medium-size rodents	46	17.8	70.0	14.5
	big-size rodents	57	22.1	150.0	38.5
Soricidae	shrew	2	0.8	12.5	0.1
<i>Passer domesticus</i>	House Sparrow	8	7.8	22.0	0.8
Passeriformes	birds	7	17.8	30.0	1.0

1985). While there were insufficient remains in the pellets to make clear identifications, the high frequency of big-size rodents in the present study was presumably due to the higher population of *T. indica* compared to other rodents in the Zabol area (M. Javidkar, unpubl. data). Most of the unidentified big-size rodents were probably *T. indica*, and the unidentified small-sized *Gerbillus* spp. were likely dominated by *Gerbillus nanus*.

In a synthesis on the Long-eared Owl, Cramp (1985) reported an average of 2.3 prey items per pellet (range 0–8; $n = 11\ 390$ pellets). Similarly, in Lithuania, Balčiauskienė et al. (2006) found 1.75 prey items per pellet. In Diyarbakir, Turkey, Long-eared Owl pellets collected from July 2000 to June 2001 contained generally 1–2 prey items per pellet (Seçkin & Coşkun 2005). In the present study, we found an average of 1.0 prey item in each pellet. Compared to other regions, the number of prey per pellet in the present study was less – simply due to the larger size of prey taken in the Zabol region.

REFERENCES

- Balčiauskienė L., Jovaišs A., Naruševičius V., Petraška A. & Skuja S. 2006. Diet of Tawny Owl (*Strix aluco*) and Long-eared Owl (*Asio otus*) in Lithuania as found from pellets. *Acta Zool. Lituanica* 16: 37–45.
- Birrer S. 2009. Synthesis of 312 studies on the diet of the Long-eared Owl *Asio otus*. *Ardea* 97: 615–624.
- Cramp S. (ed.) 1985. Handbook of the Birds of Europe, the Middle East and North Africa. The Birds of the Western Palearctic. Oxford University Press, Oxford.
- Etemad E. 1978. Mammals of Iran. Iran Department of the Environment, Tehran.
- Khaleghizadeh A. & Rokni M.-R. 2008. Pellet content of the Barn Owl *Tyto alba* in Haft-Tappeh, Khuzestan Province, Iran. Proceedings of the 15th national & 3rd International Conference of Biology, 19–21 August 2008, University of Tehran.
- Mansoori J. 2008. A field guide to the birds of Iran. Farzaneh Publishing Co., Tehran. (In Persian)
- Noori G., Arbabi T. & Khamari M. 2007a. Report on the occurrence of Long-eared Owl *Asio otus* from Zabol, Iran's border with Afghanistan–Pakistan. *Raptors Conservation* 10: 63–64. (In Russian)
- Noori G., Arbabi T. & Noori S. 2007b. Hamoon Wetland, the life of Sistan. Sepehr publication. (In Persian)
- Obuch J. & Kristin A. 2004. Prey composition of the little owl *Athene noctua* in an arid zone (Egypt, Syria, Iran). *Folia Zool.* 53: 65–79.
- Seçkin S. & Coşkun Y. 2005. Small mammals in the diet of the Long-eared Owl, *Asio otus*, from Diyarbakir, Turkey. *Zool. Middle East* 35: 102–103.
- Ziaie H. 2008. A field guide to the mammals of Iran. Iran Wildlife Center.

SAMENVATTING

In 2006–07 werden 250 braakballen van Ransuilen *Asio otus* onderzocht die afkomstig waren van verschillende winterroestplaatsen in Zabol, ZO-Iran. De uilen bleken voornamelijk knaagdieren van zo'n 150 gram te eten. Dit is in tegenstelling tot het gangbare wintervoedsel elders, dat voornamelijk kleinere zoogdieren omvat. De Indische Naaktzoolrenmuis *Tatera indica* en de Kortstaartmolrat *Nesokia indica* maakten 72,9% van de totale biomassa van het voedsel in Zabol uit. Er werden naast eerder in het voedsel van Ransuilen aangetroffen soorten ook renmuizen van de geslachten *Meriones* en *Gerbillus* aangetroffen, knaagdieren die nog niet eerder in het voedsel van Ransuilen werden gevonden.

ARDEA

TIJDSCHRIFT DER NEDERLANDSE ORNITHOLOGISCHE UNIE (NOU)

ARDEA is the scientific journal of the Netherlands Ornithologists' Union (NOU), published bi-annually in spring and autumn. Next to the regular issues, special issues are produced frequently. The NOU was founded in 1901 as a non-profit ornithological society, composed of persons interested in field ornithology, ecology and biology of birds. All members of the NOU receive ARDEA and LIMOSA and are invited to attend scientific meetings held two or three times per year.

N E T H E R L A N D S O R N I T H O L O G I S T S ' U N I O N (N O U)

Chairman – J.M. Tinbergen, Animal Ecology Group, University of Groningen, P.O. Box 14, 9750 AA Haren, The Netherlands

Secretary – P.J. van den Hout, Royal Netherlands Institute for Sea Research (NIOZ), P.O. Box 59, 1790 AB Den Burg, Texel, The Netherlands (hout@nioz.nl)

Treasurer – E.C. Smith, Ir. van Stuivenbergweg 4, 6644 AB Ewijk, The Netherlands (ekko.diny@planet.nl)

Further board members – E. Boerma, G.J. Gerritsen, J. Komdeur, J. Ouweland, G.L. Ouweneel, J.J. de Vries

Membership NOU – The 2010 membership fee for persons with a postal address in The Netherlands is €42 (or €25 for persons <25 years old at the end of the year). Family members (€9 per year) do not receive journals. Foreign membership amounts to €54 (Europe), or €65 (rest of the world). Payments to ING-bank account 285522 in the name of Nederlandse Ornithologische Unie, Sloetmarke 41, 8016 CJ Zwolle, The Netherlands (BIC: INGBNL2A and IBAN: NL36INGB0000285522). Payment by creditcard is possible. Correspondence concerning membership, payment alternatives and change of address should be sent to: Erwin de Visser, Sloetmarke 41, 8016 CJ Zwolle, The Netherlands (nou ledenadmin@gmail.com).

Research grants – The NOU supports ornithological research and scientific publications through its Huib Kluijver Fund and the 'Stichting Vogeltrekstation'. Applications for grants can be addressed to the NOU Secretary. Donations to either fund are welcomed by the NOU treasurer.

Internet – www.nou.nu

A R D E A

Editors of ARDEA – Rob G. Bijlsma, Wapse (Editor in chief); Christiaan Both, Groningen; Niels J. Dingemanse, Groningen; Dik Heg, Bern; Ken Kraaijeveld, Leiden; Kees van Oers, Heteren; Jouke Prop, Ezinge (Technical editor); Julia Stahl, Oldenburg; B. Irene Tieleman, Groningen; Yvonne I. Verkuil, Groningen

Dissertation reviews – Popko Wiersma, Groningen

Editorial address – Jouke Prop, Allersmaweg 56, 9891 TD Ezinge, The Netherlands (ardea.nou@planet.nl)

Internet – www.ardeajournal.nl. The website offers free downloads of all papers published in Ardea and forerunners from 1904 onwards. The most recent publications are available only to subscribers to Ardea and members of the NOU.

Subscription ARDEA – Separate subscription to ARDEA is possible. The 2010 subscription rates are €36 (The Netherlands), €42 (Europe), and €50 (rest of the world). Institutional subscription rates are €53, €69, and €78, respectively). Papers that were published more than five years ago can be freely downloaded as pdf by anyone through ARDEA's website. More recent papers are available only to members of the NOU and subscribers of ARDEA-online. Receiving a hard-copy with additional access to ARDEA-online costs €55 (The Netherlands and Europe), €70 (rest of the world), or €110 (institutions). Subscriptions to ARDEA-online (without receiving a hard copy) cost €40 (individuals worldwide), or €85 (institutions). Payments to ING-bank account 125347, in the name of Nederlandse Ornithologische Unie, Ir. van Stuivenbergweg 4, 6644 AB Ewijk, The Netherlands (BIC: INGBNL2A and IBAN: NL16INGB0000125347). Correspondence concerning subscription, change of address, and orders for back volumes to: Ekko Smith, Ir. van Stuivenbergweg 4, 6644 AB Ewijk, The Netherlands (ekko.diny@planet.nl).

World Owl Conference Special

Editors – David H. Johnson, Dries Van Nieuwenhuyse and James R. Duncan, in cooperation with Jouke Prop and Rob G. Bijlsma

Technical editor – Jouke Prop

Dutch summaries – Arie L. Spaans, Dries Van Nieuwenhuyse, Jouke Prop, Rob G. Bijlsma, or authors

Graphs and layout – Dick Visser

Drawings – Jos Zwarts

Cover photos – Serge Sorbi

front – Snowy Owl

back – Snowy Owl, Great Grey Owl and young Tengmalm's Owl

Production – Hein Bloem, Johan de Jong and Arnold van den Burg

© Nederlandse Ornithologische Unie (NOU), 2009

Printed by Van Denderen, Groningen, The Netherlands, December 2009