

## **Apharinodes sinensis sp. n. (Coleoptera: Staphylinidae: Pselaphinae) from China, and discovery of male wing dimorphism in Hybocephalini**

Authors: Yin, Zi-Wei, and Jiang, Ri-Xin

Source: *Revue suisse de Zoologie*, 124(1) : 9-14

Published By: Muséum d'histoire naturelle, Genève

URL: <https://doi.org/10.5281/zenodo.322660>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## ***Apharinodes sinensis* sp. n. (Coleoptera: Staphylinidae: Pselaphinae) from China, and discovery of male wing dimorphism in Hybocephalini**

Zi-Wei Yin<sup>1</sup> & Ri-Xin Jiang

Department of Biology, Shanghai Normal University, 100 Guilin Road, Shanghai, 200234, P. R. China. E-mail: [pselaphinae@gmail.com](mailto:pselaphinae@gmail.com)

**Abstract:** *Apharinodes sinensis* sp. n. is described from Yunnan, and represents the first named species of the tribe Hybocephalini Raffray from China. Male wing dimorphism is for the first time reported for a hybocephaline. An identification key to the four known *Apharinodes* species is given.

**Keywords:** Hybocephalini, *Apharinodes*, taxonomy, new species, male wing dimorphism, Asia.

### **INTRODUCTION**

The ant-loving beetle tribe Hybocephalini Raffray (Pselaphinae: Pselaphitae) comprises some 70 species grouped in 10 genera (Newton & Chandler, 1989; Besuchet 2008). Within this tribe most members are found from the tropical forested areas of Africa and Oriental region (Raffray, 1908; Newton & Chandler, 1989), with two species (of *Apharinodes* Raffray and *Stipesa* Sharp, respectively) occurring in eastern Palaearctic region (Sharp, 1874; Nomura, 1989), and one species (of *Stipesa*) from Australia (Chandler, 2001). Hybocephalines can be distinguished from those of all other tribes of the Pselaphitae by the generally compact and convex habitus, relatively robust antennae, small maxillary palpi, and more noticeably, squamous setae that may partially to entirely cover the body surface (Raffray, 1890a; Chandler, 2001). The number of the antennomeres by which the clubs are formed, relative length(s) of the abdominal segments, presence/absence and number of the abdominal paratergites, and presence/absence of elytral discal striae are frequently used to determine taxa at generic level (Raffray, 1908; Jeannel 1949a, b, 1959).

The genus *Apharinodes* Raffray is a small group currently including three species confined to East Asia (Raffray, 1911; Newton & Chandler, 1989; Nomura, 1989). Raffray (1890a) established *Apharinodes* with *A. squamosa* Raffray from Singapore (type species subsequently designated by Raffray, 1890b), and later described (Raffray, 1895) a second species *A. miranda* Raffray from the same country. Most recently, Nomura (1989)

described a third species *A. papageno* Nomura from Okinawa Island, Japan, from where additional collecting data for the same species was later added (Sugaya, 2003). The actual diversity of *Apharinodes* is expected to be much higher than currently documented. At least eight additional species from China, Malaysia and Thailand were listed in several reports on the local pselaphine fauna of these areas, though none of these has been formally named (Nomura 2000; Nomura & Idris, 2005; Nomura & Mohamed, 2008; Nomura *et al.*, 2008, 2010, 2013). *Apharinodes* has the antennal clubs composed of only the enlarged terminal antennomere (XI), the first two visible abdominal tergites (morphologically IV and V) are subequal in length, and almost the entire body surface is covered by thickened squamous setae. Such a combination of characters provides reliable means with regard to the identification of *Apharinodes* within the Hybocephalini.

Based on the third edition of the International Code of Zoological Nomenclature (ICZN, 1985), Nomura (1989) emended the gender of *Apharinodes* to masculine, and changed the suffixes of the two known species accordingly. But according to the latest (fourth) edition of the ICZN (1999), the name *Apharinodes* was originally combined with adjectival species-group names that are feminine (i.e. *squamosa* and *miranda*), thus the gender of *Apharinodes* should be remained as feminine (Article 30.1.4.4.).

Wing dimorphism is long- and well-known in insects. However, for the megadiverse polyphagan family Staphylinidae, sex-related wing dimorphism was only recently documented and discussed in detail. Thayer

(1992) firstly reported in *Omalium flavidum* Hamilton the presence of fully-winged and flightless males and flightless females, which belongs to a less common pattern of wing dimorphism. In Pselaphinae (formerly Pselaphidae, see Newton & Thayer, 1995), such pattern has been so far recorded for Batrisini (Nomura, 1991), Trichonychini (Nomura, 1988; Yin & Li, 2016), Bythinini (Kurbatov, 1994; Löbl *et al.*, 1998), and Tyrini (Nomura, 1997). In these groups, the males with normally developed hind wings tend to have larger eyes, longer elytra and metaventricle, and stronger elytral humeri. Recently, two male specimens of an undescribed *Apharinodes* were collected from southwestern China (Yunnan) exhibiting male wing dimorphism in a group of the Pselaphinae (Hybocephalini) in which such a phenomenon has never been recorded.

## MATERIAL AND METHODS

Material studied in this paper is housed in the Insect Collection of Shanghai Normal University (SNUC).

Dissected parts were preserved in Euparal on plastic slides that were placed on the same pin with the specimen. The habitus image was taken using a Canon 5D Mark III camera in conjunction with a Canon MP-E 65 mm f/2.8 1-5X Macro Lens and a Canon MT-24EX Macro Twin Lite Flash. Images of the morphological details were made using a Canon G9 camera mounted on an Olympus CX31 microscope. Zerene Stacker (version 1.04) was used for image stacking. All images were modified and grouped in Adobe Photoshop CS5 Extended.

The label data of the material are quoted verbatim, with additional notes included in parentheses. The following abbreviations are applied: AL—length of the dorsally visible part of the abdomen (posterior to elytra) along the midline; AnL—length of the antenna; AnCL—length of the antennal club; AW—maximum width of the abdomen; EL—length of the elytra along the suture; EW—maximum width of the elytra; HL—length of the head from the anterior clypeal margin to the occipital constriction; HW—width of the head across eyes; PL—length of the pronotum along the midline; PW—maximal width of the pronotum. The length of the body (BL) is a combination of HL + PL + EL + AL. Terminology for morphological structures follows Chandler (2001) with the exception that the term ‘ventrite’ replaces ‘sternite’, except prosternum.

## TAXONOMY

### *Apharinodes sinensis*, new species

Figs 1-2

**Type material:** **Holotype:** macropterous male: ‘China: Yunnan Province, Lincang City, Yun Hsien (云县), Manwan Town (漫湾镇), Caozidicun (草子地村), 24°40′27″N, 100°23′09″E, under stone, 1757 m,

9.iv.2016, Zi-Chun Xiong leg.’ (SNUC). **Paratype:** 1 apterous male, same label data as the holotype (SNUC).

**Diagnosis:** BL 2.28-2.39 mm; postocular region longer than eye (Fig. 2A); anterolateral branches (Fig. 2A-1) of squamous scales on pronotum distinct but shorter than basolateral branches (Fig. 2A-2), basolateral branches and mediobasal branch (Fig. 2A-3) separated anteriorly; male antennomere XI with large, shallow semicircular excavation on mesal surface; aedeagus (Fig. 2G-H) weakly sclerotized, median lobe with large basal capsule, endophallus with three sclerites, parameres thick and short.

**Description:** *Macropterous male* (Fig. 1A). Body length 2.39 mm. Body light reddish brown, mouthparts and tarsi lighter in color (the light color of the two males available are possibly due to immature adult status); most part of dorsal surface covered with thick squamous setae.

Head (Fig. 2A) slightly longer than wide, HL 0.53 mm, HW 0.46 mm, small vertexal and frontal fovea largely covered by setae; eyes large, each composed of about 32 facets; postocular margins (Fig. 2B) longer than eyes, roundly narrowed toward head base; antennae (Fig. 2C) robust, scapes as long as combined length of next two antennomeres, antennomeres II transversely quadrate, III to X strongly transverse, successively wider, XI largest, mesal surface with large, shallow, semicircular excavation, AnL 0.95 mm, AnCL 0.33 mm, AnCL/AnL = 0.35; maxillary palpi small and short, three-segmented, with distinct conical palpal cone; gular area flat, densely covered with setae basolaterally.

Pronotum (Fig. 2A) slightly wider than long, PL 0.48 mm, PW 0.52 mm, lateral margins nearly straight at basal half, narrowed from middle toward apex; anterolateral branches of squamous scales distinct but short, extending less mesally than basolateral branches, mediobasal branch and basolateral branches clearly separated at anterior ends. Whole surface of prosternum (Fig. 2B) covered with dense setae.

Elytra convex, wider than long, EL 0.73 mm, EW 0.88 mm, posterior margin with band of dense setae; two basal foveae covered by squamous scales; with complete discal and sutural striae. Metathoracic wings fully developed. Mesoventrite densely covered with small squamous setae; metaventricle densely covered with setae at middle, areas posterior to mesocoxae with two elongate projections that extend to metaventral posterior margin.

Abdomen subglobose, wider than long, AL 0.65 mm, AW 0.86 mm; tergite IV (first visible tergite) as long as tergite V, with pair of lateral sulci, tergites VI and VII slightly shorter than V, VIII (Fig. 2D) circular viewed posteriorly; sternite IV (second visible sternite) to VII successively shorter, VIII (Fig. 2E) shallowly emarginate at middle, IX (Fig. 2F) well-sclerotized at apex, membranous at posterior half.




Fig. 1. Dorsal habitus of *Apharinodes sinensis*. (A) Macropterous male. (B) Apterous male. Scale bars: 0.5 mm.

**Key to species of *Apharinodes* Raffray**

- 1 Body less than 2 mm; anterolateral branches of squamous scales on pronotum extending more mesal (closer to midline) than basolateral branches. (Singapore) ..... *A. squamosa* Raffray
- Body no less than 2 mm; anterolateral branches of squamous scales on pronotum shorter than basolateral branches (Figs 1, 3).....2
- 2 Male antennomere XI with large, shallow, quadrate excavation; pronotal mediobasal and basolateral branches of squamous scales fusing anteriorly (Fig. 3B). (Singapore)..... *A. miranda* Raffray
- Male antennomere XI with deep, rounded excavation (Figs 1, 2C, 3A); pronotal mediobasal and basolateral branches of squamous scales separated anteriorly (Figs 1, 2A, 3A) .....3
- 3 Excavation of male antennomere XI deeper (Fig. 3A); postocular margins slightly shorter than eye length (Nomura, 1989: fig. 1); pronotal anterolateral branches of squamous scales indistinct (Fig. 3A). (Japan: Okinawa).....  
..... *A. papageno* Nomura
- Excavation of male antennomere XI shallower (Figs 1, 2C); postocular margins longer than eye length (Fig. 2A); pronotal anterolateral branches of squamous scales distinct (Figs 1, 2A). (China: Yunnan) ..... *A. sinensis*, **sp. n.**




Fig. 2. Details of male *Apharinodes sinensis*. (A) Head dorsum and pronotum (numbers indicate branches of squamous scales: 1. anterolateral branch; 2. basolateral branch; 3. mediobasal branch). (B) Head venter and prosternum. (C) Right antenna. (D) Tergite VIII. (E) Sternite VIII. (F) Sternite IX. (G) Aedeagus, in ventral view. (H) Same, in lateral view. Scale bars: A-C = 0.3 mm; D, E, G, H = 0.1 mm; F = 0.05 mm.




Fig. 3. Dorsal habitus of *Apharinodes* species. (A) *A. papageno*, male paratype. (B) *A. miranda*, male. Scale bars: 0.5 mm.

Aedeagus (Fig. 2G-H) weakly sclerotized, length 0.38 mm; median lobe broad, with large basal capsule, broadly emarginate along apical margin; parameres short and thick, each with three thick apical setae; endophallus composed of three sclerites, left sclerite longest, curved toward right, middle sclerite shortest, membranous, right sclerite broad, weakly sclerotized, narrowed at apex.

*Apterous male* (Fig. 1B). Similar to macropterous male in general habitat, except eyes smaller, each composed of about 11 facets; elytra and metaventrite shorter; and metathoracic wings absent. Measurements: BL 2.28 mm, HL 0.50 mm, HW 0.44 mm, PL 0.47 mm, PW 0.52 mm, EL 0.65 mm, EW 0.87 mm, AL 0.66 mm, AW 0.85 mm,

AnL 0.97 mm, AnCL 0.34 mm, AnCL/AnL = 0.35, length of aedeagus 0.35 mm.

*Female*. Unknown.

**Distribution:** Southwestern China: Yunnan.

**Etymology:** The specific epithet indicating the country where the new species was collected.

#### ACKNOWLEDGMENTS

We thank Zi-Chun Xiong (Lincang, China) for collecting and offering the specimens used in this paper. Habitus pictures of *A. papageno* and *A. miranda* were taken

during the first author's visit to National Museum of Nature and Science, Tokyo in 2012. The curator of Coleoptera, Shūhei Nomura provided various help and supports. The present study was supported by grants of the National Natural Science Foundation of China (No. 31501874) and the Technology Commission of Shanghai Municipality (No.15YF1408700).

## REFERENCES

- Besuchet C. 2008. Synonymies et combinaisons nouvelles, revalidations et description de taxa nouveaux de Pselaphinae (Coleoptera: Staphylinidae). *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 81: 61-82.
- Chandler D.S. 2001. Biology, morphology and systematics of the ant-like litter beetles of Australia (Coleoptera: Staphylinidae: Pselaphinae). *Memoirs on Entomology, International* 15: vii + 560 pp.
- ICZN 1985. International Code of Zoological Nomenclature. Third Edition. *The International Trust for Zoological Nomenclature, London*. xx + 338 pp.
- ICZN 1999. International Code of Zoological Nomenclature. Fourth Edition. *The International Trust for Zoological Nomenclature, London*. xxvii + 306 pp.
- Jeannel R. 1949a. Les Pselaphides de l'Afrique Orientale (Coleoptera). *Mémoires du Muséum National d'Histoire Naturelle (N.S.)* 29: 1-226.
- Jeannel R. 1949b. Faune du Congo Belge et du Ruanda Urundi, II: Pselaphidae. *Annales du Musée du Congo Belge, Tervuren (Série 8°: Sciences Zoologiques)* 2: 1-275.
- Jeannel R. 1959. Révision des Pselaphides de l'Afrique intertropicale. *Annales du Musée Royal du Congo Belge, Tervuren (Série 8°: Sciences Zoologiques)* 75: 1-742.
- Kurbatov S.A. 1994. Les *Bryaxis* de l'Extrême-Est de la Russie (Coleoptera, Pselaphidae). *Russian Entomological Journal* 3: 39-47.
- Löbl I., Kurbatov S.A., Nomura S. 1998. On the Japanese species of *Bryaxis* (Coleoptera: Staphylinidae: Pselaphinae), with notes on allied genera and on endoskeletal polymorphy. *Species Diversity* 3: 219-269.
- Newton A.F., Chandler D.S. 1989. World catalog of the genera of Pselaphidae (Coleoptera). *Fieldiana: Zoology, (N.S.)* 53: iv + 1-93.
- Newton A.F., Thayer M.K. 1995. Protopselaphinae new subfamily for *Protopselaphus* new genus from Malaysia, with a phylogenetic analysis and review of the Omaliine Group of Staphylinidae including Pselaphidae (Coleoptera) (pp. 219-320). In: Pakaluk J. & Ślipiński S.A. (eds). Biology, phylogeny, and classification of Coleoptera: Papers celebrating the 80th birthday of Roy A. Crowson. *Muzeum i Instytut Zoologii PAN, Warszawa*. x + 1092 pp. in 2 vols.
- Nomura S. 1988. A revision of the subtribe Acetaliina (Coleoptera, Pselaphidae) from Japan, Part I. *Kontyû* 56: 251-260.
- Nomura S. 1989. Description of a new species of *Apharinodes* (Coleoptera, Pselaphidae) from Okinawa Island, Japan. *Japanese Journal of Entomology* 57: 278-282.
- Nomura S. 1991. Systematic study on the genus *Batrisoplisus* and its allied genera from Japan (Coleoptera, Pselaphidae). *Esakia* 30: 1-462.
- Nomura S. 1997. Flight of pselaphine beetles - morphological review on flight mechanism of tiny beetles [in Japanese, English title only]. *Insekutaryûmu: Ikita Konchû no Zasshi [=Insectarium]* 34: 58-66.
- Nomura S. 2000. A list of the pselaphine and protopselaphine species (Coleoptera, Staphylinidae) collected from Yunnan, southwest China in 1992-1998 (pp. 197-238). In: Aoki J.I., Yin W.-Y., Imadaté G. (eds). Taxonomical studies on the soil fauna of Yunnan Province in Southwest China. *Tokai University Press*.
- Nomura S., Idris A.G. 2005. Faunistic notes on the pselaphine species of the supertribes Goniaceritae, Pselaphitae and Clavigeritae from Malaysia and Singapore (Coleoptera: Staphylinidae: Pselaphinae). *Serangga* 10: 1-36.
- Nomura S., Mohamed M. 2008. A faunistic review of the pselaphine species (Insecta, Coleoptera, Staphylinidae) known from Borneo. *Memoirs of the National Museum of Nature and Science* 45: 13-40.
- Nomura S., Sakchoowong W., Ogata K., Chanpaisaeng J. 2008. Lists of pselaphine and protopselaphine species (Coleoptera, Staphylinidae) collected from Doi Inthanon and Khao Yai National Parks. *Report on Insect Inventory Project in Tropical Asia (TAIIV)* 2008: 265-294.
- Nomura S., Sakchoowong W., Chanpaisaeng J. 2010. A list of the pselaphine species (Insecta, Coleoptera, Staphylinidae) collected from the Kaeng Krachan National Park, West Thailand. *Bulletin of the National Museum of Nature and Science (A: Zoology)* 36: 7-25.
- Nomura S., Sakchoowong W., Maruyama M. 2013. Further study on the Pselaphine fauna (Insecta, Coleoptera, Staphylinidae) of the Kaeng Krachan National Park, West Thailand in 2010-2012. *Bulletin of the National Museum of Nature and Science (A: Zoology)* 39: 73-92.
- Raffray A. 1890a. Etude sur les Pselaphides. V. Tableaux synoptiques.- Notes et synonymie. *Revue d'Entomologie* 9: 81-172.
- Raffray A. 1890b. Etude sur les Pselaphides. VI. Diagnoses des espèces nouvelles sur lesquelles sont fondés des genres nouveaux. *Revue d'Entomologie* 9: 193-219, pls 2-3.
- Raffray A. 1895. Révision des Pselaphides des Iles de Singapour et de Penang (suite et fin). *Revue d'Entomologie* 14: 21-82, pl. 2.
- Raffray A. 1908. Coleoptera. Fam. Pselaphidae (pp. 1-487, 9 pls). In: Wytzman P. (ed.). Genera Insectorum. Fasc. 64. *V. Verteneuil & L. Desmet, Bruxelles*.
- Raffray A. 1911. Pselaphidae (pp. 1-222). In: Schenkling S. (ed.). Coleopterorum Catalogus, Pars 27. *W. Junk, Berlin*.
- Sharp D. 1874. The Pselaphidae and Scydmaenidae of Japan. *Transactions of the Entomological Society of London* 1874: 105-130.
- Sugaya H. 2003. Notes on *Apharinodes papageno* (Coleoptera, Staphylinidae, Pselaphinae) in Okinawa-jima, the Ryukyus, Japan. *Elytra* 31: 125-126.
- Thayer M.K. 1992. Discovery of sexual wing dimorphism in Staphylinidae (Coleoptera): "*Omalium*" *flavidum*, and a discussion of wing dimorphism in insects. *Journal of the New York Entomological Society* 100: 540-573.
- Yin Z.-W., Li L.-Z. 2016. A new species of *Acetalius* Sharp from eastern China (Coleoptera, Staphylinidae, Pselaphinae). *ZooKeys* 592: 93-101.