

SEASONAL VARIATION OF THE UPPER DIGESTIVE TRACT YEAST FLORA OF FERAL PIGEONS

Authors: KOCAN, R. M., and HASENCLEVER, H. F.

Source: Journal of Wildlife Diseases, 10(3) : 263-266

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-10.3.263>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

SEASONAL VARIATION OF THE UPPER DIGESTIVE TRACT YEAST FLORA OF FERAL PIGEONS

R. M. KOCAN, U.S. Bureau of Sport Fisheries and Wildlife, Patuxent Wildlife Research Center, Laurel, Maryland 20810, U.S.A.

H. F. HASENCLEVER, Laboratory of Microbiology, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, Maryland 20014, U.S.A.

Abstract: Feral pigeons were sampled over a 16-month period to determine whether their normal yeast flora varied according to season. *Candida albicans* and *Saccharomyces telluris* occurred during the entire sampling period, with *C. albicans* reaching its highest levels between August and January and *S. telluris* peaking from March through May. *Candida krusei* was present for 10 months but exhibited no predictable variation in density. *Candida tropicalis*, *C. guilliermondii* and *Geotrichum* were isolated on several occasions while *C. lusitaniae*, *C. pseudotropicalis* and *Torulopsis glabrata* were each isolated once. The high levels of infection and frequency of occurrence of some yeast species make the feral pigeon highly suspect as a carrier and disseminator of potentially pathogenic yeast.

INTRODUCTION

Pigeons have been recognized as potential public health hazards for some time, particularly in the case of ornithosis and equine encephalitis.^{1,2,6} They are also known to carry pathogenic yeasts,^{4,7} but no studies have shown to what extent they play a role as reservoirs for these organisms.

Studies by Kocan and Hasenclever⁴ on the normal mycotic flora of several species of columbids revealed that samples collected over a period of time contained different ratios of some species, while others appeared and disappeared with no apparent regularity. To determine whether this might be a cyclical phenomenon, 441 feral pigeons (*Columba livia*) were trapped over a 16-month period (excluding November and February due to poor trapping conditions) at the National Zoological Park in Washington, D.C. The male-female and adult-juvenile ratios were about even throughout the sampling period, probably the result of an active pest control program at the zoo. The number examined per month ranged between 19 and 51 (mean = 30).

MATERIALS AND METHODS

Primary yeast isolations were made by culturing throat swabs in 5 ml Sabouraud's liquid medium containing 5000 units each of penicillin and streptomycin and incubated at 37C for 48 hours. The resultant growth was transferred to Sabouraud's agar plates from which colonies were picked for identification. Identifications were made using the morphological and physiological characteristics described for these species.⁵ No attempts were made to identify other associated microbiological agents, although pox lesions were seen in about 2% of the young birds. The pigeons trapped in August were held in large flypens, fed commercial pigeon chow, and sampled along with newly trapped birds for two additional months in order to determine whether the observed changes of their yeast flora occurred as a result of endogenous changes in the birds or from changes in the local habitat or food source. The majority of the birds appeared to be in good health at the time of capture. Observations at the Zoo showed that they fed on a variety of food thrown them by visitors, but they




FIGURE 1. Frequency of occurrence of three most commonly encountered oral yeasts of feral pigeons over a 16-month period. A—*C. albicans*; T—*S. telluris*; K—*C. krusei*. Total sampled = 441; mean per month = 30 pigeons (range 19-51).

were primarily found congregated at feed troughs which contained pelletized food for the hoofed stock.

RESULTS

Candida albicans and *Saccharomyces telluris* were present for the entire sampling period while *C. krusei* appeared during 10 of the 16 months. *Candida albicans* was most prevalent from August through January while *S. telluris* was most prevalent from March through May. Both species were present in over 60% of the birds during their peaks and ranged between 20 and 60% for most of the time between peaks. *Candida krusei* exhibited no particular cyclical variation during the sampling period even though it fluctuated between 0 and 95% (Fig. 1).

Candida tropicalis occurred in 3% of the birds in January and March; *C. guilliermondii* appeared in 3 to 28% of the birds in January, May, June and September; and *Geotrichum* appeared in 4 to 28% of the birds in March, June, July, September and October. *Candida lusitanae* and *C. pseudotropicalis* appeared in June in 14 and 3%, respectively, and *Torulopsis glabrata* appeared in July in 24% of the birds.

The group of 20 pigeons which was sampled for 3 consecutive months (Aug.-Oct.) showed *C. albicans* and *C. krusei* to remain essentially the same in both captive and newly trapped pigeons while *S. telluris* steadily decreased in the captive flock (Table 1).

In order to determine whether diurnal yeast shedding might not be responsible for the observed differences in monthly isolation rates, a series of samples was taken from eight pigeons every 4 hours for 4 days. There was no indication of any change in the oral flora of these birds, indicating that the time of day samples were taken did not influence the observed incidence of infection.

DISCUSSION

The regularity of occurrence of *C. albicans*, *C. krusei* and *S. telluris* would indicate that they were a part of the normal flora of pigeons. Closer examination, however, shows that *S. telluris* gradually disappears from a population of *Columba livia* when held in captivity. The cause for *S. telluris*' continued presence under natural conditions is unknown, but appears to be related to factors in the environment which were not duplicated for the captive pigeons. A different serotype of this yeast appears in other columbid species and in particularly high numbers in white-crowned pigeons (*C. leucocephala*).³ In this case the organisms did not disappear when the birds were confined, even though their normal fruit and seed diet was changed to mixed grain.

The similarity in occurrence of *C. albicans* and *C. krusei* in captive and wild birds indicated that they fluctuate as a result of endogenous changes in the bird, rather than with external environmental factors.

TABLE 1. Comparison of yeast isolation rates for wild and captive pigeons for the same 3-month period.

| Yeast | | Aug. | Sept. | Oct. |
|--------------------|---------|------|-------|------|
| <i>C. albicans</i> | wild | 75% | 88% | 73% |
| | captive | 75% | 65% | 63% |
| <i>C. krusei</i> | wild | 60% | 0% | 4% |
| | captive | 60% | 0% | 6% |
| <i>S. telluris</i> | wild | 55% | 32% | 62% |
| | captive | 55% | 24% | 6% |

The causes for seasonal fluctuations in these three species of yeast is not understood. Since the overall incidence of infection in the pigeons does not drop very low at any one time, it is doubtful that their role as carriers or disseminators is significantly affected during any season—if such a role exists.

The six species which sporadically appeared were transient in their association with the pigeons, and were probably acquired from the local environment. This does not necessarily eliminate the pigeon as a potential disseminator of these species when they are infected.

A monthly comparison of temperature and precipitation throughout the sampling period showed a mean January temperature of 3.5C and a mean July tempera-

ture of 25.1C. These are both within 0.83C of the normal means. At no time during the study did the mean monthly temperature fall below freezing, and only trace amounts of snow fell during the sampling period. Precipitation ranged between 2.76 cm and 32.43 cm with a yearly mean of 11.6 cm. There was no correlation between incidence of yeast and climatological highs and lows.

Since environmental factors of some type appear to affect the presence of various yeasts, it could be assumed that pigeons collected in different geographical locations would have different yeast flora. Any evaluation of the pigeon's role as a carrier of these organisms would depend on the local environment as well as the time of year.

LITERATURE CITED

1. BURKHART, R. L. and L. A. PAGE. 1971. Chlamydiosis. In *Infectious and Parasitic Diseases of Wild Birds*. J. W. Davis, R. C. Anderson, L. Karstad and D. O. Trainer, (eds.) Iowa State Univ. Press, Ames, Iowa.
2. FOTHERGILL, L. D. and J. H. DINGLE. 1938. A fatal disease of pigeons caused by the virus of the eastern variety of equine encephalomyelitis. *Science* 88: 549-550.
3. HASENCLEVER, H. F. and R. M. KOCAN. 1972. Serotypes in *Saccharomyces telluris*: Their relation to source of isolation. *Inf. and Imm.* 7: 610-612.
4. KOCAN, R. M. and H. F. HASENCLEVER. 1972. Normal yeast flora of the upper digestive tract of some wild columbids. *J. Wildl. Dis.* 8: 365-368.
5. LODDER, J. 1970. *The Yeasts: A Taxonomic Study*. North Holland Pub. Co., Amsterdam and London.
6. McDIARMID, A. 1960. *Diseases of Free-living Wild Animals*. Academic Press, New York.
7. MONGA, D. P. 1972. Prevalence of pathogenic fungi in wild birds. *Indian J. Med. Res.* 60: 517-519.

Received for publication 3 October 1973