



An Infection by *Vibrio alginolyticus* in an Atlantic Bottlenose Dolphin Housed in an Open Ocean Pen

Authors: Schroeder, J. P., Wallace, J. G., Cates, M. B., Greco, S. B., and Moore, P. W. B.

Source: Journal of Wildlife Diseases, 21(4) : 437-438

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-21.4.437>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

antigens also exists, but that has not been a significant problem with the use of this test in domestic poultry.

This study was funded by the Environmental Institute of Oklahoma State University under the direction of Howard Gerald. The authors wish to thank Mr.

Gerald, the Cooperative Wildlife Research Unit of Oklahoma State University under the direction of Dr. Paul Vohs for the use of their equipment, the personnel of Washita National Wildlife Refuge for their assistance, and Dr. J. E. Williams for providing the *Salmonella* group antigens.

Journal of Wildlife Diseases, 21(4), 1985, pp. 437-438

An Infection by *Vibrio alginolyticus* in an Atlantic Bottlenose Dolphin Housed in an Open Ocean Pen

J. P. Schroeder,¹ J. G. Wallace,² M. B. Cates,¹ S. B. Greco,³ and P. W. B. Moore,¹ ¹Naval Ocean Systems Center, Hawaii Laboratory, Kailua, Hawaii 96734, USA; ²U.S. Navy Environmental and Preventive Medicine Unit No. 6, Pearl Harbor, Hawaii 96860, USA; and ³University of Hawaii, Department of Microbiology, Honolulu, Hawaii 96822, USA

An adult male Atlantic bottlenose dolphin (*Tursiops truncatus*) had a history of recurring skin problems. The dolphin weighed 159 kg and was kept in Kaneohe Bay, Hawaii, in an open ocean floating pen, 6 m square, 3.5 m deep, enclosed by 16-cm-square wire mesh. The lesions were first noted in 1975 and were always similar in size, shape and location: 2.5- to 5.0-cm-diameter oval ulcers at the anterior insertion of the pectoral fins. In 1983, the lesions appeared the most severe since 1975. The dolphin's skin was ulcerated to the musculature in the cranial insertion area of both pectoral fins and, in addition, to the subcutaneous tissue on the tip of the rostrum and on the leading edge and lateral aspects of the dorsal fin. The dermis in the region of the pectoral lesions contained a mild neutrophil infiltrate with some neutrophil migration into the overlying stratum spinosum; such characteristics are consistent with chronic, active dermatitis.

Although the dolphin had been treated under several different therapeutic plans previously, with varying degrees of success, lesions always recurred within 4 to 6 mo.

In May 1983, during a 26-day period of no treatment, the lesions became progressively larger and more inflamed than previously seen. Before cultures were taken, the lesion and surrounding area were aseptically prepared using Povidine Iodine surgical scrub, rinsed with alcohol, and allowed to dry for 60 sec. Single swab culturettes (Precision Culture C.A.T.S.® with Modified Amies Medium, Precision Dynamics Corporation, 3031 Thornton Avenue, Burbank, California 91504, USA) were used to take cultures from the lesions. Samples were transported to the laboratory on ice and were plated within 1 hr of collection.

Each sample was plated on *Salmonella-Shigella* agar, blood agar, M-Endo agar LES, Membrane Filtration Agar for recovery of *Vibrio parahaemolyticus* (Watkins et al., 1976, *Appl. Environ. Microbiol.* 32: 679-684), mannitol salt agar and

Received for publication 2 October 1984.

TCBS agar. Each culture was incubated at 37 C and observed at 24, 48 and 72 hr. Growing colonies were biochemically typed using a computer-assisted API-20E system with 2% marine salts diluent (MacDonell et al., 1982, Appl. Environ. Microbiol. 44: 423-427), and were gram-stained and tested for oxidase and catalase. The isolated organisms were then compared to known strains source of *V. alginolyticus* control cultures (American Culture Collection Strain #17750) to confirm identification.

Vibrio alginolyticus was the only pathogenic organism isolated on any of the media initially. TCBS agar was found to be the optimum medium for the recovery of the *Vibrio* species. The second and a third series of cultures, taken 6 and 11 wk into treatment, produced identical results. These cultures were again confirmed using the same two test systems and with the appropriate controls.

No bacterial growth was detected on cultures taken from the affected areas during week 13, or during three post-treatment cultural attempts. No bacterial growth occurred in the blood cultures.

The dolphin's hematologic values changed little during the active infection. Except for a blood sample at week 13 of treatment showing a 10% reduction in hematocrit and a 35% reduction in lymphocytes (from this animal's normal), the dolphin's blood counts and blood enzyme and chemistry values were within acceptable limits throughout the treatment.

Minimal antibiotic inhibitory concentrations were determined using the Sceptor® Gram Negative/MIC panel (Bethesda Biological Laboratory, 2, Cockeysville, Maryland 21030, USA). The *Vibrio* organisms were found to be sensitive to trimethoprim/sulfamethoxazole, tetracycline, gentamycin, kanamycin and chloramphenicol. Resistance was shown against carbenicillin and ampicillin.

Based on antimicrobial susceptibility

results, a therapy was initiated using 4.8 g (30 mg/kg) of Tribissen® (trimethoprim/sulfadiazine, Burroughs Wellcome Co., Kansas City, Missouri 64108, USA) given orally once daily. Antibiotic therapy continued until all cultures were negative and the wounds were healed. The total treatment period was 13 wk, and treatment was discontinued after culture results were negative. Recrudescence has not occurred for 18 mo.

We have identified *V. alginolyticus*, *V. parahaemolyticus*, *V. fluvialis*, *V. vulnificus*, *V. cholera* (non-O-antigen) and several unclassified *Vibrio* species as normal inhabitants of the sea water in which our dolphins live (unpubl. data). Some of these *Vibrios* are also routinely isolated from blowholes, pharynx and feces of the dolphins at the Hawaii facility.

Several species of *Vibrio* are known to have caused epidemics in humans in both estuarine and nearshore marine environments (Blake et al., 1980, N. Engl. J. Med. 300: 1-5). Small odontocetes are free-ranging and occur in most inshore areas of the world. Species of *Vibrio* were also identified as normal inhabitants of newly captured small cetaceans by Solangi and Dukes (1983, Nat. Mar. Fish. Serv. Final Report #NA82-GA-C-00023,300, Washington, D.C., 177 pp.), and of cetaceans maintained in captivity over long periods of time. The potential for human acquisition of infections of *Vibrio* from dolphins exists. Species of *Vibrio* may be both primary and/or opportunistic pathogens in dolphins, therefore, people who maintain and deal with dolphins or come in contact with stranded dolphins must remain constantly aware of the zoonotic potential of these organisms.

We wish to express our special appreciation to the following people for their invaluable assistance during this project: Karl Keller, J. R. Cates, G. Purdy, E. Woods, R. Lazarte, D. Reed, and R. Fujioka.