

## **Frenkelia sp. from the Brain of a Porcupine (*Erethizon dorsatum*) from Alberta, Canada**

Authors: Kennedy, Murray J., and Frelier, Paul F.

Source: Journal of Wildlife Diseases, 22(1) : 112-114

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-22.1.112>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

without cardiac alterations is a common finding in naturally infected birds (Gray, 1958, op. cit.). Other frequently encountered lesions include peritonitis and airsacculitis. In acute cases, the necrotic foci tend to be less marked (Gray, 1958, op. cit.). Histologic lesions in the liver of the wild turkey were consistent with those described by Csontos et al. (1955, Acta Vet. Hung. 5: 261–274), as reported by Gray

(1958, op. cit.). They found foci of necrosis without a marked inflammatory reaction in both liver and spleen. In this case, since *L. monocytogenes*, a known pathogen, was isolated from the affected tissue in pure culture, it seems reasonable to ascribe the lesions encountered as well as the wild turkey's death to *L. monocytogenes*.

*Journal of Wildlife Diseases*, 22(1), 1986, pp. 112–114  
© Wildlife Disease Association 1986

### ***Frenkelia* sp. from the Brain of a Porcupine (*Erethizon dorsatum*) from Alberta, Canada**

**Murray J. Kennedy**, Alberta Agriculture, Pathology Branch, 6909 116 Street, Edmonton, Alberta T6H 4P2, Canada; and **Paul F. Frelier**, Alberta Agriculture, Veterinary Services Branch, Airdrie, Alberta T0M 0B0, Canada

Cysts of *Frenkelia* spp. have been reported in several species of rodents belonging to the families Muridae, Cricetidae, and Chinchillidae (Frenkel, 1956, Ann. N.Y. Acad. Sci. 64: 215–251; Karstad, 1963, Can. Vet. J. 4: 249–251; Krampitz and Rommel, 1977, Berl. Muench. Tieraerztl. Wochenschr. 90: 17–19; Meingassner and Burtscher, 1977, Vet. Pathol. 14: 146–153; Rommel and Krampitz, 1978, Zentralbl. Veterinaermed. Reihe B 25: 273–281). To our knowledge, the present report is the first to document *Frenkelia* as a natural infection in a porcupine.

The life cycle for any *Frenkelia* spp. has only recently been determined (Rommel and Krampitz, 1975, Berl. Muench. Tieraerztl. Wochenschr. 8: 338–340; Krampitz et al., 1976, Z. Parasitenkd. 51: 7–14; Rommel et al., 1976, Z. Parasitenkd. 50: 204–205; Rommel et al., 1977, Z. Parasitenkd. 51: 139–146). Rommel and Krampitz (1975, op. cit.) determined the life cycle and developmental stages of *F.*

*glareoli* (= *F. clethrionomyobuteonis*) in the buzzard (*Buteo buteo*) definitive host and bank vole (*Clethrionomys glareolus*) intermediate host. Other researchers have added to this information, noting that the parasite is specific for the definitive host, but not the intermediate one (Rommel et al., 1976, op. cit.; Rommel and Krampitz, 1978, op. cit.; Tadros et al., 1980, Trop. Geogr. Med. 32: 86). *Buteo buteo* is also the definitive host for *Frenkelia microti*. The specific name for *Frenkelia* spp. in North America is not known (Frenkel et al., 1979, Z. Parasitenkd. 58: 115–139).

On 13 August 1984 a female porcupine, in good physical condition, was trapped near Calgary, Alberta, Canada by personnel of the Fish and Wildlife Division and submitted to the Veterinary Laboratory, Animal Health Division, Airdrie, Alberta for examination. The porcupine was kept in captivity at the Veterinary Laboratory until 18 August, when it died. The brain was removed at necropsy and preserved in 10% buffered formalin. Portions of the brain tissue were prepared for histological examination. Histological examination re-

---

Received for publication 30 May 1985.



FIGURE 1. Cysts of *Frenkelia* sp. in the brain of a porcupine. H&E. Bar = 100  $\mu$ m.




FIGURE 2. Transmission electron micrograph of the cyst wall of *Frenkelia* sp. Note the thin wall and small rounded protrusions (P).  $\times 39,312$ .






FIGURE 3. Transmission electron micrograph of *Frenkelia* sp. Note the endodyocytes (E) and septa (S).  $\times 3,015$ .

vealed deeply lobed (Fig. 1), thin-walled (Fig. 2), septate cysts (Fig. 3) characteristic of *Frenkelia microti* (Findlay and Middleton, 1932) Biocca, 1966. However, because of the apparent final host specificity of the European species, and the geographic isolation from them, it is unlikely that our *Frenkelia* is the same species. Until the life cycle has been determined, no name is proposed. In the present case, the microscopic changes were similar to those described for other *Frenkelia* spp. from wild rodents (Frenkel, 1956, op. cit.; Karstad, 1963, op. cit.; Hayden et al., 1976, Vet. Pathol. 13: 337-342; Meingassner and Burtcher, 1977, op. cit.; Geisel et al., 1978, Vet. Pathol. 15: 621-630; Geisel et al., 1979, J. Wildl. Dis. 15:

267-270) and included perivascular and meningeal mononuclear cell infiltrates and gliosis. In contrast to wild rodents, granulomatous encephalitis has been reported from naturally infected lab-reared rats (Hayden et al., 1976, op. cit.).

The effects of this parasite on the porcupine were not determined, but it is conceivable that infections with large numbers of cysts could impair the motility of the porcupine as is believed may occur in voles (Geisel et al., 1978, op. cit.), but this has not been tested.

A voucher tissue section of *Frenkelia* sp. from the brain of this porcupine was deposited in the National Museums of Canada Parasite Collection, Ottawa (NMCP 1985-0086).