

Infestations of Lice on White-tailed Deer, *Odocoileus virginianus*, in the Great Smoky Mountains National Park, Tennessee, USA

Authors: Hribar, Lawrence J., Wilson, George Butler, and Qerhardt, Reid R.

Source: Journal of Wildlife Diseases, 22(2) : 272

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-22.2.272>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Infestations of Lice on White-tailed Deer, *Odocoileus virginianus*, in the Great Smoky Mountains National Park, Tennessee, USA

Lawrence J. Hribar,¹ George Butler Wilson,² and Reid R. Gerhardt,³ Department of Entomology and Plant Pathology, University of Tennessee, Knoxville, Tennessee 37901, USA

Tricholipeurus parallelus (Osborn) and *T. lipeuroides* (Megnin) (Mallophaga: Trichodectidae) are the only species of chewing lice known from white-tailed deer, *Odocoileus virginianus* (Zimmermann) (Emerson and Price, 1981, Misc. Publ. Entomol. Soc. Am. No. 12, 72 pp.). Deer in the Cades Cove area of the Great Smoky Mountains National Park, Tennessee, USA were examined for lice from September 1983 to October 1983, and again from January 1984 to December 1984. Deer were anesthetized with 6 to 9 mg succinyl choline administered by darts. The inner surfaces of the front and back legs and groin were examined for lice. All lice were preserved in 70% ethanol and identified according to Scanlon (1960, Wildl. Dis. No. 5, 121 pp.). Immature lice were identified to genus. Voucher specimens have been deposited in the U.S. National Parasite Collection, Beltsville, Maryland 20705, USA (Accession Numbers 78933–78935).

One hundred twenty-seven deer were examined. Seventy-nine deer (62%) were infested; 51 (40%) and 36 (28%) had *T. parallelus* and *T. lipeuroides*, respectively. There was no difference between par-

TABLE 1. Prevalence of biting lice of the genus *Tricholipeurus* on white-tailed deer of the Great Smoky Mountains National Park, Tennessee, USA.

	No. deer examined	No. deer positive (%)		
		<i>T. parallelus</i>	<i>T. lipeuroides</i>	Immature lice
Bucks	51	19 (37%)	7 (14%)	3 (6%)
Does	76	32 (42%)	29 (38%)	27 (36%)

asitism of male and female deer by *T. parallelus* ($P < 0.1$, χ^2 test), but *T. lipeuroides* parasitized significantly more females than males ($P > 0.01$, χ^2 test) (Table 1).

Tricholipeurus lipeuroides was found concurrently with *T. parallelus* on eight deer, approximately 6% of those examined. Other authors have reported concurrent infestations of the two species (Watson and Anderson, 1975, Can. J. Zool. 53: 1047–1054; Samuel et al., 1980, J. Med. Entomol. 17: 15–17).

One doe which harbored both male and female adult *T. lipeuroides* as well as immature lice was also harboring two adult female *Solenopotes ferrisi* (Fahrenholz) (Anoplura: Linognathidae).

The authors wish to thank Drs. John New and Lance Durden for assistance and advice, and Mr. Greg Wathen for assistance. Various members of the Uplands Field Research Laboratory staff assisted in collecting lice. Partial funding for this investigation was provided by the Uplands Field Research Laboratory, United States National Park Service.

Received for publication 26 February 1985.

¹ Present address: Department of Zoology/Entomology, Auburn University, Auburn, Alabama 36849, USA.

² Present address: Quillen-Discher School of Medicine, East Tennessee State University, Johnson City, Tennessee 36801, USA.

³ Author to whom reprint requests should be addressed.