

GASTROINTESTINAL HELMINTHS OF THE NORTHERN BOBWHITE IN FLORIDA: 1968 AND 1983

Authors: Moore, Janice, Freehling, Michael, and Simberloff, Daniel

Source: Journal of Wildlife Diseases, 22(4) : 497-501

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-22.4.497>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

GASTROINTESTINAL HELMINTHS OF THE NORTHERN BOBWHITE IN FLORIDA: 1968 AND 1983

Janice Moore,¹ Michael Freehling,^{2,4} and Daniel Simberloff³

ABSTRACT: We collected 153 northern bobwhites (*Colinus virginianus*) over a 10-mo period from Tall Timbers Research Station near Tallahassee, Florida, USA. Five species of gastrointestinal helminths were encountered commonly (>30% prevalence): *Cyrtocaria colini*, *Raillietina cesticillus*, *R. colinia*, *Heterakis isolonche*, and *Trichostrongylus tenuis*. Other helminths included *Brachylaima* sp., *Rhabdometra odiosa*, *Mediorhynchus papillosus*, *Cheilosporura spinosa*, *Dispharynx nasuta*, *Gongylonema ingluvicola*, and *Tetrameres pattersoni*. During the intervening 15 yr since the last year-round study of this population of birds, *C. spinosa* and *T. pattersoni* have declined markedly, and *Strongyloides* sp. probably has become extinct locally. Prevalence and intensity seem more likely to fluctuate in parasite species that have complex life cycles.

INTRODUCTION

The parasites of the northern bobwhite (*Colinus virginianus*) have been studied in many parts of the host's range (Kellogg and Calpin, 1971; Kellogg et al., 1972; Kocan et al., 1979; Davidson et al., 1980, 1982; Forrester et al., 1984; Lehmann, 1984). Most studies are based on single-year surveys (but see Forrester et al., 1984). Davidson and co-workers (1980) reported the results of a year-long monthly collection of parasites from bobwhites during 1968-1969 at Tall Timbers Research Station, Leon County, Florida. (We will refer to this as the 1968 study.) We studied the same population for 10 mo (1983-1984), hereafter called the 1983 study. Here we present our observations of prevalence and intensity and compare them to those of the previous study.

MATERIALS AND METHODS

One hundred fifty-three northern bobwhites were shot and their intestinal tracts were removed quickly and frozen. When the tracts

were thawed, helminth recovery methods of Kellogg and Prestwood (1968) were used. Unless otherwise indicated, all reported helminths are adults. In addition, we examined cecal tissue under a dissecting microscope after scraping and washing. This usually yielded additional specimens of *Trichostrongylus* that were embedded in the tissue. We used plumage analysis methods of Haugen (1957) to assign bobwhites to age categories. Data were not appropriate for parametric statistics; intensities were compared with Mann-Whitney *U*-tests. Prevalences were compared with *G*-tests. For statistical tests, $\alpha = 0.05$.

Our study differed from that of 1968 in that the number of birds per month was not constant (see Tables 2, 3), the number of males and females was not equal in some months, and our collection time was not limited to the middle of the month. We collected birds in areas immediately adjacent to 1968 collection sites.

Voucher specimens from this study have been deposited in the U.S. National Parasite Collection, Beltsville, Maryland (accession numbers 78976-78981).

RESULTS AND DISCUSSION

We found 12 species of helminths (Table 1). Of these, only five species exceeded 30% prevalence: *Raillietina colinia* and *R. cesticillus* (cestodes) and *Cyrtocaria colini*, *Heterakis isolonche* (= *H. bonasae*), and *Trichostrongylus tenuis* (nematodes). The acanthocephalan *Mediorhynchus papillosus* and the trematode *Brachylaima* sp. were rare in both studies. Other parasites that were once common (>30% prevalence) have declined dramatically (e.g.,

Received for publication 5 November 1985.

¹ Department of Zoology, Colorado State University, Fort Collins, Colorado 80523, USA.

² Tall Timbers Research Station, Route 1, Box 160, Tallahassee, Florida 32312, USA.

³ Department of Biological Science, Florida State University, Tallahassee, Florida 32306, USA.

⁴ Present address: Department of Zoology, Colorado State University, Fort Collins, Colorado 80523, USA.

TABLE 1. Helminths recovered from adult and juvenile northern bobwhites from Tall Timbers Research Station, Leon County, Florida.

Helminth	Bobwhites, 1983-1984		Bobwhites, 1968-1969*	
	Adults (n = 44)	Juveniles (n = 109)	Adults (n = 120)	Juveniles (n = 65)
Trematoda				
<i>Brachylaima</i> sp.	0	3.7 (1.5) 0-2 ^b	0	0
Cestoda				
<i>Hymenolepis</i> sp.	0	0	1 (1) 1	0
<i>Raillietina cesticillus</i>	31.8 (10.2) 1-86	17.4 (17.6) 1-51	32 (14) 1-37	32 (20) 1-124
<i>Raillietina colinia</i>	54.6 (11.8) 1-40	82.6 (9.5) 1-59	43 (8) 1-98	52 (6) 1-40
<i>Rhabdometra odiosa</i>	0	0.9 (1.0) 1	3 (4) 2-8	8 (3) 1-8
Acanthocephala				
<i>Mediorhynchus papillosus</i>	0	0.9 (1.0) 1	1 (2) 2	0
Nematoda				
<i>Cheilospirura spinosa</i>	2.3 (1.0) 1	0	72 (5) 1-35	45 (3) 1-26
<i>Cyrnea colini</i>	100 (4.9) 1-19	97.3 (5.0) 1-19	72 (4) 1-21	85 (5) 1-18
<i>Dispharynx nasuta</i>	11.4 (2.8) 1-8	19.3 (1.4) 1-3	2 (1) 1	9 (1) 1
<i>Gongylonema ingluvicola</i>	0	0.9 (1.0) 1	2 (1) 1	0
<i>Heterakis isolonche</i>	100 (50.4) 1-186	99.1 (25.0) 1-125	100 (47) 3-198	97 (20) 1-86
<i>Strongyloides</i> sp.	0	0	22 (5) 1-17	5 (5) 1-10
<i>Subulura</i> sp.	0	0	1 (1) 1	2 (1) 1
<i>Tetrameres pattersoni</i>	4.6 (1.5) 1-2	0	78 (7) 1-58	71 (10) 1-65
<i>Trichostrongylus tenuis</i>	95.5 (51.6) 1-290	75.9 (72.1) 1-470	98 (59) 1-361	42 (16) 1-123

* 1968-1969 data from Davidson et al. (1980).

^b Percent prevalence (average/infected bird) range.

Cheilospirura spinosa and *Tetrameres pattersoni*). This was probably not an artifact of disparate monthly sample sizes. Our largest sample was from February, when all common adult parasites were well-represented in the 1968 study. *Stron-*

gyloides sp. appears to have become locally extinct in bobwhites from Tall Timbers.

Prevalence and intensity of the common species of helminths and comparable data from 1968 are shown in Figure 1 and

TABLE 2. Helminth parasitism in juvenile northern bobwhites from Tall Timbers Research Station, Leon County, Florida.

Helminth*	October (n = 15)	November (n = 9)	December (n = 5)	January (n = 9)	February (n = 49)
<i>Dispharynx nasuta</i>	7 (2.0) 2 ^b	0 (0) 0	20 (2.0) 2	33 (1.7) 1-2	20 (1.4) 1-3
<i>Raillietina cesticillus</i>	27 (15.3) 1-36 40 (5)	11 (51.0) 51 40 (6)	20 (1.0) 1 40 (46)	0 (0) 0 20 (20)	23 (15.2) 1-50
<i>Raillietina colinia</i>	40 (5.5) 1-26 80 (6)	89 (4.0) 2-8 30 (17)	40 (9.5) 1-18 40 (3)	89 (9.0) 2-27 40 (6)	96 (13.1) 2-59
<i>Cyrnea colini</i>	100 (6.8) 1-19 60 (8)	89 (4.4) 1-10 90 (3)	80 (5.0) 2-11 90 (4)	100 (6.1) 1-10 100 (3)	100 (4.5) 1-12

* For *Heterakis* and *Trichostrongylus*, see Figure 1.^b Percent prevalence (average/infected bird) range—1983-1984. When two lines of data are displayed, the lower line is from the 1968 study and represents percent prevalence (average/infected bird), n = 10. 1983-1984 sample sizes are indicated under the month.




FIGURE 1. Prevalence and intensity of adult *Heterakis isolonche* and *Trichostrongylus tenuis* in juvenile northern bobwhites for 9 consecutive mo, 1968–1969 and 1983–1984. In 1968–1969, $n = 10$ quail/mo. In 1983–1984, $n = 3$ (August), 8 (September), 4 (March), 6 (April); see also Table 3. (Intensity data were not available for *T. tenuis* in August and September 1983.)

Table 2 (juvenile birds) and Table 3 (adult birds). The Southeastern Cooperative Wildlife Disease Study (SCWDS) shared their 1968 data with us, and we were thus able to compare intensity and prevalence between 1968 and 1983 birds when sample size permitted (October + November and December + January for juveniles, October + November and February for adults; Table 4). Where differences were observed, 1968 values (both prevalence and intensity) usually were lower; the sole exception was the prevalence of *R. cesti-*




FIGURE 2. Prevalence and intensity of larvae of *Trichostrongylus tenuis* in juvenile northern bobwhite quail at Tall Timbers Research Station, 1983–1984.

cillus in December + January juvenile quail. This agrees with the trends for the entire sample (Table 1). In their 5-yr southern Florida study, Forrester and co-workers (1984) found statistically significant year-to-year differences in prevalence for four helminths (*C. colini*, *Ascaridia galli*, *T. tenuis* and *H. isolonche*), and significant intensity differences for *C. colini*.

Bobwhite populations have declined generally at Tall Timbers (Davidson, pers. comm.). This may have contributed to the local extinction of *Strongyloides* and decreases in *C. spinosa* and *T. pattersoni*. However, four of the five common helminths have increased in prevalence and/or intensity since 1968. This suggests that factors other than definitive host density may be influencing populations of helminths in bobwhites. Changes in the availability and susceptibility of intermediate hosts may be important in this regard. The joint decline of *C. spinosa* and *T. pattersoni*, the only two species with grasshoppers as intermediate hosts (Cram et al., 1931), is consistent with this suggestion, as is the fact that heteroxenous species exhibited more changes in intensity and prevalence than did monoxenous ones (Table 4).

TABLE 3. Helminth parasitism in adult northern bobwhites from Tall Timbers Research Station, Leon County, Florida.

Helminth	October (n = 4)	November (n = 6)	February (n = 14)
<i>Dispharynx nasuta</i>	25 (2.0) 2*	17 (2.0) 2	7 (1.0) 1
<i>Raillietina cestitillus</i>	0 20 (2)	50 (32.7) 2-86 30 (2)	21 (2.7) 1-6 50 (7)
<i>Raillietina colinia</i>	0 0 (0)	50 (9.3) 1-26 20 (1)	93 (13.5) 2-36 50 (3)
<i>Cheilospirura spinosa</i>	25 (1.0) 1 80 (5)	— 90 (3)	— 70 (3)
<i>Cyrnea colini</i>	100 (3.0) 1-7 30 (4)	100 (4.2) 1-9 60 (2)	100 (3.3) 1-7 (n = 13) 60 (4)
<i>Heterakis isolonche</i>	100 (78.6) 11-186 100 (73)	100 (25.1) 1-123 100 (45)	100 (51.7) 1-137 100 (17)
<i>H. isolonche</i> (immature)	33 (1.0) 1 (n = 3) 30 (5)	83 (1.8) 1-3 40 (2)	64 (3.4) 1-9 0 (0)
<i>Tetrameres pattersoni</i>	— 80 (9)	16.7 (2.0) 2 80 (7)	7 (1.0) 1 80 (12)
<i>Trichostrongylus tenuis</i>	75 (19.7) 9-38 90 (22)	83 (20.3) 1-49 100 (33)	100 (92.8) 14-290 100 (32)
<i>T. tenuis</i> (immature)	0	17 (1.0) 1	36 (8.2) 1-20

* Percent prevalence (average/infected bird) range—1983–1984. When two lines of data are displayed, the lower line is from the 1968 study and represents percent prevalence (average/infected bird), n = 10. 1983–1984 sample sizes are indicated under the month unless otherwise noted.

Abiotic conditions may be important also. These may contribute to the puzzling seasonal pattern exhibited by larvae of *T. tenuis* (Fig. 2). Under optimal conditions, this monoxenous nematode can develop from fresh egg to larva in 1–2 days, and once ingested, to adulthood in 4 days (Cram et al., 1931). Increases in prevalence and intensity of adult *T. tenuis* (Fig. 1, Table 2) accompanying similar increases in larvae (Fig. 2) confirmed the fact that, despite its potentially brief,

TABLE 4. Within-season comparisons of intensity and prevalence of common bobwhite quail helminths, 1968–1969 and 1983–1984.

Helminth	Intensity		Prevalence	
	Juvenile quail	Adult quail	Juvenile quail	Adult quail
<i>Cyrnea colini</i>	1968 lower ($P < 0.05$, Dec + Jan)	NS*	1968 lower ($P < 0.05$, Oct + Nov)	1968 lower ($P < 0.005$ Oct + Nov and Feb)
<i>Raillietina cestitillus</i>	1968 lower? ($P < 0.10$, Oct + Nov)	NS	1968 higher? ($P < 0.10$, Dec + Jan)	NS
<i>Raillietina colinia</i>	1968 lower? ($P < 0.10$, Dec + Jan)	NS	1968 lower? ($P < 0.10$, Dec + Jan)	1968 lower ($P < 0.025$, Feb)
<i>Heterakis isolonche</i>	NS	1968 lower ($P < 0.05$, Feb)	—	—
<i>Trichostrongylus tenuis</i>	NS	NS	NS	NS

* NS = not significant.

monoxenous life cycle, this is a seasonally occurring worm. Hon and co-workers (1978) observed similar *T. tenuis* seasonality in wild turkeys in southern Florida. Temperature and moisture have been shown to influence greatly the development of *T. tenuis* in the laboratory (Leiper, 1911) and of a congener (*T. vitrinus*) in the field (Callinan, 1979; Rose and Small, 1984). The other monoxenous helminth in this study, *H. isolonche*, exhibits a seasonal pattern similar to that of *T. tenuis* (Fig. 1), and may likewise be responsive to abiotic conditions. Although definitive host density clearly influences the population biology of bobwhite helminths (Davidson, pers. comm.), not all populations have behaved similarly over time, and other influences, such as intermediate host and abiotic factors, are implicated.

ACKNOWLEDGMENTS

We are grateful for the assistance of personnel of the Southeastern Cooperative Wildlife Disease Study, University of Georgia, which was provided under the Federal Aid in Wildlife Restoration Act (50 Stat. 917) and through Contract Nos. 14-16-0009-82-500, 14-16-0004-83-004, and 14-16-0004-84-005, Fish and Wildlife Service, U.S. Department of the Interior. W. R. Davidson, G. L. Doster, G. W. Esch and F. E. Kellogg read an earlier draft of this manuscript and provided helpful comments. We appreciate the field assistance of J. Atkinson and B. Mueller (Tall Timbers Research Station) and the technical assistance of S. Forster and J. Liberatos (Florida State University). Dr. R. Short provided laboratory space at Florida State University and Colorado State University provided computer time. This study was funded by NSF BSR-8210218 to J. Moore, D. Simberloff and R. Short.

LITERATURE CITED

- CALLINAN, A. P. L. 1979. The ecology of the free-living stages of *Trichostrongylus vitrinus*. *Int. J. Parasitol.* 9: 133-136.
- CRAM, E. B. 1931. Developmental stages of some nematodes of the Spiruroidea parasitic in poultry and game birds. U.S. Dep. Agric. Tech. Bull. No. 227, 27 pp.
- , M. F. JONES, AND E. A. ALLEN. 1931. Internal parasites and parasitic diseases of the bobwhite. *In* The Bobwhite Quail: Its Habits, Preservation and Increase, H. L. Stoddard. Charles Scribner's Sons, New York, pp. 229-313.
- DAVIDSON, W. R., F. E. KELLOGG, AND G. L. DOSTER. 1980. Seasonal trends of helminth parasites of bobwhite quail. *J. Wildl. Dis.* 16: 367-375.
- , ———, AND ———. 1982. An overview of disease and parasitism in southeastern bobwhite quail. *In* Proc. Second Nat. Bobwhite Quail Symp., F. Schitoskey, E. C. Schitoskey, and L. G. Talent (eds.). Oklahoma State University, Stillwater, pp. 57-63.
- FORRESTER, D. J., J. A. CONTI, A. O. BUSH, L. D. CAMPBELL, AND R. K. FROHLICH. 1984. Ecology of helminth parasitism of bobwhites in Florida. *Proc. Helminthol. Soc. Wash.* 51: 255-260.
- HAUGEN, A. O. 1957. Distinguishing juvenile from adult bobwhite quail. *J. Wildl. Manage.* 21: 29-32.
- HON, L. T., D. J. FORRESTER, AND L. E. WILLIAMS, JR. 1978. Helminth acquisition by wild turkeys (*Meleagris gallopavo osceola*) in Florida. *Proc. Helminthol. Soc. Wash.* 45: 211-218.
- KELLOGG, F. E., AND J. P. CALPIN. 1971. A checklist of parasites and diseases reported from the bobwhite quail. *Avian Dis.* 15: 704-715.
- , AND G. L. DOSTER. 1972. Diseases and parasites of the bobwhite. *In* Proc. First Nat. Bobwhite Quail Symp., J. S. Morrison and J. C. Lewis (eds.). Oklahoma State University, Research Foundation, Stillwater, pp. 233-267.
- , AND A. K. PRESTWOOD. 1968. Gastrointestinal helminths from wild and pen-raised bobwhites. *J. Wildl. Manage.* 32: 468-475.
- KOCAN, A. A., L. HANNON, AND J. H. EVE. 1979. Some parasitic and infectious diseases of bobwhite quail from Oklahoma. *Proc. Okla. Acad. Sci.* 59: 20-22.
- LEHMANN, V. W. 1984. Bobwhites in the Rio Grande plain of Texas. Texas A&M University Press, College Station, Texas, 371 pp.
- LEIPER, R. T. 1911. On the development and bionomics of *Trichostrongylus pergracilis*. *In* The Grouse in Health and in Disease, Lord Lovat (chairman, Committee of Inquiry on Grouse Disease). Smith, Elder and Co., London, pp. 218-234.
- ROSE, J. H., AND A. J. SMALL. 1984. Observations on the bionomics of the free-living stages of *Trichostrongylus vitrinus*. *J. Helminthol.* 58: 49-58.