

## **Serosurvey for Selected Virus Infections of Wild Carnivores in Taiwan and Vietnam**

Authors: Ikeda, Yasuhiro, Miyazawa, Takayuki, Nakamura, Kazuya, Naito, Risako, Inoshima, Yasuo, et al.

Source: Journal of Wildlife Diseases, 35(3) : 578-581

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-35.3.578>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## Serosurvey for Selected Virus Infections of Wild Carnivores in Taiwan and Vietnam

Yasuhiro Ikeda,<sup>1</sup> Takayuki Miyazawa,<sup>1</sup> Kazuya Nakamura,<sup>1</sup> Risako Naito,<sup>1</sup> Yasuo Inoshima,<sup>1</sup> Kwong-Chung Tung,<sup>2</sup> Wei-Ming Lee,<sup>2</sup> Ming-Chu Chen,<sup>3</sup> Tzong-Fu Kuo,<sup>4</sup> James A. Lin<sup>2</sup> and Takeshi Mikami<sup>1,5</sup>

<sup>1</sup> Department of Veterinary Microbiology, Graduate School of Agricultural and Life Sciences, The University of Tokyo, 1-1-1 Yayoi, Bunkyo-ku, Tokyo 113-8657, Japan; <sup>2</sup> Department of Veterinary Medicine, National Chung Hsing University, 250, Kwo-Kwang Rd., Taichung 40227, Taiwan, ROC.; <sup>3</sup> Taiwan Provincial Research Institute for Animal Health, 376, Chung-Cheng Rd., Tam-Sui, Taipei, Taiwan, ROC.; <sup>4</sup> Department of Veterinary Medicine, National Taiwan University, 142, Zou-Shan Rd., Taipei, Taiwan, ROC; and <sup>5</sup> Corresponding author (e-mail: ataka@hongo.ecc.u-tokyo.ac.jp).

**ABSTRACT:** Serum samples from two leopard cats (*Felis bengalensis*) and four Formosan gem-faced civets (*Paguma larvata taivana*) in Taiwan, September 1995, and nine leopard cats in Vietnam, August and December 1997, were examined for the prevalence of antibodies against feline parvovirus, feline herpesvirus type 1, feline calicivirus and feline immunodeficiency virus. All civets and nine of 11 leopard cats were shown to have antibodies against feline parvovirus (FPV), and FPVs were isolated from mononuclear cells in the peripheral blood of the six leopard cats.

**Key words:** Feline calicivirus, feline herpesvirus type 1, feline immunodeficiency virus, feline parvovirus, *Felis bengalensis*, gem-faced civet, leopard cat, *Paguma larvata*, serosurvey.

Significant disease syndromes in domestic cats are associated with viral infections, and several viruses of these are known to infect wild felines (Artois and Remond, 1994). In domestic cats, feline herpesvirus type 1 (FHV-1) causes rhinotracheitis (Gaskell and Povey, 1979), feline calicivirus (FCV) is responsible for stomatitis, gingivitis, and circumscribed lesions of the tongue (Wardly and Povey, 1977), feline leukemia virus (FeLV) causes several neoplastic and non-neoplastic diseases (Neil et al., 1991), feline immunodeficiency virus (FIV) induces immunosuppression (Pedersen et al., 1987; Carpenter and O'Brien, 1995) and feline parvovirus (FPV) is associated with clinical signs of gastroenteritis and panleukopenia (Parrish, 1995).

Formosan gem-faced civets (*Paguma larvata taivana*) in Taiwan and leopard cats (*Felis bengalensis*) in Asia are threatened species of the Viverridae and Felidae, respectively. Although parasites of wild felidae in Thailand have been report-

ed (Patton and Rabinowitz, 1994), little is known about the prevalence of viral diseases in Asian wild Felidae or Viverridae. In the present study, we investigated the presence of certain feline viruses in a community of leopard cats and gem-faced civets in Taiwan and Vietnam and attempted the isolation of FPV from these animals.

A list of animals examined and their assigned numbers are shown in Table 1. All 15 animals examined were apparently healthy. Blood samples of eight leopard cats, LVT1 to LVT8, were collected in Hanoi district (Vietnam; 21°05'N, 105°55'E) in 1997. LVT1 was kept in the Hanoi Zoological Garden. LVT2 and LVT3 were captured and protected in cages in a wildlife conservation center. Five free-ranging cats, LVT4 to LVT8, were captured in mountain areas near Hanoi City (Vietnam). LVT9 was a captive leopard cat, which was sold at an animal market in Ho Chi Minh City (Vietnam; 10°58'N, 106°40'E) in 1997. Two leopard cats, LTW1 and LTW2, had been captured in Nantou County (Taiwan; 24°02'N, 121°20'E) in 1994 and 1989, respectively and kept in cages as pets. Four gem-faced civets were born in a breeding cattery at Taichung County (Taiwan; 24°15'N, 121°02'E) in 1994, and kept in cages as pets. Blood samples of these six animals from Taiwan were collected in 1995.

Peripheral blood mononuclear cells (PBMCs) of the animals were isolated by Ficoll-Paque (Pharmacia LKB Biotechnology AB, Upsalla, Sweden) and stimulated by 10 µg/ml of concanavalin-A (con-A) for

3 days as reported previously (Miyazawa et al., 1989). Con-A stimulated PBMCs were maintained in RPMI1640 growth medium supplemented with 10% fetal calf serum, antibiotics and 100 units/ml of recombinant human interleukin-2.

Serum antibodies against FHV-1, FCV, and FPV were detected by indirect immunofluorescence (IF) assays using glass slides coated with the Crandell feline kidney (CRFK) cells (Crandell et al., 1973) infected with the respective viruses. The cells were fixed in cold acetone for 30 min, then incubated at 37 C for 30 min with the 100-fold diluted serum samples. After incubation, the cover slips were washed twice with phosphate-buffered saline (PBS), and incubated with 200-fold diluted goat anti-cat immunoglobulin G (whole molecule) conjugated with fluorescein isothiocyanate (Cappel, Aurora, Ohio, USA). After incubation at 37 C for 30 min, the glass slides were washed twice with PBS, mounted in buffered glycerol and examined by fluorescence microscopy. Sera from FHV-1-, FCV-, FPV- or mock-infected specific pathogen-free cats were used as positive and negative controls. Additionally, the titers of virus neutralizing (VN) antibodies against FPV in the plasma samples were determined as reported previously (Ikeda et al., 1998a). For detection of FPV antigens in the cultured PBMCs, an indirect IF assay using anti-FPV VP2 monoclonal antibody (mAb) 2D9 (Mochizuki et al., 1989) was performed as reported previously (Ikeda et al., 1998b). FeLV antigen and antibodies against FIV were checked using an ELISA method using commercial kits (Cite-Combo: IDEXX Co., Portland, Maine, USA).

Recently, we reported that the eight leopard cats in Vietnam were not infected with FIV or FeLV (Miyazawa et al., 1998). By using a commercial kit, it also was shown that the cat in Ho Chi Minh City and two cats and four civets in Taiwan did not have FeLV-specific antigens or antibodies against FIV. By using the indirect IF assay, it was demonstrated that the two

leopard cats and four civets in Taiwan had antibodies against FPV but not FHV-1 or FCV. Alternatively, the leopard cats in Vietnam were shown to have antibodies against FHV-1 (1/9), FCV (3/9) and FPV (7/9). In addition, the titers of VN antibodies against FPV were demonstrated to range from 160 to >5,120 in the four civets and 640 to >5,120 in the nine leopard cats (Table 1).

The isolation of FPV from the PBMCs of the infected domestic cats with high VN antibodies has been reported (Miyazawa, et al., 1999). For isolation of FPV from the civets or the feral cats, the Con-A-stimulated PBMCs of the all civets and 10 cats were maintained for more than 2 wk in culture. At 3 to 9 days after cultivation, the PBMCs of the six leopard cats showed cytopathic effects (CPEs) such as cell rounding and nuclear disintegration. The observed CPE's were similar to those observed in FPV-infected feline PBMCs (Ikeda et al., 1998). By the indirect IF assay using mAb 2D9, PBMCs showing CPEs from six leopard cats were shown to contain FPV-specific antigens (Table 1). In contrast, no FPV-specific antigen was detected in the other eight cultured samples.

In the present study, we found the seroprevalence of FCV, FHV-1 and FPV in leopard cats and the susceptibility of gem-faced civets to FPV. Additionally, we isolated FPV from the PBMCs of six leopard cats with high VN antibodies.

Since the number of samples is unfortunately low, it is difficult to conclude that the observed prevalence of the viruses is representative of the local population of feral cats or civets in Taiwan or Vietnam. However, these results indicate that leopard cats are susceptible to infection with FHV-1, FCV and FPV and that the civets also are susceptible to FPV infection. Infection of FHV-1 or FCV in domestic cats is known to occur by close contact between the animals. As shown in Table 1, most of the cats examined had antibodies against FPV but few of the cats had antibodies against FHV-1 or FCV, suggesting

TABLE 1. Detection of antibodies against feline calicivirus, feline herpesvirus type 1, and feline parvovirus in Formosan gem-faced civets and leopard cats in Taiwan and Vietnam.

Animal	Number	Sex	FCV <sup>a</sup> IFA	FHV-1 <sup>b</sup> IFA	FPV <sup>c</sup> IFA	FPV titer	FPV isolation
Civet	CTW1	M	—	—	+	>5,120	—
	CTW2	F	—	—	+	×2,560	—
	CTW3	F	—	—	+	×160	—
	CTW4	F	—	—	+	×5,120	—
Leopard cat	LTW1	F	—	—	+	×640	—
	LTW2	F	—	—	+	×640	+
	LVT1	M	+	+	+	>5,120	—
	LVT2	M	—	—	+	×2,560	—
	LVT3	F	—	—	+	>5,120	+
	LVT4	F	—	—	+	×2,560	+
	LVT5	?	—	—	—	<10	ND <sup>d</sup>
	LVT6	M	+	—	+	×640	+
	LVT7	M	+	—	+	×1,280	+
	LVT8	M	—	—	+	×2,560	+
	LVT9	M	—	—	—	<10	—

<sup>a</sup> Feline calicivirus.<sup>b</sup> Feline herpesvirus type 1.<sup>c</sup> Feline parvovirus.<sup>d</sup> Not done.

that the leopard cats seldom come into contact with free-roaming domestic cats in these areas. Since FPV is generally shed only for a short period of time by domestic cats but survives in the environment for up to several months (Scott, 1980), the infection of FPV may be due to the environmental exposure. The absence of FeLV antigens and relative lack of antibodies against FHV-1 or FIV in the animals suggests that these viruses may not actively circulate in the population.

In the present study, we reported the isolation of FPV from the PBMCs of the leopard cats with high VN antibodies. This observation indicates that FPV can persistently infect the lymphocytes even in the presence of high VN antibodies. Although animals that recover from FPV infection are thought to be protected for life from re-infection, it is possible that the persistent virus stimulates the host immune system for life even after the recovery. At present, however, it remains to be determined whether shedding of the virus following periods of stress occurs after clinical recovery.

Since there are few reports on isolation

of FPV from feral cats, the six FPV isolates reported here may provide insight into the FPV infection in Asian feral cats. Further studies will be necessary on biological and genetical differences between the six isolates and FPVs from domestic cats and dogs. Moreover, the method presented here may be applicable to isolate FPV from wild animals.

We thank M. Hattori for providing recombinant human IL-2-producing Ltk<sup>+</sup>IL-2. 23 cells. This work was partly supported by grants from the Ministry of Education, Science and Culture, and from the Ministry of Health and Welfare of Japan.

#### LITERATURE CITED

- ARTOIS, M., AND M. REMOND. 1994. Viral diseases as a threat to free-living wild cats (*Felis silvestris*) in Continental Europe. *Veterinary Record* 134: 651–652.
- CARPENTER, M. A., AND S. J. O'BRIEN. 1995. Coadaptation and immunodeficiency virus: lessons from the felidae. *Current Opinion in Genetics and Development* 5: 739–745.
- CRANDELL, R. A., C. G. FABRICANT, AND W. A. NELSON-REES. 1973. Development, characterization and viral susceptibility of a feline (*Felis catus*) renal cell line (CRFK). *In Vitro* (Rockville) 9: 176–185.

- GASKELL, R. M., AND R. C. POVEY. 1979. Feline rhinotracheitis: sites of virus replication and persistence in acutely and persistently infected cats. *Research in Veterinary Science* 27: 167–174.
- IKEDA, Y., T. MIYAZAWA, K. KUROSAWA, R. NAITO, S. HATAMA, C. KAI, AND T. MIKAMI. 1998a. New quantitative methods for detection of feline parvovirus (FPV) and virus neutralizing antibody against FPV using a feline T lymphoid cell line. *Journal of Veterinary Medical Science* 60: 973–974.
- IKEDA, Y., J. SHINOZUKA, T. MIYAZAWA, K. KUROSAWA, Y. IZUMIYA, Y. NISHIMURA, K. NAKAMURA, J. CAI, K. FUJITA, K. DOI, AND T. MIKAMI. 1998b. Apoptosis in feline panleukopenia virus-infected lymphocytes. *Journal of Virology* 72: 6932–6936.
- MIYAZAWA, T., T. FURUYA, S. ITAGAKI, Y. TOHYA, K. NAKANO, E. TAKAHASHI, AND T. MIKAMI. 1989. Preliminary comparisons of the biological properties of two strains of feline immunodeficiency virus (FIV) isolated in Japan with FIV Petaluma strain isolated in the United States. *Archives of Virology* 108: 131–135.
- MIYAZAWA, T., Y. IKEDA, K. MAEDA, T. HORIMOTO, Y. TOHYA, M. MOCHIZUKI, D. VU, G. D. VU, D. X. CU, K. ONO, E. TAKAHASHI, AND T. MIKAMI. 1998. Seroepidemiological survey of feline retrovirus infections in domestic and leopard cats in North Vietnam in 1997. *Journal of Veterinary Medical Science* 60: 1273–1275.
- MIYAZAWA, T., Y. IKEDA, R. NAITO, M. MOCHIZUKI, Y. TOHYA, D. VU, T. MIKAMI, AND E. TAKAHASHI. 1999b. Isolation of feline parvovirus from peripheral blood mononuclear cells of cats in northern Vietnam. *Microbiology and Immunology* 1999: In press.
- MOCHIZUKI, M., S. KONISHI, M. AJIKI, AND T. AKBOSHI. 1989. Comparison of feline parvovirus subspecific strains using monoclonal antibodies against a feline panleukopenia virus. *Japanese Journal of Veterinary Science* 51: 264–272.
- NEIL, J. C., R. FULTON, M. RIGBY, AND M. STEWART. 1991. Feline leukemia virus: generation of pathogenic and oncogenic variants. *Current Topics in Microbiology and Immunology* 171: 67–93.
- PEDERSEN, N. C., E. W. HO, M. L. BROWN, AND J. K. YAMAMOTO. 1987. Isolation of a T-lymphotropic virus from domestic cats with an immunodeficiency-like syndrome. *Science* 235: 790–793.
- PARRISH, C. R. 1995. Molecular epidemiology of parvoviruses. *Seminars in Virology* 6: 415–418.
- PATTON, S. AND A. R. RABINOWITZ. 1994. Parasites of wild felidae in Thailand: a coprological survey. *Journal of Wildlife Diseases* 30: 472–475.
- SCOTT, F. W. 1980. Virucidal disinfection on development of humoral and cellular immunity. *Cornell Veterinarian* 66: 324–332.
- WARDLEY, R. C. AND R. C. POVEY. 1977. The pathology and sites of persistence associated with three different strains of feline caliciviruses. *Research in Veterinary Science* 23: 15–19.

*Received for publication 22 June 1998.*