

HELMINTHS OF BLACK-TAILED JACK RABBITS IN NEW MEXICO 1

Author: SAMSON, KAY S.

Source: Bulletin of the Wildlife Disease Association, 4(4) : 130

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-4.4.130>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

HELMINTHS OF BLACK-TAILED JACK RABBITS IN NEW MEXICO¹

To determine if hares were reservoir hosts for helminth parasites of cattle and sheep, 25 black-tailed jack rabbits, *Lepus californicus*, were collected in Lincoln and Chaves counties, New Mexico, and examined for parasites by standard parasitological procedures. The results of these examinations are summarized in the table below.

The only parasite collected that was common to the rabbits and domestic ruminants was *Trichostrongylus colubriformis* which I recovered from the small intestines of 20 of the 25 rabbits. No parasites were recovered from the stomachs, hearts, livers, kidneys, lungs, or spleens.

The small intestines from 10 of these rabbits were cut into small pieces and digested in artificial gastric juice as described by Herlich (1956, Proc. Helm. Soc. Wash. 23:102-103). The residue was examined for nematode larvae; no larvae were found.

Although no regular examination was made for ectoparasites, the ear tick, *Dermacentor parumapertus*, was recovered from the ears of some rabbits.

Dikmans (1937, Proc. Helm. Soc. Wash. 4:65-67) described *Nematodirus neomexicanus*, a nematode parasite, from the small intestine of *Lepus californicus texianus*, collected near Albuquerque, New Mexico. To my knowledge, this is the only report of parasites of rabbits in New Mexico.

TABLE 1. Helminths of Black-Tailed Jack Rabbits.

Helminths	Number infected	Location	Worms recovered
Nematodes:			
<i>Trichostrongylus colubriformis</i>	20	Small intestine	2 to 125
<i>Nematodirus neomexicanus</i>	22	Small intestine	1 to 241
<i>Nematodirus</i> sp. (larvae)	6	Small intestine	2 to 44
<i>Dermatoxys veligera</i>	15	Large intestine	1 to 13
<i>Passalurus nonanulatus</i>	12	Large intestine	1 to 9360
<i>Micipsella brevicauda</i>	1	Abdominal cavity	1
Cestodes:			
<i>Raillietina retractilis</i>	14	Small intestine	1 to 290
<i>Mosgovoyia pectinata</i>	4	Small intestine	1 to 3

KAY S. SAMSON

USDA, ARS, Animal Disease and Parasite Research Division
 P.O. Box 3518
 Las Cruces, New Mexico 88001
 April 26, 1968

¹ This work was carried out in cooperation with the New Mexico Agricultural Experiment Station.