



**Diectophyme renale (Nematoda, Diectophymatidae)
Infection in the Crab-eating Fox (*Cerdocyon thous*) from
Brazil**

Authors: Ribeiro, Carlos Torres, Verocai, Guilherme Gomes, and
Tavares, Luiz Eduardo Roland

Source: Journal of Wildlife Diseases, 45(1) : 248-250

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-45.1.248>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

***Diectophyme renale* (Nematoda, Diectophymatidae) Infection in the Crab-eating Fox (*Cerdocyon thous*) from Brazil**

Carlos Torres Ribeiro,¹ Guilherme Gomes Verocai,^{2,3} and Luiz Eduardo Roland Tavares² ¹ Biotério de Experimentação Animal do IOC, FIOCRUZ, Rio de Janeiro, RJ, Brazil; ² Curso de Pós-Graduação em Ciências Veterinárias, Departamento de Parasitologia Animal/IV/UFRRJ, Seropédica, RJ, Brazil; ³ Corresponding author (email: verocai@gmail.com)

ABSTRACT: The purpose of this study is to report the first case of parasitism by the giant kidney worm, *Diectophyme renale* (Goeze, 1782), in the crab-eating fox, *Cerdocyon thous* (Linnaeus, 1766), in Brazil. A debilitated adult male fox was taken to the Screening Center for Wild Animals in Rio de Janeiro (CETAS/RJ) and eventually died of an unknown cause. During necropsy, a brownish-red nematode, 28.9 cm long, was found in the peritoneal cavity. The worm was identified as a male *D. renale*. This study reports the first case of diectophymatosis in the crab-eating fox, enlarging the range of wild definitive host species infected by the giant kidney worm in the Neotropical region.

Key words: Brazil, *Cerdocyon thous*, crab-eating fox, diectophymatosis, *Diectophyme renale*, helminth, nematoda, Neotropical region.

The giant kidney worm, *Diectophyme renale* (Goeze, 1782) is one of largest parasitic nematodes. Its complex life cycle requires an aquatic oligochaete or annelid as an intermediate host. Fishes and frogs are paratenic hosts that serve to infect the definitive hosts (Mace and Anderson, 1975; Anderson, 2000). Wild carnivores such as mustelids, canids, and some domesticated animals are known definitive hosts, with sporadic reports in atypical hosts which include humans (Woodhead, 1950; Anderson, 2000; Measures, 2001).

In the Neotropical region, various wild carnivore species were reported as hosts, including the maned-wolf (*Chrysocyon brachyurus*), the bush dog (*Speothos venaticus*), the coati (*Nasua nasua*), and species of grison (*Galictis cuja* and *G. vittata*; Giovannoni and Molfi, 1960; Barros et al., 1990). There have also been many reports of parasitism in domestic dogs in several Brazilian states (Costa and Lima, 1988; Kano et al., 2003; Pereira et

al., 2006). An unusual host, the noncarnivorous two-toed sloth *Choloepus didactylus*, was also reported as infected by Rocha et al. (1965). In this study, we report the first case of parasitism by the giant kidney worm, *D. renale*, in the crab-eating fox, *Cerdocyon thous* (Linnaeus, 1766), in Brazil.

An adult male crab-eating fox was taken to the Screening Center for Wild Animals in Rio de Janeiro (CETAS/RJ), which is maintained by the Institute for Environment and Renewable Natural Resources (IBAMA) and located in the Municipality of Seropédica, State of Rio de Janeiro, Southeastern Brazil. The animal was considerably debilitated and eventually died from an unknown cause. During the necropsy, a brownish-red nematode was found in the peritoneal cavity. There were no additional findings related to parasitism. The parasite was placed in saline solution and sent to the Department of Animal Parasitology of the Veterinary Institute of the Universidade Federal Rural do Rio de Janeiro, where it was fixed in alcohol-formalin-acetic acid solution. Identification was carried out using the key of Anderson and Bain (1982). The nematode was a male *D. renale* measuring 28.9 cm long with a maximum width of 3.8 mm. The specimen was deposited in the Helminthological Collection of Oswaldo Cruz Institute labeled as CHIOC no. 35568.

Helminthic parasites of the crab-eating fox have been reported (Vicente et al., 1997; Ruas et al., 2003; Santos et al., 2003, 2004; Vieira et al., 2008), but none of these studies reported diectophymatosis in this host. Because crab-eating fox consume amphibians and fishes (e.g., cichlids),

which may be possible paratenic hosts in Brazil, this mammal may be exposed to natural infections (Gatti et al., 2006; Pedó et al., 2006). However, there are no reports of Neotropical freshwater fishes harboring *D. renale* larvae (Moravec, 1998). Most reports of *D. renale* are from North America, Europe, and Asia (Mace, 1976a; Measures, 2001).

Renal parasitism by *D. renale* can cause total destruction of the renal parenchyma, atrophy and fibrosis of renal tubules, dilation of the renal pelvis, and urethral obstruction, and its occurrence in the abdominal cavity can lead to adhesions and peritonitis (Mace and Anderson, 1975; Mace 1976b; Measures and Anderson, 1985). This study enlarges the range of wild definitive host species infected by the giant kidney worm, *D. renale*, in the Neotropical region by reporting the first case of dioctophymatosis in the crab-eating fox.

The authors gratefully acknowledge the Brazilian National Council of Research and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior for partial support of this work.

LITERATURE CITED

- ANDERSON, R. C. 2000. Nematode parasites of vertebrates: Their development and transmission, 2nd Edition. CAB Publishing, London, UK. 672 pp.
- , AND O. BAIN. 1982. Keys to genera of the superfamilies Rhabditoidea, Dioctophymatoidea, Trichinelloidea and Muspiceoidea, no. 9. In CIH keys to the nematode parasites of vertebrates, R. C. Anderson, A. G. Chabaud and S. Willmott (eds.). CAB International, Wallingford, UK. pp. 15–16.
- BARROS, D. M., M. L. LORINI, AND V. G. PERSSON. 1990. Dioctophymosis in the little grison (*Galictis cuja*). *Journal of Wildlife Diseases* 26: 538–539.
- COSTA, H. M. A., AND W. S. LIMA. 1988. *Dioctophyma renale* (Goeze, 1782): Ocorrências em Minas Gerais. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia* 40: 243–244.
- GATTI, A., R. BIANCHI, C. R. X. ROSA, AND S. L. MENDES. 2006. Diet of two sympatric carnivores, *Cerdocyon thous* and *Procyon cancrivorus*, in a resting area of Espírito Santo State, Brazil. *Journal of Tropical Ecology* 22: 227–230.
- GIOVANNONI, M., AND A. O. MOLFI. 1960. *Dioctophyma renale* (Goeze, 1782) no Brasil. *Anais da Faculdade de Medicina da Universidade Federal do Paraná* 3: 99–104.
- KANO, F. S., M. T. SHIMADA, S. N. SUZUKI, S. C. OSAKI, B. C. MENARIM, F. R. V. RUTHES, AND M. A. LAIDANE-FILHO. 2003. Ocorrência de dioctofimose em dois cães no município de Guarapuava-PR. *Semina Ciências Agrárias* 24: 177–180.
- MACE, T. F. 1976a. Bibliography of giant kidney worm *Dioctophyma renale* (Goeze, 1782) (Nematoda: Dioctophymatoidea). *Wildlife Disease* No. 69, 36 pp.
- . 1976b. Lesions in mink (*Mustela vison*) infected with giant kidney worm (*Dioctophyma renale*). *Journal of Wildlife Diseases* 12: 88–92.
- , AND R. C. ANDERSON. 1975. Development of the giant kidney worm, *Dioctophyma renale* (Goeze, 1782) (Nematoda: Dioctophymatoidea). *Canadian Journal of Zoology* 53: 1552–1568.
- MEASURES, L. N. 2001. Dioctophymatosis. In *Parasitic diseases of wild animals*, W. M. Samuel, M. J. Pybus and A. A. Kocan (eds.). Iowa State University Press, Ames, Iowa. pp. 357–364.
- , AND R. C. ANDERSON. 1985. Centrarchid fish as paratenic hosts of the giant kidney worm, *Dioctophyma renale* (Goeze, 1782). *Journal of Wildlife Diseases* 21: 11–19.
- MORAVEC, F. 1998. Nematodes of freshwater fishes of the Neotropical region, 1st Edition. Academia, Prague, Czech Republic. 464 pp.
- PEDÓ, E., A. C. TOMAZZONI, S. M. HARTZ, AND A. U. CHRISTOFF. 2006. Diet of crab-eating fox, *Cerdocyon thous* (Linnaeus) (Carnivora, Canidae) in a suburban area of southern Brazil. *Revista Brasileira de Zoologia* 23: 637–641.
- PEREIRA, B. J., G. L. GIRARDELI, L. O. TRIVILIN, V. R. LIMA, L. C. NUNES, AND I. V. F. MARTINS. 2006. Ocorrência de dioctofimose em cães do município de Cachoeiro do Itapemirim, Espírito Santo, Brasil, no período de maio a dezembro de 2004. *Revista Brasileira de Parasitologia Veterinária* 15: 123–125.
- ROCHA, U. F., R. G. SERRA, AND R. GRECHI. 1965. Parasitismo por *Dioctophyma renale* (Goeze, 1782) em “preguia” *Choloepus didactylus* Linnaeus, 1758. *Revista da Faculdade de Farmácia e Bioquímica de São Paulo* 3: 325–334.
- RUAS, J. L., M. P. SOARES, N. A. R. FARIAS, AND J. G. W. BRUM. 2003. Infecção por *Capillaria hepatica* em carnívoros silvestres (*Lycalopex gymnocercus* e *Cerdocyon thous*) na região sul do Rio Grande do Sul. *Arquivos do Instituto Biológico* 70: 127–130.
- SANTOS, K. R., L. S. CATENACCI, M. M. PESTELLI, R. K. TAKAHIRA, R. S. LOPES, AND R. J. SILVA. 2003. First report of *Ancylostoma buckleyi* Le Roux and Biocca, 1957 (Nematoda: Ancylostomatidae) infecting *Cerdocyon thous* Linnaeus, 1766

- (Mammalia: Canidae) from Brazil. *Revista Brasileira de Parasitologia Veterinária* 12: 179–181.
- _____, _____, _____, _____, AND R. J. SILVA. 2004. First report of *Diphyllbothrium mansonii* (Cestoda, Diphyllbothridae) infecting *Cerdocyon thous* Linnaeus, 1766 (Mammalia, Canidae) from Brazil. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia* 56: 796–798.
- VICENTE, J. J., H. O. RODRIGUES, D. C. GOMES, AND R. M. PINTO. 1997. Nematóides do Brasil. Parte V; Nematóides de mamíferos. *Revista Brasileira de Zoologia* 14: 1–452.
- VIEIRA, F. M., J. L. LUQUE, AND L. C. MUNIZ-PEREIRA. 2008. Checklist of helminth parasites in wild carnivore mammals from Brazil. *Zootaxa* 1721: 1–23.
- WOODHEAD, A. E. 1950. Life history cycle of the giant kidney worm, *Diocetophyma renale* (Nematoda), of man and many other mammals. *Transactions of the American Microscopical Society* 69: 21–46.

Received for publication 27 March 2008.