

INVESTIGATION OF THE ENTERIC BACTERIA OF THE TESTUDINATA — I: OCCURRENCE OF THE GENERA Arizona, Citrobacter, Edwardsiella and Salmonella

Authors: JACKSON, MARGUERITE M., JACKSON, CRAWFORD G., and FULTON, MACDONALD

Source: Bulletin of the Wildlife Disease Association, 5(3) : 328-329

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-5.3.328>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

INVESTIGATION OF THE ENTERIC BACTERIA OF THE TESTUDINATA — I: OCCURRENCE OF THE GENERA *Arizona*, *Citrobacter*, *Edwardsiella* and *Salmonella*

An investigation of the enteric bacteria of wild and captive turtles was initiated early in 1968. Individuals belonging to 20 species, which represent 2 families and 10 genera have to date been found positive for the presence of H:S positive, urease negative enteric bacteria of the family Enterobacteriaceae. The genera *Arizona*, *Citrobacter*, *Edwardsiella* and *Salmonella* have been isolated.

McNeil and Hinshaw (1946, Amer. J. Vet. Res. 7(22): 62-63) were the first to report *Salmonella* in turtles. At autopsy of two Galapagos tortoises in the San Diego Zoo, *Salmonella newport* and *S. sandiego* were found extensively throughout the internal organs. A number of additional reports of the occurrence of *Salmonella* in turtles have been

made since that time. The most commonly investigated turtles have been members of the family Testudinidae, belonging to the genera *Pseudemys*, *Chrysemys* and *Testudo*. Kaufmann and Morrison (1966, Am. J. Epidem. 82(2): 364-370) summarized the literature with respect to *Salmonella* and *Arizona* in turtles, noting the public health significance. Kaplan (1957, Lab. Animal Care 7: 273-277) described a septicemia in turtles caused by *Citrobacter* (Bethesda-Ballerup Group). Jackson et al (1969, Assn. Southeastern Biologists Bull. 16(2): 55) were the first to report *Edwardsiella tarda* in turtles. Rudat et al (1966, Pathol. Microbiol. 29(5): 623-629) and Lie (1968, Arch. Hyg. Bakterirol. 152(2): 139-155) reported *Salmonella* in zoo collections in Switzerland and Germany, respectively.

Methods

Samples were collected by inserting a sterile swab directly into the cloaca. The swab was then incubated in Selenite-F enrichment medium at 35°C for 18 to 24 hours. Routine procedures and culture techniques using Desoxycholate Agar plates and Kligler's Iron Agar slants were employed to isolate H:S positive forms. Urease, ONPG (O-Nitrophenyl Beta-D-galactopyranoside), and indol tests were done on the isolates.

H:S positive, urease negative cultures were then serotyped.

Water samples were also occasionally tested from tanks holding turtles. 5 cc samples were collected with sterile pipettes and discharged into 5 cc double-strength Selenite-F enrichment medium. Turtles too small for a cloacal swab can be checked by collecting a fresh fecal pellet and discharging it into the Selenite-F medium.

Findings

Of 53 individuals designated as wild (having been in strict isolation in captivity for not more than 2 weeks and unfed), 8 (15%) were positive; of 123 captive individuals, 28 (23%) were positive. As is shown in Table 1, *Salmonella* (all forms) and *Citrobacter* (all forms) were the most commonly encountered bacteria, each being isolated from 9 species of turtles. *Arizona* was isolated from only a single turtle species.

Five individuals were positive for two of the types of bacteria tested for.

It is of ecological interest that there were several additional bacterial species present, though only a tentative identification is available as yet. Of the first 40 turtles examined, 29 (70%) were positive for *Aeromonas hydrophila*. Members of the genera *Proteus*, *Aerobacter*, *Escherichia*, *Pseudomonas*, as well as streptococci were provisionally identified

by colonial morphology and other reactions. Further work with these associated species has been deferred until the primary objectives of the study have been achieved.

It is anticipated that the accumulating data will shed light on questions regarding prevalence of infection, degree of pathogenicity, and geographic distribution.

TABLE 1. *Turtles with their associated bacteria (C = captive; W = wild)*

Arizona	<i>Salmonella bredeney</i>
<i>Chelydra serpentina</i> (C)	<i>Geochelone elegans</i> (C)
Citrobacter	<i>Salmonella infantis</i>
<i>Geochelone elegans</i> (C)	<i>Chelydra serpentina</i> (W)
Citrobacter (Bethesda-Ballerup Group)	<i>Salmonella livingstone</i>
<i>Chrysemys picta</i> (C)	<i>Terrapene carolina major</i> (C)
<i>Chrysemys picta dorsalis</i> (C)	<i>Salmonella miami</i>
<i>Geochelone denticulata</i> (C)	<i>Sternotherus minor minor</i> (W)
<i>Gopherus polyphemus</i> (C)	(3 individuals)
<i>Kinosternon leucostomum</i> (C)	<i>Salmonella newport</i>
(water sample)	<i>Pseudemys scripta</i> (C)
<i>Kinosternon subrubrum</i> (W)	<i>Salmonella thompson</i>
<i>Pseudemys floridana</i> (W)	<i>Geoemyda annulata</i> (C)
<i>Pseudemys scripta elegans</i> (C)	<i>Macrochelys temmencki</i> (C)
<i>Terrapene carolina bauri</i> (C)	<i>Pseudemys nelsoni</i> (C)
<i>Edwardsiella tarda</i>	<i>Pseudemys scripta ornata</i> (C)
<i>Chrysemys picta</i> (W)	<i>Terrapene carolina bauri</i> (C)
<i>Geoemyda annulata</i> (C)	<i>Salmonella</i> Group C ₁
<i>Pseudemys concinna suwanniensis</i> (C)	(serotype undetermined)
(3 individuals)	<i>Geochelone denticulata</i> (C)
<i>Pseudemys nelsoni</i> (C)	<i>Geochelone elegans</i> (C)
<i>Pseudemys scripta</i> (W & C)	<i>Salmonella</i> Group E
<i>Pseudemys scripta ornata</i> (C)	(serotype undetermined)
<i>Sternotherus minor minor</i> (C)	<i>Geochelone denticulata</i> (C)
<i>Terrapene carolina major</i> (C)	
<i>Terrapene ornata</i> (C)	

Acknowledgments

We gratefully acknowledge the serotyping of the cultures by R. H. Andrews, Director of Laboratories, Mississippi State Board of Health. Appreciation is expressed to the following zoo personnel who allowed us to examine specimens in their care: Robert O. Wagner, Director, Livingstone Park Zoo, Jackson, Mississippi; Robert Mattlin, Director, and Tommy O'Brien,

Curator of Reptiles, Overton Park Zoo, Memphis, Tennessee; Robert Truett, Director, Ronald Westmoreland, and James Lowery, Jimmy Morgan Zoo, Birmingham, Alabama. BBL Laboratories provided media for this investigation. This research was supported in part by MSCW Faculty Research Grand funds.

MARGUERITE M. JACKSON,
CRAWFORD G. JACKSON, JR.
and

MACDONALD FULTON,
Department of Biological Sciences,
Mississippi State College for Women,
Columbus, Mississippi 39701.