

SEASONAL CHANGES IN THE DIFFERENTIAL COUNT IN FEMORAL MARROW OF THE HARE (*Lepus europaeus* PALL.)

Authors: PUJMAN, VOJTECH, and HANUSOVA, DANIELA

Source: Journal of Wildlife Diseases, 6(1) : 69-70

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-6.1.69>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

SEASONAL CHANGES IN THE DIFFERENTIAL COUNT IN FEMORAL MARROW OF THE HARE (*Lepus europaeus* PALL.)

Hematological examinations of certain game animals showed that in partridges and hares the hemoglobin levels, erythrocyte counts and packed cell volume values increased during the winter season (Novakova, 1968. Jour. Forest. Suisse, 31-44; Pujman, 1966. Symposium on partridge, VULHM CSMS, 114-117).

Experiments were conducted to determine if seasonal changes also occurred in the femoral myelogram. Because of availability the hare was chosen as the species to be investigated. Bone marrows obtained from 9 adult (mature) hares shot in December, 15 shot in June and July, and 9 shot in August, September,

TABLE 1. Mean percentages of cells in the femoral myelogram of the hare.

	December	June July	August September October
Number of hares	9	15	9
Myeloblast	0.3	0.5	0.5
Promyelocyte	2.2	2.1	2.9
Myelocyte	2.3	3.9	8.0
Metamyelocyte	2.1	5.3	7.2
Neutrophil, Band	3.2	6.8	7.4
Neutrophil, Mature	6.9	19.2	16.0
Eosinophilic Myelocyte	0.6	0.8	1.7
Eosinophilic Metamyelocyte	0.6	0.7	1.7
Eosinophil, Band	0.2	0.5	0.3
Eosinophil, Mature	0.2	0.5	0.3
Basophilic Myelocyte	0.1	0.1	0.1
Mitotic Granulocytic Cells	0.5	0.7	0.7
Megakaryocyte	0.2	0.6	0.4
Agranulocytic and Retic. Cells	5.6	6.1	3.3
Proerythroblast	0.7	0.4	0.2
Macroblast	0.6	0.1	0.3
Basophilic Normoblast	5.2	3.3	2.6
Polychromatophil Normoblast	20.3	20.2	14.9
Orthochromatophil Normoblast	47.0	27.5	31.9
Mitotic Erythroid Cells	1.3	0.6	0.1

and October, were examined, with no regard to sex. Femoral marrow smears were obtained immediately after the shooting and stained with the panoptic stain according to Pappenheim. In each smear, 500 cells were evaluated. As illustrated in Table 1, the average differential marrow counts, obtained from hares shot in December, had significantly lower granulocytic counts ($P.05$) than those examined between June and the end of October. This decrease included all developmental forms belonging to the neutrophilic granulocyte series, commencing with the myelocyte. The percentage of eosinophils was 1.4% in the December hares, 2.5% in the early summer group and 4% in the late summer-early fall hares. The mean differential erythroid counts also are illustrated in

Table 1. All developmental forms of the erythroid series appeared more numerous in the December marrow than in the marrow of the other two groups, however, a statistically significant difference was found only in the orthochromic normoblasts. The Myeloid : Erythroid (M:E) ratio for the December group was 1:4; for the early summer group it was 1:1.27; and for the August to October group, it was 1:1.06.

The results of these preliminary examinations indicated that the relative enhancement of erythropoiesis which occurred in the femoral marrow of the December hares, might be one of the factors responsible for the observed peripheral blood changes, i.e. increased Hb and PCV values.

VOJTECH PUJMAN and

DANIELA HANUSOVA

*Pharmaceutical and Biochemical
Research Institute
Praha, Czechoslovakia*

July 31, 1969
