

## **SOME FACTORS CONTRIBUTING TO THE DEVELOPMENT OF FUNGUS INFECTION IN FRESHWATER FISH**

Author: ROTH, RENE R.

Source: Journal of Wildlife Diseases, 8(1) : 24-28

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-8.1.24>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## SOME FACTORS CONTRIBUTING TO THE DEVELOPMENT OF FUNGUS INFECTION IN FRESHWATER FISH

RENE R. ROTH, Department of Zoology, University of Western Ontario, London, Ontario.

**Abstract:** The intravenous administration of 1 mg of: cortisol, cortisone, estradiol or progesterone, or the infusion of 5 i.u./kg body weight/24<sup>h</sup> of thyroid stimulating hormone, had a facilitating effect upon fungus growth on the freshwater teleost *Catostomus commersonii commersonii* Lacépède. The same dose (1 mg) of corticosterone or testosterone or adrenocorticotrophin (5 i.u./kg/24<sup>h</sup>) did not have this facilitating effect.

The possible role of steroids in the mechanism of action of temperature in promoting infections in fish is discussed.

### INTRODUCTION

The majority of fungi involved in diseases of fish are members of the family *Saprolegniaceae*, the common "water moulds". They are characterized by a myceloid, eucarpic thallus bearing numerous reproductive cells. The most ubiquitous of the aquatic fungi, commonly attacking fish, is the eurytherm, euryhaline genus *Saprolegnia*.

Most authors agree that *Saprolegnia* is not in itself a specific disease but rather makes its appearance on individuals weakened by other maladies.<sup>15</sup> Stuart and Fuller<sup>22</sup> however, do not accept the view that the fungal infection is secondary. They state that *Saprolegnia* can act as a lethal primary parasite of healthy fish and that this genus alone has been isolated constantly from lesions of fish with ulcerative dermal necrosis (UDN).

Thus, the predisposing factors, contributing to development of fungus on freshwater fish are still controversial. One precondition appears to be the presence of bruises, wounds or abraded surfaces which provide the substratum for the growth of the fungus. But the importance of other predisposing factors such as stress<sup>23</sup> and water temperature cannot be ignored.

Since stress causes the release of stress hormones (e.g. cortisone) from the interrenal tissue<sup>23</sup> it seems possible that corti-

costeroids are involved in the etiology of fungus growth. Robertson *et al.*<sup>17</sup> showed that intraperitoneal implantation of cortisol pellets to immature rainbow trout resulted in skin infections and death. This appears to support the view that experimental administration of corticosteroids might have an effect similar to stress itself.

### MATERIALS AND METHODS

Adult white suckers (*Catostomus commersonii commersonii* Lacépède), weighing between 860 and 1150 g, were kept in large holding tanks using recycled, filtered, dechlorinated, aerated water. The temperature was held at 5 C during winter and 10 C during summer. The fish were not fed.

The operative procedures were similar to those used by Hunn *et al.*<sup>10</sup> and Mackay and Beatty.<sup>12</sup> The experimental fish were anaesthetized with 0.2 g/l M.S. 222 (tricaine methanesulfonate, Sandoz) and subsequently catheterized with a Clay-Adams PE 200 polyethylene catheter introduced into the mesonephric duct and sutured into place with monofilament nylon. A PE 50 polyethylene tubing was introduced into the caudal vein for the purpose of injecting hormonal compounds. Following cannulation and catheterization each fish was placed in a specially built lucite box provided with

This paper was presented at the Conference of the International Association of Aquatic Animal Medicine, University of Guelph, Guelph, Ontario, Canada, April 29-30, 1971.

constant temperature water. Both the catheter and the cannula were led out of the box through watertight fittings.

Some of the experimental fish were kept at 5 C (group E<sub>1</sub>) while another group was kept at 13 C (E<sub>2</sub>). They were injected once with 1 mg of one of the following pure, crystalline steroids: corticosterone, cortisone or cortisol (male and female fish), estradiol or progesterone (females only), or testosterone (males only). The particular crystalline steroid was dissolved in 0.03 ml purified ethanol and 1 ml of a mixture of: 2 parts Hickman's fish saline<sup>6</sup> and 1 part propylene-glycol, was added. This solution was injected through the caudal cannula. One ml of saline was used to wash the solution into the blood stream.

Since the average weight of the fish was 1010 g, the one mg steroid injected gave an average dose level of 0.99 mg/kg body weight (range: 0.87 - 1.16 mg/kg) and an estimated mean blood level (assuming blood volume approximately 5% of body weight) of 1980 µg% (range: 1730 - 2320 µg%).

Some of the experimental animals were treated with adrenocorticotrophin (ACTH) or thyroid stimulating hormone (TSH). ACTH or TSH were administered as continuous infusion (0.005 i.u./g body weight /24h) by means of a syringe-pump, for periods of 5 or 8 days.

There were two major groups of control fish: group A consisted of the intact fish kept in the holding tanks (group A<sub>1</sub> at 5 C and A<sub>2</sub> at 10 C); group B comprised fish submitted to the same operative procedures as the experimental animals, except that they were injected with the solvent mixture only, without added steroids. The group B fish were subdivided into smaller groups and kept at the following temperatures: 5 C (subgroup B<sub>1</sub>), 10 C (B<sub>2</sub>), 13 C (B<sub>3</sub>) and 16 C (B<sub>4</sub>).

## RESULTS

During preparatory work it was noted that most of the fish kept at temperatures exceeding 10 C developed fungus growth. Based on this experience and assuming that initially the fungus invades only the

very surface of the skin, where it is still accessible to topical treatment, each newly caught fish was dipped for 30 sec. into a trough containing a 1% solution of malachite green and 0.5 g/gal M.S. 222 (to keep the fish quiet) cooled to 5 C. This treatment was repeated once more, just prior to cannulation and catheterization.

Fungus development was absent on the fish kept intact in the holding tanks (Table 1) as well as on the control groups B<sub>1</sub>, B<sub>2</sub> and B<sub>3</sub> even 30 days after cannulation and catheterization. Group B<sub>4</sub>, however, presented fungus growth after a latency period of 10-15 days following operation. The experimental fish (E<sub>1</sub> and E<sub>2</sub>) treated with estradiol, cortisol, cortisone or progesterone as well as those treated with TSH consistently developed fungus infection after a latency period of 7-10 days. Conversely, those injected with corticosterone or testosterone or infused with ACTH did not show any sign of fungus growth.

These results were the same for fish caught ripe in April and for those caught during the summer and winter months.

## DISCUSSION

The mechanism of action of temperature in promoting infections in fish is still a matter of controversy. The majority of authors<sup>1,14,23</sup> reported a better immune response at higher than at lower water temperatures, and the rate of healing of UDN lesions was found to be faster at higher temperature.<sup>10</sup> On the other hand bacterial infections may be caused<sup>3</sup> and an epizooty accentuated<sup>13</sup> by increased water temperature. Cushing<sup>5</sup> reported that although the antibody-titer in freshwater fish increased faster at 28 C than at 15 C, the end-titer was higher at 15 C.

That the link between temperature and immunity control might be found in the adrenal cortical hormones was suggested by Bisset<sup>4</sup> who pointed out that the production of these hormones is inhibited at low temperatures. In agreement with this is the work of Fontaine and Hately<sup>9</sup> who reported that ACTH restored to normal the interrenal of the

TABLE 1. Fungus growth on white suckers kept at various water temperatures, prophylactically treated with malachite green and injected (or infused) with various hormonal compounds.

Fish group	Sex	Water temp.	Operative procedure	Presence (+) or absence (—) of fungus growth																																																					
A <sub>1</sub>	M,F	5 C	None	—																																																					
A <sub>2</sub>	M,F	10 C	None	—																																																					
Injected with solvent mixture only																																																									
B <sub>1</sub>	M,F	5 C	cannulated and catheterized	—																																																					
B <sub>2</sub>	M,F	10 C	idem	—																																																					
B <sub>3</sub>	M,F	13 C	idem	—																																																					
B <sub>4</sub>	M,F	16 C	idem	+																																																					
Infused with																																																									
<table><tr><th colspan="4">Injected with</th><th colspan="3">Infused with</th></tr><tr><th>estra-diols</th><th>cortico-sterone</th><th>cortisone</th><th>cortisol</th><th>testo-sterone</th><th>proge-sterone</th><th>TSH</th><th>ACTH</th></tr><tr><td></td><td>—</td><td>+</td><td>+</td><td>—</td><td></td><td>+</td><td>—</td></tr><tr><td>+</td><td>—</td><td>+</td><td>+</td><td></td><td>+</td><td>+</td><td>—</td></tr><tr><td></td><td>—</td><td>+</td><td>+</td><td>—</td><td></td><td>+</td><td>—</td></tr><tr><td>+</td><td>—</td><td>+</td><td>+</td><td></td><td>+</td><td>+</td><td>—</td></tr></table>											Injected with				Infused with			estra-diols	cortico-sterone	cortisone	cortisol	testo-sterone	proge-sterone	TSH	ACTH		—	+	+	—		+	—	+	—	+	+		+	+	—		—	+	+	—		+	—	+	—	+	+		+	+	—
Injected with				Infused with																																																					
estra-diols	cortico-sterone	cortisone	cortisol	testo-sterone	proge-sterone	TSH	ACTH																																																		
	—	+	+	—		+	—																																																		
+	—	+	+		+	+	—																																																		
	—	+	+	—		+	—																																																		
+	—	+	+		+	+	—																																																		
E <sub>1</sub>	M	5 C	idem																																																						
E <sub>1</sub>	F	5 C	idem																																																						
E <sub>2</sub>	M	13 C	idem																																																						
E <sub>2</sub>	F	13 C	idem																																																						

hypophysectomized eel at 16 C but not at 6 C. Robertson *et al.*<sup>17</sup> found that when the water temperature was raised from 13 to 17 C, death occurred much faster and with smaller doses in fish implanted with cortisol pellets.

In the present study temperatures exceeding 10 C appeared to make the suckers more susceptible to fungus infection than lower temperatures. This might have been the result of a disruption of the balance between host and bacteria causing infections predisposing to fungus development<sup>2</sup> or of the increased invasive activity of the fungus. Nevertheless, based on the findings mentioned above<sup>4,17</sup> and on the results of the present study, it is suggested that temperature acts mainly indirectly, through humoral factors, upon the immunological defense mechanisms of the fish. High temperatures will reduce the amount of transcortin-bound corticosteroids in fish plasma<sup>7</sup> and release free, biologically active steroids.<sup>20</sup> Thus temperature may be considered a permissive element which allows the activation of the immunodepressant interrenal corticosteroids.<sup>23</sup>

Although the glucocorticoids and particularly the 17-hydroxy-corticosteroids (17-OHCS) are traditionally known as restrainers of protein synthesis and stimulators of the catabolism of the amino acid building blocks of protein and thus of antibodies, nothing of the kind is known about estrogens or progesterone. On the contrary, estradiol is known to stimulate the synthesis of RNA and proteins in mammals. In the present study, however, estradiol administration was followed by fungus growth in the experimental fish. It seems possible that this

effect was due to the triggering of free cortisol secretion, a supposition suggested by my own finding that estradiol injections caused excretion of free cortisol in the white sucker and by the report that humans treated with estrogens showed a marked increase of plasma 17-OHCS.<sup>19</sup>

The amounts of steroids injected in the present study were pharmacological rather than physiological, resulting in an initial blood level ten times higher than that found in spawned salmon plasma.<sup>11</sup> Although this high plasma level is maintained for a short time only, substantial amounts of the injected steroids or their metabolites may be detected in various organs of these fish even after 5 days from the time of injection.<sup>14</sup> The effects elicited by the presence of these high levels of steroids may be equated with those of stress<sup>23</sup> which, through a very similar mechanism, causes metabolic changes contributing to increased susceptibility to infection and significant reduction of immune response.<sup>24</sup>

ACTH did not show any effect upon fungus growth facilitation, in the present study. Whether this was due to the relatively low dosage used or to some other factor, as e.g. the possible formation of anti-ACTH antibodies during protracted treatment, is unknown. The role of ACTH in the various fish species needs much more study.

In conclusion it may be asserted that high plasma levels of 17-OHCS, estradiol, progesterone or TSH have a permissive, facilitating effect upon the occurrence of fungus infection in the white sucker fish. It is suggested that this effect is related to the impairment of antibody formation<sup>9</sup> and the suppression of reactive tissue inflammation.

#### LITERATURE CITED

1. AMBROSIUS, H., and R. LEHMANN. 1965. Beitrage zur Immunobiologie poikilothermer Wirbeltiere. III. Der Einfluss von Adjuvanten auf die Antikörper-produktion. *Acta biol. med. German.* 14: 830-844.
2. BISSET, K. A. 1946. The effect of temperature on non-specific infections of fish. *J. Path. Bact.* 58: 251-258.
3. BISSET, K. A. 1948. The effect of temperature upon antibody production in cold-blooded vertebrates. *J. Path. Bact.* 60: 87-92.
4. BISSET, K. A. 1949. The influence of adrenal cortical hormones upon immunity in cold-blooded vertebrates. *J. Endocrinol.* 6: 99-103.

5. CUSHING, J. E. 1942. An effect of temperature upon antibody-production in fish. *J. Immun.* 45: 123-126.
6. FONTAINE, M., and J. HATEY. 1953. Recherches sur le controle hypophysaire de l'interrenal anterieur d'un poisson teleosteen, l'anguille. *C. R. Soc. Biol. Paris* 147: 217-220.
7. FREEMAN, H. C., and D. R. IDLER. 1966. Transcortin binding of cortisol in Atlantic salmon (*Salmo salar*) plasma. *Gen. Comp. Endocrinol.* 7: 37-43.
8. HOAR, W. S., and C. P. HICKMAN, JR. 1967. *A Laboratory Companion for General and Comparative Physiology*. Prentice Hall, Inc., New Jersey.
9. HODKINSON, M., and A. HUNTER. 1970. Immune response of UDN-infected salmon to *Saprolegnia*. *J. Fish Biol.* 2: 305-311.
10. HUNN, J. B., R. A. SCHOETTGER, and W. A. WILLFORD. 1968. Turnover and urinary excretion of free and acetylated MS 222 by rainbow trout, *Salmo gairdneri*. *J. Fish. Res. Bd. Can.* 25: 25-31.
11. IDLER, D. R., A. P. RONALD, and P. J. SCHMIDT. 1959. Biochemical studies on sockeye salmon during spawning migration. VII. Steroid hormones in plasma. *Can. J. Biochem. Physiol.* 37: 1227-1238.
12. MACKAY, W. C., and D. D. BEATTY. 1968. The effect of temperature on renal function in the white sucker fish, *Catostomus commersonii*. *Comp. Biochem. Physiol.* 26: 235-245.
13. MARKOFF, W., and Z. K. JATSCHEWA. 1939. *Zbl. Bakt. Abt. II, c*, 194 (quoted in Bisset, K. A. 1946).
14. POST, G. 1963. The immune response of rainbow trout (*Salmo gairdneri*) to *Aeromonas hydrophila*. Utah State Dept. Fish and Game Pub. #63-7.
15. REICHENBACH-KLINKE, H., and E. ELKAN. 1965. *The Principal Diseases of Lower Vertebrates*. Academic Press, N.Y.
16. ROBERTS, R. J. 1971. Studies on ulcerative dermal necrosis of salmonids. III. The healing process in fish maintained under experimental conditions. *J. Fish Biol.* 3: 221-224.
17. ROBERTSON, O. H., S. HANE, B. C. WEXLER, and A. R. RINFRET. 1963. The effect of hydrocortisone on immature rainbow trout. *Gen. Comp. Endocrinol.* 3: 422-436.
18. ROTH, R. R. 1969. The renal and branchial imbibition and excretory indexes in *Catostomus commersonii* treated with <sup>3</sup>H- labelled and carrier steroids. *Am. Zool.* 9: 573-574.
19. SANDBERG, A. A., and W. R. SLAUNWHITE, JR. 1959. Transcortin: a corticosteroid binding protein of plasma. II. Levels in various conditions and the effects of estrogens. *J. Clin. Invest.* 38: 1290-7.
20. SANDBERG, A. A., and W. R. SLAUNWHITE, JR. 1963. Transcortin: a corticosteroid binding protein of plasma. V. *In vitro* inhibition of cortisol metabolism. *J. Clin. Invest.* 42: 51-54.
21. SMITH, W. W. 1940. Production of antibacterial agglutinins by carp and trout at 10 C. *Proc. Soc. Exptl. Biol. Med.* 45: 726-729.
22. STUART, M. R., and H. T. FULLER. 1968. Mycological aspects of diseased Atlantic salmon. *Nature* 217: 90-92.
23. WEDEMEYER, G. 1970. The role of stress in the disease resistance of fishes. In: *A Symposium on Diseases of Fishes and Shellfishes*, Snieszko, S. F., ed. Fish. Soc. Sp. Pub. No. 5: 30-35.

Received for publication May 10, 1971