



Serologic Survey of Selected Pathogens in Free-Ranging Bengal Tigers (*Panthera tigris tigris*) in Nepal

Authors: McCauley, Deborah, Stout, Virginia, Gairhe, Kamal P., Sadaula, Amir, Dubovi, Edward, et al.

Source: Journal of Wildlife Diseases, 57(2) : 393-398

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/JWD-D-20-00046>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Serologic Survey of Selected Pathogens in Free-ranging Bengal Tigers (*Panthera tigris tigris*) in Nepal

Deborah McCauley,¹ Virginia Stout,^{1,6} Kamal P. Gairhe,² Amir Sadaula,³ Edward Dubovi,⁴ Suraj Subedi,⁵ and Gretchen E. Kaufman¹ ¹Veterinary Initiative for Endangered Wildlife, 1627 W Main St., Suite 445, Bozeman, Montana 59715, USA; ²Government of Nepal Department of National Parks and Wildlife Conservation, PO Box 860, Kathmandu, Nepal; ³National Trust for Nature Conservation, Biodiversity Conservation Center, PO Box 3712, Khumaltar, Lalitpur, Nepal; ⁴Cornell University College of Veterinary Medicine, 602 Tower Rd., Ithaca, New York 14853, USA; ⁵Government of Nepal, Ministry of Agriculture and Livestock Development, Department of Livestock Services, Veterinary Standards and Drug Regulatory Laboratory, Golfutar Main Rd., Budhanilkantha, Kathmandu, Nepal; ⁶Corresponding author (email: ginger.stout@viewwildlife.org)

ABSTRACT: Serum samples of 11 Bengal tigers (*Panthera tigris tigris*) from Chitwan National Park in Nepal, collected between 2011–17, were evaluated for the presence of antibodies to eight diseases commonly investigated in large felids. This initial serologic survey was done to establish baseline information to understand the exposure of Nepal's free-ranging tiger population to these diseases. Tiger serum samples collected opportunistically during encounters such as translocation, human conflict, and injury were placed in cold storage for later use. Frozen serum samples were assessed for feline coronavirus (FCoV), feline immunodeficiency virus, feline leukemia virus, feline herpesvirus (FHV), canine distemper virus, canine parvovirus-2 (CPV-2), leptospirosis (LEP; seven serovars), and toxoplasmosis (TOX). Six tigers were found to be positive for LEP, eight for CPV-2, five for FHV, one for FCoV, and 10 for TOX. Tigers, like other wild felids, have been exposed to these common pathogens, but further research is needed to determine the significance of these pathogens to the Nepali population.

Key words: Bengal tiger, canine parvovirus, feline coronavirus, feline herpes virus, leptospirosis, Nepal, *Panthera tigris tigris*, toxoplasma.

Asia's tiger (*Panthera tigris*) populations are in decline across their range. As of April 2016, the global wild tiger population estimation was around 3,890, occupying about 7% of its historical range, mostly found in small pockets with <25 animals (Walston et al. 2010). Threats to tigers include habitat loss or fragmentation, decrease in prey density, wildlife trade and poaching, human-wildlife conflict, and infectious diseases (Dinerstein et al. 2007). Infectious disease is recognized as an increasing concern for wild carnivore populations globally, contributing to signifi-

cant declines. Canine distemper virus (CDV) in African wild dogs (*Lycaon pictus*) and lions (*Panthera leo*) as well as feline leukemia virus (FeLV) in Iberian lynx (*Lynx pardinus*) has led to several of these declines (Roelke-Parker et al. 1996; Meli et al. 2009; Woodroffe et al. 2012).

In Nepal, the tiger population has been growing, with the most recent report of 235 individuals, an increase of 19% since 2013 (National Trust for Nature Conservation 2018). Little is known about the diseases that affect tigers in Nepal. Because there has been recent evidence of diseases such as canine distemper in tigers in other areas of Asia and Russia, it is important to determine which infectious diseases may threaten the Nepali population to minimize the potential impacts on population recovery (Goodrich et al. 2012; Gilbert et al. 2015). Using serology, we determined baseline exposure to pathogens in Bengal tigers (*Panthera tigris tigris*) in Chitwan National Park including feline coronavirus (FCoV), feline immunodeficiency virus (FIV), FeLV, feline herpesvirus (FHV), CDV, canine parvovirus-2 (CPV-2), leptospirosis (LEP; seven serovars), and toxoplasmosis (TOX).

The study was conducted in Chitwan National Park in the Terai region of south-central Nepal (Fig. 1). Blood samples from 11 adult tigers were collected by opportunistic sampling from 2011–17 when tigers were captured, translocated, or held in captivity for various injuries or illnesses. Many were human-conflict tigers and were captured near human settlements. Captures were led by the




FIGURE 1. The study area includes Chitwan National Park and surrounding regions in south-central Nepal where a serologic survey of selected pathogens in Bengal tigers (*Panthera tigris tigris*) was done. Triangles represent locations of tiger captures and sampling during 2011–17.

Nepal Department of National Parks and Wildlife Conservation and National Trust for Nature Conservation. The tigers were captured by remote injection, anesthetized according to published protocol, and aged by tooth eruption and wear (Kreeger and Arneimo 2012). Blood samples were collected and the serum was extracted by centrifugation and stored in a -20 C freezer at the National Trust for Nature Conservation, Biodiversity Conservation Center in Sauraha, Nepal.

After obtaining US (no. 16US75654B/9) and Nepal (no. 1479876) Convention on International Trade in Endangered Species of Wild Fauna and Flora permits, serum samples were transported and evaluated at the New York State Diagnostic Laboratory at Cornell University in Ithaca, New York. Serum samples were evaluated using enzyme-linked immunosorbent assay to detect antibodies to FCoV, FIV, and TOX, and antigens to FeLV (Schaefer et al. 2011); serum neutralization to detect antibodies to FHV and CDV (Sheldon et al. 2017); hemagglutination inhibition to detect anti-

bodies to CPV-2 (Carmichael et al. 1980); and a microscopic agglutination test to detect antibodies to Pomona, Hardjo, Ictero, Grip-po, Canicola, Autumnalis, and Bratislava serovars of *Leptospira interrogans* (Cole et al. 1973).

Five of eight of the pathogens that we tested for had seropositive results and all but one of the tigers were seropositive for at least one of the pathogens (Table 1). All of the tigers were adults and therefore could not be compared based on their age. Seventy-three percent (8/11) of tigers tested positive for CPV-2 antibodies (with a titer ≥ 80), 45% (5/11) for FHV antibodies (with a titer of ≥ 8), 9% (1/11) for FCoV antibodies (with a titer ≥ 1.20), 91% (10/11) for *Toxoplasma gondii* antibodies (with a titer ≥ 1.20), and 55% (6/11) positive for LEP antibodies, including all serovars (with a titer >100). The highest serovar prevalences were Autumnalis with 36% positive, Hardjo with 27% positive, and Ictero with 18% positive. All samples were seronegative for FIV, FeLV, and CDV.

TABLE 1. Diagnostic summary of the serologic survey of selected pathogens in 11 adult Bengal tigers (*Panthera tigris tigris*) surveyed in and around Chitwan National Park, Nepal from 2011–17.

Pathogen	No. seropositive tigers (n=11)			
	Male (n=9)	Female (n=2)	Total n positive	% Positive
Feline coronavirus	0	1	1	9
Feline immunodeficiency virus	0 ^a	0	0	0
Feline leukemia virus	0	0	0	0
Feline herpesvirus	3	2	5	45
Canine distemper virus	0	0	0	0
Canine parvovirus	6	2	8	73
Leptospira total	5	1	6	55
Leptospira Pomona	1	0	1	9
Leptospira Hardjo	3	0	3	27
Leptospira Ictero	2	0	2	18
Leptospira Grippo	0	0	0	0
Leptospira Canicola	1	0	1	9
Leptospira Autumnalis	3	1	4	36
Leptospira Bratislava	1	0	1	9
Toxoplasmosis	8	2	10	91

^a One animal gave an equivocal positive result = suspected positive but not confirmed.

Canine parvovirus-2 antibody prevalence was high in sampled tigers. Canine parvovirus is closely related antigenically to feline panleukopenia virus (Ikeda et al. 2002). Canine parvovirus easily spreads in the environment, causing high morbidity and mortality with clinical signs including enteritis, depression, dehydration, and anorexia. A study on leopard cats (*Prionailurus bengalensis*) in Vietnam and Taiwan found that the viral isolates were more closely related to CPV in dogs than in domestic cats; therefore, it is more likely originating from dogs (*Canis lupus familiaris*) or wild carnivores than from domestic cats (*Felis catus*; Ikeda et al. 2002). The presence of a large, free-roaming domestic dog population in Chitwan, Nepal, along with wild carnivore populations in the park, could explain the high prevalence in tigers (Massei et al. 2017). A phylogenetic analysis of the virus in tigers, domestic dogs, and wild carnivores is needed to confirm this hypothesis in Nepal.

About one third of the tigers in this study were seropositive for FHV. The most common clinical signs in large felids are upper respiratory symptoms and rhinotracheitis, which

have been described in captive felids including cheetahs (*Acinonyx jubatus*), mountain lions (*Puma concolor*), jaguars (*Panthera onca*), and leopards (*Panthera pardus*; Hofmann-Lehmann et al. 1996). In Nepal, the origin and reservoir of FHV is unknown and further research is needed to determine how the tigers were exposed. This virus may be endemic in the population and may not cause many clinical signs or mortalities, as suggested with a study of lions in East Africa (Hofmann-Lehmann et al. 1996).

Feline coronavirus occurs in domestic and wild felids around the world. Feline coronavirus can induce feline infectious peritonitis or can cause mild clinical signs of enteritis (Hofmann-Lehmann et al. 1996). One tiger in this study was seropositive for FCoV. The tiger was captured from a community forest, was geriatric, severely dehydrated, weak, and cachectic. The clinical signs exhibited by this female could possibly be due to FCoV, as the titer was elevated, but this tiger was also seropositive for CPV, FHV and TOX, and coinfections could have contributed to morbidity (Table 2).

TABLE 2. A summary of coinfections of selected pathogens and reasons for capture of 11 adult Bengal tigers (*Panthera tigris tigris*) opportunistically sampled in Chitwan National Park and surrounding areas, Nepal from 2011–17.

Tiger no.	Sample year	Sex	Reason for capture	Coinfections ^a
1	2011	Male	Human conflict	LEP, TOX
2	2011	Male	Livestock predation, injury	FHV, CPV, LEP, TOX
3	2012	Male	Injury	LEP, TOX
4	2012	Male	Human conflict	CPV, LEP, TOX
5	2013	Female	Human conflict	FCoV, FHV, CPV, TOX
6	2013	Male	Predation	FHV, CPV, TOX
7	2013	Male	Livestock predation	FHV, CPV, LEP, TOX
8	2014	Male	Livestock predation, human conflict	CPV, TOX
9	2015	Male	Unknown	Negative
10	2016	Male	Unknown	CPV, TOX
11	2017	Female	Unknown	FHV, CPV, LEP, TOX

^a LEP = leptospirosis (at least one of seven serovars); TOX = toxoplasmosis; FHV = feline herpesvirus; CPV = canine parvovirus-2; FCoV = feline coronavirus.

Almost all the tigers in this study were seropositive for TOX. *Toxoplasma gondii* is an intracellular protozoan that can infect humans and livestock, although the definitive host is domestic and wild felids (Furtado et al. 2015). Toxoplasmosis, transmitted via the fecal-oral route, is prevalent worldwide, is zoonotic, and can cause a variety of serious diseases in humans, livestock, and other wildlife (Tenter et al. 2000). *Toxoplasma gondii* is present wherever felids live, usually does not cause clinical signs, and prevalence can be up to 98% in a population (Spencer et al. 2003). Our results are consistent with other studies performed on wild felids (Goodrich et al. 2012; Furtado et al. 2015). This suggests that *T. gondii* is endemic in this population, most likely not causing clinical signs of disease but contributing to the maintenance of the pathogen in the study area.

About half of the tigers tested in this study were seropositive for at least one of the LEP serovars. *Leptospira* are Gram-negative bacteria, with small mammals considered the main reservoir, and are found most commonly in tropical areas that experience frequent flooding and high rainfall (Gamage et al. 2012). Leptospirosis is common in Nepal in domestic animals, humans, and wildlife and can be transmitted between them through

indirect or direct contact of urine (Shrestha et al. 2018). Known serovars found in humans in Nepal are Bratislava, Icterohaemorrhagiae, Autumnalis, and Sejroe (Gamage et al. 2012). One study on the *Leptospira* serovar Hardjo in large ruminants in Nepal revealed that prevalence in large ruminants in Nepal was 6.88% after the monsoon (Khanal et al. 2018). In our study of tigers, the most common serovars were Autumnalis, Hardjo, and Icterohaemorrhagiae. These serovars are important in human and livestock health, with human-livestock-wildlife interaction most likely playing a role in prevalence. It is unclear if LEP causes significant clinical disease in tigers, but exposure rates are high, most likely due to high small mammal populations, livestock predation, seasonal rain and flooding, and contaminated waterways.

Our baseline study revealed exposure of tigers to several important pathogens in Nepal, some of which are shared with humans and domestic animals. Further research is needed to determine the significance and origins of these pathogens. Although there is no known exposure to CDV, FIV, and FeLV, it is important to continue surveillance for these pathogens. It is possible that these viruses were not detected due to the small sample size available for the study. If the

population is truly naïve, then CDV and other pathogens could cause significant mortalities in the future as in other populations (Roelke-Parker et al. 1996; Meli et al. 2009; Woodroffe et al. 2012). Continued health surveillance of the tiger population in Nepal, including routine serology testing and necropsy information, is needed to support the growth of endangered populations and to develop policies to minimize morbidity and mortality from these diseases in the future.

We thank the staff at Nepal Department of National Parks and Wildlife Conservation (DNPWC), National Trust for Nature Conservation (NTNC), and Chitwan National Park for their contribution to this project in collecting and storing samples. We thank Babu Ram Lamichhane and Carl Brown for map design. We also thank Dennis and Connie Keller for supporting this project.

LITERATURE CITED

- Carmichael LE, Joubert JC, Pollock RV. 1980. Hemagglutination by canine parvovirus: Serologic studies and diagnostic applications. *Am J Vet Res* 41:784–791.
- Cole JR, Sulzer CR, Pursell AR. 1973. Improved microtechnique for the leptospiral microscopic agglutination test. *Appl Microbiol* 25:976–980.
- Dinerstein E, Loucks C, Wikramanayake E, Ginsberg J, Sanderson E, Seidensticker J, Forrest J, Bryja G, Heydlauff A, Klenzendorf S, et al. 2007. The fate of wild tigers. *BioScience* 57:508–514.
- Furtado MM, Gennari SM, Ikuta CY, de Almeida Jácomo AT, de Morais ZM, de Jesus Pena HF, de Oliveira Porfirio GE, Silveira L, Sollmann R, de Souza GO, et al. 2015. Serosurvey of smooth *Brucella*, *Leptospira* spp. and *Toxoplasma gondii* in free-ranging jaguars (*Panthera onca*) and domestic animals from Brazil. *PLoS One* 10:e0143816.
- Gamage CD, Tamashiro H, Ohnishi M, Koizumi N. 2012. Epidemiology, surveillance and laboratory diagnosis of leptospirosis in the WHO South-East Asia Region. In: *Zoonosis*, Lorenzo-Morales J, editor. InTech, Rijeka, Croatia, pp. 213–226.
- Gilbert M, Soutyrina SV, Seryodkin IV, Sulikhan N, Uphyrkina OV, Goncharuk M, Matthews L, Cleaveland S, Miquelle DG. 2015. Canine distemper virus as a threat to wild tigers in Russia and across their range. *Integr Zool* 10:329–343.
- Goodrich JM, Quigley KS, Lewis JCM, Astafiev AA, Slabi EV, Miquelle DG, Smirnov EN, Kerley LL, Armstrong DL, Quigley HB, et al. 2012. Serosurvey of free-ranging Amur tigers in the Russian Far East. *J Wildl Dis* 48:186–189.
- Hofmann-Lehmann R, Fehr D, Grob M, El Gixoli M, Packer C, Martenson JS, O'Brien SJ, Lutz H. 1996. Prevalence of antibodies to feline parvovirus, calicivirus, herpesvirus, coronavirus, and immunodeficiency virus and of feline leukemia virus antigen and the interrelationship of these viral infections in free-ranging lions in East Africa. *Clin Diag Lab Immunol* 3:554–562.
- Ikeda Y, Nakamura K, Miyazawa T, Tohya Y, Takahashi E, Mochizuki M. 2002. Feline host range of canine parvovirus: Recent emergence of new antigenic types in cats. *Emerg Infect Dis* 8:341–346.
- Khanal DR, Paudyal N, Khanal S, Prajapati M, Shrestha P, Bowen R, Acharya MP, Shrestha SP, Singh UM, Thakur RP, et al. 2018. Detection of antibodies against *Leptospira hardjo* in large ruminants of Nepal. *Acta Sci Agricult* 2:131–133.
- Kreeger TJ, Armento JM. 2012. *Handbook of wildlife chemical immobilization*. 4th Ed. Terry Kreeger, Wheatland, Wyoming, 301 pp.
- Massei G, Fooks AR, Horton DL, Callaby R. 2017. Free-roaming dogs in Nepal: Demographics, health and public knowledge, attitudes and practices. *Zoonoses Public Health* 64:29–40.
- Meli ML, Cattori V, Martinez F, Lopez G, Vargas A, Simon MA, Zorrilla I, Munoz A, Palomares F, Lopez-Bao JV, et al. 2009. Feline leukemia virus and other pathogens as important threats to the survival of the critically endangered Liberian lynx (*Lynx pardinus*). *PLoS One* 4:e4744.
- National Trust for Nature Conservation. 2018. *Nepal set to become the first country to double its tigers*. <https://www.ntnc.org.np/news/nepal-set-become-first-country-double-its-tigers>. Accessed February 2019.
- Roelke-Parker ME, Munson L, Packer C, Kock R, Cleaveland S, Carpenter M, O'Brien SJ, Pospischil A, Hofmann-Lehmann R, Lutz H, et al. 1996. A canine distemper virus epidemic in Serengeti lions (*Panthera leo*). *Nature* 379:441–445.
- Schaefer JJ, White HA, Schaaf SL, Mohammed HO, Wade SE. 2011. Modification of a commercial *Toxoplasma gondii* immunoglobulin G enzyme-linked immunosorbent assay for use in multiple animal species. *J Vet Diagn Invest* 23:297–301.
- Sheldon JD, Cushing AC, Wilkes RP, Anis E, Dubovi EJ. 2017. Serologic response to canine distemper vaccination in captive Linnaeus's two-toed sloths (*Choloepus didactylus*) after a fatal canine distemper virus outbreak. *J Zoo Wildl Med* 48:1250–1253.
- Shrestha R, McKenzie JS, Gautam M, Adhikary R, Pandey K, Koiraal P, Bahadur BC, Miller LC, Collins-Emerson J, Craig SB, et al. 2018. Determinants of clinical leptospirosis in Nepal. *Zoonoses Public Health* 65:972–983.
- Spencer JA, Higginbotham MJ, Blagburn BL. 2003. Seroprevalence of *Neospora caninum* and *Toxoplasma gondii* in captive and free-ranging nondomestic felids in the United States. *J Zoo Wildl Med* 34:246–249.

- Tenter AM, Heckeroth AR, Weiss LM. 2000. *Toxoplasma gondii*: From animals to humans. *Int J Parasitol* 30: 1217–1258.
- Walston J, Robinson JG, Bennett EL, Breitenmoser U, da Fonseca GAB, Goodrich J, Gumal M, Hunter L, Johnson A, Karanth KU, et al. 2010. Bringing the tiger back from the brink—The six percent solution. *PLoS Biol* 8:e1000485.
- Woodroffe R, Prager KC, Munson L, Conrad PA, Dubovi EJ, Mazet JAK. 2012. Contact with domestic dogs increases pathogen exposure in endangered African wild dogs (*Lycan pictus*). *PLoS One* 7:1–9.

Submitted for publication 26 March 2020.

Accepted 25 July 2020.