

## **GATA4 Deficiency Impairs Ovarian Function in Adult Mice 1**

Authors: Kyrönlahti, Antti, Vetter, Melanie, Euler, Rosemarie, Bielinska, Malgorzata, Jay, Patrick Y., et al.

Source: *Biology of Reproduction*, 84(5) : 1033-1044

Published By: Society for the Study of Reproduction

URL: <https://doi.org/10.1095/biolreprod.110.086850>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# GATA4 Deficiency Impairs Ovarian Function in Adult Mice<sup>1</sup>

Antti Kyrönlahti,<sup>3,4,9</sup> Melanie Vetter,<sup>3,4,7</sup> Rosemarie Euler,<sup>4,7</sup> Malgorzata Bielinska,<sup>4</sup> Patrick Y. Jay,<sup>4,5</sup> Mikko Anttonen,<sup>8</sup> Markku Heikinheimo,<sup>4,9</sup> and David B. Wilson<sup>2,4,6</sup>

Departments of Pediatrics,<sup>4</sup> Genetics,<sup>5</sup> and Developmental Biology,<sup>6</sup> Washington University and St. Louis Children's Hospital, St. Louis, Missouri  
Hochschule Mannheim—University of Applied Sciences,<sup>7</sup> Mannheim, Germany  
Department of Obstetrics and Gynecology<sup>8</sup> and Children's Hospital,<sup>9</sup> Biomedicum Helsinki, University of Helsinki and Helsinki University Central Hospital, Helsinki, Finland

## ABSTRACT

Transcription factor GATA4 is expressed in granulosa cells and, to a lesser extent, in other ovarian cell types. Studies of mutant mice have shown that interactions between GATA4 and its cofactor, ZFPM2 (also termed FOG2), are required for proper development of the fetal ovary. The role of GATA4 in postnatal ovarian function, however, has remained unclear, in part because of prenatal lethality of homozygous mutations in the *Gata4* gene in mice. To circumvent this limitation, we studied ovarian function in two genetically engineered mouse lines: C57BL/6 (B6) female mice heterozygous for a *Gata4*-null allele, and 129;B6 female mice in which *Gata4* is deleted specifically in proliferating granulosa cells using the Cre-loxP recombination system and *Amhr2-cre*. Female B6 *Gata4*<sup>+/-</sup> mice had delayed puberty but normal estrous cycle lengths and litter size. Compared to wild-type mice, the ovaries of gonadotropin-stimulated B6 *Gata4*<sup>+/-</sup> mice were significantly smaller, released fewer oocytes, produced less estrogen, and expressed less mRNA for the putative GATA4 target genes *Star*, *Cyp11a1*, and *Cyp19*. *Gata4* conditional knockout (cKO) mice had a more severe phenotype, including impaired fertility and cystic ovarian changes. Like *Gata4*<sup>+/-</sup> mice, the ovaries of gonadotropin-stimulated cKO mice released fewer oocytes and expressed less *Cyp19* than those of control mice. Our findings, coupled with those of other investigators, support the premise that GATA4 is a key transcriptional regulator of ovarian somatic cell function in both fetal and adult mice.

*developmental biology, fertility, gene regulation, granulosa cells, ovary, ovulation, puberty, transcription factor*

## INTRODUCTION

Transcription factor GATA4 is implicated in the development and function of the mammalian ovary [1–3]. GATA4 is expressed in pregranulosa and stroma cells of the fetal ovary

and in granulosa and theca cells of the postnatal ovary [4–6]. In adult mice, GATA4 is thought to play a role in the maintenance or maturation of granulosa cells; *Gata4* is expressed in potentially mitotic and proliferating granulosa cells but is downregulated when proliferation ceases during ovulation, apoptosis, or luteinization [4, 7]. Follicle-stimulating hormone (FSH), the gonadotropin that initiates follicular growth, increases both the expression level and the intrinsic activity of GATA4 in granulosa cells [4, 8, 9]. Conversely, loss-of-function mutations in the FSH receptor are associated with reduced ovarian expression of GATA4 in both mice [10] and humans [11]. GATA4 binds and activates genes essential for ovarian steroidogenesis, including *Star* [12, 13], *Cyp11a1* [14], and *Cyp19* (aromatase) [9, 15–18].

Analysis of genetically engineered mice has shown that interactions between GATA4 and its cofactor, ZFPM2 (also termed FOG2), are necessary for early ovarian development [3, 19]. *Zfpm2*<sup>-/-</sup> mice and *Gata4*<sup>ki/ki</sup> mice, which bear a knockin mutation that abrogates the interaction of GATA4 with ZFPM2, exhibit identical fetal ovarian phenotypes [3, 19] that include decreased expression of genes essential for early ovarian development, such as *Wnt4* [20] and *Foxl2* [21]. The ovaries of *Zfpm2*<sup>-/-</sup> and *Gata4*<sup>ki/ki</sup> fetal mice also express excess DKK1, a secreted inhibitor of the canonical WNT/β-catenin pathway, a signaling cascade critical for female gonadogenesis [3, 19].

Although GATA4 appears to be essential for proper fetal ovarian development, the role of this transcription factor in postnatal ovarian function remains unclear, in part because of prenatal lethality of homozygous mutations in the *Gata4* gene in mice [22–27]. Adult transgenic mice expressing a tetracycline-inducible small interfering RNA (siRNA) directed against GATA4 appear to have intact ovarian steroidogenesis, insofar as circulating levels of FSH and luteinizing hormone (LH) are normal [28]. Approximately 10% of GATA4 siRNA transgenic mice develop ovarian teratomas, suggesting that GATA4 functions as a tumor suppressor in this tissue [28].

In the present study, we examine the impact of GATA4 deficiency on ovarian physiology using two complementary mouse models: C57BL/6 (B6) female mice heterozygous for a germline deletion mutation in *Gata4*, and 129;B6 female mice in which *Gata4* is conditionally ablated in proliferating granulosa cells using the Cre-loxP recombination system.

## MATERIALS AND METHODS

### Experimental Mice

Procedures involving mice were approved by the Washington University Institutional Committee for Laboratory Animal Care and were conducted in

<sup>1</sup>Supported by NIH DK075618 and DK52574, Academy of Finland, and the Sigrid Juselius Foundation.

<sup>2</sup>Correspondence: David B. Wilson, Box 8208, Washington University School of Medicine, 660 S. Euclid Ave., St. Louis, MO 63110.

FAX: 314 286 2892; e-mail: wilson\_d@wustl.edu

<sup>3</sup>These authors contributed equally to the work.

Received: 21 June 2010.

First decision: 30 July 2010.

Accepted: 10 January 2011.


© 2011 by the Society for the Study of Reproduction, Inc.

This is an Open Access article, freely available through *Biology of Reproduction's* Authors' Choice option.

eISSN: 1529-7268 <http://www.biolreprod.org>

ISSN: 0006-3363

FIG. 1. Delayed puberty in B6 *Gata4*<sup>+/-</sup> female mice. **A)** Representative estrous cycles from WT and *Gata4*<sup>+/-</sup> mice. **B)** Average ages for first vaginal cornification and the onset of cyclicity. Note the significant delay in the onset of cyclicity in *Gata4*<sup>+/-</sup> mice. **C)** Proportion of time spent in each estrous stage (n = 10 for each genotype). Error bars represent SD. D, diestrus; E, estrus; M, metestrus; P, proestrus.



accordance with the National Research Council publication *Guide for Care and Use of Laboratory Animals*. All mice had free access to water and standard rodent chow and were exposed to 12L:12D photoperiods. *Gata4*<sup>+/-</sup> mice, originally produced by William Pu, were genotyped as described previously [29, 30]. These mice were backcrossed with B6 mice (Charles River Laboratories) for a minimum of 10 generations. *Gata4*<sup>Flox/Flox</sup> mice (also termed *Gata4*<sup>tm1.1Sad/J</sup>), originally generated by Stephen Duncan [27, 31], were purchased from the Jackson Laboratory and genotyped as described previously [27, 31]. The 129;B6 *Amhr2*<sup>cre/+</sup> mice (also termed B6;129S7-*Amhr2*<sup>tm3(cre)Blhr/J</sup> Mmnc), originally produced by Richard Behringer, were obtained from the Mutant Mouse Regional Resource Centers (MMRRC) and genotyped as described previously [32, 33]. *ROSA26* Flox-stop-Flox *LacZ* reporter mice (also termed B6.129S4-*Gt(ROSA)26Sor*<sup>tm1Sor/J</sup>), developed by P. Soriano [34], were obtained from the Jackson Laboratory. To generate conditional knockout (cKO) mice, *Gata4*<sup>Flox/Flox</sup> mice were mated with *Amhr2*<sup>cre/+</sup> mice, and the resultant *Gata4*<sup>Flox/+;Amhr2</sup><sup>cre/+</sup> mice were mated with *Gata4*<sup>Flox/Flox</sup> mice to produce *Gata4*<sup>Flox/Flox;Amhr2</sup><sup>cre/+</sup> mice.

### Measurement of Pubertal Events

Daily vaginal lavage was begun on the day of vaginal opening and continued for the duration of the present study [35]. Cytologic analysis was performed on air-dried, Giemsa-stained smears. Estrous cycle stage was determined by the presence of nucleated epithelial cells (NEC), cornified epithelial cells (CEC), and polymorphonuclear cells (PMN), with estrus characterized by CEC and some NEC without PMN, metestrus by CEC and PMN, diestrus by PMN with few NEC, and proestrus mainly by NEC and few CEC and PMN. Ambiguous cycle status was ascertained based on determinations for the preceding and subsequent day and were labeled as an intermediate value. The age of first vaginal cornification and the onset of estrous cyclicity were determined as described previously [35]. Cycle length was defined as the time between onsets of estrus.

### Fertility Studies

Reproductive performance was assessed by housing female mice with proven male stud mice and measuring litter size over successive pregnancies.

Because some *Gata4*<sup>+/-</sup> mice die within days of birth because of developmental defects in the heart, lungs, or diaphragm [36–38], we assessed litter size on the morning of birth.

### Tissue Isolation and Histological Analyses

At specified times, mice were killed by carbon dioxide inhalation. Blood was collect by cardiac puncture. Ovaries or uteri were harvested, fixed overnight in 4% paraformaldehyde in PBS, and embedded in paraffin for routine histology. Paraffin sections (thickness, 5  $\mu$ m) were stained with hematoxylin-and-eosin or subjected to immunoperoxidase staining with goat anti-mouse GATA4 immunoglobulin (Ig) G (sc-1237; Santa Cruz Biotechnology, Inc.) at a 1:200 dilution. The secondary antibody employed for immunoperoxidase staining was donkey anti-goat biotinylated IgG (Jackson ImmunoResearch) at a 1:1000 dilution. The avidin-biotin immunoperoxidase system (Vectastain Elite ABC Kit; Vector Laboratories, Inc.) and diaminobenzidine (Sigma-Aldrich Corp.) were used to visualize the bound antibody; slides were then counterstained with toluidine blue. For X-gal staining, frozen tissue sections (thickness, 10  $\mu$ m) were prepared after embedding mouse ovaries in O.C.T. (Tissue-Tek). Sections were fixed with 0.2% glutaraldehyde for 10 min; permeabilized with 100 mM potassium phosphate (pH 7.4), 0.02% NP-40, and 0.01% sodium deoxycholate for 5 min; and then incubated in 0.5 mg/ml of X-gal (Sigma-Aldrich Corp.) with 10 mM  $K_3[Fe(CN)_6]$ , 10 mM  $K_4[Fe(CN)_6]$ , 100 mM potassium phosphate (pH 7.4), 0.02% NP-40, and 0.01% sodium deoxycholate at 37°C overnight [25]. X-gal-stained sections were counterstained with eosin.

TABLE 1. Normal litter size in *Gata4*<sup>+/-</sup> female mice.<sup>a</sup>

Genotype	Litter no. 1	Litter no. 2	Litter no. 3
WT	6.0 $\pm$ 1.9 (n = 8)	6.6 $\pm$ 2.1 (n = 8)	7.4 $\pm$ 1.2 (n = 7)
<i>Gata4</i> <sup>+/-</sup>	5.7 $\pm$ 1.4 (n = 6)	7.0 $\pm$ 2.2 (n = 6)	6.8 $\pm$ 0.8 (n = 5)

<sup>a</sup> None of the differences was statistically significant.




FIG. 2. Ostensibly normal follicular development in the ovaries of randomly cycling adult *Gata*<sup>+/-</sup> mice. **A–C** Ovaries from 3-mo-old WT (**A** and **B**) or *Gata4*<sup>+/-</sup> (**C**) mice were subjected to immunoperoxidase staining to assess ovarian morphology and to visualize the distribution of GATA4 within this tissue. Note that GATA4 is expressed in granulosa and theca cells associated with primordial, preantral, and antral follicles of both WT and *Gata4*<sup>+/-</sup> mice. Bar = 75  $\mu$ m. **D** and **E** Histomorphometric analysis of follicle development in WT (n = 10) and *Gata4*<sup>+/-</sup> (n = 6) mice. Serial ovarian sections from 7-wk-old WT and *Gata4*<sup>+/-</sup> mice were processed, and the numbers of healthy (nonatretic; **D**) or atretic (**E**) follicles were estimated. Error bars represent SD. AnF, antral follicles; CL, corpora lutea; PF, primordial and primary follicles; PrF, preantral follicles.

### Ovarian Follicle Counts

Oocyte-containing follicles were scored as described previously [39, 40]. Briefly, ovaries were collected from mice shortly after the onset of puberty, fixed (0.34 N glacial acetic acid, 28% ethanol, and 10% formalin), embedded in paraffin, serially sectioned (thickness, 6  $\mu$ m), and stained with hematoxylin-picric acid methyl blue. The number of ostensibly healthy and atretic primordial, primary, and small preantral follicles was determined in every fifth section through the entire ovary. Primordial follicles contained a compact oocyte enveloped by a single layer of flattened granulosa cells. Primary follicles had an enlarged oocyte surrounded by a single layer of cuboidal granulosa cells. Small preantral follicles had an enlarged oocyte surrounded by two to four layers of cuboidal granulosa cells but no visible antrum. Only those follicles in which the oocyte nucleus was visible were scored, and follicles were designated atretic if the oocyte was degenerated, using criteria delineated elsewhere [39, 40]. The total number of healthy and atretic follicles per ovary was then calculated as described previously [39, 40].

### Gonadotropin and Estradiol Treatment


Immature (age, 19–25 days) mice were induced to ovulate using equine chorionic gonadotropin (eCG; 5 IU i.p.), followed by human chorionic gonadotropin (hCG; 5 IU i.p.) 48 h later. Four treatment groups were analyzed: 1) no stimulation; 2) eCG for 48 h (follicular growth); 3) eCG for 48 h, followed by hCG for 18 h (postovulation); and 4) eCG for 48 h, followed by hCG for 5 days (luteal glands). Harvested ovaries were processed for histological analysis or RNA isolation. To quantify ovulation, mice were killed 18 h after hCG administration, and oocytes were isolated from oviducts [41].

### RNA Isolation and First-Strand cDNA Synthesis

Total RNA was isolated from ovaries using TRIzol (Invitrogen). First-strand cDNA was produced with the SuperScript VILO cDNA Synthesis Kit (Invitrogen) using 700 ng of RNA as a template. An aliquot (1.5  $\mu$ l) of cDNA was subjected to real-time RT-PCR using SYBR GreenER qPCR SuperMix (Invitrogen). Conditions for all quantitative RT-PCR (qRT-PCR) reactions were optimized in a light cycler (Stratagene Mx3005), which was programmed as follows: denaturation at 95°C for 10 min, followed by 40 cycles of amplification and quantification at 95°C for 30 sec and 60°C for 1 min with single fluorescent measurement, followed by melting at 60–95°C with a heating rate of 0.1°C/sec and continuous fluorescence measurement, followed by a final cooling step to 55°C. Melting curves did not reveal any significant contamination. The relative expression of target genes was calculated using the relative standard curve method as described in *Guide to Performing Relative Quantitation of Gene Expression Using Real-Time Quantitative PCR* (Applied Biosystems) and in our prior publication [42].

To compensate for variation among runs, the target gene expression was normalized to the expression of ribosomal protein L19. Primer pairs used for qRT-PCR assays were as follows: *L19*, NM\_005084, (forward) 5'-GAAATCGCCAATGCCAACTC-3', (reverse) 5'-TCTTAGACCTGCGAG CCTCA-3', 405 bp; *Cyp19*, NM\_007810, (forward) 5'-ATGTTCTTG GAAATGCTGAACCC-3', (reverse) 5'-AGGACCTGGTATTGAAGA CGAG-3', 150 bp; *Star*, NM\_011485, (forward) 5'-GCAGCAGGCAAC CTGGTG-3', (reverse) 5'-TGATTGTCTTCGGCAGCC-3', 246 bp; *Cyp11a1*, NM\_19779.3, (forward) 5'-CGATACTTCTCATGCGAG-3', (reverse) 5'-CTTCTTCCAGGCATCTGAAC-3', 125 bp; *Gata4*, NM\_008092.3, (for-

FIG. 3. Impaired ovarian response to gonadotropins in *Gata*<sup>+/-</sup> mice. **A**) Weanling WT (solid line) or *Gata*<sup>+/-</sup> (dashed line) mice were treated with one of the following regimens: 1) no stimulation; 2) eCG for 48 h; 3) eCG for 48 h, followed by hCG for 18 h; or 4) eCG for 48 h, followed by hCG for 5 days. Ovaries were harvested and weighed ( $n = 8$  for each genotype). \* $P < 0.05$ . Error bars represent SD. **B** and **C**) Tissue sections of WT (**B**) and *Gata*<sup>+/-</sup> (**C**) ovaries 48 h after eCG stimulation. The *Gata*<sup>+/-</sup> ovary is smaller and contains fewer antral follicles. Bar = 0.3 mm. **D**) Oocyte yields after gonadotropin-induced ovulation. WT and *Gata*<sup>+/-</sup> mice were treated with eCG, followed 48 h later by hCG. Oocytes were harvested from oviducts and counted 16 h after hCG treatment. Each data point represents the oocyte yield of an individual mouse. Horizontal lines represent mean values. **E** and **F**) Sections of WT (**E**) and *Gata*<sup>+/-</sup> (**F**) uteri 48 h after eCG stimulation. Note the hypoestrogenic appearance of the heterozygous uterus. Glandular elements of the endometrial layer are less complex, and only scattered glands are detected in the stroma. No difference was found in mean body weight between the WT and *Gata*<sup>+/-</sup> mice ( $13.8 \pm 1.6$  and  $14.4 \pm 1.6$  g, respectively). Bar = 0.3 mm. **G**) Serum E<sub>2</sub> levels in WT (white bar,  $n = 7$ ) and *Gata*<sup>+/-</sup> (black bar,  $n = 7$ ) uteri 48 h after eCG stimulation. \* $P < 0.05$ . Error bars represent SD.



ward 5'-CCCTACCCAGCCTACATGG-3', (reverse) 5'-ACATATCGAGAT TGGGGTGTCT-3', 138 bp.

#### Semiquantitative RT-PCR for Detection of Cre-Mediated Recombination

First-strand cDNA was subjected to RT-PCR using a forward primer from exon 2 of the *Gata4* gene, 5'-CCCTACCCAGCCTACATGG-3', and a reverse primer from exon 7 of the same gene, 5'-GAGCTGGCCTGCGATGTCTGATG-3', and the following reaction conditions: denaturation at 95°C for 10 min; followed by 35 cycles of amplification and quantification at 95°C for 30 sec, 55°C for 1 min, and 72°C for 1 min; followed by a final cooling step to 55°C. The intact *Gata4*<sup>Flox</sup> allele gave rise to a 782-bp band, whereas the recombined allele yielded a 401-bp band, reflecting deletion of exons 3–5.

#### Hormone Measurements

Serum estradiol (E<sub>2</sub>) levels were determined using a commercial radioimmunoassay (BioCheck). Serum antimüllerian hormone (AMH) levels were measured by ELISA as described elsewhere [43].

#### In Situ Hybridization

Tissue cryosections (thickness, 10 μm) were processed and subjected to mRNA in situ hybridization as described previously [25]. The sections were incubated with  $1.2 \times 10^6$  cpm [<sup>33</sup>P]αUTP-labeled (1000–3000 Ci/mmol; Amersham) antisense riboprobe in a total volume of 80 μl. The riboprobe for mouse GATA4 was prepared as described previously [25].

#### Statistical Analyses

The effects of genotype on the age of first vaginal cornification, onset of estrous cyclicity, estrous cycle length, litter size, ovarian weight, uterine weight, steroid hormone levels, and prevalence of infertility were determined by unpaired, two-tailed *t*-tests. Statistical significance was set at  $P < 0.05$ . Two-way ANOVA was used to analyze the proportion of time in each estrous stage, follicle counts, and gene expression levels. Two-population-proportion testing was used to analyze the incidence of cystic ovarian changes. All numerical data are presented as the mean  $\pm$  SD.

## RESULTS

### Delayed Puberty but Normal Estrous Cycle Length and Litter Size in *Gata4*<sup>+/-</sup> Female Mice

We first analyzed B6 mice heterozygous for a deletion in exon 2 of the *Gata4* gene [29]. This germline mutation removes the translation start site and N-terminal activation domain of GATA4. Studies have shown that mice heterozygous for this allele, hereafter referred to as *Gata4*<sup>+/-</sup> mice, express wild-type (WT) *Gata4* mRNA at half-normal levels in heart and other tissues in which this transcription factor is expressed [29, 36].

Vaginal cytology specimens were analyzed to determine the impact of *Gata4* haploinsufficiency on the timing of two pubertal events: the first vaginal cornification and the onset of estrous cyclicity [44]. Previous studies have shown that vaginal

cornification is dependent on an increase in circulating  $E_2$ , whereas the onset of estrous cyclicity is dependent on both a preovulatory increase in plasma  $E_2$  and the ability of the pituitary to respond with a preovulatory surge of LH, which stimulates ovulation and luteinization [44]. Representative estrous cycles from WT and *Gata4*<sup>+/-</sup> females are shown in Figure 1A. The onset of vaginal cornification appeared delayed in *Gata4* heterozygotes compared to WT mice ( $50 \pm 6.8$  vs.  $45 \pm 5.5$  days,  $n = 10$  per group), although this difference did not reach statistical significance (Fig. 1B). The onset of estrous cyclicity was delayed an average of 12 days in *Gata4*<sup>+/-</sup> females compared with WT females ( $P < 0.01$ ) (Fig. 1B). Average cycle length, however, did not differ between *Gata4*<sup>+/-</sup> and WT females ( $5.3 \pm 0.5$  and  $5.4 \pm 0.4$  days, respectively), nor did *Gata4*<sup>+/-</sup> and WT mice differ in the proportion of time spent in any stage of the estrous cycle (Fig. 1C). The timing of puberty is known to correlate with mouse growth [44]. WT and *Gata4*<sup>+/-</sup> female mice had similar growth curves, implying that the delay in puberty in *Gata4*<sup>+/-</sup> females was not the result of impaired growth (data not shown). We conclude that *Gata4* is a genetic determinant of the onset of puberty in female B6 mice.

The reproductive performance of *Gata4*<sup>+/-</sup> female mice was assessed by mating 2-mo-old WT or *Gata4*<sup>+/-</sup> female mice with normal males and measuring litter size over successive pregnancies. No differences were found between WT and *Gata4*<sup>+/-</sup> mice in either litter size (Table 1) or the frequency of parturition (data not shown).

Ovaries from young adult mice were examined to determine whether *Gata4* haploinsufficiency disrupts follicular development. In both WT (Fig. 2, A and B) and *Gata4*<sup>+/-</sup> (Fig. 2C) females, nuclear GATA4 immunoreactivity was evident in theca cells and in granulosa cells of primary, preantral, and antral follicles (Fig. 2, A–C), but not in atretic follicles or corpora lutea (Fig. 2, A and C). Consistent with a published report [45], no significant differences were found between WT and *Gata4*<sup>+/-</sup> ovaries in the abundance of primordial, primary, or small preantral follicles (healthy or atretic) (Fig. 2, D and E).

#### Ovaries of *Gata4*<sup>+/-</sup> Mice Exhibit an Impaired Response to Exogenous Gonadotropins

Immature WT and *Gata4*<sup>+/-</sup> female mice were treated with eCG, which mimics the effect of FSH, followed by hCG, which mimics the effect of LH, to induce synchronized follicular growth and ovulation. Unstimulated WT and *Gata4*<sup>+/-</sup> ovaries had similar weights, whereas gonadotropin-stimulated WT ovaries weighed significantly more than their *Gata4*<sup>+/-</sup> counterparts (Fig. 3A). Histological analysis confirmed that ovaries of eCG-stimulated WT mice (Fig. 3B) were consistently larger than those of *Gata4*<sup>+/-</sup> mice (Fig. 3C).

In response to superovulation, *Gata4*<sup>+/-</sup> mice released significantly fewer oocytes into the oviducts than WT mice (Fig. 3D). Thus, *Gata4* haploinsufficiency decreases oocyte release in response to exogenous gonadotropins without impairing fertility.

Compared to WT mice, the uteri of eCG-stimulated *Gata4*<sup>+/-</sup> mice weighed significantly less ( $0.32 \pm 0.05$  [ $n = 8$ ] vs.  $0.20 \pm 0.4$  [ $n = 7$ ] mg/g body weight,  $P < 0.05$ ) and appeared to be hypoestrogenic (Fig. 3, E and F). Following eCG stimulation, serum  $E_2$  levels were significantly greater in WT mice than the *Gata4*<sup>+/-</sup> mice (Fig. 3G). In contrast, basal serum  $E_2$  levels did not differ significantly between WT ( $8.8 \pm 1.1$  pg/ml) and *Gata4*<sup>+/-</sup> mice ( $9.5 \pm 0.3$  pg/ml).

Quantitative RT-PCR was used to measure the relative expression of transcripts for *Gata4* (Fig. 4A) and three of its





FIG. 4. Reduced expression of mRNA for *Gata4* and steroidogenic factors in the ovaries of gonadotropin-stimulated *Gata4*<sup>+/-</sup> mice. Weanling WT (solid line,  $n = 5$ ) or *Gata4*<sup>+/-</sup> (dashed line,  $n = 5$ ) mice were treated with one of the following regimens: 1) no stimulation; 2) eCG for 48 h; 3) eCG for 48 h, followed by hCG for 18 h; or 4) eCG for 48 h, followed by hCG for 5 days. Ovarian RNA was subjected to qRT-PCR for *Gata4* (A), *Star* (B), *Cyp11a1* (C), and *Cyp19* (D). Error bars represent SD. \* $P < 0.05$ , \*\* $P < 0.01$ .

putative target genes, *Star* (Fig. 4B), *Cyp11a1* (Fig. 4C), and *Cyp19* (Fig. 4D), in ovaries from gonadotropin-stimulated immature WT and *Gata4*<sup>+/-</sup> mice. To confirm a reduction in *Gata4* expression in the heterozygous ovaries, we performed qRT-PCR using primers specific for the WT *Gata4* mRNA. The ovaries of unstimulated and gonadotropin-stimulated *Gata4*<sup>+/-</sup> mice contained approximately half the amount of *Gata4* mRNA (normalized to the housekeeping gene *L19*) as their WT counterparts (Fig. 4A). Basal levels of mRNA for *Star* (Fig. 4B), *Cyp11a1* (Fig. 4C), and *Cyp19* (Fig. 4D) were similar in WT and *Gata4*<sup>+/-</sup> mice. Following gonadotropin stimulation, however, the levels of *Star*, *Cyp11a1*, and *Cyp19* mRNA were significantly higher in WT than in *Gata4*<sup>+/-</sup> ovaries. The target gene most profoundly affected by *Gata4* haploinsufficiency was *Cyp19*, the key gene of estrogen

FIG. 5. *Gata4* mRNA is not expressed in the müllerian duct or adult mouse uterus. Cryosections of Embryonic Day 18.5 mouse embryo (A and B) and Embryonic Day 9.5 gravid uterus (C and D) were subjected to in situ hybridization for GATA4 using radiolabeled riboprobe. Shown are corresponding bright-field (A and C) and dark-field (B and D) photomicrographs. Note that *Gata4* mRNA is expressed in fetal ovary (ov) and in endodermal cells of the small intestine (si) and yolk sac (ys), but not in the müllerian duct (md), embryonic ectoderm (em), decidua (de), or uterus (ut). Bar = 100  $\mu$ m.



biosynthesis. Previous studies have shown that CYP19 is essential for follicular growth and coordination of the ovulatory process [46–48].

#### Conditional Deletion of *Gata4* in Granulosa Cells

To further study the physiological function of GATA4 during folliculogenesis, we used the Cre-loxP recombination system to facilitate granulosa cell-specific deletion of the *Gata4* gene. We hypothesized that conditional deletion of both *Gata4* alleles in granulosa cells would produce a more severe reproductive phenotype than germline deletion of a single allele. The 129;B6 mice homozygous for a floxed allele of *Gata4* were intercrossed with 129;B6 *Amhr2*<sup>cre/+</sup> mice. Previous studies have shown that Cre-mediated excision of the region between exons 3 and 5 of the *Gata4* gene converts the floxed allele into a recombined allele no longer capable of encoding a functional GATA4 protein [27, 31]. The *Amhr2-cre* knockin transgene has been widely used to target gene deletion in proliferating granulosa cells and in the mesenchyme-derived cells of the müllerian duct, oviduct, and uterus [33, 49–58]. Because GATA4 is not expressed in the müllerian duct (Fig. 5, A and B), oviduct [59], or adult uterus (Fig. 5, C and D), Cre-mediated deletion of the *Gata4* gene should not have a direct impact on these tissues.

*ROSA26* flox-stop-flox *lacZ* reporter (*R26R*) mice were used to assess the expression of *Amhr2-cre* within ovaries. Consistent with published reports [33, 50, 52], the *Amhr2-cre* transgene directed expression of Cre to granulosa cells in secondary follicles (Fig. 6A). Interestingly, Cre expression within secondary follicles was not uniform, as evidenced by the presence of both lacZ-positive (Fig. 6A, arrow) and lacZ-negative (Fig. 6A, arrowhead) granulosa cells. Such variegated Cre expression may contribute to the variable penetrance described below.

Intercrosses of *Gata4*<sup>Flox/Flox</sup> mice and *Gata4*<sup>Flox/+</sup>;

*Amhr2*<sup>cre/+</sup> mice yielded (*Gata4*<sup>Flox/Flox</sup>;*Amhr2*<sup>cre/+</sup>) mice, henceforth referred to as cKO mice, and *Gata4*<sup>Flox/+</sup>;*Amhr2*<sup>cre/+</sup> mice in the expected mendelian ratios. The growth of cKO mice did not differ from that of *Gata4*<sup>Flox/+</sup>;*Amhr2*<sup>cre/+</sup> mice or *Amhr2*<sup>+/+</sup> controls (data not shown).

To verify Cre-mediated deletion of the *Gata4* gene in the mouse ovary, we employed an RT-PCR strategy that distinguishes transcripts derived from the intact (*Gata4*<sup>Flox</sup>) and recombined alleles (*Gata4* <sup>$\Delta$ ex3–5</sup>). RT-PCR analysis of ovaries from *Gata4*<sup>Flox/Flox</sup>;*Amhr2*<sup>+/+</sup> mice yielded a single band corresponding to the intact allele (Fig. 6B). In contrast, analysis of RNA from the ovaries of cKO mice demonstrated both the intact *Gata4*<sup>Flox</sup> allele and the recombined *Gata4* <sup>$\Delta$ ex3–5</sup> allele (Fig. 6B). The relative intensities of the *Gata4*<sup>Flox</sup> and *Gata4* <sup>$\Delta$ ex3–5</sup> bands differed among cKO mice (Fig. 6B). Variability in the extent of Cre-mediated recombination has been observed in other granulosa cell-specific knockout mice made with *Amhr2-cre* and may reflect inefficient recombination in granulosa cells (see [52] and references therein).

To confirm reduced expression of GATA4 in the cKO ovaries, we performed qRT-PCR using primers specific for the WT GATA4 mRNA. The ratio of GATA4 mRNA expression in immature cKO versus *Gata4*<sup>Flox/Flox</sup>;*Amhr2*<sup>+/+</sup> mice was  $0.44 \pm 0.02$  ( $P < 0.001$ ). Moreover, the ratio of GATA4 mRNA expression in immature cKO versus *Gata4*<sup>Flox/+</sup>;*Amhr2*<sup>cre/+</sup> mice was  $0.72 \pm 0.02$  ( $P < 0.05$ ).

Next, we compared the patterns of GATA4 immunostaining in ovaries of postpubertal *Gata4*<sup>Flox/+</sup>;*Amhr2*<sup>cre/+</sup> (Fig. 6C) and cKO (Fig. 6D) mice. In both genotypes, GATA4 protein was detected in theca cells and in granulosa cells of primary, preantral, and antral follicles. The residual GATA4 expression in the immunohistochemical analysis presumably reflects cells in which the *Amhr2-cre* did not fully inactivate both *Gata4*<sup>Flox</sup> alleles. Nevertheless, we infer from the reproductive defects described below that GATA4 expression is silenced in enough granulosa cells to elicit a phenotype.




FIG. 6. *Amhr2-cre*-mediated deletion of *Gata4<sup>Flox</sup>* in the mouse ovary. **A** *Amhr2<sup>cre/+</sup>* mice were crossed with homozygous *R26R* mice to generate *R26R<sup>+/-</sup>;Amhr2<sup>cre/+</sup>* mice. Ovaries were cryosectioned and stained with X-gal. Note the variegated pattern of Cre activity in a secondary follicle; both lacZ-positive (arrow) and lacZ-negative (arrowhead) granulosa cells are evident. **B** Ovarian RNA was isolated from pairs of 2-mo-old control (*Gata4<sup>Flox/Flox</sup>;Amhr2<sup>+/+</sup>*) or cKO mice and subjected to RT-PCR analysis with primers that distinguish the intact floxed allele from the recombined allele lacking exons 3–5 (*Gata4<sup>Δex3-5</sup>*). Note that a transcript derived from recombined allele is seen in the cKO mice but not the control mice. The identity the third band in the cKO mice is unclear. **C** and **D** Ovaries from 7-wk-old *Gata4<sup>Flox/+</sup>;Amhr2<sup>cre/+</sup>* (**C**) or cKO (**D**) mice were sectioned and subjected to immunoperoxidase staining to visualize the distribution of GATA4 within this tissue. GATA4 immunoreactivity was seen in granulosa and theca cells of both the *Gata4<sup>Flox/+</sup>;Amhr2<sup>cre/+</sup>* and cKO mice. Bar = 75 μm. AAnF, atretic antral follicles; AnF, antral follicles; PF, primordial and primary follicles; PrF, preantral follicles.

### Reduced Fertility in *Gata4* cKO Mice

To assess the impact of conditional deletion of *Gata4* on fertility, female 2-mo-old cKO and *Gata4<sup>Flox/+</sup>;Amhr2<sup>cre/+</sup>* mice were housed with proven male studs. Both strains gave birth and raised pups to weanlings, but fertility defects were evident in the cKO females. Six of 18 (33%) cKO females failed to produce offspring, compared to only 1 of 20 (5%) *Gata4<sup>Flox/+</sup>;Amhr2<sup>cre/+</sup>* females ( $P < 0.05$ ). Even more striking, the average litter size of cKO mice was reduced ( $P < 0.005$ ) to approximately half that of *Gata4<sup>Flox/+</sup>;Amhr2<sup>cre/+</sup>* mice (Table 2).

Although cKO mice were clearly subfertile, no differences were detected in the onset of puberty or in the length of the estrous cycle among cKO mice, *Gata4<sup>Flox/+</sup>;Amhr2<sup>cre/+</sup>* mice, and *Amhr2<sup>+/+</sup>* littermate mice (data not shown). That puberty was delayed in B6 *Gata4<sup>+/-</sup>* mice (Fig. 1) but not in 129;B6 cKO or *Gata4<sup>Flox/+</sup>;Amhr2<sup>cre/+</sup>* mice might reflect background strain influences [35] or the timing/efficiency of Cre-mediated ablation. Alternatively, the B6 *Gata4<sup>+/-</sup>* mice may have extragonadal defects that delay puberty.

Ovaries of young adult *Gata4<sup>Flox/+</sup>;Amhr2<sup>cre/+</sup>* (Fig. 7A) and cKO (Fig. 7B) mice were analyzed to determine the impact of conditional ablation of *Gata4* on follicular development. On gross inspection, the ovaries of young adult *Gata4<sup>Flox/+</sup>;Amhr2<sup>cre/+</sup>* and cKO mice were indistinguishable. On histological examination, the number of primordial, primary, and small antral follicles (healthy or atretic) was similar in cKO and *Gata4<sup>Flox/+</sup>;Amhr2<sup>cre/+</sup>* ovaries (Fig. 7, C and D).

By 6 mo of age, however, differences were found in the gross and microscopic appearance of ovaries from *Gata4<sup>Flox/+</sup>;Amhr2<sup>cre/+</sup>* (Fig. 7, E and G) and cKO (Fig. 7, F and H) mice. Very large, nonhemorrhagic cysts were seen in six of seven (86%) ovaries from cKO mice and in 0 of 10 (0%) ovaries from *Gata4<sup>Flox/+</sup>;Amhr2<sup>cre/+</sup>* mice ( $P < 0.001$ ). The cysts were lined by a flattened epithelium containing scattered ciliated cells (data not shown).

Next, we measured serum AMH levels in *Gata4<sup>Flox/+</sup>;Amhr2<sup>cre/+</sup>* and cKO mice at different ages (Fig. 7I). Studies of WT mice have shown that serum levels of AMH, a protein secreted by granulosa cells within growing follicles [60], remain relatively constant until 8 mo of age and then decline

TABLE 2. Reduced litter size in B6;129 *Gata4* cKO mice.

Genotype	Litter no. 1	Litter no. 2	Litter no. 3
<i>Gata4<sup>Flox/+</sup>; Amhr2<sup>cre/+</sup></i>	8.0 ± 2.8 (n = 13) <sup>a</sup>	9.4 ± 2.0 (n = 12)	10.3 ± 2.2 (n = 9)
cKO	4.3 ± 2.7* (n = 14)	4.2 ± 2.0** (n = 12)	3.7 ± 2.4* (n = 7)

<sup>a</sup> The litter size of *Gata4<sup>Flox/+</sup>;Amhr2<sup>cre/+</sup>* mice did not differ from that of *Amhr2<sup>+/+</sup>* controls (8.6 ± 3.4 for litter no. 1, n = 5).

\*  $P < 0.005$ ; \*\*  $P < 0.001$ .






FIG. 7. Morphological and functional abnormalities in the ovaries of cKO mice. **A** and **B**) Hematoxylin-and-eosin-stained sections of ovaries from 7-wk-old  $Gata4^{Flox/+};Amhr2^{cre/+}$  and cKO mice, respectively. Bar = 200  $\mu$ m. **C** and **D**) Histomorphometric analysis of follicle development in 7-wk-old  $Gata4^{Flox/+};Amhr2^{cre/+}$  ( $n = 10$ ) and cKO ( $n = 6$ ) mice ovaries. Serial ovarian sections were processed, and the numbers of healthy (nonatretic; **D**) or atretic (**E**) follicles were estimated. **E–H**) Ovaries from 6-mo-old  $Gata4^{Flox/+};Amhr2^{cre/+}$  (**E** and **G**) and cKO (**F** and **H**) mice. The arrowheads and asterisk highlight large cysts in the cKO ovary. Bar = 200  $\mu$ m. **I**) Serum AMH levels in  $Gata4^{Flox/+};Amhr2^{cre/+}$  ( $n = 3–7$ ) and cKO ( $n = 5–7$ ) mice of varying ages. Error bars represent SD. The bars denoted by *a* differ significantly ( $P < 0.01$ ) from that denoted by *b* but not from one another.

steadily over the ensuing 10 mo [43]. This age-related decline in serum AMH correlates with decreases in both the number of growing follicles and the size of the primordial follicle pool [43], making AMH a useful marker to quantify ovarian reserve




FIG. 8. Aberrant response to superovulation in cKO mice. Immature mice were treated with eCG, followed 48 h later by hCG. Tissue samples were harvested for analysis 18 h after hCG treatment. **A** and **B**) Ovaries from eCG/hCG-treated **(A)**  $Gata4^{Flox/+};Amhr2^{cre/+}$  and **(B)** cKO mice. Note the presence of hemorrhagic follicles (arrowheads) in the cKO ovary. **C**) Oocyte yields after superovulation. Each data point represents the oocyte yield of an individual mouse. The horizontal lines represent the mean values. The sample sizes were  $Gata4^{Flox/+};Amhr2^{cre/+}$  ( $n = 11$ ) and cKO ( $n = 13$ ).

[61]. AMH levels were similar in 2.5-mo-old  $Gata4^{Flox/+};Amhr2^{cre/+}$  and cKO mice. In contrast, AMH levels were significantly lower ( $P < 0.01$ ) in 6-mo-old cKO mice. Thus, AMH levels decline prematurely in  $Gata4$  cKO mice, indicating that ovarian granulosa cell function is compromised. In light of these results, we also measured AMH levels in the serum of older B6 WT and  $Gata4^{+/-}$  mice but found no significant differences (data not shown).

#### *Gata4* cKO Mice Exhibit an Impaired Response to Exogenous Gonadotropins

Next, we assessed the response of immature  $Gata4^{Flox/+};Amhr2^{cre/+}$  and cKO mice to superovulation. No significant differences were found in the weight or gross appearance of unstimulated  $Gata4^{Flox/+};Amhr2^{cre/+}$  and cKO mice (data not shown). However, the ovaries of cKO mice treated with eCG plus hCG were significantly smaller than those of  $Gata4^{Flox/+};Amhr2^{cre/+}$  mice ( $4.8 \pm 2.0$  mg [ $n = 10$ ] and  $7.9 \pm 2.0$  mg [ $n = 8$ ], respectively;  $P < 0.05$ ) and had hemorrhagic follicular




FIG. 9. Reduced expression of mRNA for *Gata4* and steroidogenic factors in the ovaries of gonadotropin-stimulated cKO mice. Weanling *Gata4<sup>Flox/+</sup>;Amhr2<sup>cre/+</sup>* (solid line, n = 5) or cKO (dashed line, n = 5) mice were treated with one of the following regimens: 1) no stimulation; 2) eCG for 48 h; 3) eCG for 48 h, followed by hCG for 18 h; or 4) eCG for 48 h, followed by hCG for 5 days. Ovarian RNA was subjected to qRT-PCR for *Gata4* (A), *Star* (B), *Cyp11a1* (C), and *Cyp19* (D). Error bars represent SD. \* $P < 0.05$ , \*\* $P < 0.01$ .

cysts on their surface (Fig. 8, A and B). A significant decrease ( $P < 0.05$ ) was observed in the yield of oocytes from cKO mice compared with *Gata4<sup>Flox/+</sup>;Amhr2<sup>cre/+</sup>* mice (Fig. 8C), although considerable variability occurred in the response of the cKO mice. One-third of the cKO mice tested released no oocytes following superovulation. Collectively, these data show that the *Gata4* granulosa cell-specific knockout mice are subfertile and have an impaired response to gonadotropin-induced superovulation.

Quantitative RT-PCR was used to measure the relative expression of transcripts for *Gata4* (Fig. 9A) and its target genes *Star* (Fig. 9B), *Cyp11a1* (Fig. 9C), and *Cyp19* (Fig. 9D) in ovaries from gonadotropin-stimulated, immature *Gata4<sup>Flox/+</sup>;Amhr2<sup>cre/+</sup>* and cKO mice. Basal levels of mRNA for *Star*, *Cyp11a1*, and *Cyp19* were similar in *Gata4<sup>Flox/+</sup>;Amhr2<sup>cre/+</sup>* and cKO mice. Following eCG stimulation, however, the level

of *Cyp19* mRNA was significantly higher in *Gata4<sup>Flox/+</sup>;Amhr2<sup>cre/+</sup>* than in cKO ovaries. Additionally, a trend, albeit statistically insignificant, was observed toward decreased expression of *Star* and *Cyp11a1* in the ovaries of gonadotropin-stimulated cKO mice. We did not, however, detect a significant difference in the serum  $E_2$  levels of eCG-treated juvenile mice.

## DISCUSSION

Expression of GATA4 occurs in potentially mitotic and proliferating granulosa cells, but expression is lost once these cells become either terminally differentiated (from granulosa to luteal cells) or apoptotic during follicular atresia [4, 7]. On the basis of these expression patterns, it was suggested [4] that GATA4 might control genes involved in the maintenance or maturation of granulosa cells within early follicles (i.e., before ovulation). Consistent with its proposed role in granulosa cell maturation, the expression level of GATA4 in granulosa cells was subsequently shown to be enhanced by FSH, and the transcriptional activity of GATA4 was found to be induced by phosphorylation via signaling through the FSH receptor and the downstream mediators cAMP and protein kinase A [8, 9]. The abrupt decrease in GATA4 associated with ovulation or apoptosis led investigators to propose that this factor is not required for the later stages of follicular development, apoptosis, or luteinization, although GATA4 might serve to prime early follicular cells for the transition to late maturation or apoptosis [4]. In keeping with these previous observations and predictions, the present data demonstrate that germline and conditional deletions in the *Gata4* gene cause defects in granulosa cell function that impair ovarian function in adult mice.

The B6 *Gata4<sup>+/-</sup>* female mice were fertile but had delayed puberty and a blunted ovarian response to exogenous gonadotropins; the ovaries of gonadotropin-stimulated *Gata4<sup>+/-</sup>* mice were smaller, released fewer oocytes, and expressed less mRNA for the steroidogenic genes *Star*, *Cyp11a1*, and *Cyp19* (Table 3). In the absence of a fertility defect, the physiological significance of a reduced response to ovarian superstimulation is arguable. On the other hand, our data show that provocative stimulation can be used to highlight “subclinical” defects in hormonal responsiveness that shed light on the function of GATA4 in vivo. The steroidogenic gene most profoundly affected by *Gata4* haploinsufficiency was *Cyp19*, the key gene of estrogen biosynthesis and a known target for activation by GATA4 [8, 9]. Consistent with reduced *Cyp19* expression, serum  $E_2$  levels were significantly reduced in *Gata4<sup>+/-</sup>* mice (Fig. 3G), and the uteri of eCG-stimulated *Gata4<sup>+/-</sup>* mice appeared to be hypoestrogenic. Like *Gata4<sup>+/-</sup>* mice, *Cyp19* knockout mice have impaired ovulation and hypoestrogenic uteri [46–48], suggesting that the abnormal phenotype of gonadotropin-stimulated *Gata4* haploinsufficient mice may reflect, at least in part, attenuated *Cyp19* expression and a concomitant decrease in  $E_2$  production. We speculate that the delay in the onset of puberty in *Gata4<sup>+/-</sup>* female mice could reflect a slight delay in the production of  $E_2$  in these mice, although we have no direct evidence of this and cannot exclude an extragonadal effect of *Gata4* haploinsufficiency. That *Gata4<sup>+/-</sup>* female mice have a normal estrous cycle length suggests that these animals eventually achieve adequate plasma levels of  $E_2$ .

The *Gata4* cKO females have a more profound phenotype, including impaired fertility, cystic ovarian changes, and an attenuated ovulation response to exogenous gonadotropins (Table 3). As with the *Gata4* haploinsufficient mice, the

TABLE 3. Comparison of the phenotypes of the two models of GATA4 deficiency used in this study.

Parameter	<i>Gata4</i> <sup>+/-</sup>	<i>Gata4</i> <sup>Flox/Flox</sup> ; <i>Amhr2</i> <sup>cre/+</sup>
Nature of gene deletion	Germline	Conditional
Genetic background	B6 (pure)	129;B6 (mixed)
Experimental control	WT	<i>Gata4</i> <sup>Flox/+</sup> ; <i>Amhr2</i> <sup>cre/+</sup>
Onset of puberty	Delayed	Normal
Fertility	Normal	Decreased
Expression of <i>Cyp19</i> mRNA response to exogenous gonadotropins	Decreased	Decreased
Ovulation in response to exogenous gonadotropins	Decreased	Decreased
Cystic ovarian changes	Absent	Present
Serum AMH	Normal for age	Decreased for age

steroidogenic gene most profoundly affected by conditional deletion of the *Gata4* gene was *Cyp19*. That most cKO mice were subfertile rather than completely infertile might reflect an inherent limitation of using *Amhr2-cre* to drive tissue-specific recombination. As shown by studies of  $\beta$ -catenin cKO mice generated using *Amhr2-cre* [52], the importance of a given gene for ovarian function may be masked by compensatory responses that promote selective proliferation of granulosa cells in follicles that escape Cre-mediated recombination. Indeed, a hallmark of the *Gata4* cKO mice and other knockout mice generated using *Amhr2-cre* is animal-to-animal variability, which may reflect inefficient recombination in granulosa cells (see [52] and references therein).

In *Gata4* cKO, but not *Gata4*<sup>+/-</sup>, mice, serum AMH levels declined earlier than in age-matched controls. Studies have shown that AMH, which is produced by granulosa cells in growing follicles, suppresses the development of primordial follicles [62–64]. In the absence of AMH, the primordial follicle pool depletes prematurely [62]. Serum AMH levels correlate with the size of the primordial follicle pool, making it a useful serum marker of ovarian reserve in both mice and women [61]. Of note, the *Amh* gene has been shown to be a direct target for activation of GATA4 in the testis and ovary (see [1] and references therein). Thus, it remains unclear whether the decrease in serum AMH in the cKO mice reflects a decrease in ovarian reserve, reduced GATA4-dependent activation of the *Amh* promoter, or both. Regardless of the mechanism, the premature decline in AMH levels underscores that ovarian granulosa cell function is compromised in the cKO mice.

Studies by Tevosian and colleagues [3, 19, 22] have established that GATA4 and its cofactor, ZFP281, mediate early events in ovarian development. In particular, these transcriptional regulators appear to modulate the response of embryonic ovarian tissue to extracellular signals, such as the WNT/ $\beta$ -catenin pathway. We presume, based on the appearance of unstimulated ovaries from young mice, that fetal ovarian development proceeds normally in both *Gata4* haploinsufficient mice and granulosa cell-specific cKO mice. This is in accordance with the results of studies on combined haploinsufficiency of *Nr5a1* and *Gata4*, which did not reveal a genetic interaction in mouse gonadal development [45]. In the case of *Gata4* cKO mice, we speculate that fetal ovarian development is normal, because efficient Cre-mediated recombination does not take place until robust expression of *Amhr2-cre* occurs in secondary and small antral follicles [52, 65].

In summary, the present study establishes a role for GATA4 in the regulation of adult ovarian function. These results, coupled with those of other investigators [9], show that GATA4 regulates the response of postnatal ovarian tissue to gonadotropins. Similarities in the postnatal ovarian phenotypes of *Gata4* cKO mice and mice harboring granulosa cell-specific knockouts of genes in the WNT/ $\beta$ -catenin pathway [52, 57]

raise the possibility that GATA4 regulates this signaling pathway not only in fetal mice [3, 19, 22] but also in adults; further experiments will explore this possibility. In a complementary study [66], we show that GATA4 also regulates testicular function in adults; male mice in which *Gata4* was conditionally deleted in Sertoli cells using *Amhr2-cre* exhibit impaired fertility and age-dependent testicular atrophy.

## ACKNOWLEDGMENTS

We thank Christine Ratajczak for technical advice and members of the Digestive Diseases Research Center Histology Core for their assistance. We also thank Simone Wagner for assistance with the *Gata4*<sup>+/-</sup> experiments. The *Gata4*<sup>+/-</sup> mice were generously provided by William Pu (Children's Hospital, Boston, MA).

## REFERENCES

- Viger RS, Guittot SM, Anttonen M, Wilson DB, Heikinheimo M. Role of the GATA family of transcription factors in endocrine development, function, and disease. *Mol Endocrinol* 2008; 22:781–798.
- LaVoie HA. The role of GATA in mammalian reproduction. *Exp Biol Med* 2003; 228:1282–1290.
- Tevosian SG, Manuylov NL. To beta or not to beta: canonical beta-catenin signaling pathway and ovarian development. *Dev Dyn* 2008; 237:3672–3680.
- Heikinheimo M, Ermolaeva M, Bielinska M, Rahman NA, Narita N, Huhtaniemi IT, Tapanainen JS, Wilson DB. Expression and hormonal regulation of transcription factors GATA-4 and GATA-6 in the mouse ovary. *Endocrinology* 1997; 138:3505–3514.
- LaVoie HA, McCoy GL, Blake CA. Expression of the GATA-4 and GATA-6 transcription factors in the fetal rat gonad and in the ovary during postnatal development and pregnancy. *Mol Cell Endocrinol* 2004; 227:31–40.
- McCoard SA, Wise TH, Fahrenkrug SC, Ford JJ. Temporal and spatial localization patterns of Gata4 during porcine gonadogenesis. *Biol Reprod* 2001; 65:366–374.
- Laitinen MP, Anttonen M, Ketola I, Wilson DB, Ritvos O, Butzow R, Heikinheimo M. Transcription factors GATA-4 and GATA-6 and a GATA family cofactor, FOG-2, are expressed in human ovary and sex cord-derived ovarian tumors. *J Clin Endocrinol Metab* 2000; 85:3476–3483.
- LaVoie HA, Singh D, Hui YY. Concerted regulation of the porcine steroidogenic acute regulatory protein gene promoter activity by follicle-stimulating hormone and insulin-like growth factor I in granulosa cells involves GATA-4 and CCAAT/enhancer binding protein beta. *Endocrinology* 2004; 145:3122–3134.
- Kwintkiewicz J, Cai Z, Stocco C. Follicle-stimulating hormone-induced activation of Gata4 contributes in the up-regulation of Cyp19 expression in rat granulosa cells. *Mol Endocrinol* 2007; 21:933–947.
- Balla A, Danilovich N, Yang Y, Sairam MR. Dynamics of ovarian development in the FORKO immature mouse: structural and functional implications for ovarian reserve. *Biol Reprod* 2003; 69:1281–1293.
- Vaskivuo TE, Aittomaki K, Anttonen M, Ruokonen A, Herva R, Osawa Y, Heikinheimo M, Huhtaniemi I, Tapanainen JS. Effects of follicle-stimulating hormone (FSH) and human chorionic gonadotropin in individuals with an inactivating mutation of the FSH receptor. *Fertil Steril* 2002; 78:108–113.

12. Hiroi H, Christenson LK, Strauss JF III. Regulation of transcription of the steroidogenic acute regulatory protein (StAR) gene: temporal and spatial changes in transcription factor binding and histone modification. *Mol Cell Endocrinol* 2004; 215:119–126.
13. Hui YY, Lavoie HA. GATA4 reduction enhances 3',5'-cyclic adenosine 5'-monophosphate-stimulated steroidogenic acute regulatory protein messenger ribonucleic acid and progesterone production in luteinized porcine granulosa cells. *Endocrinology* 2008; 149:5557–5567.
14. Sher N, Yivgi-Ohana N, Orly J. Transcriptional regulation of the cholesterol side chain cleavage cytochrome P450 gene (CYP11A1) revisited: binding of GATA, cyclic adenosine 3',5'-monophosphate response element-binding protein and activating protein (AP)-1 proteins to a distal novel cluster of cis-regulatory elements potentiates AP-2 and steroidogenic factor-1-dependent gene expression in the rodent placenta and ovary. *Mol Endocrinol* 2007; 21:948–962.
15. Tremblay JJ, Viger RS. GATA factors differentially activate multiple gonadal promoters through conserved GATA regulatory elements. *Endocrinology* 2001; 142:977–986.
16. Bouchard MF, Taniguchi H, Viger RS. Protein kinase A-dependent synergism between GATA factors and the nuclear receptor, liver receptor homolog-1, regulates human aromatase (CYP19) PII promoter activity in breast cancer cells. *Endocrinology* 2005; 146:4905–4916.
17. Stocco C. In vivo and in vitro inhibition of cyp19 gene expression by prostaglandin F2 $\alpha$  in murine luteal cells: implication of GATA-4. *Endocrinology* 2004; 145:4957–4966.
18. Stocco C, Kwinkiewicz J, Cai Z. Identification of regulatory elements in the Cyp19 proximal promoter in rat luteal cells. *J Mol Endocrinol* 2007; 39:211–221.
19. Manuylov NL, Smagulova FO, Leach L, Tevosian SG. Ovarian development in mice requires the GATA4-FOG2 transcription complex. *Development* 2008; 135:3731–3743.
20. Vainio S, Heikkilä M, Kispert A, Chin N, McMahon AP. Female development in mammals is regulated by Wnt-4 signaling. *Nature* 1999; 397:405–409.
21. Uhlenhaut NH, Treier M. Foxl2 function in ovarian development. *Mol Genet Metab* 2006; 88:225–234.
22. Tevosian SG, Albrecht KH, Crispino JD, Fujiwara Y, Eicher EM, Orkin SH. Gonadal differentiation, sex determination and normal Sry expression in mice require direct interaction between transcription partners GATA4 and FOG2. *Development* 2002; 129:4627–4634.
23. Kuo CT, Morrissey EE, Anadappa R, Sigrist K, Lu MM, Parmacek MS, Soudais C, Leiden JM. GATA4 transcription factor is required for ventral morphogenesis and heart tube formation. *Genes Dev* 1997; 11:1048–1060.
24. Molkenin JD, Lin Q, Duncan SA, Olson EN. Requirement of the transcription factor GATA4 for heart tube formation and ventral morphogenesis. *Genes Dev* 1997; 11:1061–1072.
25. Narita N, Bielinska M, Wilson D. Cardiomyocyte differentiation by GATA-4 deficient embryonic stem cells. *Devel* 1997; 122:3755–3764.
26. Narita N, Bielinska M, Wilson DB. Wild-type endoderm abrogates the ventral developmental defects associated with GATA-4 deficiency in the mouse. *Dev Biol* 1997; 189:270–274.
27. Watt AJ, Battle MA, Li J, Duncan SA. GATA4 is essential for formation of the proepicardium and regulates cardiogenesis. *Proc Natl Acad Sci U S A* 2004; 101:12573–12578.
28. Thurisch B, Liang S, Sarioglu N, Schomburg L, Bungert J, Dame C. Transgenic mice expressing small interfering RNA against Gata4 point to a crucial role of Gata4 in the heart and gonads. *J Mol Endocrinol* 2009; 43:157–169.
29. Pu WT, Ishiwata T, Juraszek AL, Ma Q, Izumo S. GATA4 is a dosage-sensitive regulator of cardiac morphogenesis. *Dev Biol* 2004; 275:235–244.
30. Zeisberg EM, Ma Q, Juraszek AL, Moses K, Schwartz RJ, Izumo S, Pu WT. Morphogenesis of the right ventricle requires myocardial expression of Gata4. *J Clin Invest* 2005; 115:1522–1531.
31. Oka T, Maillet M, Watt AJ, Schwartz RJ, Aronow BJ, Duncan SA, Molkenin JD. Cardiac-specific deletion of Gata4 reveals its requirement for hypertrophy, compensation, and myocyte viability. *Circ Res* 2006; 98:837–845.
32. Jamin SP, Arango NA, Mishina Y, Hanks MC, Behringer RR. Requirement of Bmpr1a for mullerian duct regression during male sexual development. *Nat Genet* 2002; 32:408–410.
33. Jeyasuria P, Ikeda Y, Jamin SP, Zhao L, de Rooij DG, Themmen AP, Behringer RR, Parker KL. Cell-specific knockout of steroidogenic factor 1 reveals its essential roles in gonadal function. *Mol Endocrinol* 2004; 18:1610–1619.
34. Friedrich G, Soriano P. Promoter traps in embryonic stem cells: a genetic screen to identify and mutate developmental genes in mice. *Genes Dev* 1991; 5:1513–1523.
35. Nelson JF, Felicio LS, Randall PK, Sims C, Finch CE. A longitudinal study of estrous cyclicity in aging C57BL/6J mice: I. Cycle frequency, length and vaginal cytology. *Biol Reprod* 1982; 27:327–339.
36. Jay PY, Bielinska M, Erlich JM, Mannisto S, Pu WT, Heikinheimo M, Wilson DB. Impaired mesenchymal cell function in Gata4 mutant mice leads to diaphragmatic hernias and primary lung defects. *Dev Biol* 2007; 301:602–614.
37. Bielinska M, Jay PY, Erlich JM, Mannisto S, Urban Z, Heikinheimo M, Wilson DB. Molecular genetics of congenital diaphragmatic defects. *Ann Med* 2007; 39:261–274.
38. Rajagopal SK, Ma Q, Obler D, Shen J, Manichaikul A, Tomita-Mitchell A, Boardman K, Briggs C, Garg V, Srivastava D, Goldmuntz E, Broman KW, et al. Spectrum of heart disease associated with murine and human GATA4 mutation. *J Mol Cell Cardiol* 2007; 43:677–685.
39. Morita Y, Perez GI, Maravei DV, Tilly KL, Tilly JL. Targeted expression of Bcl-2 in mouse oocytes inhibits ovarian follicle atresia and prevents spontaneous and chemotherapy-induced oocyte apoptosis in vitro. *Mol Endocrinol* 1999; 13:841–850.
40. Tilly JL. Ovarian follicle counts—not as simple as 1, 2, 3. *Reprod Biol Endocrinol* 2003; 1:11–15.
41. Hogan B, Beddington R, Costantini F, Lacy E. *Manipulating the Mouse Embryo: A Laboratory Manual*. Cold Spring Harbor, NY: Cold Spring Harbor Press; 1994.
42. Kyronlahti A, Ramo M, Tamminen M, Unkila-Kallio L, Butzow R, Leminen A, Nemer M, Rahman N, Huhtaniemi I, Heikinheimo M, Anttonen M. GATA-4 regulates Bcl-2 expression in ovarian granulosa cell tumors. *Endocrinology* 2008; 149:5635–5642.
43. Kevenaar ME, Meerasahib MF, Kramer P, van de Lang-Born BM, de Jong FH, Groome NP, Themmen AP, Visser JA. Serum anti-mullerian hormone levels reflect the size of the primordial follicle pool in mice. *Endocrinology* 2006; 147:3228–3234.
44. Nelson JF, Karelus K, Felicio LS, Johnson TE. Genetic influences on the timing of puberty in mice. *Biol Reprod* 1990; 42:649–655.
45. Pelusi C, Zhao L, Stallings NR, Parker KL. Combined haploinsufficiency of SF-1 and GATA4 does not reveal a genetic interaction in mouse gonadal development. *Sex Dev* 2007; 1:152–160.
46. Fisher CR, Graves KH, Parlow AF, Simpson ER. Characterization of mice deficient in aromatase (ArKO) because of targeted disruption of the cyp19 gene. *Proc Natl Acad Sci U S A* 1998; 95:6965–6970.
47. Britt KL, Drummond AE, Cox VA, Dyson M, Wreford NG, Jones ME, Simpson ER, Findlay JK. An age-related ovarian phenotype in mice with targeted disruption of the Cyp19 (aromatase) gene. *Endocrinology* 2000; 141:2614–2623.
48. Toda K, Takeda K, Okada T, Akira S, Saibara T, Kaname T, Yamamura K, Onishi S, Shizuta Y. Targeted disruption of the aromatase P450 gene (Cyp19) in mice and their ovarian and uterine responses to 17 $\beta$ -estradiol. *J Endocrinol* 2001; 170:99–111.
49. Deutscher E, Hung-Chang Yao H. Essential roles of mesenchyme-derived beta-catenin in mouse mullerian duct morphogenesis. *Dev Biol* 2007; 307:227–236.
50. Gonzalez G, Behringer RR. Dicer is required for female reproductive tract development and fertility in the mouse. *Mol Reprod Dev* 2009; 76:678–688.
51. Orvis GD, Jamin SP, Kwan KM, Mishina Y, Kaartinen VM, Huang S, Roberts AB, Umans L, Huylebroeck D, Zwijsen A, Wang D, Martin JF, et al. Functional redundancy of TGF-beta family type 1 receptors and receptor-Smads in mediating anti-mullerian hormone-induced mullerian duct regression in the mouse. *Biol Reprod* 2008; 78:994–1001.
52. Hernandez Gifford JA, Hunzicker-Dunn ME, Nilson JH. Conditional deletion of beta-catenin mediated by Amhr2cre in mice causes female infertility. *Biol Reprod* 2009; 80:1282–1292.
53. Petit FG, Jamin SP, Kurihara I, Behringer RR, DeMayo FJ, Tsai MJ, Tsai SY. Deletion of the orphan nuclear receptor COUP-TFII in uterus leads to placental deficiency. *Proc Natl Acad Sci U S A* 2007; 104:6293–6298.
54. Fan HY, Shimada M, Liu Z, Cahill N, Noma N, Wu Y, Gossen J, Richards JS. Selective expression of KrasG12D in granulosa cells of the mouse ovary causes defects in follicle development and ovulation. *Development* 2008; 135:2127–2137.
55. Duggavathi R, Volle DH, Matakci C, Antal MC, Messaddeq N, Auwerx J, Murphy BD, Schoonjans K. Liver receptor homolog 1 is essential for ovulation. *Genes Dev* 2008; 22:1871–1876.
56. Ren Y, Cowan RG, Harman RM, Quirk SM. Dominant activation of the hedgehog signaling pathway in the ovary alters theca development and prevents ovulation. *Mol Endocrinol* 2009; 23:711–723.
57. Boyer A, Lapointe E, Zheng X, Cowan RG, Li H, Quirk SM, Demayo FJ,

- Richards JS, Boerboom D. WNT4 is required for normal ovarian follicle development and female fertility. *FASEB J* 2010; 24:3010–3025.
58. Pelusi C, Ikeda Y, Zubair M, Parker KL. Impaired follicle development and infertility in female mice lacking steroidogenic factor 1 in ovarian granulosa cells. *Biol Reprod* 2008; 79:1074–1083.
59. Anttonen M, Ketola I, Parviainen H, Pusa AK, Heikinheimo M. FOG-2 and GATA-4 are coexpressed in the mouse ovary and can modulate mullerian-inhibiting substance expression. *Biol Reprod* 2003; 68:1333–1340.
60. Munsterberg A, Lovell-Badge R. Expression of the mouse anti-mullerian hormone gene suggests a role in both male and female sexual differentiation. *Development* 1991; 113:613–624.
61. van Houten EL, Themmen AP, Visser JA. Anti-mullerian hormone (AMH): regulator and marker of ovarian function. *Ann Endocrinol (Paris)* 2010; 71:191–197.
62. Gruijters MJ, Visser JA, Durlinger AL, Themmen AP. Anti-mullerian hormone and its role in ovarian function. *Mol Cell Endocrinol* 2003; 211: 85–90.
63. Weenen C, Laven JS, Von Bergh AR, Cranfield M, Groome NP, Visser JA, Kramer P, Fauser BC, Themmen AP. Anti-mullerian hormone expression pattern in the human ovary: potential implications for initial and cyclic follicle recruitment. *Mol Hum Reprod* 2004; 10:77–83.
64. Durlinger AL, Gruijters MJ, Kramer P, Karels B, Ingraham HA, Nachtigal MW, Uilenbroek JT, Grootegoed JA, Themmen AP. Anti-mullerian hormone inhibits initiation of primordial follicle growth in the mouse ovary. *Endocrinology* 2002; 143:1076–1084.
65. Jorgez CJ, Klysiak M, Jamin SP, Behringer RR, Matzuk MM. Granulosa cell-specific inactivation of follistatin causes female fertility defects. *Mol Endocrinol* 2004; 18:953–967.
66. Kyronlahti A, Euler R, Bielinska M, Schoeller EL, Moley KH, Toppari J, Heikinheimo M, Wilson DB. GATA4 regulates Sertoli cell function and fertility in adult male mice. *Mol Cell Endocrinol* 2011; 333:85–95.