



**Ornithodoros sawaii (Acari: Ixodida: Argasidae)  
collected from nest soil/litter of Synthliboramphus  
antiquus, Synthliboramphuswumizusume, and nest  
soil/litter and nestlings of Hydrobatesmonorhis on  
Gugul and Sogugul Islands, Jeollanam Province,  
Republic of Korea**

Authors: Kim, Heung-Chul, Park, Chang-Uk, Kim, Miran, Kim, Yang-Mo, Yeo, Juhan, et al.

Source: Systematic and Applied Acarology, 22(11) : 1959-1969

Published By: Systematic and Applied Acarology Society

URL: <https://doi.org/10.11158/saa.22.11.13>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## ***Ornithodoros sawaii* (Acari: Ixodida: Argasidae) collected from nest soil/litter of *Synthliboramphus antiquus*, *Synthliboramphus wumizusume*, and nest soil/litter and nestlings of *Hydrobates monorhis* on Gugul and Sogugul Islands, Jeollanam Province, Republic of Korea**

HEUNG-CHUL KIM<sup>1\*</sup>, CHANG-UK PARK<sup>2\*</sup>, MIRAN KIM<sup>2</sup>, YANG-MO KIM<sup>2</sup>, JUHAN YEO<sup>2</sup>, YOUNG-SOO KWON<sup>2</sup>, SEOK-MIN YUN<sup>3</sup>, WON-JA LEE<sup>3</sup>, SUNG-TAE CHONG<sup>1</sup>, TERRY A. KLEIN<sup>4</sup> & RICHARD G. ROBBINS<sup>5,6</sup>

<sup>1</sup> 5<sup>th</sup> Medical Detachment, 168<sup>th</sup> Multifunctional Medical Battalion, 65<sup>th</sup> Medical Brigade, Unit 15247, APO AP 96205-5247, USA

<sup>2</sup> Migratory Bird Research Center, Korea National Park Research Institute, Korea National Park Service, Jinmaeul-gil, Heuksan-myeon, Sinan county, Jeollanam Province 58863, Republic of Korea

<sup>3</sup> Division of Arboviruses, Center for Immunology and Pathology, National Institute of Health, Korea Centers for Diseases Control and Prevention, Cheongju-si, Chungbuk Province 28159, Republic of Korea

<sup>4</sup> Medical Department Activity-Korea/65<sup>th</sup> Medical Brigade, Unit 15281, APO AP 96205-5281, USA

<sup>5</sup> Walter Reed Biosystematics Unit, Department of Entomology, Smithsonian Institution, MSC, MRC 534, 4210 Silver Hill Road, Suitland, MD 20746-2863, USA

<sup>6</sup> Corresponding author. E-mail: richard.g.robbins@gmail.com

\* These authors contributed equally to this work.

### **Abstract**

The 65<sup>th</sup> Medical Brigade and Medical Department Activity-Korea, in collaboration with the Migratory Birds Research Center, National Park Research Institute, conducted a migratory bird tick-borne disease surveillance program on Gugul and Sogugul Islands, two small, remote, uninhabited islands in southwestern Jeollanam Province, Republic of Korea (ROK) from 2015–2016. Ticks were collected from nest soil/litter of the Ancient Murrelet, *Synthliboramphus antiquus*, Japanese Murrelet, *Synthliboramphus wumizusume*, and Swinhoe's Storm Petrel, *Hydrobates monorhis*, using Tullgren funnels. A total of 100 *Ornithodoros sawaii* (25 females, 56 males, 17 nymphs, and 2 larvae) were collected from 2/33 (6.1%) Ancient Murrelet, 3/11 (27.3%) Japanese Murrelet, and 23/94 (24.5%) Swinhoe's Storm Petrel nest sites. In addition, 3 larvae were collected from Swinhoe's Storm Petrel nestlings. Ticks were identified morphologically and species determination was confirmed using polymerase chain reaction techniques. This is the first record of *O. sawaii* collected from nest soil/litter of the Japanese Murrelet during this bird's nesting season in the ROK.

**Key words:** Argasidae, *Ornithodoros sawaii*, *Synthliboramphus antiquus*, *S. wumizusume*, *Hydrobates monorhis*, Korea

### **Introduction**

Worldwide, there has been a resurgence of interest in zoonotic tick-borne pathogens that affect the health of wild and domestic animals, birds, and humans (Heath & Hardwick 2011, Dantas-Torres *et al.* 2012). In the Republic of Korea (ROK), tick-borne pathogenic bacteria [*e.g.*, *Ehrlichia* and *Anaplasma* (Chae *et al.* 2003, Kang *et al.* 2013), *Bartonella* (Kim *et al.* 2005, Kang *et al.* 2013), *Borrelia* (Park *et al.* 1992, Kee *et al.* 1994), and *Rickettsia* spp. (Jang *et al.* 2004, Lee *et al.* 2013,

Noh *et al.* 2017)] and viruses [*e.g.*, tick-borne encephalitis (Kim *et al.* 2009b, Ko *et al.* 2010, Yun *et al.* 2013) and severe fever with thrombocytopenia syndrome (SFTS) (Kim *et al.* 2013, Park *et al.* 2014, Yun *et al.* 2014, 2016)] manifest varying degrees of disease nationwide.

The role of migratory seabirds in the transport of exotic tick species to and from their summer breeding and winter feeding grounds has been poorly documented, in part because their breeding sites are often located on remote, uninhabited or sparsely populated islands that are usually under government protection and inaccessible to the general public. Annual migrations of migratory seabirds to and from their breeding and feeding grounds may result in the importation of exotic tick species that are transported over long distances and to diverse environments in non-endemic regions (Kohls 1957, Hughes *et al.* 1964, Amerson 1968, Nuttall 1984, Heath 1987, 2006, Hutcheson *et al.* 2005, Kawabata *et al.* 2006, Kim *et al.* 2009a, Dietrich *et al.* 2011, Kang *et al.* 2013).

With exceptions, ticks belonging to the family Ixodidae (hard ticks) are more commonly found on so-called perching birds (order Passeriformes) (Morshed *et al.* 2005), while members of the family Argasidae (soft ticks) are usually collected from the nests and burrows of non-passeriform ground-dwelling/nesting birds, including migratory seabirds (Hoogstraal *et al.* 1976, Gomez-Diaz *et al.* 2012). The genus *Ornithodoros* Koch contains the largest number of described species of soft ticks (112/193) (Guglielmone *et al.* 2010), and these are frequently associated with various seabird hosts in both the Eastern and Western Hemispheres, including the Pacific Islands (Vander Velde & Vander Velde 2013), North and South America (Denmark & Clifford 1962, Jonkers *et al.* 1973, Keirans *et al.* 1992), New Zealand (Heath 1987, 2006, Murray *et al.* 1990), Australia (Murray *et al.* 1990), Antarctica (Murray *et al.* 1990), Japan (Asanuma *et al.* 1955, Asanuma & Fukuda 1957, Asanuma 1960, 1965, Yamaguti *et al.* 1971, Kawabata *et al.* 2006), Russia (Akhundova *et al.* 1971), and Africa (Hoogstraal *et al.* 1976).

This report describes collections of *Ornithodoros sawaii* Kitaoka and Suzuki, 1973 from nest soil/litter of the Ancient Murrelet, *Synthliboramphus antiquus* (Gmelin, 1789), the Japanese Murrelet, *Synthliboramphus wumizusume* (Temminck, 1835), and both nest soil/litter and nestlings of Swinhoe's Storm Petrel, *Hydrobates monorhis* (Swinhoe, 1867), in the ROK. The collections from *S. wumizusume* nest soil/litter represent a probable new host record for *O. sawaii*.

## Materials and methods


### Survey area

The Migratory Birds Research Center, National Park Research Institute, located on Heuksan Island, Heuksan-myeon (district), Sinan-Gun (county), Jeollanam Province, ROK, conducted conservation and breeding status surveys of the Ancient Murrelet, Japanese Murrelet, and Swinhoe's Storm Petrel on Gugul and Sogugul Islands (34°04' N, 125°07' E), two small, remote, uninhabited islands that are protected National Monuments (No. 341, 13 August 1984); the islands have a total land area of 13.52 ha and 1.84 ha, respectively, and are located in Gageodo-ri, Heuksan-myeon, Sinan-gun, Jeollanam Province, 136 km west of Mokpo, a major mainland port city (Fig. 1).

### Tick collections

Tick surveys were conducted on Gugul and Sogugul Islands during the Ancient Murrelet and Japanese Murrelet breeding season (March–early May), and Swinhoe's Storm Petrel breeding season (June–October). Although the Murrelet and Storm Petrel breeding seasons do not overlap, they occupy the same nest sites. Using a small scoop, 50–100 g of soil and litter were collected from nests occupied by Ancient Murrelets and Japanese Murrelets on 22 April, 21 May 2015, and 7 May 2016, and from nests occupied by Swinhoe's Storm Petrel on 11 August, 12 September, and 15 October

2016 (Fig. 2). Soil and litter samples were placed in plastic Ziploc® bags (25 x 28 cm) that were then transferred to a Styrofoam cooler and transported to the 5<sup>th</sup> Medical Detachment, Yongsan U.S. Army Garrison, Seoul, ROK. Soil and litter samples from each nest site were placed separately inside Tullgren funnels equipped with a 52W incandescent light bulb (heat source) at the top and a collection bottle (120 ml urine specimen container) at the base containing 50 ml of 70% ethanol. After 24 hours of exposure, the material in the collection bottle was examined for arthropods. Ticks were removed and placed individually in cryovials containing 80% ethanol, labeled with a unique nest identification number, and later microscopically examined to determine developmental stage and genus using taxonomic identification keys (Kohls 1957, Yamaguti *et al.* 1971, Kitaoka & Suzuki 1973). Breeding season Swinhoe's Storm Petrel nestlings were carefully removed from their nests, identified to species, and then banded with a unique identification number. Prior to release, they were examined for ticks, particularly around the head, neck, wings and abdomen, and ticks, carefully removed with a forceps to prevent injury to the bird, were placed individually in 2 ml cryovials containing 70% EtOH and then labeled with a unique nest identification number, as described by Kim *et al.* (2016a). Two females and four males (2015 samples) and one female, three males, three nymphs, and one larva (2016 samples) were submitted to the Korea National Institute of Health for specific identification by polymerase chain reaction (PCR). Total DNA was prepared from individual ticks using a DNeasy tissue kit (Qiagen, Hilden, Germany) according to the manufacturer's instructions with minor modification and stored at -20°C until used.



**FIGURE 1.** Collection sites (○) of *Ornithodoros sawaii* from the Ancient Murrelet (*Synthliboramphus antiquus*), Japanese Murrelet (*Synthliboramphus wumizusume*) nest soil/litter, and Swinhoe's Storm Petrel (*Hydrobates monorhis*) nest soil/litter and nestlings on Gugul and Sogugul Islands, Jeollanam Province, Republic of Korea [Mokpo (●), mainland port city] (map image from <http://map.daum.net>).

#### PCR and sequencing analysis

PCR was performed using primer sets (mt-rrs1: 5-CTG CTC AAT GAT TTT TTA AAT TGC TGT GG-3 and mt-rrs2: 5-CCG GTC TGA ACT CAG ATC AAG TA-3) based on the mitochondrial 16S rDNA gene (mt-rrs) fragment previously described by Black and Piesman (1994) and Ushijima

*et al.* (2003). PCR assays were performed using 50  $\mu$ L of reaction mixture with TaKaRa ExTaq™ DNA polymerase (Takara, Shiga, Japan) at 94°C for 5 min, followed by 35 cycles for 10 sec at 94°C, 30 sec at 55°C, and then 30 sec at 72°C, with a final extension step of 5 min at 72°C. PCR products were then purified using a QIAquick® Gel Extraction Kit (Qiagen, Hilden, Germany) according to the manufacturer's instructions and sequenced after cloning into pCR®4-TOPO® plasmid (Invitrogen, Carlsbad, USA), using ABI Prism BigDye™ Terminator v3.1 Cycle Sequencing Kits and an ABI 3730xl sequencer (Applied Biosystems®, Foster City, USA) at MacroGen, Inc. (Daejeon, ROK). Sequencing results were assembled using the SeqMan program implemented in DNASTAR software (version 5.0.6; DNASTAR, Inc., Madison, WI, USA) to determine consensus sequences. The mt-rrs sequences of ticks identified in this study have been deposited in GenBank under accession numbers KY654988-KY655001.



**FIGURE 2.** Three avian hosts of *Ornithodoros sawaii*: (A) Japanese Murrelet (*Synthliboramphus wumizusume*), (B) Ancient Murrelet (*Synthliboramphus antiquus*), and (C) Swinhoe's Storm Petrel (*Hydrobates monorhis*) (photos by Chang-Uk Park, co-author).

#### Phylogenetic analysis

Sequence data were analyzed using MEGA 6.0 software (<http://www.megasoftware.net>) (Tamura *et al.* 2013). The amplified partial mt-rrs segment sequences for *Ornithodoros* spp. collected from the nesting sites were aligned and compared with previously published *Ornithodoros* spp., a process facilitated using the CLUSTALW method (Lasergene program version 5, DNASTAR, Inc.). For phylogenetic analysis, neighbor-joining (NJ) and bootstrap tests were carried out according to the Kimura 2-parameter distance method (Kimura 1980, Saitou & Nei 1987). Pairwise alignments were performed with an open-gap penalty of 15 and a gap extension penalty of 6.66. Multiple alignments were also performed using the same values. All positions containing alignment gaps and missing data were eliminated in pairwise sequence comparisons (pairwise deletion).

#### Results and Discussion

A total of 100 *Ornithodoros sawaii* (25 females, 56 males, 17 nymphs, and 2 larvae) were collected from 2/33 (6.1%) Ancient Murrelet, 3/11 (27.3%) Japanese Murrelet, and 23/94 (24.5%) Swinhoe's Storm Petrel nest sites. In addition, 3 larvae were collected from Swinhoe's Storm Petrel nestlings (Table 1).

Because the morphological identification of *Ornithodoros* spp. nymphs and adults is often unreliable, specimens collected on Gugul and Sogugul Islands were identified by PCR using partial mitochondrial 16S ribonucleic acid gene (mt-rrs) primer sets developed for the identification of soft ticks and sequence analysis with *O. sawaii* in Japan and other countries (Fig. 3). Sequence data for

*O. sawaii* collected from nest soil and litter on Gugul Island during the breeding season of the Ancient Murrelet [2 males (KOR-G1505-E2-1 and KOR-G1505-E2-2)], the Japanese Murrelet [2 females (KOR-G1505W-Z1-7 and KOR-G1505W-Z1-8) and 2 males (KOR-G1505W-Z3-3 and KOR-G1505W-Z3-4)], and Swinhoe's Storm Petrel [1 female (KOR-G1608-40-1), 2 males (KOR-G1608-2-2 and KOR-G1608-2-3), 2 nymphs (KOR-G1610-1-3 and KOR-G1610-1-4), and 1 larva (KOR-G1609-4-3)] aligned closely with *O. sawaii* sequence data submitted to GenBank (Fig. 3). Similarly, sequence data for *O. sawaii* collected from nest soil and litter on Sogugul Island during Swinhoe's Storm Petrel breeding season [1 male (KOR-S1610-13-1), and 1 nymph (KOR-S1610-2-2)] aligned closely with *O. sawaii* in the ROK (Fig. 3). Comparatively, there were just 0-2 base differences, with 99.5–100% nucleotide sequence identities between *O. sawaii* collected from Gugul and Sogugul Islands and other collection sites in Japan (Miyazaki, Shimane, Kyoto, Kutsujima, and Ishikawa Islands) (Kawabata *et al.* 2006, Takano *et al.* 2014). In Japan, *O. sawaii* has been recorded from two migratory seabirds, the Streaked Shearwater, *Calonectris leucomelas* (Temminck, 1836), and Swinhoe's Storm Petrel (Kitaoka and Suzuki 1973, 1974, Kawabata *et al.* 2006, Takano *et al.* 2014). Our survey showed that *O. sawaii* is associated with nests of the Ancient Murrelet and Japanese Murrelet, as well as Swinhoe's Storm Petrel. This is the first record of *O. sawaii* collected from nest soil/litter of the Japanese Murrelet.

**TABLE 1.** Number and stage of *Ornithodoros sawaii* collected from nest soil/litter of Ancient Murrelet (*Synthliboramphus antiquus*), Japanese Murrelet (*Synthliboramphus wumizusume*), and nest soil/litter and nestlings of Swinhoe's Storm Petrel (*Hydrobates monorhis*) on Gugul and Sogugul Islands, Jeollanam Province, Republic of Korea, 2015–2016.

Year	Collection Island	Host bird	Month	Infested/ tested soil and litter	Infestation rates (%)	<i>Ornithodoros sawaii</i>					
						F	M	N	L	L*	Total
2015	Gugul Island	Ancient Murrelet ( <i>Synthliboramphus antiquus</i> )	APR	0/5	0	0	0	0	0	0	
		Japanese Murrelet ( <i>Synthliboramphus wumizusume</i> )	MAY	2/25	8.0	0	3	2	0	5	
		Ancient Murrelet ( <i>Synthliboramphus antiquus</i> )	APR	0/1	0	0	0	0	0	0	
		Japanese Murrelet ( <i>Synthliboramphus wumizusume</i> )	MAY	3/10	30.0	13	19	3	0	35	
2016	Gugul Island	Ancient Murrelet ( <i>Synthliboramphus antiquus</i> )	MAY	0/3	0	0	0	0	0	0	
		Swinhoe's Storm Petrel ( <i>Hydrobates monorhis</i> )	AUG	12/46	26.1	8	29	1	0	38	
	Sogugul Island	Swinhoe's Storm Petrel ( <i>Hydrobates monorhis</i> )	SEP	2/12	16.7	1	0	1	2	1	5
		Swinhoe's Storm Petrel ( <i>Hydrobates monorhis</i> )	OCT	3/22	13.6	1	0	5	0	2	8
		Swinhoe's Storm Petrel ( <i>Hydrobates monorhis</i> )	OCT	6/14	42.9	2	5	5	0	12	
Total				28/138	20.3	25	56	17	2	3	103

\*Larvae collected from nestlings of Swinhoe's storm petrel, *Hydrobates monorhis*.

Japanese Murrelets are medium-sized blackish seabirds (Fig. 2A) that are endemic to regions of warm ocean currents near central and southern Japan, Korea, and Russia. Following the breeding season, these birds move to the southeastern coastal margins of Hokkaido, Honshu and Kyushu, where they overwinter. Japanese Murrelet populations are declining in many localities, particularly the Izu Islands (BirdLife International 2016a). Ancient Murrelets are medium-sized blackish seabirds that nest in colonies in rock crevices and burrow under the roots of *Carex bootiana* Hooker and Arnott (Fig. 2B). Their range extends from the Yellow Sea (islands off China and Korea),

Taiwan, Hong Kong, islands in the East Sea along the Korean Peninsula, the Russian Pacific coast and the Aleutian Islands to the Haida Gwaii (Queen Charlotte) archipelago of British Columbia, Canada, and as far south as the southern coast of California, USA (Del Hoyo *et al.* 1996, BirdLife International 2016b). On Gugul and Sogugul Islands, Japanese and Ancient Murrelets arrive concurrently at their feeding/nesting grounds one month before egg-laying from March–early May, and after rearing their chicks, they move to their coastal summer feeding grounds. Swinhoe’s Storm Petrels migrate to Gugul and Sogugul Islands and other nearby islands from early June–October and occupy the same nest sites as the Ancient and Japanese Murrelets (Lee *et al.* 2009, Park & Takeshi 2011). Swinhoe’s Storm Petrels are small seabirds that range from southern Indonesia and the Indian Ocean to northeastern Asia [Russia (south of Vladivostok), Japan, Korea, and Taiwan] (Lee *et al.* 2009, BirdLife International 2016c) (Fig. 2C).


*Ornithodoros sawaii* is closely related to *O. capensis*, a species with a broad distribution that includes North Atlantic, Pacific, and Indian Ocean islands and coastal areas of South Africa, New Zealand, and southern Australia (Kohls 1957, Heath 1987). *Ornithodoros sawaii* was first described in 1973, but in Japan it is known only from nests of the Streaked Shearwater and Swinhoe’s Storm Petrel located on the Japanese islands of Hanmya, Miyazaki, Shimane, Maizuru, Kyoto, and Ishikawa (Kitaoka & Suzuki 1973, 1974, Kawabata *et al.* 2006, Takano *et al.* 2014). In Korea, *O. sawaii* larvae were recovered from the wing and abdomen of Swinhoe’s Storm Petrel on Sogugul and Gaerin Islands (Kim *et al.* 2016a), and nymphs and adults were collected from nest soil/litter during the nesting season of the Ancient Murrelet and Swinhoe’s Storm Petrel on Chilbal Island (Kim *et al.* 2015, 2016b).

*Rickettsia* and *Borrelia* spp. have been detected in *O. sawaii* collected from the Streaked Shearwater and Swinhoe’s Storm Petrel in Japan (Kawabata *et al.* 2006, Takano *et al.* 2009). Recently, Kang *et al.* (2013) reported *Anaplasma*, *Bartonella*, and *Borrelia* spp. from *Ixodes* spp. that were collected from migratory birds during 2008–2009 on Hong Island (34°41' N, 125°11' E), located near Gugul and Sogugul Islands. Such results do not necessarily mean that migratory birds are reservoirs of these pathogens or are responsible for the introduction of exotic ticks into non-endemic areas, but they hint at the potential role of migratory birds in the dispersal of ticks and tick-borne microbial agents in northeastern Asia.

Further studies are needed to define the geographical distribution, host range, and specific pathogens associated with ticks collected from nesting sites and from resident and migratory seabirds inhabiting islands and coastal areas of the ROK.

## Acknowledgments

We thank Dr. Jong-Gil Park, Chief, Migratory Birds Research Center, National Park Research Institute, Korea National Park Service, on Heuksan Island, Jeollanam Province, ROK, for collecting nest soil and litter during seabird conservation and breeding surveys. The seabird survey was supported by “2016 Bird Survey and Research,” Korea National Park Research Institute; the Armed Forces Health Surveillance Branch-Global Emerging Infections Surveillance and Response System (AFHSB-GEIS), Silver Spring, MD, USA; and the 65<sup>th</sup> Medical Brigade, Seoul, Korea. The opinions expressed herein are those of the authors and are not to be construed as official or reflecting the views of the US Departments of the Army or Defense.



**FIGURE 3.** Phylogenetic analysis based on mt-rrs of *Ornithodoros sawaii* collected from the Ancient Murrelet (*Synthliboramphus antiquus*) and Japanese Murrelet (*Synthliboramphus wumizusume*) nest soil/litter, and Swinhoe's Storm Petrel (*Hydrobates monorhis*) nest soil/litter and nestlings. The phylogenetic trees were constructed based on NJ methods and bootstrap tests carried out according to the Kimura 2-parameter distances method. The percentage of replicate trees in which the associated taxa are clustered together in the bootstrap test (1,000 replicates) was calculated. The phylogenetic branches were supported with more than 70% bootstrap values in this analysis. The length of the bar corresponds to the degree of sequence divergence. All positions containing alignment gaps and missing data were eliminated in pairwise sequence comparisons (pairwise deletion).

## References

- Akhundova, E.D., Abushev, F.A. & Sterknova, N.N. (1971) Ectoparasites of some bird species in different regions of Azerbaijan SSR. *Materialy Simpoziuma. Itogi VI Svyazan. Simpoziuma po Izucheniyu Virusov, Ekologicheskii Svyazannykh s Ptitsami*, 7–9 Dekabrya 1971, Omsk, 128–130 (in Russian with English summary) [Translation 671 (T671) by Medical Zoology Department, United States Naval Medical Research Unit No. 3, Cairo, Egypt].
- Amerson, A.B., Jr. (1968) Tick distribution in the Central Pacific as influenced by sea bird movement. *Journal of Medical Entomology*, 5, 332–339.  
<https://doi.org/10.1093/jmedent/5.3.332>



- Asanuma, K. (1960) Host and distribution records of the soft tick, *Ornithodoros capensis* Neumann in Japan. *Japanese Journal of Sanitary Zoology*, 11, 94 (in Japanese).
- Asanuma, K. (1965) Suborder Ixodides. In: Sasa, M. (Ed.), *Mites: an Introduction to Classification, Bionomics and Control of Acarina*. Tokyo, University of Tokyo Press, pp. 101–128 (in Japanese).
- Asanuma, K. & Fukuda, S. (1957) On the life-history of a tick, *Ixodes signatus*, infesting a Black-tailed Gull (*Larus crassirostris*) in Japan. *Japanese Journal of Sanitary Zoology*, 8, 147–159.  
<https://doi.org/10.7601/mez.8.147>
- Asanuma, K., Okubo, K. & Fukuda, S. (1955) Infestation of a Black-tailed Gull (*Larus crassirostris*) with a tick, *Ixodes signatus*, in Kabushima, Aomori Prefecture, Japan. *Miscellaneous Reports of the Research Institute for Natural Resources*, 38, 85–86 (in Japanese).
- BirdLife International (2016a) *Synthliboramphus wumizusume*. The IUCN Red List of Threatened Species 2016: e.T22694899A93475195. Downloaded on 25 January 2017.
- BirdLife International (2016b) *Synthliboramphus antiquus*. The IUCN Red List of Threatened Species 2016: e.T22694896A93474893. Downloaded on 25 January 2017.
- BirdLife International (2016c) *Hydrobates monorhis*. The IUCN Red List of Threatened Species 2016: e.T22698520A93687373. Downloaded on 25 January 2017.
- Black, W.C. IV & Piesman, J. (1994) Phylogeny of hard- and soft-tick taxa (Acari: Ixodida) based on mitochondrial 16S rDNA. *Proceedings of the National Academy of Sciences USA*, 91, 10034–10038.  
<http://dx.doi.org/10.1073/pnas.91.21.10034>
- Chae, J.S., Kim, C.M., Kim, E.H., Hur, E.J., Klein, T.A., Kang, T.K., Lee, H.C. & Song, J.W. (2003) Molecular epidemiological study for tick-borne disease (*Ehrlichia* and *Anaplasma* spp.) surveillance at selected U.S. military training sites/installations in Korea. *Annals of the New York Academy of Sciences*, 990, 118–125.  
<https://doi.org/10.1111/j.1749-6632.2003.tb07349.x>
- Dantas-Torres, F., Chomel, B.B. & Otranto, D. (2012) Ticks and tick-borne diseases: a one health perspective. *Trends in Parasitology*, 28, 437–446.  
<http://dx.doi.org/10.1016/j.pt.2012.07.003>
- Del Hoyo, J., Elliott, A. & Sargatal, J. (1996) *Handbook of the Birds of the World, vol. 3: Hoatzin to Auks*. Barcelona, Spain, Lynx Edicions, 821 pp.
- Denmark, H.A. & Clifford, C.M. (1962) A tick of the *Ornithodoros capensis* group established on Bush Key, Dry Tortugas, Florida. *Florida Entomologist*, 45, 139–142.  
<http://dx.doi.org/10.2307/3492564>
- Dietrich, M., Gómez-Díaz, E. & McCoy, K.D. (2011) Worldwide distribution and diversity of seabird ticks: implications for the ecology and epidemiology of tick-borne pathogens. *Vector-Borne and Zoonotic Diseases*, 11, 453–470.  
<https://doi.org/10.1089/vbz.2010.0009>
- Gómez-Díaz, E., Morris-Pocock, J.A., González-Solís, J. & McCoy, K.D. (2012) Trans-oceanic host dispersal explains high seabird tick diversity on Cape Verde Islands. *Biology Letters*, 23, 616–619.  
<http://dx.doi.org/10.1098/rsbl.2012.0179>
- Guglielmone, A.A., Robbins, R.G., Apanaskevich, D.A., Petney, T.N., Estrada-Peña, A., Horak, I.G., Shao, R. & Barker, S.C. (2010) The Argasidae, Ixodidae and Nuttalliellidae (Acari: Ixodida) of the world: a list of valid species names. *Zootaxa*, 2528, 1–28.
- Heath, A.C.G. (1987) A review of the origins and zoogeography of tick-borne disease in New Zealand. *Tuatara*, 29, 19–29.
- Heath, A.C.G. (2006) Observations on *Ixodes eudyptidis* Maskell (Acari: Ixodidae), *Ornithodoros capensis* Neumann (Acari: Argasidae), and other tick parasites of sea birds in New Zealand. *Systematic & Applied Acarology*, 11, 131–140.  
<http://dx.doi.org/10.11158/saa.11.2.1>
- Heath, A.C.G. & Hardwick, S. (2011) The role of humans in the importation of ticks to New Zealand: a threat to public health and biosecurity. *New Zealand Medical Journal*, 124, 1–16.
- Hoogstraal, H., Kaiser, M.N. & Easton, E.R. (1976) *Ornithodoros (Alectorobius) capensis* Neumann (Ixodoidea: Argasidae) parasitizing a human and birds nesting on islands in East African lakes. *Journal of Medical Entomology*, 12, 703–704.  
<https://doi.org/10.1093/jmedent/12.6.703>
- Hughes, L.E., Clifford, C.M., Thomas, L.A., Denmark, H.A. & Philip, O.B. (1964) Isolation and characterization of a virus from soft ticks (*Ornithodoros capensis* Group) collected on Bush Key, Dry Tortugas, Florida. *American Journal of Tropical Medicine and Hygiene*, 13, 118–122.

- <https://doi.org/10.4269/ajtmh.1964.13.118>
- Hutcheson, H.J., Gorham, C.H., Machain-Williams, C., Lorono-Pino, M.A., James, A.M., Marlenee, N.L., Winn, B., Beaty, B.J. & Blair, C.D. (2005) Experimental transmission of West Nile virus (Flaviviridae: *Flavivirus*) by *Carios capensis* ticks from North America. *Vector-Borne and Zoonotic Diseases*, 5, 293–295.  
<https://doi.org/10.1089/vbz.2005.5.293>
- Jang, W.J., Kim, J.H., Choi, Y.J., Jung, K.D., Kim, Y.G., Lee, S.H., Choi, M.S., Kim, I.S., Walker, D.H. & Park, K.H. (2004) First serologic evidence of human spotted fever group rickettsiosis in Korea. *Journal of Clinical Microbiology*, 42, 2310–2313.  
<http://dx.doi.org/10.1128/JCM.42.5.2310-2313.2004>
- Jonkers, A.H., Casals, J., Aitken, T.H. & Spence, L. (1973) Soldado virus, a new agent from Trinidadian *Ornithodoros* ticks. *Journal of Medical Entomology*, 10, 517–519.  
<https://doi.org/10.1093/jmedent/10.5.517>
- Kang, J.G., Kim, H.C., Choi, C.Y., Nam, H.Y., Chae, H.Y., Chong, S.T., Klein, T.A., Ko, S.J. & Chae, J.S. (2013) Molecular detection of *Anaplasma*, *Bartonella* and *Borrelia* species in ticks collected from migratory birds from Hong-do Island, Republic of Korea. *Vector-Borne and Zoonotic Diseases*, 13, 215–225.  
<https://doi.org/10.1089/vbz.2012.1149>
- Kawabata, H., Ando, S., Kishimoto, T., Kurane, I., Takano, A., Nogami, S., Fujita, H., Tsurumi, M., Nakamura, N., Sato, F., Takahashi, M., Ushijima, Y., Fukunaga, M. & Watanabe, H. (2006) First detection of *Rickettsia* in soft-bodied ticks associated with seabirds, Japan. *Microbiology and Immunology*, 50, 403–406.  
<http://dx.doi.org/10.1111/j.1348-0421.2006.tb03807.x>
- Kee, S., Hwang, K.J., Oh, H.B., Kim, M.B., Shim, J.C., Ree, H.I. & Park, K.S. (1994) Isolation and identification of *Borrelia burgdorferi* in Korea. *Journal of the Korean Society for Microbiology*, 29, 301–310.
- Keirans, J.E., Hutcheson, H.J. & Oliver, J.H., Jr. (1992) *Ornithodoros (Alectorobius) capensis* Neumann (Acari: Ixodoidea: Argasidae), a parasite of seabirds, established along the southeastern seacoast of the United States. *Journal of Medical Entomology*, 29, 371–373.  
<https://doi.org/10.1093/jmedent/29.2.371>
- Kim, C.M., Kim, J.Y., Yi, Y.H., Lee, M.J., Cho, M.R., Shah, D.H., Klein, T.A., Kim, H.C., Song, J.W., Chong, S.T., O'Guinn, M.L., Lee, J.S., Lee, I.Y., Park, J.H. & Chae, J.S. (2005) Detection of *Bartonella* species from ticks, mites and small mammals in Korea. *Journal of Veterinary Science*, 6, 327–334.
- Kim, H.C., Ko, S.J., Choi, C.Y., Nam, H.Y., Chae, H.Y., Chong, S.T., Klein, T.A., Sames, W.J., Robbins, R.G. & Chae, J.S. (2009a) Migratory bird tick surveillance, including a new record of *Haemaphysalis ornithophila* Hoogstraal and Kohls, 1959 (Acari: Ixodidae) from Hong-do (Hong Island), Republic of Korea. *Systematic & Applied Acarology*, 14, 3–10.  
<http://dx.doi.org/10.11158/saa.14.1.1>
- Kim, H.C., Park, C.U., Park, J.H., Kwon, Y.S., Yun, S.M., Lee, W.J., Chong, S.T., Lee, I.Y., Klein, T.A. & Robbins, R.G. (2015) *Ornithodoros sawaii* Kitaoka and Suzuki (Acari: Ixodida: Argasidae) collected from nest soil and litter of *Synthliboramphus antiquus* and *Hydrobates monorhis*, on Chilbal Island, southwestern Jeollanam Province, Republic of Korea. *Systematic & Applied Acarology*, 20, 721–730.  
<http://dx.doi.org/10.11158/saa.20.7.1>
- Kim, H.C., Choi, C.Y., Kwon, Y.S., Yun, S.M., Lee, W.J., Chong, S.T., Robbins, R.G. & Klein, T.A. (2016a) *Ornithodoros sawaii* (Acari: Ixodoidea: Argasidae) larvae collected from the seabird, *Hydrobates monorhis* in the Republic of Korea. *Korean Journal of Parasitology*, 54, 233–238.  
<https://doi.org/10.3347/kjp.2016.54.2.233>
- Kim, H.C., Kwon, Y.S., Kim, M.R., Park, C.U., Yun, S.M., Lee, W.J., Chong, S.T., Klein, T.A. & Robbins, R.G. (2016b) Ticks collected from Chilbal Island, Jeollanam Province, Republic of Korea during 2014–2015. *Systematic & Applied Acarology*, 21, 1641–1650.  
<https://doi.org/10.11158/saa.21.12.5>
- Kim, K.H., Yi, J.Y., Kim, G.Y., Choi, S.J., Jun, K.I., Kim, N.H., Gyun, P.G., Kim, N.J., Lee, J.K. & Oh, M.D. (2013) Severe fever with thrombocytopenia syndrome, South Korea, 2012. *Emerging Infectious Diseases*, 19, 1892–1894.  
<http://dx.doi.org/10.3201/eid1911.130792>
- Kim, S.Y., Jeong, Y.E., Yun, S.M., Lee, I.Y., Han, M.G. & Ju, Y.R. (2009b) Molecular evidence for tick-borne encephalitis virus in ticks in South Korea. *Medical and Veterinary Entomology*, 23, 15–20.  
<http://dx.doi.org/10.1111/j.1365-2915.2008.00755.x>

- Kimura, M. (1980) A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *Journal of Molecular Evolution*, 16, 111–120.  
<http://dx.doi.org/10.1007/BF01731581>
- Kitaoka, S. & Suzuki, H. (1973) *Ornithodoros (Alectorobius) sawaii* sp. n. (Ixodoidea, Argasidae) associated with the Streaked Shearwater, *Calonectris leucomelas*, from the Amami-Oshima Islands, Japan. *National Institute of Animal Health Quarterly*, 13, 142–148.
- Kitaoka, S. & Suzuki, H. (1974) Reports of medico-zoological investigations in the Nansei Islands. Part II. Ticks and their seasonal prevalences in southern Amami-oshima. *Sanitary Entomology*, 25, 21–26 (in Japanese).  
<https://doi.org/10.7601/mez.25.21>
- Ko, S., Kang, J.G., Kim, S.Y., Klein, T.A., Kim, H.C., Chong, S.T., Sames, W.J., Yun, S.M., Ju, Y.R. & Chae, J.S. (2010) Prevalence of tick-borne encephalitis virus in ticks from southern Korea. *Journal of Veterinary Science*, 11, 197–203.  
<https://doi.org/10.4142/jvs.2010.11.3.197>
- Kohls, G.M. (1957) Acarina: Ixodoidea. *Insects of Micronesia*, 3, 85–104.
- Lee, G.K., Ko, G.N., Jegal, G.M. & Park, C.A. (2009) Impacts of introduced plants on the breeding of Swinhoe's Storm Petrels (*Oceanodroma monorhis*) and conservation activities in Shinan-gun County, Korea. *In: Chae, H.Y., Choi, C.Y. & Nam, H.Y. (Eds.), Proceeding [sic] of the 3<sup>rd</sup> International Symposium on Migratory Birds. Seabirds in Danger: Invasive Species and Conservation of Island Ecosystem [sic]*. Shinan, Korea; Publication of the National Park Migratory Birds Center, pp. 19–36.
- Lee, K.M., Choi, Y.J., Shin, S.H., Choi, M.K., Song, H.J., Kim, H.C., Klein, T.A., Richards, A.L., Park, K.H. & Jang, W.J. (2013) Spotted fever group *Rickettsia* closely related to *Rickettsia monacensis* isolated from ticks in South Jeolla Province, Republic of Korea. *Microbiology and Immunology*, 57, 487–495.  
<https://doi.org/10.1111/1348-0421.12062>
- Morshed, M.G., Scott, J.D., Fernando, K., Beati, L., Mazerolle, D.F., Geddes, G. & Durden, L.A. (2005) Migratory songbirds disperse ticks across Canada, and first isolation of the Lyme disease spirochete, *Borrelia burgdorferi*, from the avian tick, *Ixodes auritulus*. *Journal of Parasitology*, 91, 780–790.  
<https://doi.org/10.1645/GE-3437.1>
- Murray, M.D., Palma, R.L. & Pilgrim, R.L.C. (1990) Ectoparasites of Australian, New Zealand and Antarctic birds. *In: Marchant, S. & Higgins, P.J. (Eds.), Handbook of Australian, New Zealand and Antarctic Birds. Vol. 1. Ratites to Ducks*. Melbourne, Oxford University Press, pp. 1365–1374.
- Noh, Y., Lee, Y.S., Kim, H.C., Chong, S.T., Klein, T.A., Jiang, J., Richards, A.L., Lee, H.K., & Kim, S.Y. (2017) Molecular detection of *Rickettsia* species in ticks collected from the southwestern provinces of the Republic of Korea. *Parasites & Vectors*, 10, 20.  
<https://doi.org/10.1186/s13071-016-1955-x>
- Nuttall, P.A. (1984) Tick-borne viruses in seabird colonies. *Seabird*, 7, 31–41.
- Park, K.H., Lee, S.H., Won, W.J., Jang, W.J. & Chang, W.H. (1992) Isolation of *Borrelia burgdorferi*, the causative agent of Lyme disease, from *Ixodes* ticks in Korea. *Journal of the Korean Society for Microbiology*, 27, 307–312 (in Korean).
- Park, C.U. & Takeshi, O. (2011) Present breeding status of Crested Murrelets and Ancient Murrelets on Guguldo and Chilbaldo, Shinangun. *2011 Annual Report on Migratory Bird Researches, Korean National Park Research Institute*, pp. 155–179 (in Korean).
- Park, S.W., Song, B.G., Shin, E.H., Yun, S.M., Han, M.G., Park, M.Y., Park, C. & Ryou, J. (2014) Prevalence of severe fever with thrombocytopenia syndrome virus in *Haemaphysalis longicornis* ticks in South Korea. *Ticks and Tick-borne Diseases*, 5, 975–977.  
<https://doi.org/10.1016/j.ttbdis.2014.07.020>
- Saitou, N. & Nei, M. (1987) The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution*, 4, 406–425.  
<https://doi.org/10.1093/oxfordjournals.molbev.a040454>
- Takano, A., Muto, M., Sakata, A., Ogasawara, Y., Ando S., Hanaoka, N., Tsurumi, M., Sato, F., Nakamura, N., Fujita, H., Watanabe, H. & Kawabata, H. (2009) Relapsing fever spirochete in seabird tick, Japan. *Emerging Infectious Diseases*, 15, 1528–1530.  
<https://doi.org/10.3201/eid1509.0904590.7601/mez.65.13>
- Takano, A., Fujita H., Kadosaka T., Takahashi M., Yamauchi T., Ishiguro, F., Takada, N., Yano, Y., Oikawa, Y., Honda, T., Gokuden, M., Tsunoda, T., Tsurumi, M., Ando, S., Andoh, M., Sato, K. & Kawabata, H. (2014) Construction of a DNA database for ticks collected in Japan: application of molecular identification based on the mitochondrial 16S rDNA gene. *Medical Entomology and Zoology*, 65, 13–21.

<https://doi.org/10.7601/mez.65.13>

- Tamura, K., Stecher, G., Peterson, D. Filipinski A., & Kumar, S. (2013) MEGA 6. Molecular Evolutionary Genetics Analysis. Version 6.0. *Molecular Biology and Evolution*, 28, 2731–2739.  
<https://doi.org/10.1093/molbev/mst197>
- Ushijima, Y., Oliver, J. H. Jr., Keirans, J. E., Tsurumi, M., Kawabata, H., Watanabe, H. & Fukunaga, M. (2003) Mitochondrial sequence variation in *Carios capensis* (Neumann), a parasite of seabirds, collected on Torishima Island in Japan. *Journal of Parasitology*, 89, 196–198.  
[https://doi.org/10.1645/0022-3395\(2003\)089\[0196:MSVICC\]2.0.CO;2](https://doi.org/10.1645/0022-3395(2003)089[0196:MSVICC]2.0.CO;2)
- Vander Velde, N. & Vander Velde, B. (2013) Known and potential ticks and tick-borne pathogens of Micronesia. *Micronesica*, 2013-01, 1–26.
- Yamaguti, N., Tipton, V.J., Keegan, H.L. & Toshioka, S. (1971) Ticks of Japan, Korea and the Ryukyu Islands. *Brigham Young University Science Bulletin, Biological Series*, 15, 1–226.
- Yun, S.M., Lee W.G., Ryou, J.S., Yang S.C., Park, S.W., Roh, J.Y., Lee, Y.J., Park, C. & Han, M.G. (2014) Severe fever with thrombocytopenia syndrome virus in ticks collected from humans, South Korea, 2013. *Emerging Infectious Diseases*, 20, 1358–1361.  
<http://dx.doi.org/10.3201/eid2008.131857>
- Yun, S.M., Lee Y.J., Choi, W.Y., Kim, H.C., Chong, S.T., Chang, K.S., Coburn, J. M., Klein, T.A. & Lee, W.J. (2016) Molecular detection of severe fever with thrombocytopenia syndrome and tick-borne encephalitis viruses in ixodid ticks collected from vegetation, Republic of Korea, 2014. *Ticks and Tick-borne Diseases*, 7, 970– 978.  
<https://doi.org/10.1016/j.ttbdis.2016.05.003>
- Yun, S.M., Song, B.G., Choi W.Y., Park W.I., Kim, S.Y., Roh, J.Y., Ryou, J.S., Park, C. & Shin, E.H. (2013) Prevalence of tick-borne encephalitis virus in ixodid ticks collected from the Republic of Korea during 2011–2012. *Osong Public Health and Research Perspectives*, 3, 213–221.  
<https://doi.org/10.1016/j.phrp.2012.10.004>

*Submitted: 29 Aug. 2017; accepted by Lidia Chitimia-Dobler: 28 Sept. 2017; published: 3 Nov. 2017*