

Infectious disease physicians as host: *Dermacentor variabilis* (Ixoidae: Ixodidae) in a Panamanian returning from Wisconsin, United States

Authors: Bermúdez C, Sergio, Suárez, José A., Domínguez, Lillian, and Gundacker, Nathan D.

Source: Systematic and Applied Acarology, 24(10) : 1814-1816

Published By: Systematic and Applied Acarology Society

URL: <https://doi.org/10.11158/saa.24.10.2>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Correspondence

Infectious disease physicians as host: *Dermacentor variabilis* (Ixoidae: Ixodidae) in a Panamanian returning from Wisconsin, United States

SERGIO BERMÚDEZ C^{1,2*}, JOSÉ A. SUÁREZ^{1,3}, LILLIAN DOMÍNGUEZ¹ & NATHAN D. GUNDACKER⁴

¹ Instituto Conmemorativo Gorgas de Estudios de la Salud, Panamá.

² Estación Científica Coiba AIP, Ciudad del Saber, Panamá.

³ Sistema Nacional de Investigación, Panamá.

⁴ Division of Infectious Diseases, Medical College of Wisconsin, Milwaukee, US.

* Corresponding author: sbermudez@gorgas.gob.pa

ORCID: 0000-0003-1830-3133.

Dermacentor variabilis (Say, 1821) is a species of tick native of the Nearctic Eco Zone, through southern Canada, central-eastern United States, and northern Mexico, but there are also populations in western USA, Alaska and Neotropical areas of Mexico (Guzmán-Cornejo *et al.* 2016, Durden *et al.* 2016). Adult ticks are parasites of Artiodactyla and Carnivora, with the immature ticks parasitizing mainly rodents and to a lesser extent birds or reptiles (Guglielmone & Robbins 2018). *Dermacentor variabilis* is also a frequent parasite of humans and has the capacity to transmit diseases as such tularemia, rickettsiosis and can provoked paralysis (Carter *et al.* 2016, Durden *et al.* 2016, Guglielmone & Robbins 2018). Furthermore, this species is also vector of anaplasmosis and piroplasmosis, common veterinary diseases (Durden *et al.* 2016).

Although ticks have low dispersal capacity, they can be translocate to areas outside their natural range of distribution, through migratory birds, pet trade or tourists (Morshed *et al.* 2005, Goodfellow & Shaw 2005). In this sense, there are reports of *D. variabilis* exported from the USA to Australia (Halliday & Sutherst 1990), Panama (Bermúdez *et al.* 2010) and New Zealand (Heath & Hardwick 2011). Exportation seems to be highest risk in June to July, the time when the most tick activity is seen in North America (Burg 2001).

In this paper, we report the second finding of a *D. variabilis* in Panama. A 64-year-old male infectious diseases physician traveled from Milwaukee, Wisconsin, USA, to Panama, after giving a lecture on tropical medicine. The affected and his wife visited Milwaukee in May of 2019 and walked through several prairies in the peri-urban areas of the city. During these walks, both noticed several males and females *D. variabilis* crawling on their clothes and skin, although did not notice bites. Several days later in Panama, the man observed a tick on his neck. The tick was transported to the Department of Medical Entomology of the Gorgas Memorial Institute (ICGES) in Panama and morphologically identified (Briton *et al.* 1965, Yunker *et al.* 1986). The tick was examined using a stereomicroscope Leica M 205A and photographed with a camera Leica MC 170 HD. At the time of identification, it could be observed that tissue was maintained in the hypostome, which could indicate that the tick adhered and then detached. The specimen has been deposited in the Zoological Collection "Dr. Eustorgio Méndez" in ICGES (Fig. 1, 2).

The role of exotic ticks in public health is unclear. Even so, there is a potential risk to the traveler (Chen & Wilson 2009, Ramharter *et al.* 2010). Considering that a tourist could become infected during the last days of the trip, the symptoms would appear on return to the country where medical

personnel may be unfamiliar with foreign vector borne diseases. This could complicate the diagnosis of the disease and delay treatment, especially if the history of the tick bite is not reported. For example, *D. variabilis* is a vector of *Rickettsia rickettsii* and *Francisella tularensis*; while *R. rickettsii* is considered an important pathogen in Panama (Bermúdez *et al.* 2018), there are no records of tularemia in this country. Moreover, because the species could be confused with native ixodid fauna, the correct identification of the tick species is also important to determine so such exotic diseases can be considered in tourists and pets (Halliday & Sutherst 1990, Bermúdez *et al.* 2010, Heath & Hardwick 2011).



FIGURE 1. Dorsal view of *Dermacentor variabilis*.

References

- Bermúdez, S., Miranda, R., Zaldívar, Y. & Page, K. (2010) *Dermacentor variabilis* (Ixodida: Ixodidae) in Panama: report associated with tourism. *Journal of Vector Ecology*, 35, 208–209.
<https://doi.org/10.1111/j.1948-7134.2010.00079.x>
- Bermúdez, S., Domínguez, L., Suárez, J., Daza, C., Cumbreira, A. & González, J. (2018) *Presente y pasado de las rickettsiosis en Panamá*. ISBN 978-9962-699-26-2. 30 pp.
- Brinton, E.P., Beck, D.E. & Allred, D.M. (1965) *Identification of the adults, nymphs and larvae of ticks of the genus Dermacentor Koch (Ixodidae) in the western United States*. Brigham Young University Science Bulletin, pp 1–44.
- Burg, J. (2001) Seasonal activity and spatial distribution of host-seeking adults of the tick *Dermacentor variabilis*. *Medical and Veterinary Entomology*, 15, 413–421.
<https://doi.org/10.1046/j.0269-283x.2001.00329.x>
- Carter, C., Yambem, O., Carlson, T., Hickling, G., Collins, K., Jacewicz, M. & Tsao, J. (2016) Male tick bite: a rare cause of adult tick paralysis. *Neurology Neuroimmunology and Neuroinflammation* 3, e243.
<https://doi.org/10.1212/NXI.0000000000000243>

- Chen, L. & Wilson, M. (2009) Tick-borne rickettsiosis in traveler returning from Honduras. *Emerging Infectious Diseases*, 15, 1321–1323.
<https://doi.org/10.3201/eid1508.090172>
- Durden, L., Beckmen, K. & Gerlach, R. (2016) New records of ticks (Acari: Ixodidae) from dogs, cats, humans, and some wild vertebrates in Alaska: invasion potential. *Journal of Medical Entomology*, 53, 1391–1395.
<https://doi.org/10.1093/jme/tjw128>
- Goodfellow, M. & Shaw, S. (2005) Exotic diseases of dogs and cats at risk of importation to Ireland. *Irish Veterinary Journal*, 58, 271–277.
<https://doi.org/10.1186/2046-0481-58-5-271>
- Guglielmone, A. & Robbins, R. (2018) *Hard ticks (Acari: Ixodida: Ixodidae) parasitizing humans. A global overview*. Springer International Publishing, 313 pp.
- Guzmán-Cornejo, C., Robbins, R.G., Guglielmone, A.A., Montiel-Parra, G., Rivas, G. & Pérez, T.M. (2016) The *Dermacentor* (Acari, Ixodida, Ixodidae) of Mexico: hosts, geographical distribution and new records. *ZooKeys* 569, 1–22.
<https://doi.org/10.3897/zookeys.569.7221>
- Halliday, R. & Sutherst, R. (1990) An Australian record of the American dog tick *Dermacentor variabilis*, and the risk of its establishment outside North America. *Experimental and Applied Acarology*, 8, 65–70.
<https://doi.org/10.1007/BF01193382>
- Heath, A. & Hardwick, S. (2011) The role of humans in the importation of ticks to New Zealand: a threat to public health and biosecurity. *The New Zealand Medical Journal*, 124, 67–82.
- Morshed, M., Scott, J., Fernando, K., Beati, L., Mazerolle, D., Geddes, G. & Durden, L. (2005) Migratory songbirds disperse ticks across Canada, and first isolation of the Lyme disease spirochete, *Borrelia burgdorferi*, from the avian Tick, *Ixodes auritulus*. *Journal of Parasitology*, 91, 780–790.
<https://doi.org/10.1645/GE-3437.1>
- Ramharter, M., Walochnik, J., Lagler, H., Winkler, S., Wernsdorfer, W.H., Stoiser, B. & Graninger, W. (2010) Clinical and molecular characterization of a near fatal case of human babesiosis in Austria. *Journal of Travel Medicine*, 17, 416–418.
<https://doi.org/10.1111/j.1708-8305.2010.00446.x>
- Whitten, T., Demontigny, C., Bjork, J., Foss, M., Peterson, M., Scheftel, J., Neitzel, D., Sullivan, M. & Smith, K. (2019) Prevalence of *Francisella tularensis* in *Dermacentor variabilis* ticks, Minnesota, 2017. *Vector Borne and Zoonotic Diseases*, 19, 596–603.
<https://doi.org/10.1089/vbz.2018.2388>
- Yunker, C., Keirans, J., Clifford, C. & Easton, E. (1986) *Dermacentor* ticks (Acari: Ixodidae: Ixodidae) of the New World. A scanning electron microscope atlas. *Proceedings of the Entomological Society of Washington*, 88, 609–627.

Submitted: 15 Jun. 2019; accepted by Lidia Chitimia-Dobler: 17 Jul. 2019; published: 7 Oct. 2019