

## **Investigations of Koutango Virus Infectivity and Dissemination Dynamics in *Aedes aegypti* Mosquitoes**

Authors: Lobo, Jaime M. de Araújo, Christofferson, Rebecca C., and Mores, Christopher N.

Source: Environmental Health Insights, 8(s2)

Published By: SAGE Publishing

URL: <https://doi.org/10.1177/EHI.S16005>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## Supplementary Issue: Disease Vectors

### Investigations of Koutango Virus Infectivity and Dissemination Dynamics in *Aedes aegypti* Mosquitoes

Jaime M. de Araújo Lobo<sup>1,2</sup>, Rebecca C. Christofferson<sup>2</sup> and Christopher N. Mores<sup>2,3</sup>

<sup>1</sup>Project Coordinator, Agricultural Projects Agrosop, Luanda, Angola. <sup>2</sup>Department of Pathobiological Sciences, Louisiana State University, Baton Rouge, LA, USA. <sup>3</sup>Center for Experimental Infectious Disease Research, Louisiana State University, Baton Rouge, LA, USA.

**ABSTRACT:** *Aedes aegypti* has already been implicated in the emergence of dengue and chikungunya viruses in the southern US. Vector competence is the ability of a mosquito species to support transmission of an arbovirus, which is bounded by its ability to support replication and dissemination of the virus through the mosquito body to the salivary glands to be expectorated in the saliva at the time of feeding on a vertebrate host. Here, we investigate the vector competence of *A. aegypti* for the arbovirus koutango by orally challenging mosquitoes with two titers of virus. We calculated the effective vector competence, a cumulative measure of transmission capability weighted by mosquito survival, and determined that *A. aegypti* was competent at the higher dose only. We conclude that further investigation is needed to determine the infectiousness of vertebrate hosts to fully assess the emergence potential of this virus in areas rich in *A. aegypti*.

**KEYWORDS:** arbovirus, transmission, vector-borne virus, koutango, vector competence, extrinsic incubation period

**SUPPLEMENT:** Disease Vectors

**CITATION:** Lobo et al. Investigations of Koutango Virus Infectivity and Dissemination Dynamics in *Aedes aegypti* Mosquitoes. *Environmental Health Insights* 2014;8(S2) 9–13  
doi: 10.4137/EHI.S16005.

**RECEIVED:** July 31, 2014. **RESUBMITTED:** August 30, 2014. **ACCEPTED FOR PUBLICATION:** September 2, 2014.

**ACADEMIC EDITOR:** Timothy Kelley, Editor in Chief

**TYPE:** Original Research

**FUNDING:** This work was supported by the National Institutes of Health P20GM103458, Tulane University School of Public Health and Tropical Medicine, the South Louisiana Institute for Infectious Disease Research sponsored by the Louisiana Board of Regents, and the NIH/NIGMS U01GM097661. The authors confirm that the funders had no influence over the study design, content of the article, or selection of this journal.

**COMPETING INTERESTS:** Authors disclose no potential conflicts of interest.

**COPYRIGHT:** © the authors, publisher and licensee Libertas Academica Limited. This is an open-access article distributed under the terms of the Creative Commons CC-BY-NC 3.0 License.

**CORRESPONDENCE:** rcarri1@lsu.edu

Paper subject to independent expert blind peer review by minimum of two reviewers. All editorial decisions made by independent academic editor. Prior to publication all authors have given signed confirmation of agreement to article publication and compliance with all applicable ethical and legal requirements, including the accuracy of author and contributor information, disclosure of competing interests and funding sources, compliance with ethical requirements relating to human and animal study participants, and compliance with any copyright requirements of third parties.

## Introduction

*Aedes aegypti* mosquitoes originated from Africa but are now found in tropical regions around the world.<sup>1,2</sup> Urbanization particularly has contributed to the increase of *Aedes* mosquito populations, especially *A. aegypti* and *Aedes albopictus*.<sup>1,2</sup> There has been a threefold increase in urban human population densities in Africa since the 1950s, and even larger increases have occurred in Asia and the Americas.<sup>3</sup> With the related increase in contact between urban-associated *Aedes* mosquito populations, arboviruses, especially dengue virus (DENV), have established endemic transmission in these areas.<sup>4</sup>

*A. aegypti* is found in the subtropical region of the United States and in popular tropical tourist destinations, and has

been implicated in the recent chikungunya (CHIKV) outbreak in the Caribbean resulting in over 100 imported and several locally acquired cases in the US.<sup>5–7</sup> Furthermore, *A. aegypti* has been implicated as the primary vector of DENV in the southern United States.<sup>8</sup> *A. aegypti* are primarily anthrophilic, preferring to bite humans than other potential hosts, and have adapted their behavior and ecology to maximize contact with humans.<sup>9</sup> For instance, they tend to be closely associated with human domiciles and breed in (often man-made) containers near houses.<sup>9,10</sup>

Koutango virus (KOUTV) was first isolated in 1968 in Somalia, and it has been suggested that this virus is a variant of West Nile virus (WNV).<sup>11–14</sup> However, species of the main vector genus of WNV (*Culex* spp.) were not found



to be competent for KOUTV,<sup>13</sup> indicating that questions regarding the transmission cycle of this virus remain. Early observations of transovarial transmission of KOUTV and subsequent competence of emerged *A. aegypti* females suggest this as a potential vector in the transmission cycle as a real possibility<sup>15</sup> and transmission of KOUTV by *A. aegypti* viruses has been observed.<sup>15</sup> However, recent studies involving KOUTV have not focused on the mosquito, and there are still some questions regarding the potential transmission dynamics of this virus. To address this, we experimentally investigated the dynamics of KOUTV infectivity to the mosquito *A. aegypti* and determined the effective vector competence (EVC)<sup>16</sup> to characterize the threat this virus poses in areas where *A. aegypti* is present, such as in the southern United States.

Materials and Methods

**Vector competence.** *Virus.* The KOUTV strain utilized in this experiment was the KOUTV DAK Ar D 5443 received from Robert B. Tesh (CBEID-UTMB). Virus was propagated by inoculating 100 µL of viral stock into the T-75 flask of confluent Vero cells. To determine titer, a plaque assay was developed and standard curves created for quantitative reverse transcription polymerase chain reaction (qRT-PCR) as previously described.<sup>16</sup> Briefly, we utilized a SuperScript III One-step qRT-PCR kit (Invitrogen, Carlsbad, CA) with the primers and probes given in Table 1.

The presence of KOUTV viral RNA was detected via qRT-PCR using the following protocol: RT step (1 cycle) 48° C for 2 minutes and 95 °C for 2 minutes, amplification and data recording step (40 cycles) 95 °C for 15 seconds and 60 °C for 30 seconds. Primers were designed and obtained via Integrated DNA technologies with 5'-FAM fluorophore and 3'-Black-Hole quencher. Prior to offering virus to mosquitoes, virus was propagated into Vero cells. Viral standard curves and concentrations were obtained via plaque assay as described previously before the beginning of the experiment.<sup>16</sup>

**Mosquito Exposure and Detection of Disseminated Infection.** *A. aegypti* (Linnaeus, 1762) Rockefeller strain were reared from the colony at Louisiana State University School of Veterinary Medicine. Mosquitoes were orally exposed to an artificial bloodmeal 3–5 days post emergence with 10<sup>9</sup> plaque forming units (PFU)/mL or 10<sup>6</sup> PFU/mL. Feeding occurred as previously described.<sup>16</sup> Briefly, mosquitoes were reared and maintained at constant environmental parameters of 28 °C, 75–80% humidity, and a 16:8 light:dark photoperiod.<sup>16</sup>

Mosquitoes were then orally challenged using the Hemotect (Discovery Workshops, Arrington, Lancashire, UK) as in Supernatant from virus-infected cell culture was mixed with bovine blood (Hemostat, Dixon, CA) at a ratio of 2:1 and viral titers verified by qRT-PCR as in.<sup>16</sup>

After feeding, fully engorged females were placed into clean cartons and then sampled at days 3, 5, 7 and 11 post exposure (dpe) to test for dissemination status. It has been previously shown that detectable arbovirus in the legs is indicative of a fully disseminated infection; thus the salivary glands are presumed to be infected as well.<sup>17</sup> Mosquito legs were removed and analyzed separately for infection from the bodies as previously described.<sup>16,18–21</sup> RNA was extracted using the MagMax-96 kit (Ambion) on a King Fisher® nucleic acid extraction instrument according to the manufacturer's instructions (Thermo Scientific). After extraction, the samples were tested for the presence of dengue viral RNA via qRT-PCR using the following protocol: RT step (1 cycle) 48 °C for 2 minutes and 95 °C for 5 minutes; amplification and data recording step (40 cycles) 95 °C for 15 seconds and 60 °C for 30 seconds.

**Data analysis.** Vector competence was calculated as the number of disseminated infections divided by the total number of mosquitoes that fed to repletion (number exposed), EVC was calculated as in Ref. 16. All analyses and modeling was performed in R (version 3.0.1). EVC is a cumulative measure of vector competence over a range of days, bounded and weighted by the mosquito lifespan. Briefly, define the rate of change of vector competence over days as *b(N)*:

b(N) = β<sub>0</sub> + β<sub>1</sub>N

where β<sub>1</sub> = rate of change of vector competence over time. EVC, represented by the phi (ϕ) is then given by:

ϕ = ∫<sup>N</sup> p<sup>N</sup> b(N) dN

where *N* is the extrinsic incubation period, here a range of days defined by the study duration; *b(N)* is as above; and *P* = probability of daily survival. This is held constant so as to isolate the viral phenotypes and assess relative fitness.<sup>16</sup> EVC gives a measure of viral fitness while entomological and vertebrate density parameters are held.

To obtain 95% confidence intervals (CIs), we employed a bootstrapping method as in Ref. 16. Per day of study, the number of mosquitoes with disseminated infections was coded as 1 and those without as 0. We then resampled (with replacement) for *n* = 1000 iterations for a bootstrapped distribution of

Table 1. Primers and probe sequences.

FORWARD PRIMER SEQUENCE	REVERSE PRIMER SEQUENCE	PROBE SEQUENCE
ACCAGGAGGCAAGA TTTACG	CGCTTTGGTTATC CGTGTG	ACAAGAGGCAAGATTTCG CAGACCGCTGGCTGGGACACACGGATAACCAAGCG

Notes: The sequences for the primers and probe used to amplify KOUTV and CHIKV are given. All sequences are 5' → 3'.

dissemination rates (for each set of  $N$  study days). The  $b(N)$  was then calculated for each 1000 sets of dissemination over  $N$  days. EVC ( $\phi$ ) was calculated for each set of replicated data and the distribution of bootstrapped  $\phi$  was obtained (Fig. 1), from which we obtained the 95% CI.

## Results

**KOUTV vector competence.** No mosquitoes had a disseminated infection after exposure to  $10^6$  PFU/mL, indicating that at this titer, *A. aegypti* are not competent for KOUTV. However, the higher KOUTV titer exposure resulted in disseminated infections as early as 3 days post exposure (Table 2). The dissemination rates were calculated as 2.22%, 8.51%, 17.24%, and 55.56% on 3, 5, 7, and 11 days post exposure, respectively (Fig. 2).

The calculated EVC of KOUTV was 857 indicating that 85.7% of mosquitoes achieved a disseminated infection and were still alive at the end of the 11-day period of the study. The calculated 95% CI of KOUTV EVC was [0.701,1.15]. For interpretative purposes, the upper confidence limit (UCL) is asymptotically bounded by 1, because proportion has a range [0,1]. However, the function of  $\phi$  is not based on a probability density function and thus mathematically has no upper asymptote; therefore, the mathematical 97.5% quantile may exceed 1 as it did here (Fig. 2). Figure 3 shows the relationship among the raw data, the  $b(N)$  for the data, and the curve resulting from weighting  $b(N)$  by the mortality rate of the mosquito vector.

## Discussion

Vector competence is a critical determinant of the success of a transmission cycle. It is affected by both intrinsic (genetically determined virus and vector) and extrinsic (environmental and ecological) factors.<sup>22</sup> Specifically, transmission success of a virus by arthropod vectors is dependent on factors such as population density of the mosquitoes, susceptibility of


**Table 2.** *A. aegypti* vector competence for KOUTV.

DAYS POST EXPOSURE	LEGS		TOTAL TESTED
	UNINFECTED	INFECTED	
3	44	1	45
5	43	4	47
7	24	5	29
11	15	21	36
Total	153	34	187


**Note:** Proportion of mosquitoes with uninfected or infected legs after oral challenge with  $10^9$  PFU/mL on 3, 5, 7, and 11 days post exposure.

amplifying hosts, environmental temperature, and the feeding preferences and habits.<sup>23</sup> In South Florida, this mosquito species has already shown that it is competent for DENV and CHIKV, where local transmission has been reported.<sup>5,8</sup>

While we reported significant KOUTV dissemination rates in *A. aegypti* challenged with a high dose of KOUTV, it is important to consider other factors that would alter the vector competence of KOUTV. First, vector competence for DENV and CHIKV can vary significantly among populations of the same species of mosquito vector.<sup>24–26</sup> It is likely that such variability would also be seen for KOUTV across *A. aegypti* populations. Second, genetic diversity among strains of DENV (within and among serotypes) and CHIKV affects vector competence and transmission potential.<sup>26,27</sup> Likewise, this could affect KOUTV transmission, although there is a lack of data regarding KOUTV diversity. Third, extrinsic factors such as seasonality and mosquito ecology can affect vector competence.<sup>23,28,29</sup> Little is known about the interface of KOUTV and *A. aegypti* ecology and remains to be intensively investigated. Lastly, as our methods are dependent on mosquito mortality, it is important to recognize that laboratory conditions promote longevity where field-mosquito mortality is likely highly variable and influenced by many extrinsic factors.




**Figure 1.** Histogram of bootstrapped EVC values. The bootstrapped distribution of KOUTV EVC values from 1000 simulations demonstrates how the 95% CI is determined.



**Figure 2.** *A. aegypti* vector competence for KOUTV. Dissemination rates of high-dose ( $10^9$ ) KOUTV in *A. aegypti* at 3, 5, 7, and 11 days post exposure.





**Figure 3.** Relationship among the raw data, the  $b(N)$  for the data, and the curve resulting from weighting  $b(N)$  by the mortality rate of the mosquito vector. The raw data (thinner, black dotted line) connect the points experimentally determined at days 3, 5, 7, and 11 days post exposure, while the  $b(N)$  (thicker, red dashed line) is the best linear fit of the raw data. The curve (solid black line) is the fitted line weighted by the mortality of the mosquito vector ( $p^M$ ). EVC is determined by integrating to find the area under this curve.

Lastly, KOUTV reached high dissemination rates after exposure to high titer ( $10^9$  PFU/mL), but was undetected in the legs of mosquitoes challenged with a lower dose ( $10^6$  PFU/mL). This disparity in success of infection corresponding to the dose titer offered to mosquitoes indicates that the infectiousness of KOUTV may be limited to high doses. The epidemiological relevance of this finding is that only periods of high viremia will render a human (or potentially other vertebrate reservoirs) infectious. Given the relatively high dose required to infect mosquitoes and the role of infectiousness of humans in the probability of emergence, it is unlikely that KOUTV poses the same threat as other arboviruses such as dengue and chikungunya.<sup>5,8,23</sup> However, since little is known about the infectivity of KOUTV in vertebrates, it is possible that serum viremia levels reach high titers capable of transmitting to mosquitoes.

## Conclusions

*A. aegypti* pose a major health threat as they are competent for several arboviruses. Given the repeated introductions and eventual emergence of DENV and CHIKV in the southern United States, it is important to understand the potential for other viruses to emerge. Critical to our preparedness is an understanding of the potential contributors to transmission cycle(s) capable of maintaining these viruses. Here, we show that *A. aegypti* is capable of supporting KOUTV transmission at high titers. Therefore, more data are needed to determine the viremia titers of people infected with KOUTV or to determine what other potential reservoir species could support adequate viral loads.

## Author Contributions

Conceptualized and performed experiments: JML, RCC, and CNM. Analyzed data: JML and RCC. Wrote the paper: JML, RCC, and CNM. Agree with manuscript results and conclusions: JML, RCC, CNM. Jointly developed the structure and arguments for the paper: JML, RCC, CNM. Made critical revisions and approved final version: JML, RCC, CNM. All authors reviewed and approved of the final manuscript.

## REFERENCES

- Gubler DJ. Dengue and dengue hemorrhagic fever. *Clin Microbiol Rev.* 1998 Jul;11(3):480–96.
- Halstead SB. Dengue virus-mosquito interactions. *Annu Rev Entomol.* 2008; 53:273–91.
- Demographic year book, 1950–2007. In: Division S, editor: United Nations; 2009.
- Appawu M, Dadzie S, Abdul H, Asmah H, Boakye D, Wilson M, et al. Surveillance of viral haemorrhagic fevers in Ghana: entomological assessment of the risk of transmission in the northern regions. *Ghana Med J.* 2006;40(4):137–41.
- Florida Department of Health, Florida Health Series, Florida Health, 2014.
- Americas Geographic Distribution | Chikungunya virus | CDC. 2014/07/31/T17:56:30Z [Available from: <http://www.cdc.gov/chikungunya/geo/americas.html>], last updated September 17, 2014.
- Leparc-Goffart I, Nougaiere A, Cassadou S, Prat C, de Lamballerie X. Chikungunya in the Americas. *Lancet.* February 8, 2014;383(9916):514.
- Florida Department of Health, Florida Health Series, Florida Health, 2013.
- Weaver SC, Reisen WK. Present and future arboviral threats. *Antiviral Res.* 2010 Feb;85(2):328–45.
- Azil AH, Bruce D, Williams CR. Determining the spatial autocorrelation of dengue vector populations: influences of mosquito sampling method, covariables, and vector control. *J Vector Ecol.* 2014 Jun;39(1):153–63.
- Charrel RN, Brault AC, Gallian P, Lemasson JJ, Murgue B, Murri S, et al. Evolutionary relationship between Old World West Nile virus strains. Evidence for viral gene flow between Africa, the Middle East, and Europe. *Virology.* October 25, 2003;315(2):381–8.
- Prow NA, Setoh YX, Biron RM, Sester DP, Kim KS, Hobson-Peters J, et al. The West Nile-like flavivirus Koutango is highly virulent in mice due to delayed viral clearance and the induction of a poor neutralizing antibody response. *J Virol.* June 18, 2014.
- Fall G, Diallo M, Loucoubar C, Faye O, Sall AA. Vector competence of *Culex neavei* and *Culex quinquefasciatus* (Diptera: Culicidae) from Senegal for lineages 1, 2, Koutango and a putative new lineage of West Nile virus. *Am J Trop Med Hyg.* 2014 Apr;90(4):747–54.
- Butenko AM, Semashko IV, Skvortsova TM, Gromashevskii VL, Kondrashina NG. [Detection of the Koutango virus (Flavivirus, Togaviridae) in Somalia]. *Med Parazitol (Mosk).* 1986 May–Jun;3(3):65–8.
- Coz J, Valade M, Cornet M, Robin Y. [Transovarian transmission of a Flavivirus, the Koutango virus, in *Aedes aegypti* L.]. *C R Acad Sci Hebd Seances Acad Sci D.* July 5, 1976;283(1):109–10.
- Christofferson RC, Mores CN. Estimating the magnitude and direction of altered arbovirus transmission due to viral phenotype. *PLoS One.* 2011;6(1):e16298.
- Turell MJ, Gargan TP, 2nd, Bailey CL. Replication and dissemination of Rift Valley fever virus in *Culex pipiens*. *Am J Trop Med Hyg.* 1984 Jan;33(1):176–81.
- Alto BW, Lounibos LP, Mores CN, Reiskind MH. Larval competition alters susceptibility of adult *Aedes* mosquitoes to dengue infection. *Proc Biol Sci.* February 22, 2008;275(1633):463–71.
- Pesko K, Westbrook CJ, Mores CN, Lounibos LP, Reiskind MH. Effects of infectious virus dose and bloodmeal delivery method on susceptibility of *Aedes aegypti* and *Aedes albopictus* to chikungunya virus. *J Med Entomol.* 2009 Mar;46(2):395–9.
- Reiskind MH, Pesko K, Westbrook CJ, Mores CN. Susceptibility of Florida mosquitoes to infection with chikungunya virus. *Am J Trop Med Hyg.* 2008 Mar;78(3):422–5.
- Turell MJ, Dohm DJ, Mores CN, Terracina L, Wallette DL, Jr., Hribar LJ, et al. Potential for North American mosquitoes to transmit Rift Valley fever virus. *J Am Mosq Control Assoc.* 2008 Dec;24(4):502–7.
- Tabachnick WJ. Nature, nurture and evolution of intra-species variation in mosquito arbovirus transmission competence. *Int J Environ Res Public Health.* 2013 Jan;10(1):249–77.
- Christofferson RC, Mores CN, Wearing HJ. Characterizing the likelihood of dengue emergence and detection in naive populations. *Parasit Vectors.* 2014;7(1):282.

24. Pongsiri A, Ponlawat A, Thaisomboonsuk B, Jarman RG, Scott TW, Lambrechts L. Differential susceptibility of two field *aedes aegypti* populations to a low infectious dose of dengue virus. *PLoS One*. 2014;9(3):e92971.
25. Ye YH, Ng TS, Frentiu FD, Walker T, van den Hurk AF, O'Neill SL, et al. Comparative susceptibility of mosquito populations in North Queensland, Australia to oral infection with dengue virus. *Am J Trop Med Hyg*. 2014 Mar;90(3):422–30.
26. Vega-Rua A, Zouache K, Girod R, Failloux AB, Lourenco-de-Oliveira R. High level of vector competence of *Aedes aegypti* and *Aedes albopictus* from ten American countries as a crucial factor in the spread of Chikungunya virus. *J Virol*. 2014 Jun;88(11):6294–306.
27. Guo XX, Zhu XJ, Li CX, Dong YD, Zhang YM, Xing D, et al. Vector competence of *Aedes albopictus* and *Aedes aegypti* (Diptera: Culicidae) for DEN2-43 and New Guinea C virus strains of dengue 2 virus. *Acta Trop*. 2013 Dec;128(3):566–70.
28. Aziz S, Aidil RM, Nisfariza MN, Ngui R, Lim YA, Yusoff WS, et al. Spatial density of *Aedes* distribution in urban areas: A case study of breteau index in Kuala Lumpur, Malaysia. *J Vector Borne Dis*. 2014 Apr–Jun;51(2):91–6.
29. Stoddard ST, Wearing HJ, Reiner RC, Jr., Morrison AC, Astete H, Vilcarromero S, et al. Long-term and seasonal dynamics of dengue in Iquitos, Peru. *PLoS Negl Trop Dis*. 2014 Jul;8(7):e3003.