

Assessment of Navel Oranges, Clementine Tangerines, and Rutaceous Fruits as Hosts of *Bactrocera cucurbitae* and *Bactrocera latifrons* (Diptera: Tephritidae)

Authors: Mcquate, Grant T., Follett, Peter A., Liquido, Nicanor J., and Sylva, Charmaine D.

Source: International Journal of Insect Science, 7(1)

Published By: SAGE Publishing

URL: <https://doi.org/10.1177/IJIS.S20069>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Assessment of Navel Oranges, Clementine Tangerines, and Rutaceous Fruits as Hosts of *Bactrocera cucurbitae* and *Bactrocera latifrons* (Diptera: Tephritidae)

Grant T. McQuate¹, Peter A. Follett¹, Nicanor J. Liquido² and Charmaine D. Sylva¹

¹USDA-ARS, Daniel K. Inouye U.S. Pacific Basin Agricultural Research Center, Hilo, HI, USA. ²USDA-APHIS-PPQ, CPHST, Plant Epidemiology and Risk Analysis Laboratory, Honolulu, HI, USA.

ABSTRACT: Export of *Citrus* spp. fruits may require risk mitigation measures if grown in areas with established tephritid fruit fly (Diptera: Tephritidae) populations capable of infesting the fruits. The host status of *Citrus* spp. fruits is unclear for two tephritid fruit fly species whose geographic ranges have expanded in recent years: melon fly, *Bactrocera cucurbitae* (Cocquillet), and *Bactrocera latifrons* (Hendel). In no choice cage infestation studies, *B. latifrons* oviposited into intact and punctured Washington navel oranges (*Citrus sinensis* [L.] Osbeck) and Clementine tangerines (*C. reticulata* L. var. Clementine), but eggs rarely developed to the adult stage. *B. cucurbitae* readily infested intact and punctured tangerines, and to a lesser extent punctured oranges, but did not infest intact oranges. Limited cage infestation and only a single literature report of field *Citrus* spp. infestation suggest that risk mitigation of *Citrus* spp. for *B. latifrons* is not needed. Risk mitigation options of *Citrus* spp. for *B. cucurbitae*, including heat and cold treatments and systems approaches, are discussed.

KEYWORDS: citrus, host status, *Bactrocera cucurbitae*, *Bactrocera latifrons*, melon fly, risk mitigation

CITATION: McQuate et al. Assessment of Navel Oranges, Clementine Tangerines, and Rutaceous Fruits as Hosts of *Bactrocera cucurbitae* and *Bactrocera latifrons* (Diptera: Tephritidae). *International Journal of Insect Science* 2015;7 1–9 doi:10.4137/IJIS.S20069.

RECEIVED: September 10, 2014. **RESUBMITTED:** November 16, 2014. **ACCEPTED FOR PUBLICATION:** December 2, 2014.

ACADEMIC EDITOR: Helen Hull-Sanders, Editor in Chief

TYPE: Original Research

FUNDING: This research was supported by base funds allocated to the USDA-ARS, Daniel K. Inouye U.S. Pacific Basin Agricultural Research Center, Hilo, HI. The authors confirm that the funder had no influence over the study design, content of the article, or selection of this journal.

COMPETING INTERESTS: Authors disclose no potential conflicts of interest.

COPYRIGHT: © the authors, publisher and licensee Libertas Academica Limited. This is an open-access article distributed under the terms of the Creative Commons CC-BY-NC 3.0 License.

CORRESPONDENCE: grant.mcquate@ars.usda.gov

DISCLAIMER: This article reports the results of this research only. Mention of a proprietary product does not constitute an endorsement or a recommendation by the USDA for its use. USDA is an equal opportunity provider and employer.

Paper subject to independent expert blind peer review by minimum of two reviewers. All editorial decisions made by independent academic editor. Upon submission manuscript was subject to anti-plagiarism scanning. Prior to publication all authors have given signed confirmation of agreement to article publication and compliance with all applicable ethical and legal requirements, including the accuracy of author and contributor information, disclosure of competing interests and funding sources, compliance with ethical requirements relating to human and animal study participants, and compliance with any copyright requirements of third parties. This journal is a member of the Committee on Publication Ethics (COPE).

Introduction

Citrus spp. (Rutaceae family) are believed to be native to tropical and subtropical regions of Southeast Asia, but are now widely cultivated throughout the tropics and subtropics.¹ Throughout their geographic range of distribution, *Citrus* spp. fruits can be subject to infestation by a range of different tephritid (Diptera: Tephritidae) fruit fly species. Although some tephritid fruit fly species, like Mediterranean fruit fly, *Ceratitidis capitata* (Wiedemann), and Oriental fruit fly, *Bactrocera dorsalis* (Hendel), can be major pests of *Citrus* spp.,^{2,3} the host status of *Citrus* spp. is less clear for other tephritid fruit fly species, such as melon fly, *Bactrocera cucurbitae* (Cocquillet), and *Bactrocera latifrons* (Hendel). Both these species have expanded their ranges

far beyond Southeast Asia into some Pacific Islands^{4,5} and into the African continent.^{6,7} If *Citrus* spp. are hosts of these fruit fly species, then regulatory procedures would need to be developed in countries of citrus production where these fruit fly species are present in order to minimize the risk of introducing these fruit flies during export of fruits to areas where they are not established. A pest risk assessment (PRA), aimed to examine plant pest risks associated with the movement into the continental United States of *Citrus* spp. fruits (*Citrus paradisi* Macfaden, *Citrus limon* [L.] Burm. f., *Citrus aurantiifolia* [Christmann] Swingle, *C. sinensis* [L.] Osbeck, *Citrus grandis* [L.] Osbeck, and *C. nobilis* Lour. var. *deliciosa* [Ten.] Swingle) grown in Hawaii, concluded that *B. dorsalis*, *B. cucurbitae*, and



C. capitata are high-risk pests of citrus fruits, and that specific phytosanitary measures were strongly recommended to achieve quarantine security mandated by USDA-APHIS-PPQ.⁸ The Hawaii Department of Agriculture (HDOA) petitioned the USDA-Animal and Plant Health Inspection Service (APHIS) to approve a cold treatment schedule ($\leq 0.99^{\circ}\text{C}$, for 17 days, or $\leq 1.38^{\circ}\text{C}$ for 20 days) as a quarantine treatment to mitigate fruit fly infestation in *C. sinensis*.

Further data on the infestability of citrus by *B. cucurbitae* and *B. latifrons* are, however, needed to better establish the host status of *Citrus* spp. Here, we (1) present results of laboratory trials that assess the infestability of two citrus species (Washington navel oranges, *C. sinensis* (L.) Osbeck, and Clementine tangerines, *C. reticulata* L. var. Clementine) by *B. cucurbitae* and *B. latifrons*, and (2) summarize references in published literature to infestation of fruits belonging to the plant family Rutaceae by *B. cucurbitae* and *B. latifrons*. Potential regulatory procedures are discussed to mitigate the risk of introduction of fruit fly pests exported from areas with established *B. cucurbitae* and/or *B. latifrons* populations.

Materials and Methods

Insect colonies. *B. latifrons* and *B. cucurbitae* flies used in experiments were obtained from laboratory colonies at the USDA ARS Daniel K. Inouye U.S. Pacific Basin Agricultural Research Center in Hilo, HI. The *B. latifrons* colony has been maintained for over 20 years (over ~208 generations), and the *B. cucurbitae* colony has been maintained for over 36 years (over ~478 generations) with infrequent infusion of wild flies. Fruit flies used in our tests were kept in an insectary at 24–27°C, 65–70% RH, and a photoperiod of 12:12 (L:D) hours. Adults were fed water and a diet of sugar cubes and a “protein cake” consisting of three parts of sucrose, one part of protein yeast hydrolysate (Enzymatic; United States Biochemical Corporation, Cleveland, OH), and 0.5 part of torula yeast (Lake States Division, Rhinelander Paper Co., Rhinelander, WI) from the time of emergence from puparia until noon, the day before the experiment, at which time, cohorts of 50 gravid females were placed with a water wick and two sugar cubes (no “protein cake”), in 26.5 × 26.5 × 26.5 cm cubical screened cages. When fruits were added (see below), the water wick remained in the cage, but the two sugar cubes were removed. Adult flies were approximately 16–18 days old at the start of the experiments.

Bioassays. Bioassays were conducted from 24 October, 2012, to 2 August, 2013. Fruits used were randomly selected from unblemished fruits available at a local grocery store. For each bioassay, a thoroughly rinsed single harvest-mature fruit (either navel orange or tangerine) was weighed and then placed in each of the eight cages. The fruit in four of the cages was undamaged (intact), while the fruit in the other four cages was randomly punctured 50 times using a 1.0 mm diameter probe, with probes penetrating to a depth of 1.0 cm. An undamaged control fruit, known to be a good host of the fruit fly species being tested, was placed in another cage. This group of nine

concurrent cage tests is hereafter referred to as a “trial.” Trials were conducted separately for *B. latifrons* and *B. cucurbitae*. For *B. latifrons*, the control fruit was either eggplant (*Solanum melongena* L.), Anaheim pepper (*Capsicum annuum* L. var. Anaheim), or papaya (*Carica papaya* L.). For *B. cucurbitae*, the control fruit was papaya. Fruits were introduced into the cages with 50 gravid female flies at 9:00 am and removed after 24 hours. Holding conditions during the time of fruit exposure were 24–27°C, 65–70% RH, and a photoperiod of 12:12 (L:D) hours. Following fruit fly exposure, fruits were transferred to 5 L screen-topped HI-PLAS buckets (Highland Plastics, Inc., Mira Loma, CA), which held a 300 mL layer of sand on the bottom to serve as a pupariation medium. After 2 weeks, sand from the buckets was sieved, and fruits cut open to recover all pupariating larvae and pupae, which were then transferred to 7.0 cm (diameter) × 7.5 cm screened-top cups with 20 mL sand and held for adult emergence. Numbers of pupae recovered and the number of emerged adults were recorded for each fruit. For the Clementine tangerine trials, data are reported only for those bioassays where at least 10 flies were recovered from the associated control fruit.

Statistical analyses. Separate statistical analyses were performed for each fruit species (including results of both fruit fly species) with their respective controls. For each fruit species, significance of differences of pupal and adult recoveries per kilogram fruit among intact, punctured, and control fruits for both fruit fly species was tested using analysis of variance (ANOVA) on trial averages, after square root transformation ($\sqrt{\text{catch} + 0.5}$). The data for the ANOVA came from the average *B. latifrons* recovery from intact test fruits for each trial, the average *B. latifrons* recovery from punctured test fruits for each trial, the average *B. latifrons* recovery from the control fruit from each trial, the average *B. cucurbitae* recovery from intact test fruits for each trial, the average *B. cucurbitae* recovery from punctured test fruits for each trial, and the average *B. cucurbitae* recovery from the control fruit from each trial. Separate ANOVAs were performed for pupal recovery per kilogram fruit for navel oranges and for tangerines, and for adult recovery per kilogram fruit for navel oranges and for tangerines. Tukey’s honest significant difference (HSD) was used to test for mean separation. Statistical analyses were performed using JMP 10.0.0.⁹ Untransformed data are presented in the summary charts.

Literature review. References to infestation of fruits in the family Rutaceae by *B. cucurbitae* and *B. latifrons* were taken from host status summaries on host plants presented in various state, national and international host listings, as well as in scientific publications indexed in searchable databases, such as Agricola, CAB Abstracts, and Scopus. Host data also were obtained from pest interceptions reported by U.S. Federal and State governments.

Host data were classified as “field infestation data,” “laboratory infestation data,” or as “listing only” if no supporting data were provided. For field and laboratory infestation data, a summary was prepared detailing the number of fruits collected, from where the fruits were collected, the condition

of the fruits, and the level of infestation found, when this information was available.

Results

In the navel orange trials, there was a significant difference among treatments in pupal recovery per kilogram fruit ($F = 72.1$; $df = 6, 104$; $P < 0.0001$), and in adult recovery per kilogram fruit ($F = 19.7$; $df = 6, 104$; $P < 0.0001$). Pupal recovery per kilogram fruit, for both fruit fly species, was significantly greater for control fruits than for any other treatment groups. *B. cucurbitae* pupal recovery per kilogram fruit was significantly greater for punctured fruits than for intact fruits, where there was no recovery from any tested fruit. *B. cucurbitae* pupal recovery per kilogram fruit from punctured fruits was also greater than *B. latifrons* pupal recovery per kilogram fruit for either punctured or intact treatment fruits (Table 1). There was no significant difference among *B. latifrons* pupal recovery per kilogram fruit in intact or punctured fruits. Relative magnitudes of adult per kilogram recovery among the different treatment groups were comparable with the pupae per kilogram fruit recoveries, but the difference in *B. latifrons* recovery per kilogram fruit between intact control fruits and punctured navel oranges was not significantly different from the *B. cucurbitae* recovery from punctured navel oranges (Table 1).

In the Clementine tangerine trials, there was significant difference among treatments in pupal recovery per kilogram fruit ($F = 17.1$; $df = 5, 66$; $P < 0.0001$) and in adult recovery per kilogram fruit ($F = 13.5$; $df = 5, 66$; $P < 0.0001$). *B. cucurbitae* pupal recoveries per kilogram fruit from control fruits and from both intact and punctured treatment fruits were not significantly different, but were significantly greater than *B. latifrons* recovery from both intact and punctured treatment fruits. *B. latifrons* pupal recoveries per kilogram fruit from control fruits were significantly greater than *B. latifrons* recovery per kilogram fruit from both intact and punctured treatment fruits. Relative magnitudes of adult per kilogram recovery among the different treatment groups were comparable with the pupae per kilogram fruit recoveries (Table 1).

Bactrocera latifrons.

Bioassays.

Navel oranges. Out of a total of 16 trials, one trial with intact fruits and nine trials with punctured fruits had infested fruits. Considering actual fruit numbers, one of the 64 intact navel oranges and ten of the 64 punctured navel oranges were infested, whereas 100% of the control fruits (nine of the nine eggplants and seven of the seven papayas) were infested. Overall recovery from the intact and punctured navel oranges averaged 0.36 pupae/kg fruit and 5.51 pupae/kg fruit, respectively, whereas 454.1 pupae/kg fruit (eggplant) and 333.1 pupae/kg fruit (papaya) were recovered from the nine control eggplants and the seven control papayas (Table 1).

Clementine tangerines. Out of a total of 20 trials, at least 10 adult flies were recovered from control Anaheim peppers in 12 trials (Table 1). Of those 12 trials, one had infestation in

intact fruits and two had infestation in punctured fruits. Considering actual fruit numbers, one of the 48 intact Clementine tangerines and two of the 48 punctured Clementine tangerines were infested, whereas 100% of the Anaheim peppers were infested. Overall recovery from the intact and punctured tangerines averaged 0.51 pupae/kg fruit and 0.39 pupae/kg fruit, respectively, whereas 378.7 pupae/kg fruit were recovered from the 12 control peppers (Table 1).

Literature review. There are only two reports of field infestation of fruits of the plant family Rutaceae by *B. latifrons*, no reports of laboratory infestations, and ten “listing only” references (Table 2). One field infestation report is a citrus species, lime, *Citrus aurantiifolia* [Christm.] Swingle; the other is a non-citrus species, mock orange, *Murraya paniculata* [L.] Jack. Both these infestations came from extensive fruit collections in Malaysia and Thailand.¹¹ In both cases, *B. latifrons* was recovered from only one collection. The publication, however, did not report the total number of fruits included in the collection or the total number of collections made. The “listing only” references come from two *Citrus* spp.

Bactrocera cucurbitae.

Bioassays.

Navel oranges. Out of a total of 21 trials, no trials with intact fruits and 17 trials with punctured fruits had infested fruits. Considering actual fruit numbers, zero of the 84 intact navel oranges and 44 of the 84 punctured navel oranges were infested, whereas 100% of the control papayas were infested. Overall recovery from the intact and punctured navel oranges averaged 0.0 pupae/kg fruit and 84.5 pupae/kg fruit, respectively, whereas 466.7 pupae/kg fruit were recovered from the control fruits (papaya).

Clementine tangerines. Out of a total of 13 trials, at least 10 adult flies were recovered from control papayas in 12 trials. In these 12 trials, eight trials with intact fruits and eleven trials with punctured fruits had infested fruits. Considering actual fruit numbers, 15 of the 48 intact Clementine tangerines and 28 of the 48 punctured tangerines were infested, whereas 100% of the control papayas were infested. Overall recovery from the intact and punctured Clementine tangerines averaged 340.8 and 240.1 pupae/kg fruit, respectively, whereas 310.2 pupae/kg fruit were recovered from the control fruit (papaya).

Literature review. There are ten reports of field infestation (covering six species), three reports of laboratory infestation (covering three separate species), and 115 “listing only” reports (covering 13 species) of infestation of fruits of the plant family Rutaceae by *B. cucurbitae* (Table 3). Five of the six plant species for which field infestations are reported are of *Citrus* spp., while all the three reported laboratory infestations are of *Citrus* spp. Of the “listing only” references, nine are of *Citrus* spp. In *Citrus* spp. field infestation studies, adults of *B. cucurbitae* were recovered from citron, *C. medica* L.; Kaffir lime, *C. hystrix* DC; pummelo, *C. maxima* (Burm.) Merr.; tangerine, *C. reticulata* Blanco; and, sweet orange,



Table 1. Infestation of intact and punctured Washington navel oranges and Clementine tangerines relative to papayas, eggplant, and Anaheim peppers following 24 hours exposure of individual fruits to 50 gravid female *B. cucurbitae* or *B. latifrons*.

FRUIT FLY SPECIES	FRUIT ID	FRUIT STATUS	NO. TRIALS	NO. TRIALS WITH INFESTATION	TOTAL NO. FRUITS	FRUIT WEIGHT (kg)	NO. INFESTED FRUITS (%)	WEIGHT INFESTED FRUITS (kg)	AVG. NO. PUPAE RECOVERED PER kg FRUIT	AVG. NO. ADULTS RECOVERED PER kg FRUIT	AVG. % EMERGENCE
<i>B. cucurbitae</i>	Clementine	Intact	12	8	48	2.78	15 (31.2)	0.84	340.8 ^b	284.7 ^a	74.2
<i>B. cucurbitae</i>	Clementine	Punctured	12	11	48	2.92	28 (58.3)	1.59	240.1 ^{a,b}	168.2 ^a	69.8
<i>B. cucurbitae</i>	Papaya	Control	12	12	12	6.48	12 (100.0)	6.48	310.2 ^{a,b}	256.5 ^a	80.4
<i>B. latifrons</i>	Clementine	Intact	12	1	48	5.69	1 (2.1)	0.16	0.51 ^c	0.51 ^b	100.0
<i>B. latifrons</i>	Clementine	Punctured	12	2	48	5.77	2 (4.2)	0.32	0.39 ^c	0.39 ^b	100.0
<i>B. latifrons</i>	Anaheim pepper	Control	12	12	12	0.82	12 (100.0)	0.82	378.7 ^a	285.2 ^a	69.7
<i>B. cucurbitae</i>	Navel orange	Intact	21	0	84	26.28	0 (0.0)	0.00	0.0 ^c	0.0 ^c	–
<i>B. cucurbitae</i>	Navel orange	Punctured	21	17	84	25.85	44 (52.4)	13.72	84.5 ^b	24.8 ^b	17.4
<i>B. cucurbitae</i>	Papaya	Control	21	21	21	10.59	21 (100.0)	10.59	466.7 ^a	307.3 ^a	69.1
<i>B. latifrons</i>	Navel orange	Intact	16	1	64	17.98	1 (1.6)	0.26	0.36 ^c	0.060 ^c	16.7
<i>B. latifrons</i>	Navel orange	Punctured	16	9	64	18.40	10 (15.6)	2.96	5.51 ^c	2.06 ^c	37.6
<i>B. latifrons</i>	Eggplant	Control	9	9	9	1.44	9 (100.0)	1.44	454.1 ^a	388.7 ^{a,b}	85.8
<i>B. latifrons</i>	Papaya	Control	7	7	7	2.94	7 (100.0)	2.94	333.1 ^a	275.8 ^{a,b}	83.7

Notes: Numbers of pupae or adults recovered per kilogram fruit followed by the same letter are not significantly different at the $\alpha = 0.05$ level (analyses run separately for lementine fruits [with respective controls] and for navel oranges [with respective controls]).

Table 2. Summary of fruits from the plant family Rutaceae, which have been reported to be infested by *Bactrocera latifrons*.

SCIENTIFIC NAME	COMMON NAME	GRIN NO.	INFESTATION RECORD	REFERENCE CITATIONS AND INFESTATION SUMMARIES
<i>Citrus</i> spp.	Citrus species	312282	Listing Only	5,10
<i>Citrus aurantiifolia</i> (Christm.) Swingle	Lime	10683	Field Infestation	From fruit collections in Peninsular Malaysia (1986 to 1988) and in East Malaysia (Sabah and Sarawak) and Thailand (1990 to 1994), <i>B. latifrons</i> was recovered from 1 sample. No infestation rate data given. ¹¹
<i>Citrus limon</i> (L.) Burm. f.	Lemon	10732	Listing Only	5,10,12,13
<i>Citrus sinensis</i> (L.) Osbeck	Sweet orange	10782	Listing Only	5,10,12,13
<i>Murraya paniculata</i> (L.) Jack	Mock orange	24704	Field Infestation	From fruit collections in Peninsular Malaysia (1986 to 1988) and in East Malaysia (Sabah and Sarawak) and Thailand (1990 to 1994), <i>B. latifrons</i> was recovered from 1 sample. No infestation rate data given. ¹¹

Notes: Included, for each plant species, is a reference to the taxonomy of the plant species (as provided by the USDA-ARS Germplasm Repository Information Network [GRIN]), the citation of the references from which the infestation data were recovered, along with an indication as to whether the references were based on field data, laboratory data or were "listing only." In cases where there were laboratory or field infestation data, succinct summaries of the infestation data are also provided.

C. sinensis (L.) Osbeck. Infestation rates of *B. cucurbitae* in these aforementioned *Citrus* spp. were low (Table 3). For the non-citrus rutaceous fruit for which field infestation was reported, ie, limeberry, *Tripkasia trifolia* [Burm. f.] P. Wilson, the infestation rate by *B. cucurbitae* was also low. In laboratory infestation studies, *B. cucurbitae* larvae developed through pupation in sour orange, *Citrus aurantium*, and in tangerine, *C. reticulata*, but failed to develop through to pupation in lemon, *C. limon*.

Discussion

***Bactrocera latifrons*.** There has only been one report of field infestation of *B. latifrons* in *Citrus* spp., and that is from only one sample of *C. aurantiifolia* (lime) collected in Southeast Asia;¹¹ the number of collections conducted during the field host determination survey and the number of fruit samples during each collection were not specified. However, the field host survey conducted in Southeast Asia met the highest reliability of the presence of a pest in an area as defined by the ISPM 8: Determination of Pest Status in an Area,⁶¹ as the survey team included several fruit fly specialists, including the taxonomic expert for the genus *Bactrocera* (R. A. Drew). By logical extension, the field infestation record of *B. latifrons* in lime¹¹ is reliable.

No choice infestation studies reported here showed that laboratory *B. latifrons* adults can oviposit in both intact and punctured navel oranges, *C. sinensis*, but the eggs rarely develop to the adult stage. Successful adult emergence was found with both intact and punctured fruits, but only one adult fly was recovered from intact oranges (0.060 adult flies/kg fruit), while 10 flies were recovered from punctured oranges (2.06 adult flies/kg fruit), compared to an average adult recovery of 275.8 and 388.7 adults/kg fruit in control papaya and control eggplant, respectively. There is no confirmation in

the literature that citrus species such as oranges and tangerines can be natural hosts for *B. latifrons*.⁶² However, it should be noted that it can be difficult to find citrus orchards (where fruit sampling could be done to assess infestation by tephritid fruit flies) where a well-established *B. latifrons* population is present, because *B. latifrons* field populations can be best represented in pastures or recently disturbed fallow lands having wild solanaceous plants.^{5,63} The field recovery of *B. latifrons* from *C. aurantiifolia* in Southeast Asia,¹¹ combined with the observation that laboratory populations of *B. latifrons* in Hawaii can oviposit in both intact and punctured navel oranges and that the eggs can rarely develop successfully to the adult stage (Table 1), suggest that further research, especially research involving wild (field) *B. latifrons* populations interacting with intact fruits in the field, is needed to determine the host suitability of oranges and tangerines grown in Hawaii to *B. latifrons* following the guidelines specified in RSPM 30: Guidelines for the determination and designation of host status of a fruit or vegetable for fruit flies (Diptera: Tephritidae).⁶⁴

***Bactrocera cucurbitae*.** There have been reports of field infestation of *B. cucurbitae* in multiple *Citrus* spp., including *Citrus hystrix*, *C. maxima*, and *C. medica*, as well as in *C. reticulata* and *C. sinensis* (Table 3). In the no choice infestation studies reported here, significantly more *B. cucurbitae* adults than *B. latifrons* adults were recovered from both Clementine tangerines and navel oranges. Both the literature reports and the results of the no choice infestation trials support a conclusion that *Citrus* spp., in general, are better hosts for *B. cucurbitae* than for *B. latifrons*. Comparing the results of no choice infestation trials of Clementine tangerine with those of navel orange by *B. cucurbitae*, it is interesting to note that there was no significant difference in infestation rate for intact versus punctured fruits in the relatively thin-skinned

**Table 3.** Summary of fruits from the plant family Rutaceae, which have been reported to be infested by *Bactrocera cucurbitae*.

SCIENTIFIC NAME	COMMON NAME	GRIN NO.	INFESTATION RECORD	REFERENCE CITATIONS AND INFESTATION SUMMARIES
<i>Aegle marmelos</i> (L.) Corrêa	Bael	1560	Listing Only	14
<i>Casimiroa edulis</i> La Llave & Lex	White sapote	9292	Listing Only	13,15–20
<i>Citrus aurantium</i> L.	Sour orange	10684	Laboratory Infestation	Using 1st instar larvae obtained from eggs oviposited on bottle gourd (<i>Lagenaria vulgaris</i>), 49 of 100 1st instar larvae (49%) raised on orange pupated, with an average time to pupation of 8.9 days. In a separate test, 94 of 100 1st instar larvae (94%) were found to feed on pieces of orange. ²¹
			Listing Only	13,15,17–20
<i>Citrus deliciosa</i> Ten.	Italian tangerine	314340	Listing Only	18
<i>Citrus hystrix</i> DC.	Kaffir lime	10714	Field Infestation	In 1992, <i>B. cucurbitae</i> was recovered from 2 samples of <i>Citrus hystrix</i> [Thailand, Malaysia, Southern India]. Infestation rate data not given. <i>B. cucurbitae</i> individuals identified by R.A.I. Drew and D.L. Hancock. ¹¹
<i>Citrus limon</i> (L.) Burm. f.	Lemon	10732	Laboratory Infestation	Using 1st instar larvae obtained from eggs oviposited on bottle gourd (<i>Lagenaria vulgaris</i>), 49 of 100 1st instar larvae (49%) were found to feed on lemon. No larvae (0 of 100) completed growth to the point of pupation, but survived longer (4–6 days) than 1st instar larvae fed on diets of water alone or 2.5% agar gel. ²¹
			Listing Only	13,15,17–20,22,23
<i>Citrus maxima</i> (Burm.) Merr.	Pummelo	10744	Field Infestation	<i>B. cucurbitae</i> adults were recovered from infested <i>C. maxima</i> fruits, randomly collected on Penang Island, West Malaysia. No infestation rate data given. ²⁴
			Listing Only	4,13,15,19,22,25–29
<i>Citrus medica</i> L.	Citron	10745	Field Infestation	Fallen and marketable sized <i>C. medica</i> fruits were harvested over a seven week period in September–October, 1975 in Hissar, India. Fruits were cut open to check for fruit fly incidence. <i>B. cucurbitae</i> was recovered in 6 of 7 weekly collections (85.7%), with an average weekly infestation rate of 28.0%. Overall, 13 of 52 collected fruits were infested by <i>B. cucurbitae</i> . ³⁰
<i>Citrus myrtifolia</i> Raf.	Myrtle-leaf orange	10756	Listing Only	18
<i>Citrus paradisi</i> Macfad.	Grapefruit	10772	Listing Only	13,15,17–20,22,23
<i>Citrus reticulata</i> Blanco	Tangerine	10778	Field Infestation	One adult <i>B. cucurbitae</i> and 259 adult <i>B. dorsalis</i> were recovered from 10 tangerine fruits collected from the Punahou area of Honolulu (Oahu, Hawaii) in April, 1947. ³¹
			Field Infestation	<i>B. cucurbitae</i> individuals (adults?) were recovered from <i>C. reticulata</i> fruits collected between 2005–2007 in Benin, with infestation rate falling in the range of 1–25 <i>B. cucurbitae</i> per kg fruit. No data presented on the number of fruits collected, the weight of fruits collected or the percentage infestation of collected fruits. ⁶
			Laboratory Infestation	In captivity, female melon flies laid eggs on cut fruits of <i>C. reticulata</i> . The eggs hatched out and the development of the larvae proceeded to continue normally through pupation. No data presented on methods used for the lab infestation or the resulting infestation rate. ³²
			Listing Only	13–15,18,19,22,27,33–36



Table 3. (Continued).

SCIENTIFIC NAME	COMMON NAME	GRIN NO.	INFESTATION RECORD	REFERENCE CITATIONS AND INFESTATION SUMMARIES
<i>Citrus sinensis</i> (L.) Osbeck	Sweet orange	10782	Field Infestation	Adult <i>B. cucurbitae</i> have been reared from oranges, but these fruits do not serve regularly as <i>B. cucurbitae</i> hosts. "Only in rare instances does the melon fly attack them, and then only slightly." No infestation rate data presented. ³⁷
			Field Infestation	In 1910, a few oranges provided by a farmer from Kaimuki (Oahu, Hawaii) were placed in a breeding jar from which mostly <i>Drosophila</i> spp. were recovered, but also one adult melon fly. ³⁸
			Field Infestation	About 10% of orange fruits recovered in the vicinity of the University of Agriculture in Faisalabad, Pakistan, were infested by <i>B. cucurbitae</i> . ³⁹
			Field Infestation	<i>B. cucurbitae</i> individuals (adults?) were recovered from <i>C. sinensis</i> fruits collected between 2005–2007 in Benin and in Burkina Faso, with infestation rate falling in the range of 1–25 <i>B. cucurbitae</i> per kg fruit. No data presented on the number of fruits collected, the weight of fruits collected or the percentage infestation of collected fruits. ⁶
			Listing Only	4,13–15,17–20,22,23,25–28,31–36,40–48
<i>Citrus</i> spp.	Citrus species	312282	Listing Only	13,14,16,17,20–22,31,34,35,42,49–59
<i>Citrus vulgaris</i> Risso		102860	Listing Only	18
<i>Clausena lansium</i> (Lour.) Skeels	Wampi	10811	Listing Only	13,15–20
<i>Triphasia trifolia</i> (Burm. f.) P. Wilson	Limeberry	40476	Field Infestation	13 of 29 samples (44.8%) of <i>T. trifolia</i> fruits made in Rota, Marianas Islands, between 1959–1963, were infested by <i>B. cucurbitae</i> and/or <i>B. dorsalis</i> . A total of seven <i>B. cucurbitae</i> adults were recovered from a total of 13,729 fruits. ⁶⁰
			Listing Only	4,13,18

Notes: Included, for each plant species, is a reference to the taxonomy of the plant species (as provided by the USDA-ARS Germplasm Repository Information Network [GRIN]), the citation of the references from which the infestation data were recovered, along with an indication whether the references were based on field data, laboratory data, or were "listing only." In cases where there were laboratory or field infestation data, succinct summaries of the infestation data are also provided.

Clementine tangerines, while the infestation rate in the relatively thicker skinned navel oranges was significantly higher in punctured fruits than in intact fruits, where no infestation was observed. This suggests that fruit damage may be an important factor in *B. cucurbitae* infestation in navel oranges in the field. Overall, both published infestation data and the results of the no choice infestation trials reported here indicate that movement of both Clementine tangerines and navel oranges from Hawaii to the continental US will require appropriate risk mitigation measures.

Risk mitigation options for tephritid fruit flies in oranges and tangerines. Risk mitigation measures are not needed for *B. latifrons* because no field infestation by *B. latifrons* of citrus fruits like Clementine tangerines and navel oranges has been reported to date, but the laboratory studies reported here indicate that further field studies are

needed, especially studies involving exposure of fruits on trees to gravid wild females. However, because both field and laboratory data show that *B. cucurbitae* can infest citrus fruits like Clementine tangerines and navel oranges, shipment of citrus fruits out from areas where *B. cucurbitae* is present will require risk mitigation measures. At present, there is one postharvest treatment that could currently be used, and several options are available for systems approaches. An irradiation quarantine treatment could be used based on research that established a 150 Gy minimum absorbed dose as a generic treatment dose for postharvest disinfestation of tephritid fruit flies in fruits and vegetables.^{65,66}

Use of high-temperature forced-air is another possible tephritid fruit fly disinfestation treatment. A high-temperature forced-air disinfestation treatment using four temperature stages was developed that successfully disinfested



color-break to half-ripe papaya (*Carica papaya* L.) of *C. capitata* (Wiedemann), *B. dorsalis* (Hendel), and *B. cucurbitae*, whether the fruit flies were introduced as eggs or as third instars.⁶⁷

Alternatively, cold treatment is frequently used to control fruit flies in citrus.⁶⁸ A cold-based disinfestation treatment was shown to be effective for carambola (*Averrhoa carambola* L.) against *C. capitata* (Wiedemann), *B. dorsalis* (Hendel), and *B. cucurbitae* (Coquillett).⁶⁹ However, further research would be needed to demonstrate that the established temperature range would effectively disinfest tephritid fruit flies from citrus fruits. Assessment of whether established cold treatment schedules (for *C. capitata* or *Anastrepha ludens* [Loew]) would be effective for *Bactrocera zonata* (Saunders), the peach fruit fly, was recently tested in order to develop a cold treatment to safely permit shipment of oranges from Egypt (where *B. zonata* is established) to other localities where *B. zonata* is not present. In that research, *B. zonata* was found to be more cold tolerant than *C. capitata* but less cold tolerant than *A. ludens*, so treatment schedules previously developed for *A. ludens* were determined to provide quarantine security for oranges that might be infested by *B. zonata*.⁷⁰

Finally, quarantine security could be sought through a systems approach incorporating a series of risk-reducing steps. Adequate quarantine security might not be achieved by each individual step, but could be achieved when multiple steps are applied sequentially. As an example, a systems approach could incorporate a cold treatment while using an alternative security measure, such as low prevalence,⁷¹ or a pest-free production area.^{72,73} In a recently established systems approach to permit the shipping of Sharwil avocados (a poor host of *B. dorsalis*) from Hawaii to the continental US,^{74,75} a grower compliance agreement requires multiple risk-reducing steps including limiting the dates over which fruits can be picked, protecting picked fruits from exposure to *B. dorsalis* adults, monitoring *B. dorsalis* field population levels, and application of a protein bait spray if *B. dorsalis* trap catch, monitored weekly, exceeds a specified level.

Acknowledgment

We thank Steven A. Brown for technical assistance with this study. We also thank Victoria Yokoyama, Dorothy Alontaga, and two anonymous reviewers who contributed to manuscript improvement through reviews of an earlier draft of this manuscript.

Author Contributions

Conceived and designed the experiments: GTM, PAF, CDS. Analyzed the data: GTM, CDS. Wrote the first draft of the manuscript: GTM, NJL. Contributed to the writing of the manuscript: GTM, PAF, NJL, CDS. Agree with manuscript results and conclusions: GTM, PAF, NJL, CDS. Jointly developed the structure and arguments for the paper: GTM, PAF, NJL. Made critical revisions and approved final version: GTM, PAF, NJL, CDS. All authors reviewed and approved the final manuscript.

REFERENCES

- Purseglove JW. *Tropical Crops. Dicotyledons*. London, England: Longman Scientific and Technical; 1968.
- Martinez-Ferrer MT, Campos JM, Fibla M. Field efficacy of *Ceratitis capitata* (Diptera: Tephritidae) mass trapping technique on clementine groves in Spain. *J Appl Entomol*. 2012;136:181–190.
- Han P, Wang X, Niu C-Y, Dong Y-C, Zhu J-Q, Desneux N. Population dynamics, phenology, and overwintering of *Bactrocera dorsalis* (Diptera: Tephritidae) in Hubei Province, China. *J Pest Sci*. 2011;84:289–295.
- Hollingsworth RG, Vagalo M, Tsatsia F. Biology of melon fly, with special reference to the Solomon Islands. In: Allwood AJ, Drew RAI, eds. *Management of Fruit Flies in the Pacific, A Regional Symposium. ACLAR Proceedings no. 76*. Canberra, Australia: Australian Centre for International Agricultural Research; 1997:140–144.
- Liquido NJ, Harris EJ, Dekker LA. Ecology of *Bactrocera latifrons* (Diptera: Tephritidae) populations: host plants, natural enemies, distribution, and abundance. *Ann Entomol Soc Am*. 1994;87:71–84.
- Vayssières JF, Rey JY, Traore L. Distribution and host plants of *Bactrocera cucurbitae* in West and Central Africa. *Fruits*. 2007;62:391–396.
- Mziray HA, Makundi RH, Mwatawala M, Maerere A, De Meyer M. Host use of *Bactrocera latifrons*, a new invasive tephritid species in Tanzania. *J Econ Entomol*. 2010;103:70–76.
- HDOA (Hawaii Department of Agriculture). *Movement of Citrus Fruits, Citrus spp., From Hawaii into Other Regions of the United States: A Qualitative, Pathway-Initiated Pest Risk Assessment*. Honolulu, HI: Hawaii Department of Agriculture; 2000.
- SAS Institute Inc. *JMP 10.0.0*. Cary, NC: SAS Institute, Inc; 2012.
- Yunus A, Ho HT. *List of Economic Pests, Host Plants, Parasites and Predators in West Malaysia (1920–1978)*, Bulletin No. 153. Kuala Lumpur, Malaysia: Malaysia Ministry of Agriculture; 1980:150–153.
- Allwood AL, Chinajariyawong A, Drew RAI, Allwood AJ. Host plant records for fruit flies (Diptera: Tephritidae) in Southeast Asia. *Raffles B Zool*. 1999;7:1–92.
- Vijaysegaran S. The current situation of fruit flies in Peninsular Malaysia. In: Vijaysegaran S, Ibrahim AG, eds. *Fruit Flies in the Tropics, Proceedings of the 1st International Symposium*. Kuala Lumpur: Malaysian Agricultural Research and Development Institute and Malaysian Plant Protection Society; 1991:125–139.
- White IM, Elson-Harris MM. *Fruit Flies of Economic Significance; Their Identification and Bionomics*. Wallingford, UK: CAB International; 1992.
- Kapoor VC. Indian Tephritidae with their recorded hosts. *Orient Insects*. 1970; 4(2):207–251.
- Holbrook FR. *Fruit Fly Host List*. Letter to Maehler KL, Plant Quarantine Division. Berkeley, CA: USDA-ARS, Entomology Research Division, Hawaii Fruit Fly Investigations, Honolulu, HI; 1967.
- Isnadi S. The distribution of *Dacus* spp. in the Indonesian Archipelagos. In: Vijaysegaran S, Ibrahim AG, eds. *Fruit Flies in the Tropics, Proceedings of the 1st International Symposium*. Kuala Lumpur: Malaysian Agricultural Research and Development Institute and Malaysian Plant Protection Society; 1991:99–107.
- Oakley RG. Part III fruit flies (Tephritidae). *Manual of Foreign Plant Pests*. Washington, D.C.: United States Department of Agriculture, Agricultural Research Administration, Bureau of Entomology and Plant Quarantines; 1950:198–199.
- USDA. *Melon Fly, Dacus cucurbitae*. Beltsville, MD: Biological Assessment Support Staff, National Program Planning Staff, Plant Protection and Quarantine; 1986.
- USDA-APHIS-PPQ. *Action Plan: Melon Fly Dacus cucurbitae* Coquillett. Washington, DC: USDA-APHIS-PPQ; 1984.
- USDA-APHIS-PPQ. *Pests Not Known to Occur in the United States or of Limited Distribution, No.33: Melon Fly*. Hyattsville, MD: USDA-APHIS-PPQ; 1983.
- Rajamannar N. Growth, orientation and feeding behaviour of the larva of melon fly, *Dacus cucurbitae* Coq., on various plants. *Proc Indian Natl Sci Acad B Biol Sci*. 1962;28:133–142.
- Cantrell BK. *National Contingency Plan for Response to an Incursion of Melon Fly Bactrocera cucurbitae* (Coquillett) (Diptera: Tephritidae) in Australia. Gordon, NSW: Horticultural Research & Development Corporation; 1999.
- Margosian ML, Bertone CA, Borchert DM, Takeuchi Y. *Identification of Areas Susceptible to the Establishment of Fifty-Three Bactrocera spp. (Diptera: Tephritidae: Dacinae) in the United States*. Manhattan, KS: USDA-APHIS-PPQ; 2009.
- Tan KH, Lee SL. Species diversity and abundance of *Dacus* (Diptera: Tephritidae) in five ecosystems of Penang, West Malaysia. *B Entomol Res*. 1982;72: 709–716.
- Botha J, Reeves A, Hardie D. *Melon Fruit Fly (Bactrocera cucurbitae) With Reference to Other Fruit Fly Species of Importance to the Cucurbit Industry*, Fact Sheet no. 5/204. Perth: Western Australia Department of Agriculture; 2004.
- CAB International. *Crop Protection Compendium—Report—Bactrocera cucurbitae (Melon Fly)*. Wallingford, UK: CAB International; 2007.
- Dhillon MK, Singh R, Naresh JS, Sharma HC. The melon fruit fly, *Bactrocera cucurbitae*: a review of its biology and management. *J Insect Sci*. 2005;5:1–16.
- Government of Western Australia. *Pest/Host Database*. South Perth, Australia: Department of Agriculture and Food, Government of Western Australia; 2010.
- Singh A, Sardana HR, Chaurasia V, Stonehouse J. *IMFFI Knowledge Review: Publications on Indian Fruit Fly Ecology, Infestation and Management*. Pusa, New Delhi: Project Integrated Management of Fruit Flies in India; 2004.



30. Gupta JN, Verma AN. Screening of different cucurbit crop for the attack of the melon fruit fly, *Dacus cucurbitae* Coquillett (Diptera: Tephritidae). *Haryana Agr Univ J Res*. 1978;7:78–82.
31. McBride OC, Tanada YA. A revised list of host plants of the melon fly in Hawaii. *Proc Hawaii Entomol Soc*. 1949;13:411–421.
32. Chawla SS. Some critical observations on the biology of the melon fly *Dacus cucurbitae* Coquillett (Diptera: Trypetidae). *Res B Panjab Univ*. 1966;17:105–109.
33. CDFA (California Department of Food & Agriculture). *Exotic Fruit Fly Regulatory Response Manual, Section 5, Appendix A: Fruit Fly Host Materials List. Melon Fruit Fly (Bactrocera cucurbitae)*. Sacramento, CA: CDFA; 2001.
34. Narayanan ES, Batra HN. *Fruit Flies and Their Control*. New Delhi: Indian Council of Agricultural Research; 1960:1–67.
35. Syed RA. *Studies on Trypetids and Their Natural Enemies in West Pakistan. V. Dacus (Strumeta) cucurbitae* Coquillett. Vol 14. Rawalpindi, Pakistan: Commonwealth Institute of Biology Control Technical Bulletin; 1971:63–75.
36. USDA-APHIS. Part 301. Domestic Quarantine Notices. Subpart—Melon Fruit Fly. United States Code of Federal Regulations 7 CFR 310.97. Restrictions on interstate movement of regulated articles. 301.97-2. Regulated articles. United States Government Printing Office, Washington, DC; 2008. Available at: <http://www.gpo.gov/fdsys/pkg/CFR-2008-title7-vol5/pdf/CFR-2008-title7-vol5-sec301-97-2.pdf>. Viewed May 28, 2014.
37. Back EA, Pemberton CE. The Melon Fly, Bulletin no. 643, Washington, D.C.: Government Printing Office, United States Department of Agriculture; 1918.
38. Ehrhorn EM. Report of superintendent of entomology. *Hawaii For Agric*. 1910;7:336–338.
39. Inayatullah C, Khan L, Mohsin A-U, Haq M-U. Relationship between fruit infestation and the density of melon fruit fly adults and puparia. *Pak J Zool*. 1993;25:201–202.
40. Back EA, Pemberton CE. The melon fly in Hawaii. *USDA B*. 1917;491:1–64.
41. Heppner JB. *Larvae of fruit flies. V Dacus cucurbitae (Melon fly) Diptera: Tephritidae*. Florida: Florida Department of Agriculture and Consumer Services; 1989. [Entomology Circular No. 315:1–2].
42. Lall BS. Studies on the biology and control of melon fly *Dacus cucurbitae* Coq. *Pesticides*. 1975;9:31–36.
43. NAPPO (North American Plant Protection Organization)—PAS (Phytosanitary Alert System). *Bactrocera cucurbitae* (Coquillett). 2014. Available at: <http://www.pestalert.org/viewArchPestAlert.cfm?rid=56>. Viewed May 1, 2014.
44. Severin HHP, Severin HC, Hartung WI. The ravages, life history, weights of stages, natural enemies and methods of control of melon fly. *Ann Entomol Soc Am*. 1914;7:178–207.
45. USDA-ARS. Insects not known to occur in the United States. Melon fly (*Dacus cucurbitae* (Coq.)). *Coop Econ Insect Rep*. 1959;9(19):367–368.
46. Weems HV. *Major Fruit Flies of the World*. Gainesville, Florida: Florida Department of Agriculture, Division of Plant Industry; 1967. [Leaflet No. 3:1–4].
47. Weems HV. *Melon Fly (Dacus cucurbitae) Coquillett (Diptera: Tephritidae)*. Gainesville, Florida: Florida Department of Agriculture and Consumer Services; 1964:1–2. [Entomology Circular 29].
48. Weems HV, Heppner JB, Fasulo TR. Featured Creatures. Melon fly, *Bactrocera cucurbitae* (Coquillett) (Insecta: Diptera: Tephritidae). Publication no. EENY-199, University of Florida, Gainesville, Florida; 2001. Available at: http://entomology.ifas.ufl.edu/creatures/fruit/tropical/melon_fly.htm. Viewed May 1, 2014. Revised March 2012.
49. Agrawal N, Mathur YK. The fruit fly problem associated with its cultivated crops in India and its control. *Fruit Flies in the Tropics, Proceedings of the 1st International Symposium*. Kuala Lumpur: Malaysian Agricultural Research and Development Institute and Malaysian Plant Protection Society; 1991:140–151.
50. Baloch UK. Pakistan. In: *Citrus Pest Problems and Their Control in the Near East*, FAO Plant Production and Protection Paper no. 135. Rome, Italy: Food and Agriculture Organization of the United Nations; 1996:111–126.
51. Gopalan M. Insecticidal control of the fruit fly *Dacus cucurbitae* Coq. on bitter gourd (*Momordica charantia*). *SI Hort*. 1977;25:43–44.
52. HDOA (Hawaii Department of Agriculture). *Distribution and Host Records of Agricultural Pests and Other Organisms in Hawaii. Bactrocera cucurbitae* (Coquillett). Honolulu, Hawaii, USA: Plant Industry Division, Hawaii Department of Agriculture; 2006.
53. Kapoor VC. Dacines (Diptera: Tephritidae: Dacine) of the Indian subcontinent. In: Vijaysegaran S, Ibrahim AG, eds. *Fruit Flies in the Tropics*, Proceedings of the 1st International Symposium. Kuala Lumpur: Malaysian Agricultural Research and Development Institute and Malaysian Plant Protection Society; 1991:204–209.
54. Kapoor VC. 2.1 *Dacus cucurbitae* Coquillett. In: Robinson AS, Hooper G, eds. *World Crop Pests: Fruit Flies, Their Biology, Natural Enemies and Control*. Vol 3A. Oxford: Elsevier; 1989:60.
55. Kapoor VC, Agarwal ML. Fruit flies and their increasing host plants in India. In: Cavalloro R, ed. *Proceedings of the CEC/IOBC International Symposium*. Rotterdam: A. A. Balkema; 1983:252–257.
56. Lall BS. Vegetable pests. In: Pant NC, ed. *Entomology in India, 1938–1963*. New Delhi: Entomological Society of India; 1964:187–211.
57. Luck RF, Gumpf DJ, Morse JG. A summary of citrus pest problems in the Near East. In: Baloch UK, ed. *Citrus pest problems and their control in the Near East. FAO Plant Production and Protection Paper no. 135*. Rome, Italy: Food and Agriculture Organization of the United Nations; 1996:309–349.
58. Ramadan MM, Messing RH. A survey for potential biocontrol agents of *Bactrocera cucurbitae* (Diptera: Tephritidae) in Thailand. *Proc Hawaii Entomol Soc*. 2003;36:115–122.
59. Vargas RI, Long J, Miller NW, et al. Releases of *Psytalia fletcheri* (Hymenoptera: Braconidae) and sterile flies to suppress melon fly (Diptera: Tephritidae) in Hawaii. *J Econ Entomol*. 2004;97:1531–1539.
60. Nakagawa S, Faria GJ, Urago T. Newly recognized hosts of the oriental fruit fly, melon fly, and Mediterranean fruit fly. *J Econ Entomol*. 1968;61:339–340.
61. FAO. *Determination of Pest Status in an Area. International Standards for Phytosanitary Measures (ISPM) Publication No. 8, Secretariat of the International Plant Protection Convention*. Rome, Italy: Food and Agriculture Organization of the United Nations; 1998. Available at: https://www.ippc.int/sites/default/files/documents/1323945129_ISPM_08_1998_En_2011-11-29_Refor.pdf. Accessed March 24, 2014.
62. Aluja M, Mangan RL. Fruit fly (Diptera: Tephritidae) host status determination: critical conceptual, methodological, and regulatory considerations. *Annu Rev Entomol*. 2008;53:473–502.
63. McQuate GT, Bokonon-Ganta AH, Peck SL. Population biology and prospects for suppression of the solonaceous fruit fly, *Bactrocera latifrons* (Diptera: Tephritidae). *Proc Hawaii Entomol Soc*. 2007;39:111–115.
64. NAPPO. *NAPPO RSPM 30: Guidelines for the determination and designation of host status of a fruit or vegetable for fruit flies (Diptera: Tephritidae)*. Ottawa, Canada: The Secretariat of the North American Plant Protection Organization; 2008.
65. Follett PA. Generic radiation quarantine treatments: the next steps. *J Econ Entomol*. 2009;102:1399–1406.
66. Follett PA, Armstrong JW. Revised irradiation doses to control melon fly, Mediterranean fruit fly, and oriental fruit fly (Diptera: Tephritidae) and a generic dose for tephritid fruit flies. *J Econ Entomol*. 2004;97:1254–1262.
67. Armstrong JW, Hansen JD, Hu BKS, Brown SA. High-temperature forced-air quarantine treatment for papayas infested with tephritid fruit flies (Diptera: Tephritidae). *J Econ Entomol*. 1989;82:1667–1674.
68. Willink E, Gastaminza G, Salvatore A, et al. Quarantine cold treatments for *Ceratitris capitata* and *Anastrepha fraterculus* (Diptera: Tephritidae) for citrus in Argentina: Conclusions after 10 years of research. *Fruit Flies of Economic Importance: From Basic to Applied Knowledge, Proceedings of the 7th International Symposium on Fruit Flies of Economic Importance*. Salvador, Brazil: Sociedade Brasileira para o Progresso da Ciência (SBPC); 2008:285–293.
69. Armstrong JW, Silva ST, Shishido VM. Quarantine cold treatment for Hawaiian carambola fruit infested with Mediterranean fruit fly, melon fly, or Oriental fruit fly (Diptera: Tephritidae) eggs and larvae. *J Econ Entomol*. 1995;88:683–687.
70. Hallman GJ, Myers SW, Taret G, Fontenot EA, Vreyson MJB. Phytosanitary cold treatment for oranges infested with *Bactrocera zonata* (Diptera: Tephritidae). *J Econ Entomol*. 2013;106:2336–2340.
71. FAO. *Requirements for the Establishment of Areas of Low Pest Prevalence. International Standards for Phytosanitary Measures (ISPM) Publication No. 22. Secretariat of the International Plant Protection Convention*. Rome, Italy: Food and Agriculture Organization of the United Nations; 2005. Available at: https://www.ippc.int/sites/default/files/documents/1323946136_ISPM_22_2005_En_2011-11-29_Refor.pdf. Accessed March 24, 2014.
72. FAO. *Requirements for the Establishment of Pest Free Areas. International Standards for Phytosanitary Measures (ISPM) Publication No. 4. Secretariat of the International Plant Protection Convention*. Rome, Italy: Food and Agriculture Organization of the United Nations; 1995. Available at: https://www.ippc.int/sites/default/files/documents/1367570788_ISPM_04_1995_En_2011-12-01_Refor.pdf. Accessed March 24, 2014.
73. FAO. *Requirements for the Establishment of Pest Free Places of Production and Pest Free Production Sites. International Standards for Phytosanitary Measures (ISPM) Publication No. 10. Secretariat of the International Plant Protection Convention*. Rome, Italy: Food and Agriculture Organization of the United Nations; 1999. Available at: https://www.ippc.int/sites/default/files/documents/1323945204_ISPM_10_1999_En_2011-11-29_Refor.pdf. Accessed March 24, 2014.
74. USDA-APHIS. Interstate movement of Sharwil avocados from Hawaii. Final Rule by the Animal and Plant Health Inspection Service. *Fed Regist*. 2013;78(177):56129–56132.
75. Follett PA, Vargas RI. A systems approach to mitigate oriental fruit fly risk in ‘Sharwil’ avocados exported from Hawaii. *Acta Hort*. 2010;880:439–445.