



Mechanism of Cytoplasmic mRNA Translation

Authors: Browning, Karen S., and Bailey-Serres, Julia

Source: The Arabidopsis Book, 2015(13)

Published By: The American Society of Plant Biologists

URL: <https://doi.org/10.1199/tab.0176>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

First published on April 24, 2015: e0176. doi: 10.1199/tab.0176

Mechanism of Cytoplasmic mRNA Translation

Karen S. Browning^{a,c} and Julia Bailey-Serres^{b,c}

^a Department of Molecular Biosciences and Institute for Cell and Molecular Biology, University of Texas at Austin, Austin TX 78712-0165

^b Department of Botany and Plant Sciences and Center for Plant Cell Biology, University of California, Riverside, CA, 92521 USA

^c Both authors contributed equally to this work

Address correspondence to kbrowning@cm.utexas.edu or serres@ucr.edu

Protein synthesis is a fundamental process in gene expression that depends upon the abundance and accessibility of the mRNA transcript as well as the activity of many protein and RNA-protein complexes. Here we focus on the intricate mechanics of mRNA translation in the cytoplasm of higher plants. This chapter includes an inventory of the plant translational apparatus and a detailed review of the translational processes of initiation, elongation, and termination. The majority of mechanistic studies of cytoplasmic translation have been carried out in yeast and mammalian systems. The factors and mechanisms of translation are for the most part conserved across eukaryotes; however, some distinctions are known to exist in plants. A comprehensive understanding of the complex translational apparatus and its regulation in plants is warranted, as the modulation of protein production is critical to development, environmental plasticity and biomass yield in diverse ecosystems and agricultural settings.

INTRODUCTION

Plant growth and function requires highly regulated spatial and temporal regulation of gene expression. The decoding of the mRNA into a polypeptide chain (protein) by the ribosome is a key step in the regulatory continuum from gene to protein to phenotype. The process of translation requires many RNAs, the messenger RNA (mRNA) transcript, transfer RNAs (tRNAs) and the ribosomal RNAs (rRNAs) of the ribosome as well as scores of soluble protein factors that function either as individual proteins or in multi-subunit complexes. The decoding of mRNA into a polymer of amino acids is a very ancient process, such that the machine for this process, the two-subunit ribosome, is conserved across all forms of life on Earth. The three basic phases of the process of translation – initiation, elongation and termination – are also generally conserved. Consequentially, many components of the complex apparatus involved are recognizable across phyla, especially across the plant, fungal and animal kingdoms. Despite this conservation of the basic chemistry and process of protein synthesis, nature has evolved many ways of starting the first phase, known as initiation. The second phase known as elongation, in which additional amino acids are covalently added to the polypeptide, and the third phase known as termination that completes the process are much more preserved across all kingdoms. Eubacteria have three proteins known as initiation factors to unite the mRNA, the initiator tRNA (usually tRNA^{Met}) and the

small subunit of the ribosome and assemble them with the large subunit of the ribosome to commence the elongation process. *Archaea* and eukaryotes have expanded this machinery to include 10 or more proteins or protein complexes, although *Archaea* lack the eIF3 and eIF4 families found in eukaryotes. In addition, a number of “flourishes” have been added to the nuclear-encoded eukaryotic mRNA such as a 5'-m⁷GpppN cap structure at the 5' end and a stretch of adenine residues, the poly(A) tail at the 3' end. These features are added in the nucleus during transcription and are important in transcript stability during the journey from nucleus to cytoplasm and the lifetime of the mRNA. The translational apparatus has evolved to use these added mRNA features to facilitate the process of initiation in the cytoplasm. It is also thought that the role of the extended cohort of initiation factors in eukaryotes is to participate in exquisitely complicated schemes to regulate the process. What could be more important to a cell than the synthesis of the proteins that catalyze the chemistry of metabolism to make the energy for cell growth, division and function? It is therefore not surprising that the plant translational apparatus and its regulation varies from other eukaryotic organisms due to the specialized cellular biochemistry, developmental complexity and environmental plasticity that confers survival and reproduction centered around the capture of light energy and the conversion to chemical energy, i.e., photosynthesis. Another chapter of *The Arabidopsis Book* evaluates the regulation of translation of cytoplasmic mRNAs (Roy and von Arnim, 2013). Organellar mRNA translation (chloroplast and mitochondria) and its coordi-

nation with cytoplasmic translation is beyond the scope of this chapter; therefore, the reader is referred to recent reviews (Gonzalez and Giegé, 2014; Janska and Kwasniak, 2014; Tiller and Bock, 2014). Here, we detail the process of cytoplasmic translation, its machinery and regulation in plant cells as a drama that occurs in several acts.

DELIVERY OF THE SCRIPT: FROM PRE-mRNA TO QUALITY-CHECKED CYTOSOLIC mRNA

Following the selection of the transcript start site and polymerization of approximately the first 20 nucleotides of the pre-mRNA by RNA polymerase II, the 5'-end of the nascent transcript is modified by the addition of a 5'-m⁷GpppN-cap structure. This event augments subsequent steps in pre-mRNA biogenesis including intron removal by the spliceosome (Izaurralde et al., 1994) and the cleavage event that marks the 3'-end and site of poly(A) addition (Cooke and Alwine, 1996; Hunt, 2011). Mechanisms of 5'-cap addition in plants are not well studied, but are thought to resemble those of other eukaryotes. The 5'-cap provides protection for the mRNA until it is removed by the decapping machinery and subsequent degradation occurs in a 5' to 3' manner (Jiao et al., 2008). As will be discussed, the cap structure also plays a definitive role in the selection of an mRNA for translation. At the 3' end of the pre-mRNAs, the process of cleavage and polyadenylation has both highly conserved eukaryotic and plant-specific features (Hunt, 2011). The 3'-poly(A) addition site of an individual gene transcript can vary, with ~25% of *Arabidopsis thaliana* genes displaying multiple 3' cleavage sites (Wu et al., 2011). This heterogeneity in the 3' untranslated region (3'UTR) is likely to impact mRNA stability as well as translation. Also pertinent to translation can be features of the 5'-leader sequence prior to the initiation codon of the protein-encoding open reading frame (ORF), referred to as the 5' untranslated region (5'UTR) or 5' leader. Sequences or secondary structures within the 5'UTR can predispose a transcript to distinct translational regulation (Arribere and Gilbert, 2013), as can the presence of short upstream ORFs (uORFs) (Roy and von Arnim, 2013). High-throughput mRNA sequencing (mRNA-seq) has further expanded appreciation for transcript isoform variants that arise due to selection of the site of transcript initiation and variation in intron selection that are regulated in environmental and developmental contexts (Yamamoto et al., 2009). Of these two, variation in intron removal appears to be more prevalent, but both lead to further diversity and potential regulation of protein expression (Filichkin et al., 2010; Li et al., 2010; Reddy et al., 2013).

The mechanism of constitutive intron splicing of plant pre-mRNAs is generally similar to the pathway detailed in yeast and mammals (Reddy, 2007; Koncz et al., 2012; Reddy et al., 2013). An aspect of this process is the recording of splicing events by binding of an exon junction complex (EJC) 20-30 nt upstream of the site of intron removal. There is modulation of intron removal through regulation of the selection of alternative splice sites and intron retention, affecting upwards of 60% of plant mRNAs during development or due to environmental influences (Filichkin et al., 2010; Wu et al., 2011; Filichkin and Mockler, 2012; Kalyna et al., 2012; Marquez et al., 2012; Syed et al., 2012; Leviatan

et al., 2013; Staiger and Brown, 2013). Alternative splicing and intron retention events have numerous consequences, ranging from the generation of transcript isoforms that encode distinct proteins or are differentially regulated at the level of message stability, transport, localization or translation. When transcription or splicing produces a transcript containing a premature termination codon, typically upstream of an EJC, the mRNA is targeted for nonsense mediated decay (NMD) after the first round of translation (Reddy et al., 2013). Those transcripts that survive the pioneering round of translation are templates for protein synthesis until they are targeted for degradation or sequestered into translationally inactive complexes and removed from the "cast of actors" in the drama that is translation.

ACT 1: INITIATION OF TRANSLATION

The most well studied aspect of translation in eukaryotes is the initiation phase, which is by far considered currently to be the predominant level of regulation. Initiation of translation of a cytosolic mRNA utilizes both the 5'-m⁷GpppN-cap and the 3'-poly(A) tail with initiation factors that specifically recognize these features to start the process of initiation of translation. Baker's yeast (*Saccharomyces cerevisiae*), has provided a genetic treasure trove for structural and functional insight of the highly interactive initiation machinery. Comparative studies have shown that the machinery and their functions are highly conserved, although there are some interesting differences across the spectrum of eukaryotes. In fact, there are remarkable tales of diversity in the machinery that are unique to various organisms and ecological niches (Hernández and Vazquez-Pianzola, 2005; Hernández et al., 2012). Several recent reviews on translation provide mechanistic and structural details of translation derived with *S. cerevisiae* and mammalian systems (Sonenberg and Hinnebusch, 2009; Jackson et al., 2010; Lorsch and Dever, 2010; Hinnebusch, 2011; Aitken and Lorsch, 2012; Dever and Green, 2012; Hernández et al., 2012; Hershey et al., 2012; Hinnebusch and Lorsch, 2012; Valasek, 2012; Voigts-Hoffmann et al., 2012; Lomakin and Steitz, 2013; Hinnebusch, 2014; Mead et al., 2014; Merrick and Harris, 2014). Translation in plants has been reviewed with different emphases in the past five years (Bailey-Serres et al., 2009; Muench et al., 2012; Muñoz and Castellano, 2012; Echevarría-Zomeño et al., 2013; Browning, 2014; Gallie, 2014), and several historical reviews provide the back story (Browning, 1996; Bailey-Serres, 1999; Kawaguchi and Bailey-Serres, 2002; Browning, 2004; Gallie, 2007). As will be described, the translational machinery of plants resembles that of *S. cerevisiae* and mammals. Because plants have unique biological activities, such as photosynthesis and the capacity to respond to stresses *in situ*, they have evolved translational control mechanisms relevant to their needs. This chapter will outline the process of initiation, elongation and termination as largely derived from detailed studies in *S. cerevisiae* and mammals, but will include specific aspects of the plant apparatus where known. Our knowledge of plant translation is based largely on the *Arabidopsis thaliana* accession Col-0 (referred to here as Arabidopsis) and the *in vitro* system derived from the germ (embryo) of hexaploid bread wheat (*Triticum aestivum*). Undoubtedly there will be myriad dif-

ferences within the plant kingdom, not only in the translational apparatus but modulation of protein synthesis as is needed in particular ecological niches and environmental circumstances.

THE ACTORS: THE BASIC MACHINERY OF INITIATION

The current estimate of the number of “basic” initiation factors of eukaryotes is >16 and growing (see Table 1). The unified nomenclature for these proteins/complexes includes five categories for the different aspects of the initiation process (Safer, 1989). These are the eukaryotic initiation factors (eIF) 1 to 6. Table 1 includes several additional proteins that are now being considered part of the translational machinery, but have yet to be evaluated in plants, although in many cases a plant ortholog has been recognized. Initiation factor nomenclature is challenging, especially the names of the initiation factors which have evolved within eukaryotic phyla. eIFs include single or multi-subunit protein complexes that are distinguished by complex number (*i.e.*, eIF2, 3, 4) and Roman letters or Greek letters (*i.e.*, eIF2A or eIF2 α), each of which is a different complex or protein. Some proteins were originally designated eIFs because of their ability to stimulate rabbit reticulocyte *in vitro* translation. For example, the eIF2C family turns out to correspond to the Argonautes that participate in RNA-mediated gene silencing (Chen, 2010). Why the addition of an AGO was found to stimulatory is unknown, but AGO1 copurifies with membrane-associated ribosomes during microRNA (miRNA)-mediated translational inhibition in *Arabidopsis* (Li et al., 2013b).

Table 1 presents the current nomenclature of proteins with known functional activities in mRNA translation in the model plant *Arabidopsis thaliana*. More than one functional gene encodes most of these factors or factor subunits. Therefore, there are multiple isoforms of each factor or factor subunit, which could accumulate in a distinct quantitative, spatial or temporal manner which may have functional consequence.

A schematic of the initiation process is shown in Figure 1 and emphasizes where plant translation is known to differ from that of yeast or mammals. In the next section we introduce and present details about the initiation factors and their interactions with other actors, mRNA, tRNAs and the ribosome, the *prima donna*. The order of the description of members of these acting troupes corresponds to the sequence of their appearance on the stage with the 40S ribosomal subunit. The 40S ribosome/associated factors and mRNA/associated factors are then joined by the 60S ribosome to form the functional 80S ribosome complex for elongation of the polypeptide. The 60S subunit possesses the peptidyl transferase activity to join together amino acids as directed by the codon sequence of the mRNA. After the introduction of the initiation factors, we consider the first act of protein synthesis: the sequence of events that culminates in initiation of polypeptide synthesis. The second and third acts, elongation and termination of protein synthesis will introduce several new performers (*i.e.*, eEFs, eIF6, RACK1, ABCE1, and eRFs). There are also two *encore* events that have garnered attention in recent years that involve efficient recycling of ribosomes for maintained translation of an mRNA or in some cases re-initiation at a downstream open reading frame on an mRNA (von Arnim et al., 2014). There are also side shows

of translation including mRNA turnover (see section on “Curtains for some mRNAs”) and protein degradation (for recent reviews in this series see Callis, 2014; Choi et al., 2014).

eIF2 GROUP AND tRNA^{MET}

This group of factors functions in formation of the ternary complex comprised of Met-tRNA^{Met}•eIF2•GTP and the exchange of GDP for GTP from eIF2•GDP by eIF2B (also called guanine nucleotide exchange factor or GEF). The eight known proteins of this group have challenging names (Table 1). eIF2A (not to be confused with eIF2 α or eIF2B α) and eIF2D (not to be confused with eIF2B δ) are new members of the eIF2 group in animals (Komar et al., 2012). The genes appear to be conserved in plants, but their role in plant initiation is not currently known.

Also to be considered along with this group is the initiator Met-tRNA^{Met}, which has a very specific role in the selection of the correct initiation codon (AUG). This tRNA does not function in elongation and can be distinguished from Met-tRNA^{Met} used for addition of internal methionine residues. Initiator tRNAs have evolved several strategies for this role and avoiding interaction with elongation factor EF1A (reviewed in Koltitz and Lorsch, 2010). Plants and fungi appear to use a strategy of modification of a certain nucleotide in the T-loop with a large O-ribosylphosphate moiety to prevent interaction of initiator Met-tRNA^{Met} with eEF1A.

eIF2 and Ternary Complex Formation

eIF2 is among the most studied of the translation initiation factors. The primary role of this heterotrimeric complex in both eukaryotes and *Archaea* is to bring the Met-tRNA^{Met} and GTP to the ribosome, a task performed by the single polypeptide factor IF2 in eubacteria (reviewed in Schmitt et al., 2010). The eIF2 complex is composed of three subunits, designated eIF2 α , eIF2 β and eIF2 γ . eIF2 α and eIF2 β interact with eIF2 γ which forms the core of the complex and also contains the GTP nucleotide binding site. This factor has structural similarity to other GTP binding factors such as elongation factors EF-Tu (prokaryotic) or eEF1A (eukaryotic) (Schmitt et al., 2010). A zinc-binding domain is present in eIF2 β that is similar to one found in eIF5 and is proposed to play a role during start codon recognition (Nanda et al., 2013). The major binding site of eukaryotic Met-tRNA^{Met} appears to be the eIF2 γ subunit and there is less contribution to binding of Met-tRNA^{Met} by eIF2 α than eIF2 β in eukaryotes; whereas in *Archaea* eIF2 α and eIF2 β comprise the major binding site of Met-tRNA^{Met} (Schmitt et al., 2010). eIF2 and associated proteins have been purified from wheat germ and biochemically studied (Benne et al., 1980; Lax et al., 1982; Osterhout et al., 1983; Seal et al., 1983; Shaikhin et al., 1992; Benkowski et al., 1995a, b). Structural and functional similarity of plant eIF2 to yeast and mammalian eIF2 are expected, although there could be plant specific molecular interactions of the subunits or Met-tRNA^{Met} given that plant eIF2 β lacks the third poly-lysine region in the N-terminal domain found in other eukaryotic eIF2 β subunits (Metz and Browning, 1997).

Table 1. Initiation Factors of Arabidopsis

Factor	Mr^a	Function	Arabidopsis Gene^f
eIF1	12,600	Formation of and scanning by pre-initiation complex; start site selection; controls GTP activating protein activity of eIF5	At4g27130, At5g54760, At5g54940, At1g54290
eIF1A (eIF4C ^b)	17,600	Formation of and scanning by pre-initiation complex; start site selection	At5g35680, At2g04520
eIF2		Forms ternary complex with GTP and Met-tRNA; Binds Met-tRNA, to 40S subunit; GTPase activity in presence of eIF5	
α	42,000	Target for GCN2 kinase in plants	At2g40290, At5g05470
β	38,000		At5g20920, At5g01940, At3g07920
γ	50,000		At1g04170, At4g18330
eIF2A		Unknown in plants; in mammals participates in IRES mediated initiation	At1g73180
eIF2B ^c		Recycles eIF2•GDP to eIF2•GTP; unknown if similar function in plants	
α	42-65,000		At1g53880, At1g72340, At1g53900
β	43,800		At3g07300
γ	49,000		At5g19485
δ	37-73,000		At5g38640, At1g48970, At2g44070
ε	80,000		At3g02270, At2g34970, At4g18300
eIF3	13 subunits	Formation of and scanning by pre-initiation complex; start site selection; binding of mRNA to PIC; prevention of pre-mature 60S ribosome association	
a	114,000		At4g11420
b	85,000		At5g27640, At5g25780
c	103,000		At3g56150, At3g22860
d	67,000		At4g20980, At5g44320
e ^d	52,000		At3g57290
f	32,000		At2g39990
g	36,000		At3g11400, At5g06000
h	38,000	Required for efficient re-initiation of main open reading frame of mRNAs with upstream open reading frames	At1g10840
i ^e	36,000		At2g46280, At2g46290
j	25,000		At1g66070, At5g37475
k	26,000		At4g33250
l	60,200		At5g25754, At5g25757
m	50,000		At3g02200, At5g15610
eIF4A	47,000	ATP-dependent unwinding of mRNA Binds mRNA to 40S subunit	At3g13920, At1g54270
eIF4B	58,000	ATP-dependent unwinding of mRNA Binds mRNA to 40S subunit	At3g26400, At1g13020

Phosphorylation of eIF2

Mammals possess four eIF2 kinases: (HRI, heme-regulated inhibitor; PKR, double stranded RNA-dependent kinase; PERK, PKR-like ER kinase; GCN2, general control non-derepressible-2 kinase). All phosphorylate a conserved serine residue (Ser51) in mammalian eIF2 α that inhibits initiation of translation in response to various stresses. eIF2B (see below) cannot easily dissociate from phosphorylated eIF2 α and therefore guanine nucleotide exchange is inhibited depleting available eIF2 for ternary complex formation (reviewed in Donnelly et al., 2013). Despite early reports of a “PKR-like” activity in virus-infected plants (Hiddinga et al., 1988; Langland et al., 1995; Langland et al., 1996; Chang et al., 1999), no specific kinase could be purified and the sequence of a putative PKR ortholog is absent from plant genomes (Immanuel et al., 2012). GCN2 is therefore the only recognizable plant eIF2 α kinase at this time and targets a similar serine residue in plant eIF2 α (Halford et al., 2004; Byrne et al., 2012; Li et al., 2013a; Wang et al., 2014). Other eIF2 α kinases may exist,

but have yet to be described. GCN2 was identified in yeast in response to nutrient deprivation, particularly amino acid or purine starvation (Hinnebusch, 2005), but it is induced by other stresses (e.g., UV, osmotic and oxidative stress) and functions similarly in mammals (Donnelly et al., 2013). The general amino acid control pathway in yeast is controlled by the transcription factor GCN4 which activates transcription of numerous genes in many biosynthetic pathways in response to nutrient deprivation (Hinnebusch, 2005). The translation of yeast *GCN4* mRNA, utilizes four short upstream open reading frames (uORFs) in the 5' leader sequence to control expression of the ORF encoding GCN4.

GCN2 kinase, which is activated during nutrient deprivation by sensing tRNAs that are unchanged (i.e., low amino acid levels), phosphorylates eIF2 α , preventing its interaction with eIF2B for guanine nucleotide recycling. The amount of available ternary complex falls and protein synthesis initiation is inhibited. When the levels of ternary complex are high, initiation occurs at the first AUG in the 5' leader and elongation and termination precede; subsequent reinitiation events are likely at uORFs 2-4 and therefore initiation at the

Table 1. (continued)

Factor	Mr ^a	Function	Arabidopsis Gene ^f
eIF4F		Complex of eIF4G and eIF4E; ATP-dependent unwinding of mRNA; Binds mRNA to 40S subunit	
eIF4G	188,000	Interaction with eIF4A, eIF4B, eIF5, eIF4E	At3g60240
eIF4E	26,000	Binds to eIF4G and m ⁷ G cap on mRNA	At4g18040, At1g29590, At1g29550
eIFiso4F		Complex of eIFiso4G and eIFiso4E; Plant specific isoform of eIF4F; ATP-dependent unwinding of mRNA; Binds mRNA to 40S subunit	
eIFiso4G	84,000	Interaction with eIF4A, eIF4B, eIF5, eIF4E	At5g57870, At2g24050
eIFiso4E	22,500	Binds to eIFiso4G and m ⁷ G cap on mRNA	At5g35620
eIF5	48,600	Joining of 60S subunit; GTPase activating protein	At1g77840, At1g36730
eIF5B	121-142,000	Positions Met-tRNA _i at AUG with eIF1A	At1g76810, At1g21160
eIF5C	47,000	eIF5 “mimic protein” also called 5MP1 or BZW2; regulates eIF2 function by being both a mimic and competitor for eIF5; role unclear in plants	At5g36230, At1g65220
eIF6	26,000	Prevents association of 60S and 40S subunits	At3g55620, At2g39820
PABP	~60-74,000	Binds poly A on mRNA; interacts with eIF4G	At2g23350, At4g34110, At1g22760, At1g71770, At3g16380, At1g49760, At2g36660, At1g34140, At5g65250, At5g65250
4E2 (nCBP ^b , 4EHP)	25,700	Unclear in plants	At5g18110

^a Approximate molecular weight based on TAIR9 data.

^b Prior nomenclature used in literature.

^c Hypothetical genes based on similarity to mammalian eIF2B subunits; protein complex has not been isolated from plant source and shown to function as GDP exchange factor.

^d Also known as INT6

^e Also known as TRIP1 (TGF-beta receptor interacting protein)

^f Links to various data bases using the Arabidopsis Gene Identifier can be found at <http://browning.cm.utexas.edu/arabidopsis/fiat>

AUG of the GCN4 coding region is limited. However, when ternary complex is low, reinitiation at uORFs 2-4 is less likely and the 40S ribosomes continue to scan, acquire ternary complex, reach the AUG of the GCN4 ORF and commence synthesis of GCN4 (Hinnebusch, 2005). The process also involves the transient maintenance of eIF3 association with the ribosome as it translates the first of the four

uORFs under starvation conditions (Szamecz et al., 2008). This is an exquisitely complex regulatory system in yeast for sensing and response to nutrient status through translational control.

Arabidopsis GCN2 kinase complements yeast GCN2 kinase suggesting some aspects of the yeast general amino acid control (GAAC) mechanisms may be conserved in plants (Zhang et




Figure 1. Overview of the steps of translation initiation in the cytoplasm of plants.

Once the mRNA has been exported from the nucleus into the cytoplasm it interacts with a cap-binding complex (eIF4F or plant-specific eIFiso4F) at the 5' end and PABP at the 3' end. Additional factors, eIF4A and eIF4B are recruited to the mRNA to promote ATP-dependent unwinding of secondary structure prior to interaction with the 43S PIC (pre-initiation complex). The 43S PIC is formed from the 40S ribosome and its associated factors eIF1, eIF1A, eIF3 and eIF5. eIF1, eIF1A, eIF3 and eIF5 form the multi-factor complex (MFC), although it is not clear if this assembles prior to interaction with the ribosome or on the ribosome. Addition of ternary complex (TC) of eIF2•Met-tRNA_i•GTP completes the preparation of the PIC. This then engages the mRNA and its associated factors (eIF4F/eIFiso4F, eIF4A, eIF4B, PABP) to form the 48S scanning complex (open conformation), which functions to scan the 5' untranslated region of the mRNA in the 5' to 3' direction in order to select the AUG (i.e., A/GXXA_nUGG). Once the AUG is selected the 48S complex switches to the closed conformation securing the Met-tRNA_i in the P-site and ejecting eIF1. eIF5B•GTP binds to the 48S ribosome complex and the process for joining with the 60S subunit commences, and eIFs 6, 5, 3, and 2 exit the 48S ribosome. Completion of joining of the 60S ribosome results in the hydrolysis of eIF5B•GTP and its release along with eIF1A. The now functional 80S ribosome may now start peptide elongation (see Figure 2). The role of eIF2B in guanine nucleotide recycling of eIF2 in plants is unclear at this time (shown with a green line). Note that the factors and ribosomal subunits are not to scale.

al., 2003). Herbicides (*i.e.*, chlorosulfuron, glyphosate) that inhibit amino acid synthesis and thus induce amino acid starvation result in induction of GCN2 and phosphorylation of eIF2 α (Zhang et al., 2008b). GCN2 also functions in response to purine starvation, UV irradiation, wounding, hormones, cold shock (Lageix et al., 2008), cadmium stress (Sormani et al., 2011b), amino acid metabolism and sulfur signaling (Byrne et al., 2012), but evidently not virus infection (Zhang et al., 2008b). The phosphorylation of eIF2 α in response to purine starvation was correlated with reduced large polysome complexes, suggesting that it can generally inhibit initiation (Lageix et al., 2008; Sormani et al., 2011b). To date, there is no direct evidence that GCN2 regulates ternary complex formation in plants or plays a role in translation of mRNAs with uORFs.

eIF2 α and eIF2 β subunits were reported to be targets of CK2 (formerly casein kinase II) *in vitro*, but the role of phosphorylation of these subunits *in vivo* is not known (Dennis and Browning, 2009; Dennis et al., 2009). Interestingly, none of the subunits of eIF2 were reported to be phosphorylated in a study of the effects of light/dark on the phosphoproteome of the translational apparatus (Boex-Fontvieille et al., 2013). There is still much to learn about the control of protein synthesis in plants in response to various types of stresses and to what level eIF2 subunit phosphorylation regulates the process.

In plants, “eIF2B or, not 2B, that is the question”

As described above, one of the major mechanisms used by yeast and particularly mammals for the regulation of initiation of translation is the phosphorylation of a single conserved serine residue on eIF2 α . This phosphorylation event prevents eIF2B from dissociating from eIF2 during recycling of GDP for GTP, prohibiting the formation of a new ternary complex. This inability to recycle GDP/GTP effectively shuts down initiation in the absence of ternary complex formation (reviewed in Donnelly et al., 2013).

An eIF2B-like activity has not been purified from a plant source, although genes with similarity to mammalian eIF2B subunits are encoded by Arabidopsis (see Table 1) and phosphopeptides have been reported for the eIF2B γ and eIF2B δ subunits (Boex-Fontvieille et al., 2013), suggesting that the proteins are expressed and modified.

Evidence that eIF2B recycling may not be necessary, and thus phosphorylation of eIF2 α may not have as strong an inhibitory effect on translation, is the report that binding of GDP to eIF2 is only about 10-fold higher than that of GTP in comparison to the ~100-fold difference for mammalian eIF2 (Shaikhin et al., 1992). Thus the requirement for eIF2B recycling may be less in plants allowing for continued translation even in the presence of eIF2 α phosphorylation. There is a need for further studies to corroborate biochemically what is known about plant GCN2 kinase, its role in eIF2 α phosphorylation and eIF2B (*i.e.* if it exists as a complex) activity in GDP recycling and global translational activity. It will also be intriguing to decipher the cross-talk that occurs between pathways that activate GCN2 and the Target of Rapamycin (TOR) pathway, which is likely involved in sensing sucrose and other nutrients in plants and regulating translation of certain mRNAs (Immanuel et al., 2012; Robaglia et al., 2012).

eIF3 Group

The sole performer of this group, eIF3, is a complex of six subunits (a, b, c, l, g, j) in yeast, but 13 subunits (a-m) in higher eukaryotes including plants (Browning et al., 2001). The principal role of eIF3 is to bind to the 40S subunit to participate in the formation of the 43S pre-initiation complex (PIC) comprised of the 40S subunit, and the ternary complex (eIF2•GTP•tRNA^{Met}), along with the factors eIF1, eIF5 and eIF1A. eIF3 acts as a bridge to facilitate binding of mRNA with its associated factors, eIF4F, eIF4A, eIF4B and Poly(A) binding protein (PABP), with the PIC, forming a 48S scanning complex. Recent cryo-EM and structural data for eIF3 suggest that it “hugs” the 40S ribosome with contacts that span the mRNA entry and exit sites (Hashem et al., 2013; Liu et al., 2014).

eIF3

The eIF3 complex shares “architectural” similarities to other large complexes, the 26S proteasome lid and the COP9/signalosome that are collectively known as PCI complexes (Pick et al., 2009). The eight proteins that form the octamer core of each of these complexes share motifs known as PCI and MPN domains. The characteristic composition of these PCI complexes are six subunits with PCI domains and two with MPN domains (Pick et al., 2009). Higher eukaryotic eIF3 subunits a, c, e, and l all have PCI domains, k and m have structurally related winged helix domains (Zhou et al., 2005), and f and h have MPN domains. The PCI/MPN containing subunits form the “octamer” core that is similar to that found in the proteasome lid (Querol-Audi et al., 2013). The remaining subunits, b, d and g have RNA Recognition Motif (RRM) domains (Cuchalova et al., 2010); subunits b and i have WD40 domains and eIF3g has a zinc-binding domain (Hinnebusch and Lorsch, 2012; Valasek, 2012; Voigts-Hoffmann et al., 2012; Hashem et al., 2013; Querol-Audi et al., 2013). Recent cryo-EM reconstructions of mammalian eIF3 and the 43S PIC indicate that eIF3 has five lobes and the PCI/MPN octamer forms the functional core of eIF3 (Siridechadilok et al., 2005; Khoshnevis et al., 2012; Hashem et al., 2013; Querol-Audi et al., 2013). The placement of the eIF3 subunits and their contacts with the ribosome and other initiation factors awaits further structural data, but a picture is beginning to emerge at the molecular level (Wilson and Doudna Cate, 2012; Hashem et al., 2013; Liu et al., 2014). Yeast eIF3 is also implicated in termination of protein synthesis, termination codon read-through and ribosome reinitiation suggesting that we still have much to learn about this multi-functional factor and its roles during translation in all organisms (Pisarev et al., 2007; Beznosková et al., 2013).

Plant eIF3 (wheat and Arabidopsis) has been purified and its biochemical analysis suggests strong similarity both in number of subunits and sequence to mammalian eIF3 (Checkley et al., 1981; Lauer et al., 1985; Heuffer et al., 1988; Burks et al., 2001). Subunits eIF3m and eIF3l were first described in plant eIF3 (Burks et al., 2001) and subsequently identified in mammalian eIF3. Biochemical and yeast-two hybrid analysis have implicated some of the Arabidopsis eIF3 subunits in association with the 26S proteasome and COP9 signalosome complexes or subunits (Karniol

et al., 1998; Yahalom et al., 2001; Kim et al., 2004; Paz-Aviram et al., 2008). Both the 26S proteasome and COP9 signalosome play roles in protein turnover. These interactions with eIF3 or its subunits suggest that there may be additional unknown functions for some of the eIF3 subunits or some sort of communication between these large PCI complexes to coordinate various aspects of the cellular dramas of translation and protein degradation (Kim et al., 2001; von Arnim and Chamovitz, 2003).

eIF3 plays a pivotal role in initiation of translation via its interactions with numerous factors as well as the ribosome. It is therefore not surprising that only few Arabidopsis eIF3 subunit mutants have been reported. Five of the subunits for eIF3 (see Table 1) are encoded by a single gene in Arabidopsis (a, e, f, h, k) and eight are encoded by two genes (b, c, d, g, i, j, l, m). Mutations in the single genes for eIF3e or eIF3f cause male gametophytic lethality (Yahalom et al., 2008; Xia et al., 2010). Genotypes that only express *eIF3f* in pollen indicated that the absence of eIF3f is also embryo lethal (Xia et al., 2010). Partial loss of function alleles of the single *eIF3h* gene are viable but display multiple developmental defects, including reduced male gamete transmission. Interestingly, eIF3k appears to be non-essential under normal growth conditions (Tiruneh et al., 2013).

eIF3 and initiation/reinitiation

Several eIF3 subunits are implicated in translation of mRNAs with unusual 5' leader sequences. These include viral mRNAs and plant transcripts with that have one or more uORF upstream or overlapping with the main protein coding ORF (mORF) (See also, Reinitiation involving uORFs section).

Early insight into the nuanced roles of Arabidopsis eIF3 subunits came from the discovery of its function in the initiation of the 35S cauliflower mosaic virus (CaMV) mRNAs during infection. The 5' leader of this viral mRNA is long, highly structured and contains several short ORFs. A viral encoded protein called TAV (transactivation/viroplasm) interacts with eIF3 through eIF3g to retain eIF3 on ribosomes and promote reinitiation at the initiation codon of the first long viral ORF (Park et al., 2001; Park et al., 2004). Another initiation factor, eIF4B is also involved in the TAV-mediated reinitiation process (see below). A host protein called RISP (reinitiation supporting protein) was described that supports reinitiation during CaMV infection and interacts with eIF3 through the eIF3a and eIF3c subunits (Thiebauld et al., 2009). TAV binding to the TOR kinase was shown to be critical for the reinitiation event (Schepetilnikov et al., 2011). This suggests that eIF3 can be essential in reinitiation, in a manner exploited by CaMV and possibly other viruses and connects translation to the TOR signaling pathway in plants.

eIF3 is also important in translation of endogenous plant mRNAs that possess a uORF. Remarkably, over 30% of the protein coding mRNAs of Arabidopsis possess a 5' leader with one or more uORF. Of these, ~1% encode a peptide that is evolutionarily conserved among angiosperms (CPuORFs) (Jorgensen and Dorantes-Acosta, 2012). The presence of a uORF generally reduces the level of translation of the mORF, due to efficient initiation of translation of the uORF and limited initiation at the mORF. DNA microarray analysis of mRNA present on polysomes isolated from *eIF3h* mutants suggested that eIF3h is necessary for reini-

tiation on mRNAs with uORFs (Kim et al., 2004; Kim et al., 2007; Roy et al., 2010; Zhou et al., 2010a; Zhou et al., 2014a). Numerous studies document uORFs that regulate mORF translation in plants, including several that exert their regulation based on metabolite availability (Roy and von Arnim, 2013). uORF-containing mRNAs of Arabidopsis include those encoding the S class of bZIP transcription factors and Auxin Response Factors (ARFs) (Kim et al., 2004; Nishimura et al., 2005; Rahmani et al., 2009). Both eIF3h and the 60S ribosomal protein RPL24 are required for efficient reinitiation of translation of the *AtbZIP11* (*ATB2*) and *ARF* mORF (Kim et al., 2004; Nishimura et al., 2005; Kim et al., 2007; Roy et al., 2010; Zhou et al., 2010a; Zhou et al., 2014a), although global analyses of polysomal RNA do not strongly support a role of RPL24 in this process (Tiruneh et al., 2013).

The reinitiation downstream of the uORFs of *ARF3* mRNA is mediated by auxin as well as TOR kinase (Schepetilnikov et al., 2013). Auxin triggers TOR activation, followed by its association with polysomes, where it phosphorylates ribosomal protein S6 kinase (i.e., AtS6K1) rendering it active to phosphorylate eIF3h, evidently after dissociation from the ribosome. This discovery makes a direct link to auxin-mediated signaling through plant TOR/S6K1 and the translational apparatus needed to reinitiate translation of ARF mRNAs possessing uORFs and provides unequivocal evidence that TOR contributes to gene-specific translational control in plants (Schepetilnikov et al., 2013).

Functional characteristics of other eIF3 subunits are emerging. The overexpression of *eIF3g* in wheat appears to enhance tolerance to drought and other abiotic stresses (Singh et al., 2007; Singh et al., 2013). Interestingly, the only monoclonal antibody to wheat eIF3 subunits that showed any inhibitory activity *in vitro* was to eIF3g and it inhibited mRNA binding to 40S ribosomes *in vitro* (Lauer et al., 1985; Heufler et al., 1988). This observation suggests that eIF3g facilitates mRNA binding to 40S ribosomes. Further, eIF3g was shown to be involved in reinitiation events required to translate GCN4 mRNA in yeast (Cuchalova et al., 2010) and in the reinitiation of CaMV in plants (Park et al., 2001; Park et al., 2004). It can be speculated that eIF3g has a role(s) in reinitiation events through direct interaction with the mRNA and ribosome, but more work will be needed to further establish the function(s) of the eIF3g and other eIF3 subunits during initiation and reinitiation events.

Many observations suggest that there is "cross-talk" between the PCI/MPN complexes in the ribosome-mediated synthesis and proteasome-mediated degradation of proteins. Since these complexes share structural similarity in many of their subunits (Pick et al., 2009), it will be illuminating to figure out the structural and regulatory role of these subunits/complexes and their interactions, from their opening acts in synthesis to the "death scene" in degradation of proteins.

Regulation of eIF3 through Phosphorylation

Several subunits of eIF3 are reportedly phosphorylated by highly specific kinases. As mentioned, this includes TOR-regulated phosphorylation of eIF3h by AtS6K1 (Schepetilnikov et al., 2013). eIF3i was identified as a target of brassinosteroid insensitive receptor kinase (BRI1) and was shown to co-immunoprecipitate with BRI1 (Jiang and Clouse, 2001; Ehsan et al., 2005). This suggests a con-

nection between brassinosteroid signaling and eIF3 function, although the effect of the phosphorylation of eIF3i on eIF3 function is not yet known. The brassinosteroid signaling pathway has been proposed to have similarities to TGF- β signaling in mammals. The TGF- β kinase targets the eIF3i subunit, suggesting that there may be conserved signaling pathways and regulation between plants and animals, albeit repurposed in the individual phyla.

Additional pleiotropic kinases such as CK2 (Mulekar and Huq, 2013) have been shown to be active against several plant initiation factors (eIF2 α , eIF2 β , eIF5) *in vitro*, including multiple phosphorylation sites in eIF3c (Dennis and Browning, 2009). An *in vivo* phosphoproteome study of the light/dark response identified multiple subunits of AtelF3 (b, c, d) as phosphorylation targets (Boex-Fontvieille et al., 2013), including eIF3c sites that are comparable to those identified as CK2 substrates in wheat (Dennis and Browning, 2009).

Clearly there is complicated regulation of translation through multiple eIF3 subunit phosphorylation events and it will be necessary to identify the various kinases and their roles in eIF3 phosphorylation regulation at the molecular level in plants and how it compares with regulation in other organisms.

eIF4 Group

This group of factors interacts with the mRNA and facilitates its binding to the 43S PIC (Jackson et al., 2010; Valasek, 2012; Hinnebusch, 2014). Within this group are the cap-binding complexes, including eIF4F (all eukaryotes) and eIFiso4F (plant-specific). The individual subunits of these complexes are designated as the cap-binding proteins, eIF4E or eIFiso4E, and the large scaffolding proteins, eIF4G or eIFiso4G. Other members of this group are eIF4A, a DEAD box RNA helicase and eIF4B, a RNA binding protein. Both eIF4A and eIF4B are single polypeptides and interact with the large subunits of eIF4F and eIFiso4F. Another member of this group is eIF4H in mammals; however, a comparable factor has not been identified in yeast or plants. Although PABP is not an official member of this group, it will be considered here as it binds to the poly(A) tail of mRNA and interacts with other eIF4 group members during binding of the mRNA to the ribosome. Additional RNA helicases have been identified as having roles in initiation such as yeast DED1 or mammalian DHX29. These have not yet been formally designated as “eIFs” but eventually may be added to the “cast of characters” (Jackson et al., 2010). Plants have comparable RNA helicases (Bush et al., 2009), but their role in plant translation has not been elucidated and they may also have other specific roles in post-transcriptional processes. For example, the nuclear-localized AtelF4AIII was shown to function in nuclear pre-mRNA/mRNA movement during hypoxia (Koroleva et al., 2009a; Koroleva et al., 2009b) and the DEAD box helicase AtRH57 appears to be involved in rRNA processing in response to glucose and abscisic acid (Hsu et al., 2013).

eIF4A

This was the first RNA helicase to be identified and has been called “the godfather of helicases” (Rogers et al., 2002). A number of reviews on eIF4A and the DEAD/DEAH family of helicases

summarize its role in the initiation of translation in mammalian and yeast systems (Webster et al., 1991; Parsyan et al., 2011; Andreou and Klostermeier, 2013; Linder and Fuller-Pace, 2013; Marintchev, 2013; Putnam and Jankowsky, 2013). Despite being the founding member of the DEAD box helicases, so named for a conserved amino acid motif (DEAD), eIF4A is the “outlier” in the family. It possesses a minimal helicase core but lacks additional accessory domains found in other helicases (Andreou and Klostermeier, 2013, 2014). eIF4As are highly conserved proteins, and based on sequence similarity plant eIF4A is likely to share structural and mechanistic details with eIF4A from yeast or mammals.

eIF4A is a non-processive bi-directional RNA dependent ATPase that functions locally to unwind short duplexes and lacks any specificity for RNA sequence (Marintchev, 2013). It has two RecA domains that in the presence of RNA and ATP come together to form the “closed” catalytically active conformation in a dumbbell-like shape (Meng et al., 2014). eIF4G and eIF4B binding to eIF4A favors the closed conformation and likely stimulates P_i release and/or nucleotide exchange from eIF4A (see eIF4G below, reviewed in Marintchev, 2013). It is thought that eIF4A complexed with eIF4G and eIF4B interacts with the 5' end of the mRNA to relax secondary structure in an ATP-dependent manner in preparation for the binding of the 43S PIC. eIF4A further functions to remove secondary structures and/or RNA binding proteins during the scanning of the 5' leader by the PIC (Parsyan et al., 2011; Marintchev, 2013; Andreou and Klostermeier, 2014).

Biochemical studies of wheat eIF4A suggest that it is similar to mammalian and yeast eIF4A (Lax et al., 1986; Abramson et al., 1988; Balasta et al., 1993; Bi et al., 2000). Studies of Arabidopsis eIF4A during the cell cycle led to the suggestion that proliferating cells display high canonical eIF4A association with the cap binding complex whereas quiescent cells may have other types of RNA helicases associated with the cap-binding complexes (Bush et al., 2009). Overexpression of pea (*Pisum sativum*) *eIF4A* (PDH45) resulted in increased resistance to salt stress in rice (*Oryza sativa*) and tobacco (*Nicotiana tabacum*), suggesting a role in stress responses (Tajrishahi et al., 2011; Sahoo et al., 2012). A T-DNA insertion in one of the two *eIF4A* gene paralogs of *Brachypodium distachyon* resulted in a slow-growing, dwarf phenotype that could be partly reversed by heterologous expression of Arabidopsis *eIF4A-1* (Vain, et al, 2011). This phenotype is similar to a T-DNA insertion mutant in Arabidopsis *eIF4A-1* (Vain et al., 2011), one of the three genes encoding this protein (Table 1).

Phosphorylation of eIF4A

There is proteomic evidence that cytoplasmic eIF4A1/2 of Arabidopsis associates with the cyclin dependent kinase CDKA; however, an effect on eIF4A function by CDKA has not been demonstrated (Hutchins et al., 2004). eIF4A is rapidly phosphorylated in response to hypoxia in maize (*Zea mays*), but the relevant kinase or sites were not identified (Webster et al., 1991). Wheat eIF4A was also shown to be phosphorylated at an apparent single site in response to heat shock (Gallie et al., 1997). The recent phosphoproteomic analysis of the light/dark transition shows that *At-eIF4A1*, 2 and 3 gene products are phosphorylated (Boex-Fontvieille et al., 2013).

eIF4B

This is the only initiation factor that lacks a high degree of sequence similarity between yeast, mammals and plants. It is largely accepted that eIF4B functions as a RNA binding protein and enhances the helicase activity of eIF4A, presumptively by augmenting both ATP and RNA binding (Hinnebusch and Lorsch, 2012). In yeast, eIF4B binds eIF4G and induces a conformational change that in turn promotes the binding of eIF4A and increases its RNA helicase activity (Park et al., 2012). Yeast eIF4B has also been shown to bind to the 40S ribosomal protein RPS20 near the mRNA entry site, which may facilitate interaction of eIF4A (Walker et al., 2012; Zhou et al., 2014b) and recent data suggest a mechanism in yeast for eIF4B to promote association of eIF4A with eIF4F (Park et al., 2012). eIF4B is not an essential gene in yeast, but its deletion produces a cold sensitive phenotype (Altmann et al., 1993).

Extensive biochemical and kinetic characterization of the interactions of wheat eIF4B, eIF4A, eIF4G/eIFiso4G and PABP confirm that the interactions are similar to other organisms (reviewed in Gallie, 2014; Le et al., 1997; Bi and Goss, 2000; Bi et al., 2000; Khan and Goss, 2005; Cheng and Gallie, 2006, 2007; Cheng et al., 2008; Khan et al., 2008; Khan et al., 2009; Mayberry et al., 2009; Cheng and Gallie, 2010; Yumak et al., 2010; Khan and Goss, 2012; Cheng and Gallie, 2013). Wheat eIF4B has two tandem domains for interaction with eIF4A and PABP separated by a RNA binding domain in addition to binding domains for eIF4G/iso4G and eIF3g. These tandem binding domains are the most highly conserved parts of plant eIF4B. This suggests that eIF4B may interact with more than one molecule of eIF4A or PABP at a time during initiation events. Further, the interactions with eIF4G/iso4G and eIF4A appear to be specific to plant eIF4B, suggesting divergent evolution of this factor from other eukaryotes (Gallie, 2014).

eIF4B also assists eIF3 in CaMV infection and the TAV-mediated reinitiation on the 35S transcript. Specifically, eIF4B interacts with eIF3g to form a stable 43S PIC in plant cells. Upon binding of the 60S ribosomal subunit at the final step of initiation, eIF4B and eIF3 are released and therefore not found in polysomes of cells not infected by CaMV. However, TAV keeps eIF3 associated with the ribosome, as eIF3 is found in polysomes of infected cells (Park et al., 2004). The overexpression of eIF4B prevents association of TAV with translating ribosomes by competing with TAV for binding to eIF3g (Park et al., 2004). These exploitations of the translation system by a virus suggest that the role of eIF4B may not be solely in stimulation of the helicase activity of eIF4A and eIF4F on mRNA.

Arabidopsis has two forms of eIF4B (eIF4B1 and eIF4B2), which similarly support *in vitro* translation (Mayberry et al., 2009). Interestingly, a heterozygous Arabidopsis T-DNA activation tagging line that showed necrotic lesions symptomatic of programmed cell death overexpresses *eIF4B2* and lines homozygous for a disruption of this gene were embryo lethal (Gaussand et al., 2011). Ectopic overexpression of *eIF4B2* recapitulated the necrotic phenotype, leading to the conclusion that too much eIF4B can cause disruptions in gene expression that trigger programmed cell death. These data suggest a fundamental role for eIF4B in the process of initiation in plants (Gaussand et al., 2011).

Phosphorylation of eIF4B

Plant eIF4B is a target of phosphorylation by CK2 and possibly additional kinases (Gallie et al., 1997; Dennis and Browning, 2009). Arabidopsis eIF4B isoforms show differential responses to the isoforms of CK2 (Dennis and Browning, 2009) suggesting that eIF4B activity could potentially be modulated by distinct CK2s. Support for eIF4B phosphorylation was found in the light to dark phosphoproteome (Boex-Fontvieille et al., 2013) and wheat eIF4B was shown to respond to heat shock by dephosphorylation at multiple sites (Gallie et al., 1997). The effect of phosphorylation of plant eIF4B on its various activities needs further study in light of the recent work showing that phosphorylation of mammalian eIF4B and eIF4G influence the formation of an eIF4F/eIF4A/eIF4B complex and stimulate interaction with eIF3 and the 43S PIC (Dobrikov et al., 2012).

The divergence in eIF4B protein sequences between kingdoms suggest that there may be wide latitude in evolutionary constraints and unique mechanisms for regulation for this factor by phosphorylation (Hernández et al., 2010) leaving much to explore for the role of this “nonconserved” factor.

eIF4F and eIFiso4F

The role of eIF4F is to bind to the “cap” on the 5' end of the mRNA. This is a guanine residue methylated in the 7 position and attached to the first residue of the mRNA through a 5' to 5' linkage (m⁷GpppX). This reverse linkage helps to protect the mRNA, as the decapping of mRNA is one of the first enzymatically-driven steps in a major mRNA degradation pathway (Li and Kiledjian, 2010; Milac et al., 2014) and likely plays roles in plant gene expression during stress and other developmental pathways (Zhang et al., 2013). The m⁷G structure is recognized by the cap-binding protein eIF4E. eIF4E binds to the scaffold protein eIF4G to form the two-subunit complex called eIF4F, which is conserved in higher eukaryotes. eIF4G helps to assemble eIF4A, eIF4B and PABP on the mRNA to prepare it for binding to the 43S PIC (via interaction of eIF3 with eIF4G and eIF4B), in preparation for scanning of the 5' leader for an initiation codon (see below).

One distinction between the plant translational apparatus and that of other eukaryotes is the presence of a second cap-binding protein complex that differs from the canonical eIF4F. This two-protein complex, eIFiso4F, is comprised of eIFiso4G and eIFiso4E (Allen et al., 1992; van Heerden and Browning, 1994; Patrick and Browning, 2012). The cap binding proteins, eIF4E and eIFiso4E, have ~50% amino acid sequence similarity and form distinct and specific complexes with their respective binding partners, eIF4G and eIFiso4G (Mayberry et al., 2011). However, in the absence of the correct binding partner, mixed complexes will form that function in translation *in vitro* (Mayberry et al., 2011).

The larger scaffold subunits (eIF4G and eIFiso4G) share similarity in the C-terminal half with the eIF4E binding site and two HEAT (Huntington, elongation factor 3, protein phosphatase 2A, and the yeast TOR1 kinase) domains; however, the N-terminal half of eIF4G is absent in eIFiso4G (Patrick and Browning, 2012). eIFiso4F appears to be necessary for proper growth and development of Arabidopsis, as deletion of both *eIFiso4G* genes results

in slow and stunted growth, pale green rosettes and significant reproductive defects (Lellis et al., 2010). Based on phylogenetic analyses, eIFiso4G appeared in basal plant lineages before eIFiso4E and most likely formed a complex with eIF4E, whereas eIFiso4E emerged around the time flowering plants evolved (Patrick and Browning, 2012).

The role of the eIF4 factors is to bind the mRNA and prepare it for association with the 43S PIC. The current model for this process has the 5' cap binding to eIF4F through the eIF4E subunit. eIF4G (or eIFiso4G) serves as the scaffold for assembly of eIF4A, eIF4B and PABP (presumably binding the poly(A) tail and circularizing the mRNA at some time) as well as an interaction site for eIF3 (Jackson et al., 2010; Hinnebusch and Lorsch, 2012; Valasek, 2012). Although a structure for an eIF4F complex has not been determined, there are structures for portions of eIF4G in complex with eIF4E or eIF4A and considerable domain mapping for eIF4G from several organisms (Marintchev and Wagner, 2005; Marintchev et al., 2009; Dobrikov et al., 2012; Park et al., 2012), including plant eIF4G and eIFiso4G (Cheng and Gallie, 2010; Cheng and Gallie, 2013).

eIF4G and eIFiso4G

eIF4Gs have one to three HEAT domains depending upon the organism: yeast has one, plants have two, and mammals have three suggesting considerable evolution of the machinery (Hernández and Vazquez-Pianzola, 2005; Hernández et al., 2010; Hernández et al., 2012). HEAT domains are comprised of a series of alpha helices that form a coiled solenoid-like structure and provide surfaces for interaction with proteins and mRNA (Marintchev and Wagner, 2005; Valasek, 2012). Among these interactions are binding to eIF4A, eIF4B, eIF3 and PABP, which all function to bring the mRNA to the 40S subunit to initiate the scanning process (Hinnebusch and Lorsch, 2012).

The two eIF4A Rec A (N-terminal, C-terminal) domains interact with the middle section of the yeast HEAT domain or with the HEAT-1 and HEAT-2 domains of mammalian eIF4G (Marintchev et al., 2009; Hilbert et al., 2011). eIF4G promotes the transition from an "open" to a "closed" form of eIF4A based on crystal structures (Schütz et al., 2008; Marintchev et al., 2009). The model in yeast has been further refined by monitoring conformational changes in solution to include a "half-closed" intermediate for eIF4A that is stabilized upon binding of eIF4G (Hilbert et al., 2011). The closed state conformation of eIF4A is stimulated by ATP and RNA binding. It is also proposed that the closed form has a groove between the two interfaces of eIF4A and eIF4G that make the nucleotide binding site of eIF4A more accessible to ATP (Hilbert et al., 2011). The oscillations between the half-closed and closed forms may drive the release of ADP/P_i and rebinding of ATP to maintain mRNA binding affinity during scanning (Hilbert et al., 2011).

Plant eIFiso4G and eIF4G domains have been mapped and shown to have similar types of interactions with eIF4A, eIF4B and PABP, but have some differences from the mammalian or yeast interactions (Gallie, 2014). Further structural details will be required to understand how similar or different these plant factor interactions are to other eukaryotes.

An alternative to the current "script" of mRNA binding and unwinding by associated factors prior to binding to the 43S PIC, is that the eIF4 factors assemble on the 43S PIC and then recruit and unwind the mRNA, feeding it directly into the scanning complex (Hinnebusch and Lorsch, 2012). This model has many elements that explain some of the biochemical data but will require further testing not only in yeast but also in mammals and plants to determine if this is an accurate depiction of the scene within the cell.

"4F or, not 4F, that is the question"

The presence of two eIF4F complexes in plants raises the question of whether they have distinct biological activities. eIF4F promotes translation of reporter mRNAs with more secondary structure better than eIFiso4F in a cell-free translation system derived from wheat germ (Gallie and Browning, 2001). In addition, cellular mRNAs were shown to have varying levels of dependence upon eIF4F or eIFiso4F, as well as eIF4B, for optimal translation (Mayberry et al., 2009). These results suggest that plant eIFiso4F and eIF4F may have evolved selective abilities to promote or otherwise regulate translation of specific mRNA populations. What advantage distinct eIF4Fs provide to plants is not yet clear, but they may have pleiotropic roles in the synthesis of proteins involved in plant-specific functions, such as photosynthesis, cellulose biosynthesis, *etc.* eIFiso4F was implicated in the specific regulation of translation of an enzyme involved in the synthesis of chlorophyll (Chen et al., 2014) which fits with the observed pale green phenotype of Arabidopsis plants lacking eIFiso4G (Lellis et al., 2010).

Ultimately, structures of plant eIF4F or eIFiso4F in association with the 43S PIC will be needed to determine if there are functional differences in assembly of the initiation complexes in plants as compared to yeast and mammalian systems.

The cast of cap-binding proteins: eIF4E, eIFiso4E and 4EHP

All higher plants have three forms of the cap-binding proteins, eIF4E, eIFiso4E (plant-specific) and 4EHP (4E homologous protein) which all bind to m⁷GTP-Sepharose (Ruud et al., 1998; Patrick et al., 2014; Kropiwnicka et al., 2015). eIF4E and eIFiso4E are both class 1 cap-binding proteins (Joshi et al., 2005) and form the eIF4F and eIFiso4F complexes with their respective subunits, eIF4G and eIFiso4G (Mayberry et al., 2011; Patrick and Browning, 2012). 4EHP is termed a class 2 cap-binding protein (Joshi et al., 2005) and does not appear to function in canonical translation but has been implicated in regulatory roles for mRNAs during animal development in association with proteins that are not considered part of the canonical translational apparatus (Rom et al., 1998; Rhoads, 2009; Morita et al., 2012). The role of plant 4EHP (previously termed nCBP in plants for "novel" cap-binding protein) is unknown, although it has shown modest ability to stimulate translation with wheat eIFiso4G and appears to bind m⁷GTP more tightly than other cap-binding proteins (Ruud et al., 1998). There is more to learn about this unusual cap-binding protein and its role(s) in various cellular processes as capped non-coding and small RNAs are discovered and their functions elucidated.

It is also worth mentioning that there is a nuclear cap-binding protein complex (CBP20/CBP80) that shares structural and ancestral similarity to the eIF4E/eIFiso4E cap-binding proteins (CBP20) and eIF4G/eIFiso4G large subunits (CBP80). This complex does not function directly in translation but is important in pre-mRNA splicing, miRNA processing and export of mRNA in plants (Hugouvieux et al., 2001; Papp et al., 2004; Marintchev and Wagner, 2005; Topisirovic et al., 2011; Rogers and Chen, 2013; Gonatopoulos-Pournatzis and Cowling, 2014). There are many unanswered questions about the timing and location of exchange of CBP20 and eIF4E/eIFiso4E from the cap of mRNAs as it transitions from the nucleus to the cytoplasm.

The canonical type 1 plant cap-binding proteins, eIF4E and eIFiso4E, form tight complexes with their respective eIF4G subunits at the nanomolar to sub-nanomolar level (Mayberry et al., 2011). It is unlikely that these complexes readily dissociate given the tight binding affinity. They also do not appear to be regulated by any of the pathways associated with mammalian eIF4E that require dissociation/reassociation (see 4E Binding Proteins below). The crystal structure of wheat eIF4E shows similarities to both mammalian and yeast eIF4E, but revealed an intermolecular disulfide bridge between two plant-specific cysteine residues (Cys-113 and Cys151) that form under oxidizing conditions (Monzingo et al., 2007). This leads to the hypothesis that eIF4E could function as a redox sensor at the level of translational initiation. A constitutively reduced mutant (C113S) and oxidized forms of eIF4E bound m⁷GTP with a modest 1.5x difference in K_{off} values in NMR (nuclear magnetic resonance) solution studies (Monzingo et al., 2007). Further studies using mass spectrometry and a lysine-specific chemical probe indicated structural changes occurred upon altering the redox state and support the hypothesis of a redox sensor or switch for eIF4E (O'Brien et al., 2013). It remains a tantalizing possibility that the oxidation state of eIF4E (and/or eIFiso4E) may modulate cap binding in a redox-sensing manner in plants during retrograde signaling from the chloroplast to nucleus or other processes that could regulate the level of translation in response to the redox state of the cell.

Arabidopsis thaliana has three genes for the class 1 eIF4E (*eIF4E*, *eIF4E1b*, *eIF4E1c*) and one gene for eIFiso4E (Patrick and Browning, 2012; Patrick et al., 2014). *eIF4E* is the most highly expressed; the other two eIF4E genes (*eIF4E1b*, *eIF4E1c*) correspond to a tandem duplication event within the *Brassicaceae* with evidence that only *eIF4E1b* generates a transcript (Patrick and Browning, 2012) (see Table 1 for accession numbers). A number of forms of eIF4E have evolved in other organisms with special functions such as recognition of tri-methylated caps or tissue-specific regulation; however, some of these proteins have lost either their ability to bind m⁷G or eIF4G poisoning them as potential inhibitors or repressors of initiation (Joshi et al., 2005; Rhoads, 2009). eIF4E1b and eIF4E1c from *Arabidopsis* were found to bind eIF4G and to m⁷G-affinity resin and thus function biochemically *in vitro* as canonical cap-binding proteins (Patrick et al., 2014; Kropiwnicka et al., 2015). However, the double mutant *eIF4E eIFiso4E* is lethal suggesting that eIF4E1b and eIF4E1c are not able to replace eIF4E *in vivo* (Callot and Gallois, 2014; Patrick et al., 2014).

Several laboratories made the simultaneous discovery that mutations in the genes encoding eIFiso4E or eIF4E confer naturally occurring virus resistance and prevent viral replication (Jiang

and Laliberte, 2011; Wang and Krishnaswamy, 2012). Plants and many of their RNA viruses have co-evolved and a significant portion of this evolution appears to center on the use of eIF4F and eIFiso4F subunits by viruses for replication. Interestingly, neither *Arabidopsis eIF4E1b* nor *eIF4E1c* are recognized as virus resistance genes in contrast to dozens of examples for *eIF4E* and *eIFiso4E* (Robaglia and Caranta, 2006; Charron et al., 2008; Moury et al., 2013). Likely, other plants have multiple *eIF4E/eIFiso4E* genes, some of which may have evolutionary advantages for specific functions during stress or development.

The viral 5' linked protein (VPg) of potyviruses was found to interact directly with eIF4E or eIFiso4E, as well as other translation factors (eIF4G, eIFiso4G, PABP, eIF4A, eEF1A) (Jiang and Laliberte, 2011; Wang and Krishnaswamy, 2012). A number of mutations in subunits of cap-binding complexes interfere with virus reproduction, yet these mutations do not appear to compromise host protein synthesis. This suggests that it is not the protein synthesis activity *per se* that confers virus resistance, but some other aspect that has yet to be discovered. There is interest in using these genes to engineer better virus resistance for economically important crops (Wang and Krishnaswamy, 2012; Moury et al., 2013; Kim et al., 2014a) and it is likely that virus/host co-evolution has shaped the roles of these initiation factors in plants (Moury et al., 2013).

Positive strand plant viral RNAs recruit eIF4F and/or eIFiso4F using structural elements in the 3' UTRs termed 3' cap-independent translation enhancers (3'CITES) (reviewed in Simon and Miller, 2013). These varied structural RNA elements appear to function as "cap substitutes" through direct interaction with eIF4F or eIFiso4F (or both). Plant viral 3'CITES are unlike the internal ribosome entry site (IRES) elements associated with animal and insect viruses that use internal initiation as their hallmark and vary in their initiation factor requirement (Komar et al., 2012; Jackson, 2013). Plant viral 3'CITES typically make use of RNA-RNA interactions that base pair a portion of the 3'CITE with a 5' UTR loop ("kissing loop") in a manner reminiscent of the 5' to 3' interactions of the canonical initiation process involving the 5' cap and 3' poly(A) tail (Simon and Miller, 2013). Additionally, some plant viral RNA 3' UTRs use molecular mimicry by folding into structures that resemble tRNAs recruiting ribosomes directly (Simon and Miller, 2013). There is much to be learned from these interesting structures and the co-evolution of viruses and eIF4F/eIFiso4E host proteins. Since viruses are adept at co-opting host systems and using them to their advantage, it is likely that at least some host mRNAs may have elements similar to a 3'CITE.

eIF4E-Binding Proteins: Are they actors on the plant stage?

A major form of regulation of mammalian translation is through 4E binding proteins (4EBPs) that are regulated via phosphorylation by mammalian TOR (mTOR) in the PI3K-Akt signaling pathway that responds to many types of stress and environmental cues (Carrera, 2004; Richter and Sonenberg, 2005; Hernández et al., 2010). Phosphorylated mammalian 4EBPs dissociate from eIF4E, allowing it to interact with eIF4G to form a functional complex, whereas unphosphorylated 4EBPs bind to eIF4E and sequester it from interaction with eIF4G, thereby limiting eIF4F/

cap-dependent initiation (Carrera, 2004; Richter and Sonenberg, 2005). Plants have a functional TOR system that senses metabolic states (Ren et al., 2012; Robaglia et al., 2012; Xiong and Sheen, 2012; Caldana et al., 2013; Dobrenel et al., 2013; Xiong et al., 2013; Xiong and Sheen, 2013, 2014), but appear to lack 4EBPs that regulate eIF4F complex formation (Verma and Chatterjee, 2009). Given the sub/nanomolar binding of the plant eIF4F and eIFiso4F subunits to each other, it seems unlikely that these complexes will dissociate once formed (Mayberry et al., 2011). Plants have proteins with canonical eIF4E binding sites that have been shown to bind to eIF4E or eIFiso4E, such as lipoxygenase 2 and BTF3 (beta subunit of the nascent polypeptide-associated complex (Freire et al., 2000; Freire, 2005); however, the role(s) of these protein interactions with plant cap-binding proteins is still unclear.

Phosphorylation of eIF4G/iso4G and eIF4E/iso4E

Although there are reports of multiple isoelectric states of wheat eIF4F/iso4F subunits, the effect on activity and the kinases involved have not been identified (Gallie et al., 1997). Neither eIF4F nor eIFiso4F subunits were targets of CK2, unlike eIF4B (Dennis and Browning, 2009). Only eIF4G was found to be phosphorylated in the phosphoproteome of the light to dark transition (Boex-Fontvieille et al., 2013). Given the importance of phosphorylation of eIF4E, 4E-BP and eIF4G in mammalian systems, it remains to be discovered if plants have evolved a different system for regulation of these subunits through phosphorylation, redox-sensitive structure regulation or other means.

Poly(A) Binding Protein

Although not an "official" initiation factor, PABP binds to the 3' poly(A) tail of the mRNA and interacts with eIF4G and eIF4B, suggesting that the mRNA may be circularized at least transiently during initiation (Park et al., 2011). In mammals, PABPs have extensive roles in the nucleus and cytoplasm in mRNA processing, translation and degradation, as well as a role in miRNA-mediated processes (reviewed in Goss and Kleiman, 2013). Higher eukaryotes have multiple genes for PABP. In the case of *X. laevis*, PABP gene products have been shown to function similarly in translation, but are distinctly required for development indicating there may be mRNAs whose processing, expression or degradation requires a specific PABP (Gorgoni et al., 2011).

Plants have an extensive family of genes encoding PABP with considerable protein sequence diversity. The eight PAB genes in *A. thaliana* fall into four phylogenetic sub-groups with varying tissue specific expression (Le and Gallie, 2000; Belostotsky, 2003). In general, Arabidopsis and other plant PABPs have four RRM domains that consist of two α -helices and four anti-parallel β -sheets. A separately folded C-terminal domain called PABC is composed of 4 to 5 α -helices and contains a PABC interaction motif (PAM2) for protein-protein interaction. The solution structure of wheat PABC has a similar fold to the mammalian PABC domain and also contains the PABC-Interacting Motif (PAM-2) protein interaction domain (Siddiqui et al., 2007). *A. thaliana* PABC/PAM2

was shown to have multiple binding partners, several of which interfere with *in vitro* translation or are implicated in RNA metabolism (Wang and Grumet, 2004; Bravo et al., 2005; Siddiqui et al., 2007). Some plant PABPs have been reported to have two instead of four RRMs, suggesting specifically evolved functions for these proteins (Belostotsky, 2003). It is also well established that there can be multiple PABPs bound to the poly(A) tail of the mRNA at any given time, thus leading to a diversity of possible PABP molecules on one transcript, each perhaps recruiting different binding partners via the PAM2 interface. Adding further complexity is post-translational modification by phosphorylation, which also affects PABP's interactions with binding partners (Le et al., 2000).

Extensive biochemical analysis of wheat PABP has shown that the presence of eIF4G or eIF4B enhances its RNA binding activity and in turn the presence of PABP increases the affinity of the eIF4F complex for the cap and stimulates the ATPase and RNA helicase activities of eIF4A, eIF4F/eIFiso4F and eIF4B (reviewed in Gallie, 2014; Le et al., 1997; Wei et al., 1998; Bi and Goss, 2000; Luo and Goss, 2001; Khan and Goss, 2005; Cheng and Gallie, 2010). It has been further shown that plant eIF4G has an additional PABP binding domain that binds eIF4B in a competitive and mutually exclusive manner (Cheng and Gallie, 2010; Cheng and Gallie, 2013). This second domain is absent in mammals and yeast PABP. PABP is also implicated in viral replication (Smith and Gray, 2010) and plant PABP was shown to interact with the reverse transcriptase of turnip mosaic virus (TuMV) (Dufresne et al., 2008) and with the 3'UTR of tobacco etch virus (TEV) to promote internal initiation (Khan et al., 2008; Khan et al., 2009; Yumak et al., 2010; Iwakawa et al., 2012).

There are still many questions about the role of PABP in initiation, its interactions with various initiation factors and other proteins, such as the PAM2-domain containing Early Responsive to Dehydration 15 (ERD15) family members (Aalto et al., 2012). PABP function likely extends beyond initiation. For example, an Arabidopsis *pab2 pab8* loss-of-function mutant maintained translation of late-embryogenesis mRNAs in young seedlings, leading to the suggestion that PABP contributes to turnover of abundant seed transcripts during early seedling development (Tiruneh et al., 2013).

eIF1 Group

This group is involved in the stimulation and assembly of the 43S PIC and includes eIF1 (called SUI1 in yeast) and eIF1A (formerly known as eIF4C). Both of these small factors (~12-17 kDa) are single polypeptides and conserved across all eukaryotes. eIF1 has structural similarity to the initiation factor (IF)3 C-terminal domain in prokaryotes and eIF1A is the functional equivalent to prokaryotic IF1 (Valasek, 2012).

eIF1 and eIF1A

eIF1 binds near the peptidyl (P)-site of the 40S subunit and precludes Met-tRNA_{Met} bound to eIF2•GTP (the ternary complex) from being fully engaged in the peptidyl (P)-site of the 43S PIC until the initiation codon is accurately identified (Nanda et al., 2013; Martin-Marcos et al., 2014). eIF1A has an interesting struc-

ture with a folded central region that binds in the acceptor (A)-site of the 40S subunit; however, its N- and C terminal tails are unstructured and extend into the P-site. Similar to eIF1, eIF1A participates in preventing full P-site engagement of the Met-tRNA^{Met} until the correct initiation codon is identified (Nanda et al., 2013).

Recombinant wheat eIF1 was shown to function in formation of a multifactor complex (MFC) *in vitro* similar to that found in yeast and mammals (Asano et al., 2000; Dennis et al., 2009; Hinnebusch and Lorsch, 2012; Sokabe et al., 2012; Hinnebusch, 2014). The MFC, consisting of eIF1, eIF2, eIF3 and eIF5, presumably helps to organize these factors prior to binding to the 43S PIC. It also appears to stabilize binding of the ternary complex to the 40S subunit. Plant eIF1 interacts directly with eIF5 and the N-terminal domain of eIF3c (Dennis et al., 2009). On the other hand, eIF1A binds to the 40S subunit independently of the MFC. eIF1A purified from wheat germ substitutes biochemically for rabbit reticulocyte eIF1A suggesting a highly conserved function (Timmer et al., 1993). Overexpression of *eIF1* and *eIF1A* have been reported to improve salt tolerance in plants (Rausell et al., 2003; Latha et al., 2004; Diedhiou et al., 2008; Sun and Hong, 2013), suggesting a role in stress acclimation.

eIF5 Group

Two members (eIF5, eIF5B) of this group function in the selection of the start site and engagement of codon-anticodon base pairing, whereas the third (eIF5A), functions in elongation. All three group members, eIF5, eIF5A and eIF5B are found in plants and other eukaryotes, suggesting ancient origin and conserved functions. eIF5 has GTPase activating protein function. eIF5B resembles prokaryotic IF2, but does not bind Met-tRNA^{Met} and functions to promote binding of the 60S subunit (subunit joining) which in turn promotes GTP hydrolysis and release of eIF5B as the last step of the initiation process. eIF5A (nee eIF4D), although initially designated an initiation factor, is the “imposter” in the group and promotes the peptidyl transferase reactions of poly-prolyl residues during elongation (Gutierrez et al., 2013) Given the role for eIF5A in elongation, it has been proposed to rename this factor eEF5 (Dever et al., 2014) (See section on Elongation Factors and Table 2).

eIF5

This protein promotes hydrolysis of GTP bound to the ternary complex during start site recognition (Jennings and Pavitt, 2010; Aitken and Lorsch, 2012; Hinnebusch and Lorsch, 2012; Valasek, 2012; Nanda et al., 2013; Hinnebusch, 2014). eIF5 has two functional domains, N-terminal (NTD) and C-terminal (CTD) with a linker connection (Conte et al., 2006; Wei et al., 2006). An “arginine finger” (Arg-15) required for GTPase activity is positioned near the N-terminus in an unstructured region. This unstructured region is free to interact with the GTP-binding region of eIF2γ to promote GTP hydrolysis. In addition, the NTD has structural similarity to eIF1, which may play a role in events during initiation codon selection (Nanda et al., 2013). The eIF5 CTD contains a HEAT domain that interacts with eIF1, the NTD of eIF3c, and N-terminal tail of eIF2β to stabilize the MFC mentioned earlier that

is formed by these factors (Nanda et al., 2013). In addition, the CTD of yeast eIF5 interacts with an unstructured region of eIF4G that is proposed to promote binding of the mRNA to the 43S PIC and assist in scanning and subsequent release of eIF1 (Singh et al., 2012). Whether this occurs with mammalian or plant eIF4G is not known. eIF5 has been shown to be released from mammalian PIC with eIF2•GDP and appears to have a role as GDP dissociation inhibitor during recycling of eIF2•GDP by eIF2B (Jennings et al., 2013). eIF5 from plants has received little attention, except studies that show that the wheat factor functions in the formation of a MFC in a manner enhanced by phosphorylation of members of the MFC by CK2 (Dennis et al., 2009). One of the *in vitro* CK2 sites of eIF5 was confirmed in the light/dark phosphoproteome (Boex-Fontvieille et al., 2013).

eIF5B

As the structural homolog of eubacterial IF2, eIF5B carries out a similar function in eukaryotes (Allen and Frank, 2007). Upon recognition of the initiation codon, a series of events including the release of eIF1 from its position near the P-site and hydrolysis of the GTP bound to ternary complex, leads to conformation changes in eIF2 (Allen and Frank, 2007). At this point eIF5B•GTP binds to the complex via contacts with the C-terminal tail of eIF1A and may stabilize the binding of the Met-tRNA^{Met} in the P-site (Pisareva and Pisarev, 2014). The binding of eIF5B•GTP likely displaces many of the associated factors (eIF2, eIF3 and eIF5) to open a surface for 60S subunit attachment. Hydrolysis of eIF5B•GTP is promoted by the GTPase activating activity of the 60S subunit, triggering release of eIF5B•GDP and eIF1A as the newly formed 80S complex is established (Allen and Frank, 2007). Mutations in mammalian eIF5B show this factor may play multiple roles during initiation *in vitro* (Pisareva and Pisarev, 2014). Pea eIF5B has been biochemically characterized as a heat stable protein with potential properties of a chaperone and binds to GTP as expected (Rasheedi et al., 2010; Suragani et al., 2011).

eIF6 Group

Designated as an initiation factor and the sole performer of this group, eIF6 interacts with the 60S subunit and functions in the prevention of premature association of the 60S ribosomal subunit with the 43S PIC. It also has a role in the assembly of the 40S and 60S ribosomal subunits (Miluzio et al., 2009; Brina et al., 2011). eIF6 was first discovered in wheat germ as a ribosome disassociation factor that bound 60S ribosomes (Russell and Spremulli, 1978, 1979, 1980) and subsequently identified in yeast (Si et al., 1997) and mammals (Valenzuela et al., 1982; Raychaudhuri et al., 1984).

In sequenced plant genomes, eIF6 is typically encoded by multiple genes; in the case of Arabidopsis a single eIF6 gene (*AteIF6A*) is broadly expressed and a second displays more regional and regulated transcript accumulation (*AteIF6B*) (Guo et al., 2011a). The role of eIF6 in subunit joining involves its interaction with the conserved ribosome-associated protein receptor of activated C kinase 1 (RACK1) (see section below on RACK1).

Table 2. Elongation and Termination Factors of *Arabidopsis*

Factor	M _r ^a	Function	Arabidopsis Gene ^b
eEF1A (EF-1α ^b)	52,000	Bind aminoacyl-tRNA and GTP	At1g07920, At1g07930, At1g07930, At1g07940, At5g60390
eEF1B		Recycle eEF1A·GDP	
α ^c	24,000		At5g12110, At5g19510
β ^d	28,000		At1g30230, At2g18110
γ	46,000		At1g09640, At1g57720
eEF2 ^e	92,000	Translocation	At1g56070, At3g12915
eEF5 ^f (eIF4D, eIF5A) ^g	17,200	Elongation of poly-proline/glycine regions	AT1g26630, AT1g69410, AT1g13950
eRF1	49,000	Termination/peptide release	At5g47880, At1g12920, At3g26618
eRF3	60,500	Termination/peptide release	At1g18070
ABCE1/RLI1 ^g	68,000	Ribosome recycling	At3G13640, At4g19210 ^g

^a Approximate molecular weight based on TAIR9 data.

^b Prior nomenclature used in literature.

^c The old *Artemia* and mammalian designation was EF-1δ. The old plant designation was EF-1β.

^d The old *Artemia* and mammalian designation was EF-1β. The old plant designation was EF-1β'.

^e A conserved histidine residue is post-translationally modified to a diphthamide.

^f Originally designated eIF4D and later eIF5A; now known to participate in elongation and renamed eEF5; contains the unique amino acid hypusine.

^g We recommend At3G13640 be called *ABCE1A* and At4g19210 be called *ABCE1B* rather than *ABCE1* and *ABCE2*, respectively.

^h Links to various data bases using the Arabidopsis Gene Identifier can be found at <http://browning.cm.utexas.edu/arabidopsis/flat>

eIF6 also functions in ribosome biogenesis and the transport of ribosomal subunits from the nucleolus to the cytoplasm in a process that requires phosphorylation of a CK1 site that is conserved in *AteIF6A*, but lost in *AteIF6B* mutants (Guo et al., 2011a).

The Ribosome: the *Prima Donna*

The peptidyl transferase reaction is catalyzed by the ribosome, a highly evolutionarily conserved macromolecular complex of two subunits that is comprised of RNA and proteins (Yusupova and Yusupov, 2014). Without exception, the role of the small subunit of the ribosome is to decode the mRNA whereas the large subunit catalyzes the peptidyl transferase reaction (peptide bond formation). Decoding involves the A-, P- and exit (E)-site positions transiently occupied by tRNAs as they bring amino acids to transfer to the growing polypeptide chain and exit empty to be recharged.

Subunit composition

Ribosomes, their subunits, and rRNAs are measured in Svedberg (S) units corresponding to their sedimentation coefficient measured by ultracentrifugation. When joined together, cytosolic ribosomes of higher eukaryotes (including plant 40S and 60S subunits) sediment at 80S. The 40S subunit is formed with 18S rRNA

and small ribosomal proteins (RPSs) and the 60S by the 5S, 5.8S, and 25-28S rRNAs and large ribosomal proteins (RPLs). The subunits of lower eukaryotes have the same four rRNA molecules and are therefore slightly smaller. The ribosomes of bacteria, mitochondria and plastids are significantly smaller, typically consisting of a 30S small subunit with a 16S rRNA and a 50S large subunit with 23S and 5S rRNAs and no 5.8S rRNA. The rRNAs of eukaryotes possess several expanded segments and variable regions relative to their bacterial counterparts. It was postulated that these expansion regions are associated with the more complex control of translation. Accompanying the rRNA distinctions are eukaryote-specific RPs, all of which are encoded in higher plants (Barakat et al., 2001). The remaining RPs fall into two groups of either eubacterial (found across kingdoms) or archaea/eukaryote-specific origin (Armache et al., 2010b, a). A total of 54 RPs are recognized in eubacteria, 79 in yeast, and 79- 80 RPs in higher eukaryotes. With the exception of the RPs that form a flexible lateral stalk of the large subunit, all are present in a single copy per ribosome (Yusupova and Yusupov, 2014).

Ribosome architecture

High-resolution crystal structure analyses confirm pronounced conservation of the three-dimensional structure of ribosomes between eubacterial, archaebacterial and eukaryotic kingdoms

(Klinge et al., 2012; Voigts-Hoffmann et al., 2012; Yusupova and Yusupov, 2014). The larger mass of eukaryotic ribosomes has presented a greater challenge for obtaining crystals with high-quality diffraction characteristics. Consequentially, insights into eukaryote-specific structural features of ribosomes have been gleaned from cryo-EM structural analyses, including models at 38 Å (Verschoor et al., 1996) and at <10 Å resolution for translating wheat, yeast, insect, and mammalian ribosomes (Armache et al., 2010b, a). These models provide a wealth of insight into the position and structure of rRNAs and RPs.

Eukaryotic-specific RPs are located at several key positions within animal, yeast and wheat ribosomes, including the sites associated with decoding and tRNA binding, and mRNA exit on the 40S subunit. RPs specific to both subunits interact with the eukaryote-specific factor eIF3. The cryo-EM studies of insect and mammalian ribosomes emphasize the presence of rRNA expansion regions on the outer periphery that dynamically form RP-rRNA and rRNA-rRNA interactions (Anger et al., 2013). The wheat ribosome is more like that of yeast with a less extensive outer rRNA-protein layer. Further studies of plant ribosome architecture are needed to better appreciate kingdom-specific features as well as structural variations that might be associated with specific RP isoforms.

Interest in plant ribosomes, as well as other organisms, centers around several questions: Do ribosomes play specific roles in global translational activity or the translation of individual mRNAs? Is there ribosome heterogeneity due to differences in protein isoform composition or modification and if so, what is its function(s)? Are ribosome biogenesis and protein synthesis tightly regulated as a means of energy conservation and does management of these investments pace growth during the diurnal cycle or under abiotic environment? Is a threshold level of ribosomes necessary for cellular and organismal homeostasis?

Cytosolic ribosomal proteins

Eighty RP gene families of two to five paralogous members were identified in the Arabidopsis genome by comparison to the amino acid sequences of animal, yeast and *Archaea* RPs (Barakat et al., 2001) (see Table 3). The vast majority of RPs are basic in charge ($pI > 8.0$) and ≤ 45 kDa in mass. There are, however, a handful of conserved acidic RPs ($pI < 5$). Eukaryotes have two RP gene families that encode small acidic phosphoproteins that dimerize and bind the larger RPP0 to form a flexible lateral stalk structure on the large ribosomal subunit. Higher plants have a RPP1 and RPP2 family, as well as a third acidic RP called RPP3 (Szick et al., 1998; Chang et al., 2005). This stalk structure promotes eEF2 binding and GTP hydrolysis in yeast (Gonzalo and Reboud, 2003) and is important in the binding of ribosome inactivating proteins, such as the ricin toxin (Li et al., 2013c).

Several mass spectrophotometric studies have explored the proteome of Arabidopsis ribosomes. Ribosomes purified by differential centrifugation yielded 31-33 of the putative 40S RPs and 48 of the putative 60S RPs, respectively (Chang et al., 2005; Giavalisco et al., 2005; Carroll et al., 2008; Piques et al., 2009; Turkina et al., 2011; Carroll, 2013). An analysis performed on ribosomes captured by immunopurification, thereby limiting con-

tamination by organellar ribosomes and other co-sedimenting complexes, confirmed products from 204 of the estimated 232 functional RP genes of *A. thaliana* based on two or more prototypic peptides (Hummel et al., 2012). This corresponded to 64 of the 80 putative RP families and included RACK1. Of these, 74 Arabidopsis RPs were positioned in a high-resolution 80S ribosome structure map relative to the rRNA structure (Armache et al., 2010b).

Ribosome heterogeneity

The biogenesis of a ribosome requires coordinated synthesis of rRNAs and RPs. In Arabidopsis, the 18S, 5.8S and 25S rRNA are encoded by the 45S rDNA genic repeat that is tandemly duplicated over 500 times on the short arms of chromosome 2 and 4, whereas the pre-5S rRNA is encoded by repetitive pericentromeric regions on chromosomes 3, 4 and 5 (Layat et al., 2012). The 18S, 5.8S and 25S rRNA precursor of Arabidopsis is sometimes referred to as the 35S pre-rRNA. The transcription of the Arabidopsis 45S rDNA units by RNA polymerase I is regulated by direct binding of TOR to the promoter region located in the intergenic region (spacer) between the 25S and 18S genes (Ren et al., 2011). The subsequent pre-rRNA processing pathway is a complicated pathway involving cleavage and nucleotide modification events. Synthesis of 5S rRNA by RNA polymerase III is regulated by mTOR in animals (Kantidakis et al., 2010); however, the regulation of plant 5S rRNA synthesis is not well characterized.

Arabidopsis RP genes are significantly co-regulated at transcriptional and post-transcriptional levels during development, in response to various stimuli and in some translational apparatus mutants. The coordinate transcriptional and posttranscriptional regulation of many RP genes reflects the presence of common *cis*-regulatory elements in gene promoters and features of mRNA 5'UTRs (Kawaguchi et al., 2004; Branco-Price et al., 2005; McIntosh and Bonham-Smith, 2006; Nicolai et al., 2006; Tiruneh et al., 2013; Wang et al., 2013). Close inspection of Arabidopsis RP transcript accumulation data indicate that there is more than one RP transcription network regulated in response to carbon and nitrogen availability, as well as other environmental inputs (Sormani et al., 2011a; Wang et al., 2013). RP gene co-regulation was also noted when monitoring polysome-associated transcript levels. A coordinate shift of Arabidopsis RP mRNAs out of polysome complexes was observed in response to a number of environmental stresses (Kawaguchi et al., 2004; Branco-Price et al., 2005; McIntosh and Bonham-Smith, 2006; Nicolai et al., 2006; Branco-Price et al., 2008; Pyl et al., 2012; Tiruneh et al., 2013; Wang et al., 2013). Conversely, coordinated up-regulation of the translation of a large majority of Arabidopsis RP mRNAs was recorded in two mutants of the translational apparatus (*eif3h* and *rpl24b*) (Tiruneh et al., 2013). Co-regulation of RP gene transcript accumulation and translation has also been noted in *Chlamydomonas* (Schmollinger et al., 2014).

The stereotype of a ribosome is that all are the same, but there is some evidence of specialized differences in cytosolic ribosomes of Arabidopsis and other plants that may contribute to the regulation of translation (Horiguchi et al., 2012; Hummel et al., 2012). Ribosome heterogeneity is predicted to be the con-

Table 3. Cytosolic Ribosomal Protein Genes of *Arabidopsis*^a

Small Subunit Genes			Large Subunit Genes		
Protein Family Name	Gene Names	Arabidopsis Gene Identifier of Gene Family Members	Protein Family Name	Gene Names	Arabidopsis Gene Identifier of Gene Family Members
Sa	<i>RPSaA</i>	At1g72370	P0	<i>RPP0A</i>	At2g40010
	<i>RPSaB</i>	At3g04770		<i>RPP0B</i>	At3g09200
S2	<i>RPS2A</i>	At1g58380		<i>RPP0C</i>	At3g11250
	<i>RPS2B</i>	At1g59359	P1	<i>RPP1A</i>	At1g01100
	<i>RPS2C</i>	At2g41840		<i>RPP1B</i>	At4g00810
	<i>RPS2D</i>	At3g57490		<i>RPP1C</i>	At5g47700
	<i>RPS2E</i>	At1g58684		<i>RPP1D</i>	At5g24510
	<i>RPS2F</i>	At1g58983	P2	<i>RPP2A</i>	At2g27720
S3	<i>RPS3A</i>	At2g31610		<i>RPP2B</i>	At2g27710
	<i>RPS3B</i>	At3g53870		<i>RPP2C</i>	At3g28500
	<i>RPS3C</i>	At5g35530	<i>RPP2D</i>	At3g44590	
S3a	<i>RPS3aA</i>	At3g04840	<i>RPP2E</i>	At5g40040	
	<i>RPS3aB</i>	At4g34670	P3	<i>RPP3A</i>	At4g25890
S4	<i>RPS4A</i>	At2g17360		<i>RPP3B</i>	At5g57290
	<i>RPS4B</i>	At5g07090	L3	<i>RPL3A</i>	At1g43170
	<i>RPS4D</i>	At5g58420		<i>RPL3B</i>	At1g61580
S5	<i>RPS5A</i>	At2g37270	L4	<i>RPL4A</i>	At3g09630
	<i>RPS5B</i>	At3g11940		<i>RPL4D</i>	At5g02870
S6	<i>RPS6A</i>	At4g31700	L5	<i>RPL5A</i>	At3g25520
	<i>RPS6B</i>	At5g10360		<i>RPL5B</i>	At5g39740
S7	<i>RPS7A</i>	At1g48830	L6	<i>RPL6A</i>	At1g18540
	<i>RPS7B</i>	At3g02560		<i>RPL6B</i>	At1g74060
	<i>RPS7C</i>	At5g16130		<i>RPL6C</i>	At1g74050
S8	<i>RPS8A</i>	At5g20290	L7	<i>RPL7A</i>	At1g80750
	<i>RPS8B</i>	At5g59240		<i>RPL7B</i>	At2g01250
S9	<i>RPS9B</i>	At5g15200	<i>RPL7C</i>	At2g44120	
	<i>RPS9C</i>	At5g39850	<i>RPL7D</i>	At3g13580	
S10	<i>RPS10A</i>	At4g25740	L7a	<i>RPL7aA</i>	At2g47610
	<i>RPS10B</i>	At5g41520		<i>RPL7aB</i>	At3g62870
	<i>RPS10C</i>	At5g52650	L8	<i>RPL8A</i>	At2g18020
S11	<i>RPS11A</i>	At3g48930		<i>RPL8B</i>	At3g51190
	<i>RPS11B</i>	At4g30800		<i>RPL8C</i>	At4g36130
	<i>RPS11C</i>	At5g23740	L9	<i>RPL9B</i>	At1g33120
S12	<i>RPS12A</i>	At1g15930		<i>RPL9C</i>	At1g33140
	<i>RPS12C</i>	At2g32060		<i>RPL9D</i>	At4g10450
S13	<i>RPS13A</i>	At3g60770	L10	<i>RPL10A</i>	At1g14320
	<i>RPS13B</i>	At4g00100		<i>RPL10B</i>	At1g26910
S14	<i>RPS14A</i>	At2g36160		<i>RPL10C</i>	At1g66580
	<i>RPS14B</i>	At3g11510	L10a	<i>RPL10aA</i>	At1g08360
	<i>RPS14C</i>	At3g52580		<i>RPL10aB</i>	At2g27530
S15	<i>RPS15A</i>	At1g04270		<i>RPL10aC</i>	At5g22440
	<i>RPS15B</i>	At5g09490	L11	<i>RPL11A</i>	At2g42740
S15	<i>RPS15A</i>	At1g04270		<i>RPL11B</i>	At3g58700
	<i>RPS15B</i>	At5g09490		<i>RPL11C</i>	At4g18730
	<i>RPS15B</i>	At5g09490		<i>RPL11D</i>	At5g45775
S15	<i>RPS15A</i>	At1g04270	L12	<i>RPL12A</i>	At2g37190
	<i>RPS15B</i>	At5g09490		<i>RPL12B</i>	At3g53430
	<i>RPS15B</i>	At5g09490		<i>RPL12C</i>	At5g60670

continued

Table 3. (continued)

Small Subunit Genes			Large Subunit Genes		
Protein Family Name	Gene Names	Arabidopsis Gene Identifier of Gene Family Members	Protein Family Name	Gene Names	Arabidopsis Gene Identifier of Gene Family Members
S15a	<i>RPS15C</i>	At5g09500	L13	<i>RPL13B</i>	At3g49010
	<i>RPS15D</i>	At5g09510		<i>RPL13C</i>	At3g48960
	<i>RPS15E</i>	At5g43640		<i>RPL13D</i>	At5g23900
	<i>RPS15F</i>	At5g63070	L13a	<i>RPL13aA</i>	At3g07110
	<i>RPS15aA</i>	At1g07770		<i>RPL13aB</i>	At3g24830
	<i>RPS15aB b</i>	At2g19720		<i>RPL13aC</i>	At4g13170
	<i>RPS15aC</i>	At2g39590		<i>RPL13aD</i>	At5g48760
S16	<i>RPS15aD</i>	At3g46040	L14	<i>RPL14A</i>	At2g20450
	<i>RPS15aE b</i>	At4g29430		<i>RPL14B</i>	At4g27090
	<i>RPS15aF</i>	At5g59850	L15	<i>RPL15A</i>	At4g16720
S17	<i>RPS16A</i>	At2g09990		<i>RPL15B</i>	At4g17390
	<i>RPS16B</i>	At3g04230	L17	<i>RPL17A</i>	At1g27400
	<i>RPS16C</i>	At5g18380		<i>RPL17B</i>	At1g67430
S18	<i>RPS17A</i>	At2g04390	L18	<i>RPL18A</i>	At2g47570
	<i>RPS17B</i>	At2g05220		<i>RPL18B</i>	At3g05590
	<i>RPS17C</i>	At3g10610		<i>RPL18C</i>	At5g27850
	<i>RPS17D</i>	At5g04800	L18a	<i>RPL18aB</i>	At2g34480
S19	<i>RPS18A</i>	At1g22780		<i>RPL18aC</i>	At3g14600
	<i>RPS18B</i>	At1g34030	L19	<i>RPL19A</i>	At1g02780
	<i>RPS18C</i>	At4g09800		<i>RPL19B</i>	At3g16780
S20	<i>RPS19A</i>	At3g02080		<i>RPL19C</i>	At4g02230
	<i>RPS19B</i>	At5g15520	L21	<i>RPL21A</i>	At1g09590
	<i>RPS19C</i>	At5g61170		<i>RPL21C</i>	At1g09690
S21	<i>RPS20A</i>	At3g45030		<i>RPL21E</i>	At1g57660
	<i>RPS20B</i>	At3g47370	L22	<i>RPL21G</i>	At1g57860
	<i>RPS20C</i>	At5g62300		<i>RPL22A</i>	At1g02830
S22	<i>RPS21A</i>	At3g53890		<i>RPL22B</i>	At3g05560
	<i>RPS21B</i>	At3g53890	L23	<i>RPL22C</i>	At5g27770
S23	<i>RPS21C</i>	At5g27700		<i>RPL23A</i>	At1g04480
	<i>RPS23A</i>	At3g09680	L23a	<i>RPL23B</i>	At2g33370
S24	<i>RPS23B</i>	At5g02960		<i>RPL23C</i>	At3g04400
	<i>RPS24A</i>	At3g04920	L24	<i>RPL23aA</i>	At2g39460
S25	<i>RPS24B</i>	At5g28060		<i>RPL23aB</i>	At3g55280
	<i>RPS25A</i>	At2g16360		<i>RPL24A</i>	At2g36620
	<i>RPS25B</i>	At2g21580	L26	<i>RPL24B</i>	At3g53020
S26	<i>RPS25D</i>	At4g34555		<i>RPL24C</i>	At2g44860
	<i>RPS25E</i>	At4g39200	L27	<i>RPL26A</i>	At3g49910
	<i>RPS26A</i>	At2g40510		<i>RPL26B</i>	At5g67510
S27	<i>RPS26B</i>	At2g40590	L27a	<i>RPL27A</i>	At2g32220
	<i>RPS26C</i>	At3g56340		<i>RPL27B</i>	At3g22230
	<i>RPS27A</i>	At2g45710		<i>RPL27C</i>	At4g15000
S28	<i>RPS27B</i>	At3g61110	L28	<i>RPL27aA</i>	At1g12960
	<i>RPS27D</i>	At5g47930		<i>RPL27aB</i>	At1g23290
	<i>RPS27C</i>	At3g56340		<i>RPL27aC</i>	At1g70600
S29	<i>RPS28A</i>	At2g19730	L29	<i>RPL28A</i>	At2g19730
	<i>RPS28B</i>	At4g29410		<i>RPL28C</i>	At4g29410
S30	<i>RPS29A</i>	At3g06700		<i>RPL29A</i>	At3g06700

Table 3. (continued)

Small Subunit Genes			Large Subunit Genes		
Protein Family Name	Gene Names	Arabidopsis Gene Identifier of Gene Family Members	Protein Family Name	Gene Names	Arabidopsis Gene Identifier of Gene Family Members
S27a	<i>RPS27aA</i>	At1g23410	L30	<i>RPL29B</i>	At3g06680
	<i>RPS27aB</i>	At2g47110		<i>RPL30A</i>	At1g36240
	<i>RPS27aC</i>	At3g62250		<i>RPL30B</i>	At1g77940
S28	<i>RPS28A</i>	At3g10090	L31	<i>RPL30C</i>	At3g18740
	<i>RPS28B</i>	At5g03850		<i>RPL31A</i>	At2g19740
	<i>RPS28C</i>	At5g64140		<i>RPL31B</i>	At4g26230
S29	<i>RPS29A</i>	At3g43980	L32	<i>RPL31C</i>	At5g56710
	<i>RPS29B</i>	At3g44010		<i>RPL32A</i>	At4g18100
	<i>RPS29C</i>	At4g33865	L34	<i>RPL32B</i>	At5g46430
	<i>RPS30A</i>	At2g19750		<i>RPL34A</i>	At1g26880
S30	<i>RPS30B</i>	At4g29390	L35	<i>RPL34B</i>	At1g69620
	<i>RPS30C</i>	At5g56670		<i>RPL34C</i>	At3g28900
RACK1 ^c	<i>RACK1A</i>	At1g18080	L35a	<i>RPL35A</i>	At3g09500
	<i>RACK1B</i>	At1g48630		<i>RPL35B</i>	At2g39390
	<i>RACK1C</i>	At3g18130		<i>RPL35C</i>	At3g55170
				<i>RPL35D</i>	At5g02610
			L36	<i>RPL35aA</i>	At1g07070
				<i>RPL35aB</i>	At1g41880
				<i>RPL35aC</i>	At1g74270
			L36a	<i>RPL35aD</i>	At3g55750
				<i>RPL36A</i>	At2g37600
				<i>RPL36B</i>	At3g53740
			L37	<i>RPL36C</i>	At5g02450
				<i>RPL36A</i>	At3g23390
				<i>RPL36aB</i>	At4g14320
			L37a	<i>RPL37A</i>	At1g15250
				<i>RPL37B</i>	At1g52300
				<i>RPL37C</i>	At3g16080
			L38	<i>RPL37aB</i>	At3g10950
				<i>RPL37aC</i>	At3g60245
			L39	<i>RPL38A</i>	At2g43460
				<i>RPL38B</i>	At3g59540
			L40	<i>RPL39A</i>	At2g25210
				<i>RPL39B</i>	At3g02190
				<i>RPL39C</i>	At4g31985
			L41	<i>RPL40A</i>	At2g36170
				<i>RPL40B</i>	At3g52590
			L41	<i>RPL41C</i>	At2g40205
				<i>RPL41D</i>	At3g08520
				<i>RPL41E</i>	At3g11120
				<i>RPL41G</i>	At3g56020

^a RP genes that have not been identified as pseudogenes (Barakat et al., 2001; Hummel et al., 2012; The Arabidopsis Information Resource, October 2014). Proteomic studies using 1D and 2D mass spectrometry have obtained evidence for the product of a least one paralog of each RP family in purified ribosomes, with the exception of the small (~3.5 kDa) and highly basic RPL41 (reviewed by Carroll, 2013).

^b Two *RPS15a* paralogs encode RPs that assemble into mitochondrial and not cytosolic ribosomes (Carroll et al., 2008).

^c Three *RACK1* paralogs encode a protein stably associated with the ribosome; RACK1s also function in signaling outside of the ribosome.

sequence of ancestral genome duplication and subsequent neofunctionalization of members of some RP gene families as well as regulated post-translational modification of some RPs. For example, transcripts of Arabidopsis *RP* paralogs are regulated by environmental inputs including carbon, phosphate and metals (Hummel et al., 2012; Wang et al., 2013). *RP* transcript levels are also differentially regulated between cell types (e.g., (Mustroph et al., 2009)). Other sources of ribosome heterogeneity include N-terminal methionine removal, N-terminal acetylation, methylation of lysine and proline residues and phosphorylation of serine and threonine residues of RPs (Bailey-Serres and Freeling, 1990; Bailey-Serres et al., 1997; Szick-Miranda and Bailey-Serres, 2001; Williams et al., 2003; Chang et al., 2005; Carroll et al., 2008; Turkina et al., 2011; Hummel et al., 2012; Boex-Fontvieille et al., 2013; Carroll, 2013). The functional consequence of ribosome heterogeneity is largely unknown, but may provide for a complex regulatory network that impacts translation at the global or mRNA specific level.

Ribosomal protein phosphorylation

The most well studied phosphorylated RP is the 40S subunit protein RPS6, which is modified at multiple serine residues at its C-terminus. This region of the protein extends into the mRNA exit channel of the ribosome (Anger et al., 2013). In mammals, RPS6 is phosphorylated by p70S6k, whose activity is mediated by the mTOR kinase which also phosphorylates other proteins that regulate translation (Zoncu et al., 2011). The direct impact of RPS6 phosphorylation on mammalian translation may be negligible, but provides an effective readout for p70S6k activity in actively dividing cells, which promotes efficient translation of mRNAs with a polypyrimidine track at their 5' end (5'TOP). This feature is a characteristic of mRNAs encoding RPs and a number of core translation factors in mammals, but there is only limited evidence of functional 5'TOPs in plants (Jiménez-López et al., 2011). In Arabidopsis, RPS6 phosphorylation is promoted during the day (Turkina et al., 2011; Boex-Fontvieille et al., 2013) and in response to auxin and cytokinin (Turck et al., 2004), but is repressed by hypoxia (Chang et al., 2005). In monocots, RPS6 is phosphorylated in embryos during germination (Beltrán-Peña et al., 2002) and its phosphorylation rapidly declines following hypoxia, heat shock and singlet oxygen treatment (Williams et al., 2003; Khandal et al., 2009). The RPS6 kinase of Arabidopsis, AtS6K1, is important in regulating cell division and growth (Henriques et al., 2010; Shin et al., 2012). New data suggest a non-ribosomal function of RPS6 in epigenetic regulation of rDNA transcription in Arabidopsis (Kim et al., 2014c). Further clarification will require mutational analyses to determine if RPS6 phosphorylation is biologically significant or simply a hallmark of S6K activity in plants. Clearly there are many layers of S6K regulation yet to be explored and explained at the molecular level.

Most RPs assemble into pre-ribosomes in the nucleolus at a stoichiometry of one copy per ribosome. By contrast, the acidic proteins RPP1 and RPP2 complex in the cytoplasm with one another and then assemble onto the ribosome (Gonzalo and Reboud, 2003). The acidic proteins can be absent or present in multiple copies on the ribosome. Plant ribosomes possess a re-

lated third plant-specific protein called RPP3 (Szick et al., 1998). A source of heterogeneity of plant ribosomes is developmentally and environmentally regulated by modulation of RPP1, 2 and 3 levels and phosphorylation status (Bailey-Serres et al., 1997; Szick-Miranda and Bailey-Serres, 2001; Turkina et al., 2011; Boex-Fontvieille et al., 2013). The biological relevance of P-protein phosphorylation also deserves additional investigation.

Phosphoproteomic studies have also provided insight into the modulation of RP phosphorylation. In a non-targeted phosphoproteomics study, one or more isoforms of Arabidopsis RPs displayed distinct patterns of phosphorylation according to availability of CO₂ and light (Boex-Fontvieille et al., 2013). This included significant quantitative differences in phosphorylation state of RPS6A, RPS6B, RPL13D and RPL14A. Clearly further work on the functional consequences of RP phosphorylation is needed to better understand the complex interplay of signaling with protein modification and translational control in plants.

RACK1, A Ribosome Interacting Player

RACK1 is an interesting protein that is soluble, plasma-membrane-associated or ribosome-bound via interactions with the 18S rRNA and several 40S RPs (Valasek, 2012). RACK1 is stably associated with the 40S ribosomal subunit of Arabidopsis (Chang et al., 2005; Giavalisco et al., 2005; Carroll et al., 2008; Piques et al., 2009; Turkina et al., 2011; Hummel et al., 2012; Carroll, 2013), as in other eukaryotes. This protein has a seven bladed β -propeller domain that allows it to act as a scaffold, in a manner homologous to the heterotrimeric G protein G β subunit. RACK1 is involved in diverse signaling events as well as in translation (Gandin et al., 2013a). Its roles in mammals and yeast include recruitment of the protein kinase C that phosphorylates eIF6 to promote its release from the 60S. This event must take place before the joining of the 40S and 60S subunits can occur at the end of the initiation phase. Interestingly, Arabidopsis eIF6A/B appear to have both lost this protein kinase C phosphorylation site and only eIF6A retains a CK1 phosphorylation site (Guo et al., 2011a). RACK1 may have additional functions associated with protein synthesis, including recruitment of eIF3, regulation of RP synthesis, and promotion of the turnover of improperly folded nascent proteins (Gandin et al., 2013b). These distinct roles may involve different kinases recruited to the RACK1 scaffold. RACK1 was connected to the activity of miRNA in humans, *C. elegans* and Arabidopsis (Speth et al., 2013).

There are three *RACK1* paralogs in Arabidopsis. All three Arabidopsis *RACK1*s are detected in ribosomes (Chang et al., 2005; Giavalisco et al., 2005; Carroll et al., 2008; Piques et al., 2009; Turkina et al., 2011; Hummel et al., 2012; Carroll, 2013), functionally complement a CPC2/RACK mutant of yeast, and interact with eIF6 (Guo et al., 2011a). Interestingly, single and multiple *RACK1* mutants cause a variety of developmental abnormalities and enhance responsiveness to abscisic acid (ABA) (Guo et al., 2009; Guo et al., 2011b). Double mutants of *rack1a rack1b* are hypersensitive to anisomycin, an inhibitor of peptide elongation and display slightly reduced levels of 80S ribosomes under normal growth conditions and following ABA treatment (Guo and Chen, 2008). The multiple locations and functions of RACK1 makes interpretation of the double mutant phenotypes difficult. For exam-

ple, RACK1s participates in pre-miRNA processing via interaction with SERRATE, a partner of DICER-LIKE 1 and the nuclear-cap binding complex (CBP20/80) in Arabidopsis (Speth et al., 2013; Raczynska et al., 2014). This apparent nuclear role of RACK1 contrasts to its involvement with miRNAs in *C. elegans*, which is at the level of recruitment of the Ago2-miRNA silencing complex to polysomes for translational inhibition (Jannot et al., 2011). An unresolved question is whether ribosome-associated RACK1 functions in ALTERED MERISTEM PROGRAM 1 (AMP1)/AGO1/miRNA-mediated translation repression in Arabidopsis (Li et al., 2013b). In summary, plant RACK1 is ribosome-associated and functions in a conserved manner but also has extra-ribosomal functions that may be plant-specific.

Ribosomal protein mutants

Over 20 *RP* gene mutants of Arabidopsis have been characterized (reviewed by Byrne, 2009; Horiguchi et al., 2012; Roy and von Arnim, 2013). Often, single or multiple loss-of-function mutations for individual RPs result in embryo-lethality or pleiotropic developmental phenotypes affecting organ size or shape. These include asymmetric or pointed first leaves and reduced rosette size. In many cases *RP* mutants display phenotypes related to defects in auxin-mediated processes. At the mechanistic level, there are at least four possible causes of *RP* mutant phenotypes: (1) insufficient ribosomes affecting mRNAs equally or specifically (2) non-functional ribosomes, (3) a requirement for a distinct ribosome form for translation of specific mRNAs, or (4) an extra-ribosomal function of the protein (reviewed by Horiguchi et al., 2012).

Ribosome insufficiency, for example, could arise when reduced levels of a specific RP limits the biogenesis of a ribosomal subunit (Horiguchi et al., 2012). Phenotypes associated with *RP* mutants, such as smaller plant rosette size could be the consequence of reduced ribosome biogenesis. The synthesis of both subunits is tightly coordinated within the nucleolus, culminating in export of individual subunits to the cytoplasm. The co-expression of multiple *RP* gene paralogs in the same cell-type may limit ribosome insufficiency. However, reduction in a core ribosome component could limit overall ribosome numbers, rather than just the stoichiometry of an individual subunit. This could be important, since ribosome levels may be tightly regulated to meter the amount of energy consumed in translation at specific developmental states (*i.e.*, rapidly dividing versus differentiated cells) or under non-favorable environmental conditions. Indeed, the defects in cellular expansion of Arabidopsis leaves, a prominent phenotype of *RP* and ribosome biogenesis mutants, might be attributed to global reduction of ribosomes (Roy and von Arnim, 2013).

In the second scenario, *RP* mutants may not disrupt ribosome biogenesis but the subunits or complexes that form might be defective in overall activity or a specific function. This could occur if individual RPs have a specific role in translation. For example, studies of RPL24 indicated its importance in translation of mRNAs with small uORFs. RPL24 was also shown to be important in the intricate regulation of initiation of translation on CaMV 35S mRNA (reviewed by Roy and von Arnim, 2013). Levels of RPL4 and RPL5 were recognized as critical for translation of

uORF-containing mRNAs encoding proteins important for auxin responses (Rosado and Raikhel, 2010; Rosado et al., 2012).

The third possibility is that ribosome heterogeneity, due to the product of a specific *RP* gene paralog, is necessary for translation of a sub-set of transcripts. This last concept was detailed by Horiguchi et al. (2012), but definitive examples of *RP* gene paralogs of distinct function remain limited. A possible example is provided by the Arabidopsis *RPL10* paralogs, which appear to have non-redundant functions in male gametophyte development (Falcone Ferreyra et al., 2010; Falcone Ferreyra et al., 2013). However, it is necessary to rule out the possibility that RPL10 may have an extra-ribosomal function, as shown for a number of RPs in diverse eukaryotes (Warner and McIntosh, 2009; Xue and Barna, 2012).

Extra-ribosomal function has been suggested for several Arabidopsis RPs as well as RACK1. An example of an extra-ribosomal function is the proposed role played by RPS6 in the regulation of transcription of rDNA and some *RP* gene transcripts in Arabidopsis (Kim et al., 2014c). This function involves direct interaction of non-phosphorylated RPS6 with Histone Deacetylase 2B (HD2B), which suppresses rDNA transcription. It was hypothesized that phosphorylation of free RPS6 could reduce HD2B inhibition, thereby promoting rDNA transcription or processing. If regulated by TOR as proposed, this could place ribosome biogenesis and translational regulation under unified control (Kim et al., 2014c).

To move forward in our understanding of plant ribosomes there needs to be further consideration of whether limitation, excess, or modification of individual RPs modulates ribosome biogenesis or impacts translation of specific mRNAs. The use of gene silencing constructs equipped with inducible promoters or targeted gene editing as well as examinations limited to specific cells may aid in this challenge. Significant advancements would be generated by further structural analyses of 80S ribosomes or subunit-initiation/elongation factor complexes of plant ribosomes.

Ribosomes and energy for their synthesis

The biogenesis and activity of ribosomes requires a considerable investment in energy to power the synthesis of RPs and rRNAs. In rapidly dividing cells, the synthesis of rRNAs and RPs necessary for ribosome biogenesis may utilize more than half of all cellular energy, as each amino acid addition to a nascent peptide consumes at least four NTP molecules: (2 ATP for generating each amino-acyl tRNA and 2 GTP for each elongation event) (Figure 2). For ribosome biogenesis there is the additional energy outlay for rRNA synthesis. It therefore is not surprising that both ribosome biogenesis and global levels of translation are central to energy management (Piques et al., 2009; Pyl et al., 2012; Pal et al., 2013). Situations that limit ATP and GTP availability such as hypoxia, unanticipated darkness and extended nighttime limit *RP* mRNA translation in Arabidopsis (Branco-Price et al., 2008; Piques et al., 2009; Pal et al., 2013). In seedlings, cytosolic *RP* mRNAs account for ~10% of total cellular mRNA (Branco-Price et al., 2008). These transcripts are stable but rapidly translationally repressed during hypoxia due to sequestration in aggregates of oligouridylylate binding protein 1C




Figure 2. Overview of the steps of plant cytoplasmic translation elongation and termination cycles.

The elongating ribosome binds the incoming eEF1A•GTP•aa-tRNA in the A-site. If there is a match between the codon and anticodon of the tRNA, GTP hydrolysis occurs and eEF1A•GDP exits. Peptide bond formation occurs at the peptidyl transferase site; this reaction is mediated by the ribosome. eEF2•GTP binds and hydrolysis of GTP promotes translocation of the mRNA by three nucleotides, moving the now empty tRNA into the E-site, the newly elongated peptide•tRNA into the P-site, generating an empty A-site ready to accept another eEF1A•GTP•aa-tRNA. eEF1A•GDP requires the action of eEF1B to exchange GDP for GTP. eEF2 does not require a guanine exchange factor to acquire another GTP molecule. It is clear that each step of elongation is expensive in energy. Two GTP are required in the ribosome during elongation and each incoming eEF1A•GTP•aa-tRNA requires the functional equivalent of two ATP molecules to activate the amino acid and add it to the tRNA acceptor arm site. The AMP formed in this process requires two ATP to regenerate back to ATP; thus even though one ATP is consumed in the aminoacylation reaction, two ATP are ultimately consumed. The arrival of the termination codon in the A-site triggers the binding of the eRF1•eRF3•GTP complex into the A-site. Upon GTP hydrolysis, the eRF3•GDP is released. ABCE1 binds at the A-site to the remaining eRF1, promoting release of the polypeptide. Subsequent ATP hydrolysis by ABCE1 dissociates the ribosomal subunits, releasing ABCE1, eRF3, the deacetylated tRNA in the P-site and the mRNA. Note that the factors and ribosomal subunits are not to scale.

(UBP1C) (Sorenson and Bailey-Serres, 2014). This sequestration is quickly reversed upon reoxygenation, facilitating energy management during stress events.

RP mRNA translation in animals is largely mediated by mTOR activity, the presence of a 5'TOP, and the RNA binding protein La-related protein 1 (Thoreen et al., 2012; Tcherkezian et al., 2014). A characteristic of many other mammalian mRNAs that require mTOR activity is extensive secondary structure within their 5' leader sequences. In the case of higher plants, it is not clear yet if coordinate regulation of *RP* mRNA translation is TOR or 5'TOP regulated. Arabidopsis *RP* mRNAs typically have GC-rich untranslated leaders (Kawaguchi et al., 2004; Branco-Price et al.,

2005) and some have termini reminiscent of 5'TOPs. Whether or not a specific mRNA sequence or feature (i.e. structure) is involved, the manipulation of TOR levels in Arabidopsis influences overall levels of polysomes (Deprost et al., 2007). RNA binding proteins may also be a factor, as translation of a large number of *RP* transcripts was enhanced at subfreezing temperatures (4°C) by the RNA chaperone Cold Shock Protein 1, which has double-stranded RNA helicase activity (Juntawong et al., 2013). In sum, ribosome biogenesis is highly regulated due to the energy investment in both the transcription of rRNA and translation of *RP* mRNAs. Further study is needed to clarify the connections of these processes to TOR and S6K regulated energy sensing.

THE DRAMA OF INITIATION

The actors introduced in the previous section, initiation factors eIF1, eIF1A, eIF2, eIF3, and eIF5, come together on the 40S subunit to form the 43S PIC. This complex interacts with the mRNA and its associated factors (eIF4s) to form a 48S complex that is competent to search in the 5' to 3' direction for the initiation codon using the ATP-dependent scanning model proposed by Kozak (Kozak, 1978, 1980). The selection of the initiation codon depends upon its nucleotide sequence context and possibly other features in the mRNA. Upon selection of the initiation codon, a series of molecular events occurs that transforms the open scanning form of the 48S scanning complex to the closed form that is ready to engage the 60S subunit (Asano, 2014). This completes the initiation phase with an elongation competent 80S ribosome at the start of the desired ORF.

Formation of the 43S Pre-Initiation Complex

The 40S subunit, ternary complex (eIF2•GTP•Met-tRNA_i^{Met}), eIF1, eIF1A and eIF3 interact to form the 43S PIC. The ternary complex, eIF1, eIF1A, eIF3 and eIF5 can also form the MFC prior to interaction with the small ribosomal subunit. MFC formation in yeast promotes assembly and stability of the 43S PIC (Hinnebusch et al., 2007). The MFC has been shown *in vitro* to form in yeast (Asano et al., 2000), mammals (Sokabe et al., 2012) and plants (Dennis et al., 2009), suggesting that these protein interactions are part of a conserved mechanism.

Models for mRNA Binding of the eIF4s

The canonical model begins with mRNA interacting with the cap-binding complex through the eIF4E subunit, which is complexed with eIF4G. eIF4G then serves as the scaffold for assembly of eIF4A, PABP and eIF4B. This mRNA-factor complex is thought to then unwind the mRNA in an ATP-dependent manner for interaction with the 43S PIC. Since eIF4A is not a processive helicase, it is not clear exactly how an unwound region is initiated and maintained to enable ribosome binding. A more recent model of initiation (Aitken and Lorsch, 2012), places the eIF4s directly on the 43S PIC. In this scenario, the mRNA is subsequently recruited and unwound on the 40S subunit directly channeling the transcript. eIF4F/eIF4A could be bound to the mRNA (via the 5' cap or other RNA binding regions on eIF4G) as a "chaperone" which facilitates its interaction with 43S PIC associated factors such as eIF4B, eIF3 and/or eIF5. This model would explain a number of protein-protein interactions that are known to occur in yeast (e.g. eIF4G-eIF3, eIF4G-eIF5, eIF4B-40S subunit). However, some of these interactions have yet to be shown to occur in other eukaryotes (e.g. eIF4G-eIF5) and further biochemical analysis is needed to confirm the myriad of protein-protein interactions in the 43S PIC•mRNA/eIF4s complex and when/where they occur during the initiation process. Whichever model(s) proves to be true, the key aspect of the process is the relaxation of secondary structure in the 5' region of the transcript to facilitate ribosome binding, scanning, and eventually initiation codon recognition.

Start Site Recognition

Once the 43S PIC associates at the 5' end of the mRNA, scanning proceeds from 5' to 3' until a start codon is selected (Kozak, 1986; Asano, 2014). This is facilitated by binding of eIF1 to the "open" or scanning form of the 43S PIC that is stabilized by contacts with the N-terminal tail of eIF2β (Nanda et al., 2013). The zinc-binding domain in the C-terminus of eIF5 lies in close proximity to eIF1 and displaces the zinc-binding domain of eIF2β. This close proximity allows the N-terminal tail of eIF5 with its arginine finger required to interact with eIF2 and stimulate the GTPase activity of eIF2γ. The N-terminal tail of eIF5 prevents P_i release from the eIF2 ternary complex from this "open" conformation during the scanning process. When the scanning complex encounters the initiation codon in the suitable context in the P-site of the 40S subunit, the formation of the codon/anticodon base pair promotes full engagement of the Met-tRNA_i^{Met}. This disrupts eIF1 interaction with N-terminal tail of eIF2β and results in eIF1 release. The eIF2β N-terminal tail then interacts with the C-terminal domain of eIF5. The N-terminal domain of eIF5 is then able to interact with the C-terminal tail of eIF1A, which promotes the release of P_i from the ternary complex resulting in scanning arrest at a suitable initiation codon and conversion to a "closed" PIC (Asano, 2014; Hinnebusch, 2014; Saini et al., 2014).

A suitable initiation codon context is an A residue and to a lesser extent a guanine (G) residue at position -3 and a G residue at position +4 relative to the A₁UG codon of the mRNA. It is thought that nucleotides surrounding the start codon help to engage the closed PIC conformation as the AUG codon is recognized (Hinnebusch and Lorsch, 2012; Asano, 2014; Hinnebusch, 2014). The A at position -3 corresponds to the first nucleotide of the E-site of the 40S subunit and is occupied by eIF2 as the codon-anticodon interaction is established in the P-site. In Arabidopsis plants exposed to dehydration stress, transcripts that were better associated with polysomes during the stress were enriched in A nucleotides just upstream of the start codon (Kawaguchi and Bailey-Serres, 2005). A recent study of Arabidopsis 5'UTRs further showed that the positions of A residues in the -1 to -5 region from the AUG were highly correlated with translational efficiency and uracil (U) residues in the same region were negatively correlated (Kim et al., 2014b). These results suggest that the region 5' to the AUG in Arabidopsis strongly influences translational efficiency in plants (Kim et al., 2014b). mRNAs with a 5'UTR that was shorter than average (125 nt) and had low potential for secondary structure formation had higher levels of ribosome occupancy. Consistently, the G+C nucleotide content of the 5'UTR was inversely correlated with translational activity during a variety of environmental stresses including dehydration, hypoxia and darkness (Branco-Price et al., 2005; Kawaguchi and Bailey-Serres, 2005; Juntawong and Bailey-Serres, 2012).

Non-AUG codons

There are also rare examples of initiation at non-AUG codons on Arabidopsis mRNAs, including *AGAMOUS*, *FCA* and *POLyG*. The latter encodes a RNA polymerase targeted to the plastid or mitochondrion based on the AUG selected (Riechmann et al.,

1999; Wamboldt et al., 2009; Simpson et al., 2010). Mutational studies that evaluated the ramifications of initiation codon context, secondary structure, 5'UTR length and presence of uORFs on the rate of initiation on the protein coding ORFs of plants have provided insight into use of an CUG triplet as a functional initiation codon for the *FCA* transcript (Simpson et al., 2010). Advances in nucleotide-level resolution of ribosome position (Liu et al., 2013; Juntawong et al., 2014) and *in vivo* secondary structure (Ding et al., 2014) are likely to yield additional examples of non-AUG initiation and information on the surrounding mRNA region that will provide new insight into flexibility in start site selection in plants.

Reinitiation involving uORFs

The presence of one or more short uORFs that precedes a mORF presents a special situation to the scanning ribosome. Based on the characterization of *GCN4* mRNA translation in yeast, the length of the uORFs, the spacing between the uORFs and the mORF, and specific mRNA sequence features contribute to the subtle regulation of subsequent reinitiation events that determine amount of *GCN4* synthesized (Valasek, 2012). In plants, the amino acid sequence of the uORF peptide can also contribute to the translational regulation (Rahmani et al., 2009; Jorgensen and Dorantes-Acosta, 2012; Roy and von Arnim, 2013; von Arnim et al., 2014). In the case of mRNAs with multiple ORFs (polycistronic), ribosomes will initiate in the normal manner at the first AUG in a suitable context and elongation will proceed. When the termination codon is encountered, the termination process that dissociates the ribosome subunits is likely to occur. If the uORF is short there may be lingering association of eIF3 with the 40S subunit and the reassembly of the MFC may occur (Asano, 2014). In plants, this process is enhanced when the eIF3h subunit is present and phosphorylated by S6K in a TOR kinase-dependent manner (Schepetilnikov et al., 2013).

Assembly of the 80S ribosome, the final scene of initiation

Upon formation of the “closed” PIC, mammalian eIF5 is released in complex with eIF2•GDP. eIF5's role at this point is as a GDP dissociation inhibitor for eIF2•GDP until eIF2B is able to stimulate the replacement of GDP with GTP. The function of mammalian eIF2B is crucial as it allows eIF2•GDP to exchange for GTP and acquire a new Met-tRNA^{Met} for participation in another round of initiation (Jennings and Pavitt, 2010; Jennings et al., 2013); however, as described above it is not clear what the role and importance of plant eIF2B are at this time. The release of eIF5/eIF2•GDP opens the surface of the 40S for binding of the 60S subunit, whereas eIF5B•GTP facilitates the 60S ribosome joining through interactions with the C-terminal tail of eIF1A, and then eIF5B•GDP readily dissociates from the complex.

eIF1A plays a central role in the entire process of initiation. First, through its interactions with the eIF1 C-terminal tail to stabilize the “open” PIC by preventing Met-tRNA^{Met} to fully engage in the P-site. Second, upon arrival at the correct AUG, the C-terminal tail of eIF1A is displaced and interacts with eIF5 to promote

the release of P_i generated by hydrolysis of eIF2•GTP by eIF5. Lastly, eIF1A facilitates the formation of the 80S ribosome and is the last initiation factor to exit after eIF5B•GTP. Thus, the 80S ribosome positioned at the correct initiation codon is now ready to move on to the next act, elongation.

ACT 2: ELONGATION

Translational elongation is an evolutionarily conserved progression of ribosome catalyzed polypeptide formation through mRNA decoding (see Figure 2 and Table 2). Once the subunit joining is complete with the Met-tRNA^{Met} in the P-site of the ribosome, the second codon in the A-site awaits interaction with the anticodon of an aminoacyl (aa)-tRNA coupled to eEF1A•GTP (Dever and Green, 2012). Appropriate codon-anticodon interactions at the A-site will stimulate the peptidyl transferase reaction that generates a peptide bond between the Met-RNA_i^{Met} and the aa-tRNA, leaving a deacylated tRNA^{Met} in the P-site. The subsequent translocation of the mRNA by one codon shifts the peptidyl-tRNA to the P-site and the deacylated tRNA to the E-site, freeing the A-site for the next appropriate aa-tRNA and continuation of the cycle (Dever and Green, 2012; Doerfel et al., 2013). Translocation is facilitated by eEF2•GTP binding and GTP hydrolysis.

The principal cast for this process includes the aa-tRNAs, eukaryotic elongation factor (eEF)1A (homolog of bacterial EF-Tu), eEF1B (homolog of bacterial EF-Ts), eEF2 (homolog of bacterial EF-G) and the ribosome. The role of eEF5 (nee eIF5A) in the elongation process is emerging, and like its prokaryotic homolog (EF-P) appears to involve elongation of amino acid sequences enriched in runs of proline and/or glycine (Doerfel et al., 2013; Gutierrez et al., 2013; Ude et al., 2013). Aminoacyl tRNA synthetases (aa-synthetases) participate backstage. These enzymes are encoded by a large family of nuclear genes in Arabidopsis that couple cognate tRNAs to their amino acids to form acetylated tRNA (aa-tRNA) in an ATP dependent reaction.

THE ACTORS IN ELONGATION

eEF1A

eEF1A is the ortholog of bacterial elongation factor-Tu (EF-Tu). This factor forms a ternary complex with GTP and an aa-tRNA, which it delivers to the peptidyl transferase center when the corresponding codon is present in the A-site. Initial loose binding is followed by a recognition event that involves the hydrolysis of GTP and structural rearrangements of the tRNA, eEF1A and the ribosome. eEF1A•GDP is then released for recycling by eEF1B, a complex with GEF activity that recovers eEF1A•GTP (described below).

eEF1A is a highly abundant protein that may constitute up to 1% of the total protein in a cell. The protein is encoded by four paralogs in Arabidopsis. Seed endosperm of the maize *opaque 2* mutant has increased levels of eEF1A from multiple genes and is associated with improved lysine content (Lopez-Valenzuela et al., 2003; Lopez-Valenzuela et al., 2004). Interestingly, eEF1A has functions and interactions outside of its role in translation includ-

ing association with cytoskeleton (which may reflect an association of the translation process with cytoskeleton anchors), nuclear export, proteolysis, apoptosis and viral propagation (Browning, 1996; Sasikumar et al., 2012). This factor is also known to participate in processes including export of tRNAs from the nucleus and the targeting of damaged and misfolded proteins to the proteasome (Sasikumar et al., 2012). EF1A is also reported to have interactions with the tombusvirus replication complex and the 3' tRNA-like structure of turnip yellow mosaic virus (Matsuda et al., 2004; Li et al., 2009).

eEF1B

In prokaryotes, EF-Tu•GDP cannot recycle the GDP for GTP without assistance from EF-Ts. Similarly, eEF1A requires an exchange factor, eEF1B. In contrast to the single polypeptide EF-Ts, eEF1B is a complex of proteins that varies in complexity from eukaryote to eukaryote. eEF1B has three components in plants, eEF1B α , eEF1B β and eEF1B γ (Table 2), two in yeast and three in mammals (Sasikumar et al., 2012). In some organisms eEF1B also includes a valyl tRNA synthetase. Very little is known specifically about eEF1B from plants other than it has been purified from wheat germ (Lauer et al., 1984) and was shown to play a role in viral replication (Sasvari et al., 2011; Hwang et al., 2013), as have eEF1A (Matsuda et al., 2004; Li and Nagy, 2011) and cap-binding complex subunits (Wang and Krishnaswamy, 2012).

Both eEF1A and eEF1B are post-translationally modified by phosphorylation involving several kinases and by methylation, which may influence various activities such as interaction with actin (Lopez-Valenzuela et al., 2003). Presumably these modifications reflect highly complex mechanisms of regulation in eukaryotes (Le Sourd et al., 2006; Sasikumar et al., 2012). Phosphorylation of elongation factors under photosynthetic control was not reported for Arabidopsis (Boex-Fontvieille et al., 2013).

eEF2

eEF2 is the functional equivalent to EF-G of prokaryotes. Peptide bond formation occurs rapidly following acceptance of the aa-tRNA into the A-site of the peptidyl transferase center within the large ribosomal subunit. This region is largely comprised of rRNA and is highly conserved between prokaryotic and eukaryotic ribosomes, indicating that the process of peptide bond formation is quite ancient (Dever and Green, 2012). After the peptide bond is formed in the peptidyl transferase reaction catalyzed by the ribosome, it is necessary to move the now uncharged tRNA from the P-site into the E-site, freeing the A-site for the next incoming aa-tRNA•eEF1A•GTP. A GTP•eEF2 complex binds to the ribosome and its GTP hydrolysis promotes the movement of the mRNA•tRNA•tRNA hybrid forward by three nucleotides, coinciding with the movement of the P-site deacylated tRNA to the E-site and ejection of the deacylated-tRNA from the E-site.

eEF2 is post-translationally modified at a conserved histidine residue to diphthamide. This modification makes eEF2 the target for ADP-ribosylation by diphtheria-like toxins (Ortiz et al., 2006; Zhang et al., 2008a). The biological significance of this unusual modifica-

tion is unknown despite its conservation across all eukaryotes and *Archaea*. Wheat eEF2 was shown to have this modification as evidenced by ADP-ribosylation by diphtheria toxin (Lauer et al., 1984). eEF2 is also a substrate for phosphorylation by the Ca²⁺/calmodulin-dependent eEF2 kinase (eEF2K), which reduces eEF2 association with the ribosome. The phosphorylation site in mammals is a conserved threonine residue (T56). The mammalian AMP kinase and mTOR-signaling pathways converge to inhibit Ca²⁺-dependent eEF2K activity, thereby limiting translational elongation under nutrient limiting conditions (Leprivier et al., 2013). Conversely, hypoxia in mammals promotes eEF2K phosphorylation and accumulation. Because eEF2 phosphorylation is Ca²⁺-regulated, it is thought to regionally fine-tune protein synthesis, such as in dendrites of activated neurons (Heise et al., 2014). When purified wheat germ eEF2 was phosphorylated with a mammalian Ca²⁺/calmodulin-dependent kinase, its activity in the *in vitro* translation system was reduced (Smailov et al., 1993). Although a plant eEF2 kinase has not been recognized, phosphoproteomic analyses focused on translation factors detected P-Ser558 of Arabidopsis eEF2 (Guillaume Tcherkez, personal communication). This site is conserved relative to Ser595 of mammals. Interestingly, phosphorylation of Ser595 by cyclin A in mammals promotes phosphorylation of Thr56 by eEF2K (Hizli et al., 2013). The finding that the *At*eEF2 (*LOS1*) is important for protein synthesis at low temperatures (Guo et al., 2002) hints that regulation of eEF2 activity is relevant to cold acclimation and likely other stress conditions.

eEF5 (nee eIF5A/eIF4D)

eEF5 was formerly known as eIF5A/eIF4D due to its initial report as a stimulator of Met-puromycin synthesis *in vitro*, a model assay for initiation. eEF5 was later recognized as a facilitator of elongation (Nanda et al., 2009) and confirmed as the functional and structural equivalent of elongation factor P (EF-P) of eubacteria. Both EF-P and eEF5 are involved in the efficient elongation of proteins with runs of proline or glycine residues (Doerfel et al., 2013; Gutierrez et al., 2013; Ude et al., 2013). Why certain combinations of amino acids pose difficulties during elongation is not fully understood, but at least for the imino acid proline (lacking the hydrogen at the amino group) it may be due to structural constraints introduced in the peptide backbone by its presence.

eEF5 has features that distinguish it from EF-P and is truncated on its C-terminus relative to EF-P, which limits its contacts with the 60S subunit (Gutierrez et al., 2013). eEF5 is the only eukaryotic protein known to be post-translationally modified with hypusine, a modification derived from spermidine, which is required for its activity. Similarly, prokaryotic EF-P is modified by β -lysylation, also a spermidine derivative (Allen and Frank, 2007; Bullwinkle et al., 2013). The hypusine/ β -lysine modification is postulated to help eEF5 to engage the ribosome and bring proline residues into closer proximity in the peptidyl transferase center for peptide bond formation (Gutierrez et al., 2013). Given this proposed function, it may be informative to evaluate the density of ribosomes in regions of mRNAs enriched in proline codons in genotypes that vary in eEF5 abundance and hypusination. eEF5 also appears to be modified by phosphorylation in the light/dark transition (Boex-Fontvieille et al., 2013).

A number of studies of plant eEF5 have indicated a role in stress responses (abiotic, pathogen, iron deficiency), growth and development (Wang et al., 2003; Chou et al., 2004; Hopkins et al., 2008; Ma et al., 2010; Lan and Schmidt, 2011; Wang et al., 2012). eEF5 was found to be associated with eEF2 in pumpkin phloem (Ma et al., 2010). In Arabidopsis, an *eEF5* paralog (reported as *eIF5A-2*) is necessary for cytokinin-mediated promotion of protoxylem development in seedling roots through genetic interaction with Cytokinin Response 1 (CRE1), a histidine kinase that binds cytokinin and the phosphotransferase AHP6, that negatively regulates signaling by cytokinin (Ren et al., 2013). Given that these processes appear unrelated to translation, eEF5 may have additional biological function(s) in plants; alternatively, these may represent downstream outcomes due to translation defects involving proteins with poly-prolyl or glycyl residues whose translation may depend upon this factor.

Compared to initiation in plant translation, there has been less work on the elongation process and its factors. Whether there will be aspects of elongation or its control that are specific to plants await further discovery.

ACT 3, THE FINALE: FACTORS AND EVENTS OF TERMINATION

Elongation ends when translocation places one of the three stop codons (UAA, UGA, or UAG) into the A-site of the ribosome. This initiates the termination phase, which ends with disengagement of the peptide from the ribosome (Dever and Green, 2012; Jackson et al., 2012). There are two eukaryotic release factors (eRF1 and eRF3, see Table 2 and Fig. 2) in plants, the same as in mammals. The fate of the translation complex upon termination is still lacking in molecular details. Termination is usually followed by ribosome release, but may be followed by reinitiation of translation after a short coding sequence (i.e., a uORF) (Roy and von Arnim, 2013). In the special case of premature termination at a nonsense codon (a termination codon 5' of an EJC), degradation of the transcript occurs via the NMD pathway (see below).

Prior to the final “act”, a complex of release factors (RF) and GTP must form in the cytosol in preparation for interaction with the ribosome (see Table 2). eRF3•GTP binds to eRF1 which acts as a GTP dissociation inhibitor. When the elongating ribosome arrives at a stop codon on the mRNA, the eRF3•GTP•eRF1 complex is recruited to the A-site, preventing further entry of eEF1A•aa-tRNA complexes. Unlike prokaryotic termination factors that have specificity for one or more termination codons, the eukaryotic ternary complex of eRF3•GTP•eRF1 recognizes all three stop codons (UAA, UGA, UAG) by a little known mechanism (reviewed in Dever and Green, 2012; Jackson et al., 2012).

THE ACTORS IN TERMINATION

eRF1 and eRF3

eRF1 is evolutionarily related to bacterial RF1 and RF2, class 1 RFs. The structure of these proteins resembles a tRNA, allowing the N-terminal domain of the RF to dock in the A-site and directly

interact with the stop codon (Dever and Green, 2012; Jackson et al., 2012). The high fidelity of this binding coupled with the GTPase activity of eRF3 promotes the peptidyl tRNA hydrolysis necessary to release the polypeptide from the P-site and from the ribosome (Dever and Green, 2012; Jackson et al., 2012). At the structural level, the N-terminal region of eRF1 binds the stop codon, the middle domain enters the peptidyl transferase center where it promotes the hydrolytic release of the polypeptide, whereas the C-terminal region interacts with eRF3. eRF3's GTPase activity is necessary both to increase the rate of peptide hydrolysis by eRF1 and the efficiency of termination (Dever and Green, 2012; Jackson et al., 2012). The structure of the eRF3•GTP•eRF1 ternary complex on the ribosome was determined, revealing a number of features that suggest it has a very similar GTPase activation mechanism to the prokaryotic aa-tRNA•EF•Tu•GTP complex (des Georges et al., 2014). eRF1 is retained after termination and is important for recycling of the ribosome by recruiting the ABCE1/RIL1 protein (see below and Fig. 2) which functions with eIF6 in the dissociation of the ribosome into subunits for recycling (Pisarev et al., 2010). Yeast eRF1 has been shown to have additional functions that affect the cytoskeleton and cell cycle (Valouev et al., 2002) and appears to “moonlight” as observed for the eEFs (Le Sourd et al., 2006; Sasikumar et al., 2012)

Arabidopsis has three *AteRF1* genes that encode functional RFs (Chapman and Brown, 2004). The overexpression of *At-eRF1-1* resulted in the silencing of *AteRF1-1* and to some extent *AteRF1-2* and *AteRF1-3* causing a phenotype known as *broomhead* (altered spacing between inflorescence stems cause a broom-like appearance) and is associated with perturbations in cell division and cell elongation (Petsch et al., 2005). *AteRF1-2* mRNA levels are induced by high glucose levels and *AteRF1-2* overexpression lines display increased glucose-mediated repression of germination (Zhou et al., 2010b). These genotypes are also hypersensitive to paclobutrazol, an inhibitor of gibberellin biosynthesis, as well as abscisic acid. Consistently, T-DNA insertion mutants of *AteRF1-2* showed resistance to gibberellin synthesis inhibitors during germination. It is not yet clear if the role *AteRF1-2* plays in glucose sensing or phytohormone responses reflects its role in termination or some other cellular role. It is important to determine if the *broomhead* and other phenotypes associated with eRF1 mutants in Arabidopsis are related to translation or other processes. Interestingly, a mutant (*Or*) that produces an orange color in cauliflower heads (inflorescence meristems) due to an increase in β -carotene and encodes, a protein shown to interact with eRF1-2. The *Or* mutant displays altered petiole elongation and other developmental alterations suggesting a role for termination in developmental programs (Zhou et al., 2010c). Much more needs to be learned about plant termination and its actors.

NONSENSE MEDIATED mRNA DECAY: CURTAINS FOR SOME mRNAs

In special cases, termination can trigger mRNA decay (Belostotsky and Sieburth, 2009). This mechanism, termed nonsense mediated decay (NMD), provides quality control of mRNAs as they transit from the nucleus to active translation complexes. The

process provides continuity between the nuclear process of intron splicing and cytoplasmic translation. A key feature in the process is the EJC, which is deposited just 5' of exon-exon junctions following splicing, and serves as a talisman in the pioneering (first) round of translation of an mRNA. Following translational initiation, the elongating ribosome is thought to displace many of the RNA binding proteins bound to the mRNA as it translocates from codon to codon. If an EJC lies 3' of a stop codon or the transcript has an unusually long 3' UTR (>300 bp), then eRF3, responsible for termination and release of the nascent polypeptide, associates with UPF1 (helicase up-frameshift 1), a protein needed to initiate NMD. Two other proteins required for this process, UPF2 and UPF3, bind to the EJC after splicing. mRNAs with an EJC 3' of the termination codon properly position UPF1-3 such that the destruction of the mRNA is triggered (Chang et al., 2007). NMD functions similarly in plants based on the presence of orthologs of NMD components and evidence of NMD coupled to the turnover of mRNAs with premature termination codons (Kerényi et al., 2008; Reddy et al., 2013). The targeting of alternatively spliced transcripts with premature termination codons for NMD provides an example of a mechanism coupled to translation that modulates mRNA abundance in response to environmental cues (Kalyna et al., 2012). Thus the half-life of an mRNA can be intimately entwined with its translation.

The many factors, complexes and processes involved in the temporal and spatial regulation of mRNA decay in plants have received limited attention until quite recently (Maldonado-Bonilla, 2014). It is important to understand the connection between translation and decay processes, including miRNA-mediated translational inhibition and mRNA turnover (Li et al., 2013b; Rogers and Chen, 2013).

RECYCLING: IS THERE AN ENCORE?

The emerging view is that termination is followed by "recycling", efficient reuse of the ribosome. At this point in the translational process the 80S ribosome, mRNA and tRNA-polypeptide chain are still coupled. This requires that the two subunits of the ribosome dissociate and eRF1 as well as the deacylated tRNA be released. In prokaryotes ribosome recycling is promoted by EF-G and a dedicated ribosome recycling factor (RRF), present only in prokaryotes. Currently, it is thought that an essential protein, ATP-binding cassette E (ABCE1)/RNAse L INHIBITOR 1 (RLI1), conserved in eukaryotes and *Archaea*, promotes polypeptide release and ribosome recycling (Pisarev et al., 2010) in a process that requires ATP hydrolysis (Dever and Green, 2012; Jackson et al., 2012). Recent structural studies show that following eRF3•GDP release, ABCE1 binds to eRF1 and within the ribosome (Preis et al., 2014). This binding is associated with a dramatic conformational change in eRF3 that positions its central domain in the peptidyl transferase center, where it catalyzes the release of the polypeptide.

Other factors may be important in recycling of ribosomes on cytosolic mRNAs of eukaryotes. First, the proximity between the 3' and 5' ends of the message, fostered by the presumed interaction between PABP and eIF4s (Jackson et al., 2010; Valasek, 2012), may enable loosely associated 40S subunits to reform a PIC and recommence the initiation phase. There is, however,

some debate about the importance of PABP/eIF4G interactions in the closed-loop mRNA model (Afonina et al., 2014). Studies with yeast and mammals (Dever and Green, 2012) point to a role of ABCE1/RLI1 in recruiting the MFC to the 40S subunit once the ribosome is dissociated. Interestingly, there is evidence from mammals that if eIF3, eIF1, eIF1A, and eIF2•tRNA^{Met} remain associated with the 40S subunit after termination then bidirectional scanning by the 40S or 80S complex occurs. Such a scenario would enable initiation at AUGs of downstream or upstream open reading frames (i.e., uORFs) that precede the ORF encoding the functional protein (Skabkin et al., 2013). The clever use of mimicry of tRNA shapes in some plant viral 3' UTRs serves to recruit or recycle ribosomes, suggesting that recycling may be a common cellular event.

THE ACTORS IN RECYCLING

In addition to eRF1, the ABC-type ATPase ABCE1 is a key player in ribosome recycling. The *Arabidopsis* genome encodes two *ABCE1* genes, which are characterized by an N-terminal Fe-S cluster and two nucleotide binding domains. A point mutation in a *ABCE1/RLI1* ortholog in *Cardamine hirsuta*, a relative of *Arabidopsis*, converts the highly lobed leaf into a simple leaf and causes other downstream phenotypes (Kougioumoutzi et al., 2013). These findings suggest that ABCE1 plays an important role in numerous cellular developmental processes. Developmental dysfunctions including alterations in auxin homeostasis are quite frequent for mutants affecting ribosome biogenesis as described above. But caution is needed in interpreting these results, as it remains to be shown if ABCE1 has other roles or the efficiency of ribosome recycling is critical for development. Other proteins that act in the recycling of the translational apparatus in segue from termination to a new initiation event are unknown, with the exception of eIF6 which promotes subunit dissociation.

FUTURE PROSPECTS

The ease of isolation of mRNA and methods for global analyses of mRNA abundance has resulted in intense research on gene regulation in plants and other eukaryotes. Although, transcriptional regulation is frequently presumed to be the default mechanism that modulates steady-state transcript abundance, regulation that occurs at post-transcriptional levels including the processes that determine mRNA maturation, transport, stabilization, turnover and, in particular, translation have become increasingly apparent. In plants, these processes all contribute to dynamics in quantity, location and function of the gene product and are not readily discerned from steady-state transcript data. Technologies that enable the isolation of mRNAs associated with polysomes, such as translating ribosome affinity purification (TRAP) have helped to illuminate translational regulation of individual transcripts, particularly in *Arabidopsis* (Zanetti et al., 2005). Resolution of dynamics in mRNA translation will be enhanced by the ability to identify the position and frequency of ribosomes as they transit gene transcripts. This "ribosome profiling" strategy has been applied to examine changes in ribosome distribution along *Arabidopsis*

mRNAs in seedlings upon illumination-triggering photomorphogenesis and during hypoxia (Liu et al., 2013; Juntawong et al., 2014). Translational dynamics occur in response to environmental stress, metabolites, and over the course of development (reviewed by Roy and von Arnim, 2013) and as a means for overall regulation of cellular energy over the diurnal cycle (Pal et al., 2013; Sulpice et al., 2014). Translational regulation may also be important in the tolerance of polyploidy in plants, as a comparison of total and polysomal mRNAs in the allopolyploid *Glycine dolichocarpa* indicated that selective translation contributes to dominance of expression of specific homoeologous genes as well as physically linked genes (Coate et al., 2014).

Further studies of the translational apparatus is needed, including the soluble factors, ribosomes and the cadre of RNA binding proteins that act as stagehands to fine-tune translational regulation. The use of genetic approaches to dissect the roles of the apparatus will most likely benefit from inducible constructs that reduce endogenous transcript levels or produce isoforms with specific features at controlled levels. In addition to a focus on endogenous mRNAs, the study of the performances of plant viruses in translation can be helpful. In the end, the data generated over the next decade will provide key insights, but is likely to also raise more enigmas. There are currently many questions about plant translation that are unanswered:

- What is the role of eIF2 phosphorylation by GCN2 in regulating translation and are there other eIF2 kinases that might regulate global levels of translation?
- Is there a plant version of eIF2B and what is its function?
- What is the role of the plant-specific eIFiso4F and why did it evolve?
- Are there other specific differences in plant initiation complexes compared to other eukaryotes?
- What are the molecular interactions of the initiation factors with the ribosomes? Do they differ from other eukaryotes?
- What factors besides eIF3h and the ribosome are important in uORF translation?
- Is ribosome heterogeneity of biological significance?
- Do specific ribosomal proteins regulate translation of individual gene transcripts or cohorts of mRNAs during development or in response to environmental cues?
- What is the role of nutrient availability and TOR in ribosome biogenesis (including rRNA synthesis, *RP* mRNA transcription and translation), and other processes of translation?
- What are the signals from chloroplast to nucleus that regulate coordinated synthesis of nuclear encoded photosynthetic proteins?
- What RNA sequences or structures and RNA binding proteins contribute to differential translation, targeting, stability and trafficking of mRNAs?
- What mechanisms sequester mRNAs into untranslatable pools, and how do they regain their ribosome loading?
- What are the levels of interaction between chromatin, transcription, nuclear processing, translation, and mRNA turnover involving NMD, miRNA or general decay mechanisms?

As these questions are answered we will acquire a greater appreciation of the multi-dimensional and integrated performance within the cell nucleus and cytoplasm that culminate in the highly regulated “action drama” of protein synthesis in plants.

ACKNOWLEDGMENTS

We thank members of the Browning and Bailey-Serres laboratories who for more than twenty years have contributed to the advancement in understanding the importance of translational regulation and mechanisms of translation in plants. Particularly, we thank Albrecht von Arnim, Maureen Hummel and Angel Syrett for comments on this manuscript. We also thank Angel Syrett for the artwork. Our research related to translation is supported by the National Science Foundation (MCB1052530 and S-0000335) to K.S.B. and (IOS-1121626 and IOS-1238243) to J.B.-S.

REFERENCES

- Aalto, M.K., Helenius, E., Kariola, T., Pennanen, V., Heino, P., Hörak, H., Puzörjova, I., Kollist, H., and Palva, E.T. (2012). ERD15—an attenuator of plant ABA responses and stomatal aperture. *Plant Sci.* **182**, 19-28.
- Abramson, R.D., Browning, K.S., Dever, T.E., Lawson, T.G., Thach, R.E., Ravel, J.M., and Merrick, W.C. (1988). Initiation factors that bind mRNA: a comparison of mammalian factors with wheat germ factors. *J. Biol. Chem.* **263**, 5462-5467.
- Afonina, Z.A., Myasnikov, A.G., Shirokov, V.A., Klaholz, B.P., and Spirin, A.S. (2014). Formation of circular polyribosomes on eukaryotic mRNA without cap-structure and poly(A)-tail: a cryo electron tomography study. *Nuc. Acids Res.* **42**, 9461-9469.
- Aitken, C.E., and Lorsch, J.R. (2012). A mechanistic overview of translation initiation in eukaryotes. *Nat. Struct. Mol. Biol.* **19**, 568-576.
- Allen, G.S., and Frank, J. (2007). Structural insights on the translation initiation complex: ghosts of a universal initiation complex. *Mol. Microbiol.* **63**, 941-950.
- Allen, M.L., Metz, A.M., Timmer, R.T., Rhoads, R.E., and Browning, K.S. (1992). Isolation and sequence of the cDNAs encoding the subunits of the isozyme form of wheat protein synthesis initiation factor 4F. *J. Biol. Chem.* **267**, 23232-23236.
- Altmann, M., Müller, P.P., Wittmer, B., Ruchti, F., Lanker, S., and Trachsel, H. (1993). A *Saccharomyces cerevisiae* homologue of mammalian translation initiation factor 4B contributes to RNA helicase activity. *EMBO J.* **12**, 3997-4003.
- Andreou, A.Z., and Klostermeier, D. (2013). The DEAD-box helicase eIF4A: paradigm or the odd one out? *RNA Biol.* **10**, 19-32.
- Andreou, A.Z., and Klostermeier, D. (2014). eIF4B and eIF4G jointly stimulate eIF4A ATPase and unwinding activities by modulation of the eIF4A conformational cycle. *J. Mol. Biol.* **426**, 51-61.
- Anger, A.M., Armache, J.P., Berninghausen, O., Habeck, M., Subklewe, M., Wilson, D.N., and Beckmann, R. (2013). Structures of the human and *Drosophila* 80S ribosome. *Nature* **497**, 80-85.
- Armache, J.P., Jarasch, A., Anger, A.M., Villa, E., Becker, T., Bhushan, S., Jossinet, F., Habeck, M., Dindar, G., Franckenberg, S., Marquez, V., Mielke, T., Thomm, M., Berninghausen, O., Beatrix, B., Söding, J., Westhof, E., Wilson, D.N., and Beckmann, R. (2010a). Localization of eukaryote-specific ribosomal proteins in a 5.5-Å cryo-EM map of the 80S eukaryotic ribosome. *Proc. Natl. Acad. Sci. USA* **107**, 19754-19759.
- Armache, J.P., Jarasch, A., Anger, A.M., Villa, E., Becker, T., Bhushan, S., Jossinet, F., Habeck, M., Dindar, G., Franckenberg, S., Marquez, V., Mielke, T., Thomm, M., Berninghausen, O., Beatrix, B., Söding, J., Westhof, E., Wilson, D.N., and Beckmann, R. (2010b). Cryo-EM structure and rRNA model of a translating eukaryotic 80S ribosome at 5.5-Å resolution. *Proc. Natl. Acad. Sci. USA* **107**, 19748-19753.
- Arribere, J.A., and Gilbert, W.V. (2013). Roles for transcript leaders in translation and mRNA decay revealed by transcript leader sequencing.

- Genome Res. **23**, 977-987.
- Asano, K.** (2014). Why is start codon selection so precise in eukaryotes? *Translation* **2**, e28387.
- Asano, K., Clayton, J., Shalev, A., and Hinnebusch, A.G.** (2000). A multifactor complex of eukaryotic initiation factors, eIF1, eIF2, eIF3, eIF5, and initiator tRNA^{Met} is an important translation initiation intermediate *in vivo*. *Genes & Dev.* **14**, 2534-2546.
- Bailey-Serres, J.** (1999). Selective translation of cytoplasmic mRNAs in plants. *Trends Plant Sci.* **4**, 142-148.
- Bailey-Serres, J., and Freeling, M.** (1990). Hypoxic stress-induced changes in ribosomes of maize seedling roots. *Plant Physiol.* **94**, 1237-1243.
- Bailey-Serres, J., Sorenson, R., and Juntawong, P.** (2009). Getting the message across: cytoplasmic ribonucleoprotein complexes. *Trends Plant Sci.* **14**, 443-453.
- Bailey-Serres, J., Vangala, S., Szick, K., and Lee, C.H.** (1997). Acidic phosphoprotein complex of the 60S ribosomal subunit of maize seedling roots. Components and changes in response to flooding. *Plant Physiol.* **114**, 1293-1305.
- Balasta, M.L., Carberry, S.E., Friedland, D.E., Perez, R.A., and Goss, D.J.** (1993). Characterization of the ATP-dependent binding of wheat germ protein synthesis initiation factors eIF-(iso)4F and eIF-4A to mRNA. *J. Biol. Chem.* **268**, 18599-18603.
- Barakat, A., Szick-Miranda, K., Chang, I.F., Guyot, R., Blanc, G., Cooke, R., Delseny, M., and Bailey-Serres, J.** (2001). The organization of cytoplasmic ribosomal protein genes in the *Arabidopsis* genome. *Plant Physiol.* **127**, 398-415.
- Belostotsky, D.A.** (2003). Unexpected complexity of poly(A)-binding protein gene families in flowering plants: Three conserved lineages that are at least 200 million years old and possible auto- and cross-regulation. *Genetics* **163**, 311-319.
- Belostotsky, D.A., and Sieburth, L.E.** (2009). Kill the messenger: mRNA decay and plant development. *Curr. Opin. Plant Biol.* **12**, 96-102.
- Beltrán-Peña, E., Aguilar, R., Ortíz-López, A., Dinkova, T.D., and De Jiménez, E.S.** (2002). Auxin stimulates S6 ribosomal protein phosphorylation in maize thereby affecting protein synthesis regulation. *Physiol Plant* **115**, 291-297.
- Benkowski, L.A., Ravel, J.M., and Browning, K.S.** (1995a). mRNA binding properties of wheat germ protein synthesis initiation factor 2. *Biochem. Biophys. Res. Commun.* **214**, 1033-1039.
- Benkowski, L.A., Ravel, J.M., and Browning, K.S.** (1995b). Development of an *in vitro* translation system from wheat germ that is dependent upon the addition of eukaryotic initiation factor 2. *Anal. Biochem.* **232**, 140-143.
- Benne, R., Kasperaitis, M., Voorma, H.O., Ceglaz, E., and Legocki, A.B.** (1980). Initiation factor eIF-2 from wheat germ: purification, functional comparison to eIF-2 from rabbit reticulocytes and phosphorylation of its subunits. *Eur. J. Biochem.* **104**, 109-117.
- Beznosková, P., Cuchalová, L., Wagner, S., Shoemaker, C.J., Gunišová, S., von der Haar, T., and Valášek, L.S.** (2013). Translation initiation factors eIF3 and HCR1 control translation termination and stop codon read-through in yeast cells. *PLoS Genet.* **9**, e1003962.
- Bi, X.P., and Goss, D.J.** (2000). Wheat germ poly(A)-binding protein increases the ATPase and the RNA helicase activity of translation initiation factors eIF4A, eIF4B, and eIF-iso4F. *J. Biol. Chem.* **275**, 17740-17746.
- Bi, X.P., Ren, J.H., and Goss, D.J.** (2000). Wheat germ translation initiation factor eIF4B affects eIF4A and eIFiso4F helicase activity by increasing the ATP binding affinity of eIF4A. *Biochemistry* **39**, 5758-5765.
- Boex-Fontvieille, E., Daventure, M., Jossier, M., Zivy, M., Hodges, M., and Tcherkez, G.** (2013). Photosynthetic control of *Arabidopsis* leaf cytoplasmic translation initiation by protein phosphorylation. *PLoS One* **8**, e70692.
- Branco-Price, C., Kawaguchi, R., Ferreira, R.B., and Bailey-Serres, J.** (2005). Genome-wide analysis of transcript abundance and translation in *Arabidopsis* seedlings subjected to oxygen deprivation. *Ann. Bot.* **96**, 647-660.
- Branco-Price, C., Kaiser, K.A., Jang, C.J., Larive, C.K., and Bailey-Serres, J.** (2008). Selective mRNA translation coordinates energetic and metabolic adjustments to cellular oxygen deprivation and reoxygenation in *Arabidopsis thaliana*. *Plant J.* **56**, 743-755.
- Bravo, J., Aguilar-Henonin, L., Olmedo, G., and Guzman, P.** (2005). Four distinct classes of proteins as interaction partners of the PABC domain of *Arabidopsis thaliana* poly(A)-binding proteins. *Mol. Genet. Genomics* **272**, 651-665.
- Brina, D., Grosso, S., Miluzio, A., and Biffo, S.** (2011). Translational control by 80S formation and 60S availability: the central role of eIF6, a rate limiting factor in cell cycle progression and tumorigenesis. *Cell Cycle* **10**, 3441-3446.
- Browning, K.** (2014). Cytoplasm: Translational apparatus. *The Plant Sciences. Molecular Biology: Springer Reference 2014-02-13 03:17:23 UTC*.
- Browning, K.S.** (1996). The plant translational apparatus. *Plant Mol. Biol.* **32**, 107-144.
- Browning, K.S.** (2004). Plant translation initiation factors: it is not easy to be green. *Biochem. Soc. Trans.* **32**, 589-591.
- Browning, K.S., Gallie, D.R., Hershey, J.W., Hinnebusch, A.G., Maitra, U., Merrick, W.C., and Norbury, C.** (2001). Unified nomenclature for the subunits of eukaryotic initiation factor 3. *Trends Biochem. Sci.* **26**, 284.
- Bullwinkle, T.J., Zou, S.B., Rajkovic, A., Hersch, S.J., Elgmal, S., Robinson, N., Smil, D., Bolshan, Y., Navarre, W.W., and Ibbá, M.** (2013). (R)- β -lysine-modified elongation factor P functions in translation elongation. *J. Biol. Chem.* **288**, 4416-4423.
- Burks, E.A., Bezerra, P.P., Le, H., Gallie, D.R., and Browning, K.S.** (2001). Plant initiation factor 3 subunit composition resembles mammalian initiation factor 3 and has a novel subunit. *J. Biol. Chem.* **276**, 2122-2131.
- Bush, M.S., Hutchins, A.P., Jones, A.M., Naldrett, M.J., Jarmolowski, A., Lloyd, C.W., and Doonan, J.H.** (2009). Selective recruitment of proteins to 5' cap complexes during the growth cycle in *Arabidopsis*. *Plant J.* **59**, 400-412.
- Byrne, E.H., Prosser, I., Muttucumar, N., Curtis, T.Y., Wingler, A., Powers, S., and Halford, N.G.** (2012). Overexpression of GCN2-type protein kinase in wheat has profound effects on free amino acid concentration and gene expression. *Plant Biotechnol. J.* **10**, 328-340.
- Byrne, M.E.** (2009). A role for the ribosome in development. *Trends Plant Sci.* **14**, 512-519.
- Caldana, C., Li, Y., Leisse, A., Zhang, Y., Bartholomaeus, L., Fernie, A.R., Willmitzer, L., and Giavalisco, P.** (2013). Systemic analysis of inducible target of rapamycin mutants reveal a general metabolic switch controlling growth in *Arabidopsis thaliana*. *Plant J.* **73**, 897-909.
- Callis, J.** (2014). The ubiquitination machinery of the ubiquitin system. *The Arabidopsis Book* **12**, e0174.
- Callot, C., and Gallois, J.L.** (2014). Pyramiding resistances based on translation initiation factors in *Arabidopsis* is impaired by male gametophyte lethality. *Plant Signal Behav* **9**, e27940.
- Carrera, A.C.** (2004). TOR signaling in mammals. *J. Cell Sci.* **117**, 4615-4616.
- Carroll, A.J.** (2013). The *Arabidopsis* cytosolic ribosomal proteome: From form to function. *Front. Plant Sci.* **4**, 32.
- Carroll, A.J., Heazlewood, J.L., Ito, J., and Millar, A.H.** (2008). Analysis

- of the *Arabidopsis* cytosolic ribosome proteome provides detailed insights into its components and their post-translational modification. *Mol. Cell Proteomics* **7**, 347-369.
- Chang, I.F., Szick-Miranda, K., Pan, S., and Bailey-Serres, J.** (2005). Proteomic characterization of evolutionarily conserved and variable proteins of *Arabidopsis* cytosolic ribosomes. *Plant Physiol.* **137**, 848-862.
- Chang, L.Y., Yang, W.Y., Browning, K., and Roth, D.** (1999). Specific *in vitro* phosphorylation of plant eIF2 α by eukaryotic eIF2 α kinases. *Plant Mol. Biol.* **41**, 363-370.
- Chang, Y.F., Imam, J.S., and Wilkinson, M.F.** (2007). The nonsense-mediated decay RNA surveillance pathway. *Annu. Rev. Biochem.* **76**, 51-74.
- Chapman, B., and Brown, C.** (2004). Translation termination in *Arabidopsis thaliana*: characterisation of three versions of release factor 1. *Gene* **341**, 219-225.
- Charron, C., Nicolai, M., Gallois, J.L., Robaglia, C., Moury, B., Palloix, A., and Caranta, C.** (2008). Natural variation and functional analyses provide evidence for co-evolution between plant eIF4E and potyviral VPg. *Plant J.* **54**, 56-68.
- Checkley, J.W., Cooley, L.L., and Ravel, J.M.** (1981). Characterization of initiation factor eIF-3 from wheat germ. *J. Biol. Chem.* **256**, 1582-1586.
- Chen, X.** (2010). Small RNAs - secrets and surprises of the genome. *Plant J.* **61**, 941-958.
- Chen, Z., Jolley, B., Caldwell, C., and Gallie, D.R.** (2014). Eukaryotic translation initiation factor eIFiso4G is required to regulate violaxanthin de-epoxidase expression in *Arabidopsis*. *J. Biol. Chem.* **289**, 13926-13936.
- Cheng, S., and Gallie, D.R.** (2006). Wheat eukaryotic initiation factor 4B organizes assembly of RNA and eIFiso4G, eIF4A, and PABP. *J. Biol. Chem.* **281**, 24351-24364.
- Cheng, S., and Gallie, D.R.** (2007). eIF4G, eIFiso4G, and eIF4B bind the poly(A)-binding protein through overlapping sites within the RNA recognition motif domains. *J. Biol. Chem.* **282**, 25247-25258.
- Cheng, S., and Gallie, D.R.** (2010). Competitive and noncompetitive binding of eIF4B, eIF4A, and the poly(A) binding protein to wheat translation initiation factor eIFiso4G. *Biochemistry* **49**, 8251-8265.
- Cheng, S., and Gallie, D.R.** (2013). Eukaryotic initiation factor 4B and the poly(A)-binding protein bind eIF4G competitively. *Translation* **1**, e24038.
- Cheng, S., Sultana, S., Goss, D.J., and Gallie, D.R.** (2008). Translation initiation factor 4B homodimerization, RNA binding, and interaction with poly(A)-binding protein are enhanced by zinc. *J. Biol. Chem.* **283**, 36140-36153.
- Choi, C.M., Gray, W.M., Mooney, S., and Hellmann, H.** (2014). Composition, Roles, and Regulation of Cullin-Based Ubiquitin E3 Ligases. *The Arabidopsis Book* **12**, e0175.
- Chou, W.C., Huang, Y.W., Tsay, W.S., Chiang, T.Y., Huang, D.D., and Huang, H.J.** (2004). Expression of genes encoding the rice translation initiation factor, eIF5A, is involved in developmental and environmental responses. *Physiol. Plant* **121**, 50-57.
- Coate, J.E., Bar, H., and Doyle, J.J.** (2014). Extensive translational regulation of gene expression in an allopolyploid (*Glycine dolichocarpa*). *Plant Cell* **26**, 136-150.
- Conte, M.R., Kelly, G., Babon, J., Sanfelice, D., Youell, J., Smerdon, S.J., and Proud, C.G.** (2006). Structure of the eukaryotic initiation factor (eIF) 5 reveals a role common to several translation factors. *Biochemistry* **45**, 4550-4558.
- Cooke, C., and Alwine, J.C.** (1996). The cap and the 3' splice site similarly affect polyadenylation efficiency. *Mol. Cell Biol.* **16**, 2579-2584.
- Cuchalova, L., Kouba, T., Herrmannova, A., Danyi, I., Chiu, W.L., and Valasek, L.** (2010). The RNA recognition motif of eukaryotic translation initiation factor 3g (eIF3g) is required for resumption of scanning of posttermination ribosomes for reinitiation on GCN4 and together with eIF3i stimulates linear scanning. *Mol. Cell Biol.* **30**, 4671-4686.
- Dennis, M.D., and Browning, K.S.** (2009). Differential phosphorylation of plant translation initiation factors by *Arabidopsis thaliana* CK2 holoenzymes. *J. Biol. Chem.* **284**, 20602-20614.
- Dennis, M.D., Person, M.D., and Browning, K.S.** (2009). Phosphorylation of plant translation initiation factors by CK2 enhances the *in vitro* interaction of multifactor complex components. *J. Biol. Chem.* **284**, 20615-20628.
- Deprost, D., Yao, L., Sormani, R., Moreau, M., Leterreux, G., Nicolai, M., Bedu, M., Robaglia, C., and Meyer, C.** (2007). The *Arabidopsis* TOR kinase links plant growth, yield, stress resistance and mRNA translation. *EMBO Rep.* **8**, 864-870.
- des Georges, A., Hashem, Y., Unbehaun, A., Grassucci, R.A., Taylor, D., Hellen, C.U., Pestova, T.V., and Frank, J.** (2014). Structure of the mammalian ribosomal pre-termination complex associated with eRF1•eRF3•GDPNP. *Nucl. Acids Res.* **42**, 3409-3418.
- Dever, T.E., and Green, R.** (2012). The elongation, termination, and recycling phases of translation in eukaryotes. *Cold Spring Harb. Perspect. Biol.* **4**, 55-70.
- Dever, T.E., Gutierrez, E., and Shin, B.S.** (2014). The hypusine-containing translation factor eIF5A. *Crit. Rev. Biochem. Mol. Biol.* **49**, 413-425.
- Diedhiou, C.J., Popova, O.V., Dietz, K.J., and Goldack, D.** (2008). The *SUI*-homologous translation initiation factor eIF-1 is involved in regulation of ion homeostasis in rice. *Plant Biol. (Stuttg)* **10**, 298-309.
- Ding, Y., Tang, Y., Kwok, C.K., Zhang, Y., Bevilacqua, P.C., and Assmann, S.M.** (2014). *In vivo* genome-wide profiling of RNA secondary structure reveals novel regulatory features. *Nature* **505**, 696-700.
- Dobrenel, T., Marchive, C., Azzopardi, M., Clément, G., Moreau, M., Sormani, R., Robaglia, C., and Meyer, C.** (2013). Sugar metabolism and the plant target of rapamycin kinase: a sweet operatOR? *Front. Plant Sci.* **4**, 93.
- Dobrikov, M.I., Dobrikova, E.Y., and Gromeier, M.** (2012). Dynamic regulation of the translation initiation helicase complex by mitogenic signal transduction to eIF4G. *Mol. Cell Biol.* **33**, 937-946.
- Doerfel, L.K., Wohlgemuth, I., Kothe, C., Peske, F., Urlaub, H., and Rodnina, M.V.** (2013). EF-P Is essential for rapid synthesis of proteins containing consecutive proline residues. *Science* **339**, 85-88.
- Donnelly, N., Gorman, A.M., Gupta, S., and Samali, A.** (2013). The eIF2 α kinases: their structures and functions. *Cell Mol. Life Sci.* **70**, 3493-3511.
- Dufresne, P.J., Ubalijoro, E., Fortin, M.G., and Laliberte, J.F.** (2008). *Arabidopsis thaliana* class II poly(A)-binding proteins are required for efficient multiplication of turnip mosaic virus. *J. Gen. Virol.* **89**, 2339-2348.
- Echevarría-Zomeño, S., Yáñez, E., Fernández-Bautista, N., Castro-Sanz, A.B., Ferrando, A., and Castellano, M.M.** (2013). Regulation of translation initiation under biotic and abiotic stresses. *Int. J. Mol. Sci.* **14**, 4670-4683.
- Ehsan, H., Ray, W.K., Phinney, B., Wang, X., Huber, S.C., and Clouse, S.D.** (2005). Interaction of *Arabidopsis* BRASSINOSTEROID-INSENSITIVE 1 receptor kinase with a homolog of mammalian TGF- β receptor interacting protein. *Plant J.* **43**, 251-261.
- Falcone Ferreyra, M.L., Pezza, A., Biarc, J., Burlingame, A.L., and Casati, P.** (2010). Plant L10 ribosomal proteins have different roles during development and translation under ultraviolet-B stress. *Plant Physiol.* **153**, 1878-1894.
- Falcone Ferreyra, M.L., Casadevall, R., Luciani, M.D., Pezza, A., and Casati, P.** (2013). New evidence for differential roles of L10 ribosomal

- proteins from *Arabidopsis*. *Plant Physiol.* **163**, 378-391.
- Filichkin, S.A., and Mockler, T.C.** (2012). Unproductive alternative splicing and nonsense mRNAs: a widespread phenomenon among plant circadian clock genes. *Biol. Direct* **7**, 20.
- Filichkin, S.A., Priest, H.D., Givan, S.A., Shen, R., Bryant, D.W., Fox, S.E., Wong, W.K., and Mockler, T.C.** (2010). Genome-wide mapping of alternative splicing in *Arabidopsis thaliana*. *Genome Res.* **20**, 45-58.
- Freire, M.A.** (2005). Translation initiation factor (iso) 4E interacts with BTF3, the β -subunit of the nascent polypeptide-associated complex. *Gene* **345**, 271-277.
- Freire, M.A., Tourneur, C., Granier, F., Camonis, J., El Amrani, A., Browning, K.S., and Robaglia, C.** (2000). Plant lipoxygenase 2 is a translation initiation factor-4E-binding protein. *Plant Mol. Biol.* **44**, 129-140.
- Gallie, D.R.** (2007). Translational control in plants and chloroplasts. In *Translational Control in Biology and Medicine*, M.B. Mathews, N. Sonenberg, and J.W.B. Hershey, eds (Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press), pp. 747-774.
- Gallie, D.R.** (2014). The role of the poly(A) binding protein in the assembly of the cap-binding complex during translation initiation in plants. *Translation* **2**, e959378.
- Gallie, D.R., and Browning, K.S.** (2001). eIF4G functionally differs from eIFiso4G in promoting internal initiation, cap-independent translation, and translation of structured mRNAs. *J. Biol. Chem.* **276**, 36951-36960.
- Gallie, D.R., Le, H., Caldwell, C., Tanguay, R.L., Hoang, N.X., and Browning, K.S.** (1997). The phosphorylation state of translation initiation factors is regulated developmentally and following heat shock in wheat. *J. Biol. Chem.* **272**, 1046-1053.
- Gandin, V., Senft, D., Topisirovic, I., and Ronai, Z.A.** (2013a). RACK1 function in cell motility and protein synthesis. *Genes Cancer* **4**, 369-377.
- Gandin, V., Gutierrez, G.J., Brill, L.M., Varsano, T., Feng, Y., Aza-Blanc, P., Au, Q., McLaughlan, S., Ferreira, T.A., Alain, T., Sonenberg, N., Topisirovic, I., and Ronai, Z.A.** (2013b). Degradation of newly synthesized polypeptides by ribosome-associated RACK1/c-Jun N-terminal kinase/eukaryotic elongation factor 1A2 complex. *Mol. Cell Biol.* **33**, 2510-2526.
- Gaussand, G.M.D.J., Jia, Q., Van der Graaff, E., Lamers, G.E., Franz, P.F., Hooykaas, P.J.J., and De Pater, S.** (2011). Programmed cell death in the leaves of the *Arabidopsis* spontaneous necrotic spots (sns-D) mutant correlates with increased expression of the eukaryotic translation initiation factor eIF4B2. *Front. Plant Sci.* **2**, 9.
- Giavalisco, P., Wilson, D., Kreitler, T., Lehrach, H., Kloese, J., Gobom, J., and Fucini, P.** (2005). High heterogeneity within the ribosomal proteins of the *Arabidopsis thaliana* 80S ribosome. *Plant Mol. Biol.* **57**, 577-591.
- Gonatopoulos-Pournatzis, T., and Cowling, V.H.** (2014). Cap-binding complex (CBC). *Biochem. J.* **457**, 231-242.
- Gonzalez, D.H., and Giegé, P.** (2014). Biogenesis of the oxidative phosphorylation machinery in plants. From gene expression to complex assembly. *Front. Plant Sci.* **5**, 225.
- Gonzalo, P., and Reboud, J.P.** (2003). The puzzling lateral flexible stalk of the ribosome. *Biol. Cell* **95**, 179-193.
- Gorgoni, B., Richardson, W.A., Burgess, H.M., Anderson, R.C., Wilkie, G.S., Gautier, P., Martins, J.P., Brook, M., Sheets, M.D., and Gray, N.K.** (2011). Poly(A)-binding proteins are functionally distinct and have essential roles during vertebrate development. *Proc. Natl. Acad. Sci. USA* **108**, 7844-7849.
- Goss, D.J., and Kleiman, F.E.** (2013). Poly(A) binding proteins: are they all created equal? *Wiley Interdiscip. Rev. RNA* **4**, 167-179.
- Guo, J., and Chen, J.G.** (2008). RACK1 genes regulate plant development with unequal genetic redundancy in *Arabidopsis*. *BMC Plant Biol.* **8**, 108.
- Guo, J., Jin, Z., Yang, X., Li, J.F., and Chen, J.G.** (2011a). Eukaryotic initiation factor 6, an evolutionarily conserved regulator of ribosome biogenesis and protein translation. *Plant Signal. Behav.* **6**, 766-771.
- Guo, J., Wang, J., Xi, L., Huang, W.D., Liang, J., and Chen, J.G.** (2009). RACK1 is a negative regulator of ABA responses in *Arabidopsis*. *J. Exp. Bot.* **60**, 3819-3833.
- Guo, J., Wang, S., Valerius, O., Hall, H., Zeng, Q., Li, J.F., Weston, D.J., Ellis, B.E., and Chen, J.G.** (2011b). Involvement of *Arabidopsis* RACK1 in protein translation and its regulation by abscisic acid. *Plant Physiol.* **155**, 370-383.
- Guo, Y., Xiong, L., Ishitani, M., and Zhu, J.K.** (2002). An *Arabidopsis* mutation in translation elongation factor 2 causes superinduction of CBF/DREB1 transcription factor genes but blocks the induction of their downstream targets under low temperatures. *Proc. Natl. Acad. Sci. USA* **99**, 7786-7791.
- Gutierrez, E., Shin, B.S., Woolstenhulme, C.J., Kim, J.R., Saini, P., Buskirk, A.R., and Dever, T.E.** (2013). eIF5A promotes translation of polyproline motifs. *Mol. Cell* **51**, 35-45.
- Halford, N.G., Hey, S., Jhurrea, D., Laurie, S., McKibbin, R.S., Zhang, Y., and Paul, M.J.** (2004). Highly conserved protein kinases involved in the regulation of carbon and amino acid metabolism. *J. Exp. Bot.* **55**, 35-42.
- Hashem, Y., des, G.A., Dhote, V., Langlois, R., Liao, H.Y., Grassucci, R.A., Hellen, C.U., Pestova, T.V., and Frank, J.** (2013). Structure of the mammalian ribosomal 43S preinitiation complex bound to the scanning factor DHX29. *Cell* **153**, 1108-1119.
- Heise, C., Gardoni, F., Culotta, L., di Luca, M., Verpelli, C., and Sala, C.** (2014). Elongation factor-2 phosphorylation in dendrites and the regulation of dendritic mRNA translation in neurons. *Front. Cell Neurosci.* **8**, 35.
- Henriques, R., Magyar, Z., Monardes, A., Khan, S., Zalejski, C., Orelana, J., Szabados, L., de la Torre, C., Koncz, C., and Bögre, L.** (2010). *Arabidopsis* S6 kinase mutants display chromosome instability and altered RBR1-E2F pathway activity. *EMBO J.* **29**, 2979-2993.
- Hernández, G., and Vazquez-Pianzola, P.** (2005). Functional diversity of the eukaryotic translation initiation factors belonging to eIF4 families. *Mech. Dev.* **122**, 865-876.
- Hernández, G., Altmann, M., and Lasko, P.** (2010). Origins and evolution of the mechanisms regulating translation initiation in eukaryotes. *Trends Biochem. Sci.* **35**, 63-73.
- Hernández, G., Proud, C.G., Preiss, T., and Parsyan, A.** (2012). On the Diversification of the Translation Apparatus across Eukaryotes. *Comp. Funct. Genomics* **2012**, 256848.
- Hershey, J.W., Sonenberg, N., and Mathews, M.B.** (2012). Principles of translational control: an overview. *Cold Spring Harb. Perspect. Biol.* **4**, 1-10.
- Heuffer, C., Browning, K.S., and Ravel, J.M.** (1988). Properties of the subunits of wheat germ initiation factor 3. *Biochim. Biophys. Acta.* **951**, 182-190.
- Hiddinga, H.J., Crum, C.J., Hu, J., and Roth, D.A.** (1988). Viroid-induced phosphorylation of a host protein related to a dsRNA-dependent protein kinase. *Science* **241**, 451-453.
- Hilbert, M., Kebbel, F., Gubaev, A., and Klostermeier, D.** (2011). eIF4G stimulates the activity of the DEAD box protein eIF4A by a conformational guidance mechanism. *Nucl. Acids Res.* **39**, 2260-2270.
- Hinnebusch, A.G.** (2005). Translational regulation of GCN4 and the general amino acid control of yeast. *Annu. Rev. Microbiol.* **59**, 407-450.
- Hinnebusch, A.G.** (2011). Molecular mechanism of scanning and start codon selection in eukaryotes. *Microbiol. Mol. Biol. Rev.* **75**, 434-467.

- Hinnebusch, A.G.** (2014). The scanning mechanism of eukaryotic translation initiation. *Annu. Rev. Biochem.* **83**, 779-812.
- Hinnebusch, A.G., and Lorsch, J.R.** (2012). The mechanism of eukaryotic translation initiation: New insights and challenges. *Cold Spring Harb. Perspect. Biol.* **4**, 29-54.
- Hinnebusch, A.G., Dever, T.E., and Asano, K.** (2007). Mechanism of translation initiation in the yeast *Saccharomyces cerevisiae*. In *Translational Control in Biology and Medicine*, M.B. Mathews, N. Sonenberg, and J.W.B. Hershey, eds (Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press), pp. 225-268.
- Hizli, A.A., Chi, Y., Swanger, J., Carter, J.H., Liao, Y., Welcker, M., Ryzanov, A.G., and Clurman, B.E.** (2013). Phosphorylation of eukaryotic elongation factor 2 (eEF2) by cyclin A-cyclin-dependent kinase 2 regulates its inhibition by eEF2 kinase. *Mol. Cell Biol.* **33**, 596-604.
- Hopkins, M.T., Lampi, Y., Wang, T.W., Liu, Z., and Thompson, J.E.** (2008). Eukaryotic translation initiation factor 5A is involved in pathogen-induced cell death and development of disease symptoms in *Arabidopsis*. *Plant Physiol.* **148**, 479-489.
- Horiguchi, G., Van Lijsebettens, M., Candela, H., Micol, J.L., and Tsukaya, H.** (2012). Ribosomes and translation in plant developmental control. *Plant Sci* **191-192**, 24-34.
- Hsu, Y.F., Chen, Y.C., Hsiao, Y.C., Wang, B.J., Lin, S.Y., Cheng, W.H., Jauh, G.Y., Harada, J.J., and Wang, C.S.** (2013). AtRH57, a DEAD-box RNA helicase, is involved in feedback inhibition of glucose-mediated abscisic acid accumulation during seedling development and additively affects pre-ribosomal RNA processing with high glucose. *Plant J.* **77**, 119-135.
- Hugouvieux, V., Kwak, J.M., and Schroeder, J.I.** (2001). An mRNA cap binding protein, ABH1, modulates early abscisic acid signal transduction in *Arabidopsis*. *Cell* **106**, 477-487.
- Hummel, M., Cordewener, J.H., de Groot, J.C., Smekens, S., America, A.H., and Hanson, J.** (2012). Dynamic protein composition of *Arabidopsis thaliana* cytosolic ribosomes in response to sucrose feeding as revealed by label free MSE proteomics. *Proteomics* **12**, 1024-1038.
- Hunt, A.G.** (2011). RNA regulatory elements and polyadenylation in plants. *Front. Plant Sci.* **2**, 109.
- Hutchins, A.P., Roberts, G.R., Lloyd, C.W., and Doonan, J.H.** (2004). *In vivo* interaction between CDKA and eIF4A: a possible mechanism linking translation and cell proliferation. *FEBS Lett.* **556**, 91-94.
- Hwang, J., Oh, C.S., and Kang, B.C.** (2013). Translation elongation factor 1B (eEF1B) is an essential host factor for tobacco mosaic virus infection in plants. *Virology* **439**, 105-114.
- Immanuel, T.M., Greenwood, D.R., and MacDiarmid, R.M.** (2012). A critical review of translation initiation factor eIF2 α kinases in plants-regulating protein synthesis during stress. *Funct. Plant Biol.* **39**, 717-735.
- Iwakawa, H.O., Tajima, Y., Taniguchi, T., Kaido, M., Mise, K., Tomari, Y., Taniguchi, H., and Okuno, T.** (2012). Poly(A)-binding protein facilitates translation of an uncapped/nonpolyadenylated viral RNA by binding to the 3' untranslated region. *J. Virol.* **86**, 7836-7849.
- Izaurralde, E., Lewis, J., McGuigan, C., Jankowska, M., Darzynkiewicz, E., and Mattaj, I.W.** (1994). A nuclear cap binding protein complex involved in pre-mRNA splicing. *Cell* **78**, 657-668.
- Jackson, R.J.** (2013). The current status of vertebrate cellular mRNA IR-ESs. *Cold Spring Harb. Perspect. Biol.* **5**, a011569.
- Jackson, R.J., Hellen, C.U., and Pestova, T.V.** (2010). The mechanism of eukaryotic translation initiation and principles of its regulation. *Nat. Rev. Mol. Cell Biol.* **11**, 113-127.
- Jackson, R.J., Hellen, C.U., and Pestova, T.V.** (2012). Termination and post-termination events in eukaryotic translation. *Adv. Protein Chem. Struct. Biol.* **86**, 45-93.
- Jannot, G., Bajan, S., Giguère, N.J., Bouasker, S., Banville, I.H., Piquet, S., Hutvagner, G., and Simard, M.J.** (2011). The ribosomal protein RACK1 is required for microRNA function in both *C. elegans* and humans. *EMBO Rep.* **12**, 581-586.
- Janska, H., and Kwasniak, M.** (2014). Mitochondrial regulation of OXPHOS biogenesis in plants. *Front. Plant Sci.* **5**, 79.
- Jennings, M.D., and Pavitt, G.D.** (2010). eIF5 is a dual function GAP and GDI for eukaryotic translational control. *Small GTPases* **1**, 118-123.
- Jennings, M.D., Zhou, Y., Mohammad-Qureshi, S.S., Bennett, D., and Pavitt, G.D.** (2013). eIF2B promotes eIF5 dissociation from eIF2•GDP to facilitate guanine nucleotide exchange for translation initiation. *Genes Dev.* **27**, 2696-2707.
- Jiang, J., and Laliberte, J.F.** (2011). The genome-linked protein VPg of plant viruses—a protein with many partners. *Curr. Opin. Virol.* **1**, 347-354.
- Jiang, J.R., and Clouse, S.D.** (2001). Expression of a plant gene with sequence similarity to animal TGF- β receptor interacting protein is regulated by brassinosteroids and required for normal plant development. *Plant J.* **26**, 35-45.
- Jiao, Y., Riechmann, J.L., and Meyerowitz, E.M.** (2008). Transcriptome-wide analysis of uncapped mRNAs in *Arabidopsis* reveals regulation of mRNA degradation. *Plant Cell* **20**, 2571-2585.
- Jiménez-López, S., Mancera-Martínez, E., Donayre-Torres, A., Rangel, C., Uribe, L., March, S., Jiménez-Sánchez, G., and Sánchez de Jiménez, E.** (2011). Expression profile of maize (*Zea mays* L.) embryonic axes during germination: translational regulation of ribosomal protein mRNAs. *Plant Cell Physiol.* **52**, 1719-1733.
- Jorgensen, R.A., and Dorantes-Acosta, A.E.** (2012). Conserved peptide upstream open reading frames are associated with regulatory genes in angiosperms. *Front. Plant Sci.* **3**, 191.
- Joshi, B., Lee, K., Maeder, D.L., and Jagus, R.** (2005). Phylogenetic analysis of eIF4E-family members. *BMC Evol. Biol.* **5**, 48.
- Juntawong, P., and Bailey-Serres, J.** (2012). Dynamic Light Regulation of Translation Status in *Arabidopsis thaliana*. *Front. Plant Sci.* **3**, 66.
- Juntawong, P., Sorenson, R., and Bailey-Serres, J.** (2013). Cold shock protein 1 chaperones mRNAs during translation in *Arabidopsis thaliana*. *Plant J.* **74**, 1016-1028.
- Juntawong, P., Girke, T., Bazin, J., and Bailey-Serres, J.** (2014). Translational dynamics revealed by genome-wide profiling of ribosome footprints in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* **111**, E203-212.
- Kalyna, M., Simpson, C.G., Syed, N.H., Lewandowska, D., Marquez, Y., Kusenda, B., Marshall, J., Fuller, J., Cardle, L., McNicol, J., Dinh, H.Q., Barta, A., and Brown, J.W.** (2012). Alternative splicing and non-sense-mediated decay modulate expression of important regulatory genes in *Arabidopsis*. *Nucl. Acids Res.* **40**, 2454-2469.
- Kantidakis, T., Ramsbottom, B.A., Birch, J.L., Dowding, S.N., and White, R.J.** (2010). mTOR associates with TFIIC, is found at tRNA and 5S rRNA genes, and targets their repressor Maf1. *Proc. Natl. Acad. Sci. USA* **107**, 11823-11828.
- Karniol, B., Yahalom, A., Kwok, S., Tsuge, T., Matsui, M., Deng, X.W., and Chamovitz, D.A.** (1998). The *Arabidopsis* homologue of an eIF3 complex subunit associates with the COP9 complex. *FEBS Lett.* **439**, 173-179.
- Kawaguchi, R., and Bailey-Serres, J.** (2002). Regulation of translational initiation in plants. *Curr. Opin. Plant Biol.* **5**, 460-465.
- Kawaguchi, R., and Bailey-Serres, J.** (2005). mRNA sequence features that contribute to translational regulation in *Arabidopsis*. *Nucl. Acids Res.* **33**, 955-965.
- Kawaguchi, R., Girke, T., Bray, E.A., and Bailey-Serres, J.** (2004). Differential mRNA translation contributes to gene regulation under non-stress and dehydration stress conditions in *Arabidopsis thaliana*. *Plant*

- J. **38**, 823-839.
- Kerényi, Z., Mérai, Z., Hiripi, L., Benkovics, A., Gyula, P., Lacomme, C., Barta, E., Nagy, F., and Silhavy, D.** (2008). Inter-kingdom conservation of mechanism of nonsense-mediated mRNA decay. *EMBO J.* **27**, 1585-1595.
- Khan, M.A., and Goss, D.J.** (2005). Translation initiation factor (eIF)4B affects the rates of binding of the mRNA m⁷G cap analogue to wheat germ eIFiso4F and eIFiso4F•PABP. *Biochemistry* **44**, 4510-4516.
- Khan, M.A., and Goss, D.J.** (2012). Poly(A)-binding protein increases the binding affinity and kinetic rates of interaction of viral protein linked to genome with translation initiation factors eIFiso4F and eIFiso4F•4B complex. *Biochemistry* **51**, 1388-1395.
- Khan, M.A., Yumak, H., and Goss, D.J.** (2009). Kinetic mechanism for the binding of eIF4F and tobacco etch virus internal ribosome entry site RNA: effects of eIF4B and poly(A)-binding protein. *J. Biol. Chem.* **284**, 35461-35470.
- Khan, M.A., Yumak, H., Gallie, D.R., and Goss, D.J.** (2008). Effects of poly(A)-binding protein on the interactions of translation initiation factor eIF4F and eIF4F•4B with internal ribosome entry site (IRES) of tobacco etch virus RNA. *Biochim. Biophys. Acta.* **1779**, 622-627.
- Khandal, D., Samol, I., Buhr, F., Pollmann, S., Schmidt, H., Clemens, S., Reinbothe, S., and Reinbothe, C.** (2009). Singlet oxygen-dependent translational control in the *tigrina-d.12* mutant of barley. *Proc. Natl. Acad. Sci. USA* **106**, 13112-13117.
- Khoshnevis, S., Hauer, F., Milón, P., Stark, H., and Ficner, R.** (2012). Novel insights into the architecture and protein interaction network of yeast eIF3. *RNA* **18**, 2306-2319.
- Kim, B.H., Cai, X., Vaughn, J.N., and Von Arnim, A.G.** (2007). On the functions of the h subunit of eukaryotic initiation factor 3 in late stages of translation initiation. *Genome Biol.* **8**, R60.
- Kim, J., Kang, W.H., Hwang, J., Yang, H.B., Dosun, K., Oh, C.S., and Kang, B.C.** (2014a). Transgenic *Brassica rapa* plants over-expressing eIF(iso)4E variants show broad-spectrum Turnip mosaic virus (TuMV) resistance. *Mol. Plant Pathol.* **15**, 615-626.
- Kim, T., Hofmann, K., von Arnim, A.G., and Chamovitz, D.A.** (2001). PCI complexes: pretty complex interactions in diverse signaling pathways. *Trends Plant Sci.* **6**, 379-386.
- Kim, T.H., Kim, B.H., Yahalom, A., Chamovitz, D.A., and Von Arnim, A.G.** (2004). Translational regulation via 5' mRNA leader sequences revealed by mutational analysis of the *Arabidopsis* translation initiation factor subunit eIF3h. *Plant Cell* **16**, 3341-3356.
- Kim, Y., Lee, G., Jeon, E., Sohn, E.J., Lee, Y., Kang, H., Lee, D.W., Kim, D.H., and Hwang, I.** (2014b). The immediate upstream region of the 5'-UTR from the AUG start codon has a pronounced effect on the translational efficiency in *Arabidopsis thaliana*. *Nucl. Acids Res.* **42**, 485-498.
- Kim, Y.K., Kim, S., Shin, Y.J., Hur, Y.S., Kim, W.Y., Lee, M.S., Cheon, C.I., and Verma, D.P.** (2014c). Ribosomal protein S6, a target of rapamycin, is involved in the regulation of rRNA genes by possible epigenetic changes in *Arabidopsis*. *J. Biol. Chem.* **289**, 3901-3912.
- Klinge, S., Voigts-Hoffmann, F., Leibundgut, M., and Ban, N.** (2012). Atomic structures of the eukaryotic ribosome. *Trends Biochem. Sci.* **37**, 189-198.
- Kolitz, S.E., and Lorsch, J.R.** (2010). Eukaryotic initiator tRNA: finely tuned and ready for action. *FEBS Lett.* **584**, 396-404.
- Komar, A.A., Mazumder, B., and Merrick, W.C.** (2012). A new framework for understanding IRES-mediated translation. *Gene* **502**, 75-86.
- Koncz, C., Dejong, F., Villacorta, N., Szakonyi, D., and Koncz, Z.** (2012). The spliceosome-activating complex: molecular mechanisms underlying the function of a pleiotropic regulator. *Front. Plant Sci.* **3**, 9.
- Koroleva, O.A., Brown, J.W., and Shaw, P.J.** (2009a). Localization of eIF4A-III in the nucleolus and splicing speckles is an indicator of plant stress. *Plant Signal. Behav.* **4**, 1148-1151.
- Koroleva, O.A., Calder, G., Pendle, A.F., Kim, S.H., Lewandowska, D., Simpson, C.G., Jones, I.M., Brown, J.W., and Shaw, P.J.** (2009b). Dynamic behavior of *Arabidopsis* eIF4A-III, putative core protein of exon junction complex: fast relocation to nucleolus and splicing speckles under hypoxia. *Plant Cell* **21**, 1592-1606.
- Kougioumoutzi, E., Cartolano, M., Canales, C., Dupré, M., Bramsiepe, J., Vlad, D., Rast, M., Dello Iorio, R., Tattersall, A., Schnittger, A., Hay, A., and Tsiantis, M.** (2013). *SIMPLE LEAF3* encodes a ribosome-associated protein required for leaflet development in *Cardamine hirsuta*. *Plant J.* **73**, 533-545.
- Kozak, M.** (1978). How do eucaryotic ribosomes select initiation regions in messenger RNA? *Cell* **15**, 1109-1123.
- Kozak, M.** (1980). Evaluation of the "scanning model" for initiation of protein synthesis in eucaryotes. *Cell* **22**, 7-8.
- Kozak, M.** (1986). Point mutations define a sequence flanking the AUG initiator codon that modulates translation by eukaryotic ribosomes. *Cell* **44**, 283-292.
- Kropiwnicka, A., Kuchta, K., Lukaszewicz, M., Kowalska, J., Jemielity, J., Ginalski, K., Darzynkiewicz, E., and Zuberek, J.** (2015). Five eIF4E isoforms from *Arabidopsis thaliana* are characterized by distinct features of cap analogs binding. *Biochem. Biophys. Res. Commun.* **456**, 47-52.
- Lageix, S., Lanet, E., Pouch-Pelissier, M.N., Espagnol, M.C., Robaglia, C., Deragon, J.M., and Pelissier, T.** (2008). *Arabidopsis* eIF2 α kinase GCN2 is essential for growth in stress conditions and is activated by wounding. *BMC. Plant Biol.* **8**, 134.
- Lan, P., and Schmidt, W.** (2011). The enigma of eIF5A in the iron deficiency response of *Arabidopsis*. *Plant Signal. Behav.* **6**, 528-530.
- Langland, J.O., Jin, S., Jacobs, B.L., and Roth, D.A.** (1995). Identification of a plant-encoded analog of PKR, the mammalian double-stranded RNA-dependent protein kinase. *Plant Physiol.* **108**, 1259-1267.
- Langland, J.O., Langland, L.A., Browning, K.S., and Roth, D.A.** (1996). Phosphorylation of plant eukaryotic initiation factor-2 by the plant-encoded double-stranded RNA-dependent protein kinase, pPKR, and inhibition of protein synthesis *in vitro*. *J. Biol. Chem.* **271**, 4539-4544.
- Latha, R., Salekdeh, G.H., Bennett, J., and Swaminathan, M.S.** (2004). Molecular analysis of a stress-induced cDNA encoding the translation initiation factor, eIF1, from the salt-tolerant wild relative of rice, *Porteresia coarctata*. *Funct. Plant Biol.* **31**, 1035-1042.
- Lauer, S.J., Browning, K.S., and Ravel, J.M.** (1985). Characterization of initiation factor 3 from wheat germ. 2. Effects of polyclonal and monoclonal antibodies on activity. *Biochemistry* **24**, 2928-2931.
- Lauer, S.J., Burks, E., Irvin, J.D., and Ravel, J.M.** (1984). Purification and characterization of three elongation factors, EF-1 α , EF-1 β and EF-2, from wheat germ. *J. Biol. Chem.* **259**, 1644-1648.
- Lax, S.R., Osterhout, J.J., and Ravel, J.M.** (1982). Factors from wheat germ that enhance the activity of eukaryotic initiation factor eIF-2: Isolation and characterization of Co-eIF2 β . *J. Biol. Chem.* **257**, 8233-8237.
- Lax, S.R., Browning, K.S., Maia, D.M., and Ravel, J.M.** (1986). ATPase Activities of Wheat Germ Initiation Factors 4A, 4B and 4F. *J. Biol. Chem.* **261**, 15632-15636.
- Layat, E., Sáez-Vásquez, J., and Tourmente, S.** (2012). Regulation of Pol I-transcribed 45S rDNA and Pol III-transcribed 5S rDNA in *Arabidopsis*. *Plant Cell Physiol.* **53**, 267-276.
- Le, H., and Gallie, D.R.** (2000). Sequence diversity and conservation of the poly(A)-binding protein in plants. *Plant Sci.* **152**, 101-114.
- Le, H., Browning, K.S., and Gallie, D.R.** (2000). The phosphorylation state of poly(A)-binding protein specifies its binding to poly(A) RNA and its interaction with eukaryotic initiation factor (eIF) 4F, eIFiso4F, and eIF4B. *J. Biol. Chem.* **275**, 17452-17462.

- Le, H., Tanguay, R.L., Balasta, M.L., Wei, C.C., Browning, K.S., Metz, A.M., Goss, D.J., and Gallie, D.R. (1997). Translation initiation factors eIF-iso4G and eIF-4B interact with the poly(A)-binding protein and increase its RNA binding activity. *J. Biol. Chem.* **272**, 16247-16255.
- Le Sourd, F., Boulben, S., Le Bouffant, R., Cormier, P., Morales, J., Belle, R., and Mulner-Lorillon, O. (2006). eEF1B: At the dawn of the 21st century. *Biochim. Biophys. Acta.* **1759**, 13-31.
- Lellis, A.D., Allen, M.L., Aertker, A.W., Tran, J.K., Hillis, D.M., Harbin, C.R., Caldwell, C., Gallie, D.R., and Browning, K.S. (2010). Deletion of the eIFiso4G subunit of the *Arabidopsis* eIFiso4F translation initiation complex impairs health and viability. *Plant Mol. Biol.* **74**, 249-263.
- Leprivier, G., Remke, M., Rotblat, B., Dubuc, A., Mateo, A.R., Kool, M., Agnihotri, S., El-Naggar, A., Yu, B., Somasekharan, S.P., Faubert, B., Bridon, G., Tognon, C.E., Mathers, J., Thomas, R., Li, A., Barokas, A., Kwok, B., Bowden, M., Smith, S., Wu, X., Korshunov, A., Hielscher, T., Northcott, P.A., Galpin, J.D., Ahern, C.A., Wang, Y., McCabe, M.G., Collins, V.P., Jones, R.G., Pollak, M., Delattre, O., Gleave, M.E., Jan, E., Pfister, S.M., Proud, C.G., Derry, W.B., Taylor, M.D., and Sorensen, P.H. (2013). The eEF2 kinase confers resistance to nutrient deprivation by blocking translation elongation. *Cell* **153**, 1064-1079.
- Leviatan, N., Alkan, N., Leshkowitz, D., and Fluhr, R. (2013). Genome-wide survey of cold stress regulated alternative splicing in *Arabidopsis thaliana* with tiling microarray. *PLoS One* **8**, e66511.
- Li, M.W., AuYeung, W.K., and Lam, H.M. (2013a). The GCN2 homologue in *Arabidopsis thaliana* interacts with uncharged tRNA and uses *Arabidopsis* eIF2 α molecules as direct substrates. *Plant Biol.* **15**, 13-18.
- Li, P., Ponnala, L., Gandotra, N., Wang, L., Si, Y., Tausta, S.L., Kebrom, T.H., Provart, N., Patel, R., Myers, C.R., Reidel, E.J., Turgeon, R., Liu, P., Sun, Q., Nelson, T., and Brutnell, T.P. (2010). The developmental dynamics of the maize leaf transcriptome. *Nat. Genet.* **42**, 1060-1067.
- Li, S., Liu, L., Zhuang, X., Yu, Y., Liu, X., Cui, X., Ji, L., Pan, Z., Cao, X., Mo, B., Zhang, F., Raikhel, N., Jiang, L., and Chen, X. (2013b). MicroRNAs inhibit the translation of target mRNAs on the endoplasmic reticulum in *Arabidopsis*. *Cell* **153**, 562-574.
- Li, X.P., Kahn, P.C., Kahn, J.N., Grela, P., and Tumer, N.E. (2013c). Arginine residues on the opposite side of the active site stimulate the catalysis of ribosome depurination by ricin A chain by interacting with the P-protein stalk. *J. Biol. Chem.* **288**, 30270-30284.
- Li, Y., and Kiledjian, M. (2010). Regulation of mRNA decapping. *Wiley Interdiscip. Rev. RNA* **1**, 253-265.
- Li, Z., and Nagy, P.D. (2011). Diverse roles of host RNA binding proteins in RNA virus replication. *RNA Biol.* **8**, 305-315.
- Li, Z., Pogany, J., Panavas, T., Xu, K., Esposito, A.M., Kinzy, T.G., and Nagy, P.D. (2009). Translation elongation factor 1A is a component of the tombusvirus replicase complex and affects the stability of the p33 replication co-factor. *Virology* **385**, 245-260.
- Linder, P., and Fuller-Pace, F. (2013). Looking back on the birth of DEAD-box RNA helicases. *Biochim. Biophys. Acta.* **1829**, 750-755.
- Liu, M.J., Wu, S.H., Wu, J.F., Lin, W.D., Wu, Y.C., Tsai, T.Y., and Tsai, H.L. (2013). Translational landscape of photomorphogenic *Arabidopsis*. *Plant Cell* **25**, 3699-3710.
- Liu, Y., Neumann, P., Kuhle, B., Monecke, T., Schell, S., Chari, A., and Ficner, R. (2014). Translation initiation factor eIF3b contains a nine-bladed β -propeller and interacts with the 40S ribosomal subunit. *Structure* **22**, 923-930.
- Lomakin, I.B., and Steitz, T.A. (2013). The initiation of mammalian protein synthesis and mRNA scanning mechanism. *Nature* **500**, 307-311.
- Lopez-Valenzuela, J.A., Gibbon, B.C., Holding, D.R., and Larkins, B.A. (2004). Cytoskeletal proteins are coordinately increased in maize genotypes with high levels of eEF1A. *Plant Physiol.* **135**, 1784-1797.
- Lopez-Valenzuela, J.A., Gibbon, B.C., Hughes, P.A., Dreher, T.W., and Larkins, B.A. (2003). eEF1A isoforms change in abundance and actin-binding activity during maize endosperm development. *Plant Physiol.* **133**, 1285-1295.
- Lorsch, J.R., and Dever, T.E. (2010). Molecular view of 43S complex formation and start site selection in eukaryotic translation initiation. *J. Biol. Chem.* **285**, 21203-21207.
- Luo, Y.J., and Goss, D.J. (2001). Homeostasis in mRNA initiation - Wheat germ poly(A)-binding protein lowers the activation energy barrier to initiation complex formation. *J. Biol. Chem.* **276**, 43083-43086.
- Ma, Y., Miura, E., Ham, B.K., Cheng, H.W., Lee, Y.J., and Lucas, W.J. (2010). Pumpkin eIF5A isoforms interact with components of the translational machinery in the cucurbit sieve tube system. *Plant J.* **64**, 536-550.
- Maldonado-Bonilla, L.D. (2014). Composition and function of P bodies in *Arabidopsis thaliana*. *Front. Plant Sci.* **5**, 201.
- Marintchev, A. (2013). Roles of helicases in translation initiation: A mechanistic view. *Biochim. Biophys. Acta.* **1829**, 799-809.
- Marintchev, A., and Wagner, G. (2005). eIF4G and CBP80 share a common origin and similar domain organization: implications for the structure and function of eIF4G. *Biochemistry* **44**, 12265-12272.
- Marintchev, A., Edmonds, K.A., Marintcheva, B., Hendrickson, E., Oberer, M., Suzuki, C., Herdy, B., Sonenberg, N., and Wagner, G. (2009). Topology and regulation of the human eIF4A/4G/4H helicase complex in translation initiation. *Cell* **136**, 447-460.
- Marquez, Y., Brown, J.W., Simpson, C., Barta, A., and Kalyna, M. (2012). Transcriptome survey reveals increased complexity of the alternative splicing landscape in *Arabidopsis*. *Genome Res* **22**, 1184-1195.
- Martin-Marcos, P., Nanda, J.S., Luna, R.E., Zhang, F., Saini, A.K., Cherkasova, V.A., Wagner, G., Lorsch, J.R., and Hinnebusch, A.G. (2014). Enhanced eIF1 binding to the 40S ribosome impedes conformational rearrangements of the preinitiation complex and elevates initiation accuracy. *RNA* **20**, 150-167.
- Matsuda, D., Yoshinari, S., and Dreher, T.W. (2004). eEF1A binding to aminoacylated viral RNA represses minus strand synthesis by TYMV RNA-dependent RNA polymerase. *Virology* **321**, 47-56.
- Mayberry, L.K., Allen, M.L., Dennis, M.D., and Browning, K.S. (2009). Evidence for variation in the optimal translation initiation complex: plant eIF4B, eIF4F, and eIF(iso)4F differentially promote translation of mRNAs. *Plant Physiol.* **150**, 1844-1854.
- Mayberry, L.K., Allen, M.L., Nitka, K.R., Campbell, L., Murphy, P.A., and Browning, K.S. (2011). Plant cap-binding complexes eukaryotic initiation factors eIF4F and eIFiso4F: Molecular specificity of subunit binding. *J. Biol. Chem.* **286**, 42566-42574.
- McIntosh, K.B., and Bonham-Smith, P.C. (2006). Ribosomal protein gene regulation: what about plants? *Can. J. Bot.* **84**, 342-362.
- Mead, E.J., Masterton, R.J., von der Haar, T., Tuite, M.F., and Smales, C.M. (2014). Control and regulation of mRNA translation. *Biochem. Soc. Trans.* **42**, 151-154.
- Meng, H., Li, C., Wang, Y., and Chen, G. (2014). Molecular dynamics simulation of the allosteric regulation of eIF4A protein from the open to closed state, induced by ATP and RNA substrates. *PLoS One* **9**, e86104.
- Merrick, W.C., and Harris, M.E. (2014). Control not at initiation? Bah, humbug! *EMBO J.* **33**, 3-4.
- Metz, A.M., and Browning, K.S. (1997). Assignment of the β -subunit of wheat eIF2 by protein and DNA sequence analysis and immunoanalysis. *Arch. Biochem. Biophys.* **342**, 187-189.
- Milac, A.L., Bojarska, E., and Wypijewska Del Nogal, A. (2014). Decapping scavenger (DcpS) enzyme: Advances in its structure, activity and

- roles in the cap-dependent mRNA metabolism. *Biochim. Biophys. Acta.* **839**, 452-462.
- Miluzio, A., Beugnet, A., Volta, V., and Biffo, S.** (2009). Eukaryotic initiation factor 6 mediates a continuum between 60S ribosome biogenesis and translation. *EMBO Rep.* **10**, 459-465.
- Monzingo, A.F., Dhaliwal, S., Dutt-Chaudhuri, A., Lyon, A., Sadow, J.H., Hoffman, D.W., Robertus, J.D., and Browning, K.S.** (2007). The structure of eukaryotic translation initiation factor-4E from wheat reveals a novel disulfide bond. *Plant Physiol.* **143**, 1504-1518.
- Morita, M., Ler, L.W., Fabian, M.R., Siddiqui, N., Mullin, M., Henderson, V.C., Alain, T., Fonseca, B.D., Karashchuk, G., Bennett, C.F., Kabuta, T., Higashi, S., Larsson, O., Topisirovic, I., Smith, R.J., Gingras, A.C., and Sonenberg, N.** (2012). A novel 4EHP-GIGYF2 translational repressor complex is essential for mammalian development. *Mol. Cell Biol.* **32**, 3585-3593.
- Moury, B., Charron, C., Janzac, B., Simon, V., Gallois, J.L., Palloix, A., and Caranta, C.** (2013). Evolution of plant eukaryotic initiation factor 4E (eIF4E) and potyvirus genome-linked protein (VPg): A game of mirrors impacting resistance spectrum and durability. *Infect. Genet. Evol.* **27**, 472-480.
- Muench, D.G., Zhang, C., and Dahodwala, M.** (2012). Control of cytoplasmic translation in plants. *Wiley Interdiscip. Rev. RNA* **3**, 178-194.
- Mulekar, J.J., and Huq, E.** (2013). Expanding roles of protein kinase CK2 in regulating plant growth and development. *J. Exp. Bot.* **65**, 2883-2893.
- Mustroph, A., Zanetti, M.E., Jang, C.J., Holtan, H.E., Repetti, P.P., Galbraith, D.W., Girke, T., and Bailey-Serres, J.** (2009). Profiling translomes of discrete cell populations resolves altered cellular priorities during hypoxia in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* **106**, 18843-18848.
- Muñoz, A., and Castellano, M.M.** (2012). Regulation of translation initiation under abiotic stress conditions in plants: Is it a conserved or not so conserved process among eukaryotes? *Comp. Funct. Genomics* **2012**, 406357.
- Nanda, J.S., Saini, A.K., Muñoz, A.M., Hinnebusch, A.G., and Lorsch, J.R.** (2013). Coordinated movements of eukaryotic translation initiation factors eIF1, eIF1A, and eIF5 trigger phosphate release from eIF2 in response to start codon recognition by the ribosomal preinitiation complex. *J. Biol. Chem.* **288**, 5316-5329.
- Nanda, J.S., Cheung, Y.N., Takacs, J.E., Martin-Marcos, P., Saini, A.K., Hinnebusch, A.G., and Lorsch, J.R.** (2009). eIF1 controls multiple steps in start codon recognition during eukaryotic translation initiation. *J. Mol. Biol.* **394**, 268-285.
- Nicolai, M., Roncato, M.A., Canoy, A.S., Rouquié, D., Sarda, X., Freyssinet, G., and Robaglia, C.** (2006). Large-scale analysis of mRNA translation states during sucrose starvation in *Arabidopsis* cells identifies cell proliferation and chromatin structure as targets of translational control. *Plant Physiol.* **141**, 663-673.
- Nishimura, T., Wada, T., Yamamoto, K.T., and Okada, K.** (2005). The *Arabidopsis* STV1 protein, responsible for translation reinitiation, is required for auxin-mediated gynoceium patterning. *Plant Cell* **17**, 2940-2953.
- O'Brien, J.P., Mayberry, L.K., Murphy, P.A., Browning, K.S., and Brodbelt, J.S.** (2013). Evaluating the conformation and binding interface of cap-binding proteins and complexes via ultraviolet photodissociation mass spectrometry. *J. Proteome Res.* **12**, 5867-5877.
- Ortiz, P.A., Ulloque, R., Kihara, G.K., Zheng, H., and Kinzy, T.G.** (2006). Translation elongation factor 2 anticodon mimicry domain mutants affect fidelity and diphtheria toxin resistance. *J. Biol. Chem.* **281**, 32639-32648.
- Osterhout, J.J., Lax, S.R., and Ravel, J.M.** (1983). Factors from wheat germ that enhance the activity of eukaryotic initiation factor eIF-2: Isolation and characterization of Co-eIF-2 α . *J. Biol. Chem.* **258**, 8285-8289.
- Pal, S.K., Liput, M., Piques, M., Ishihara, H., Obata, T., Martins, M.C., Sulpice, R., van Dongen, J.T., Fernie, A.R., Yadav, U.P., Lunn, J.E., Usadel, B., and Stitt, M.** (2013). Diurnal changes of polysome loading track sucrose content in the rosette of wild-type *Arabidopsis* and the starchless pgm mutant. *Plant Physiol.* **162**, 1246-1265.
- Papp, I., Mur, L.A., Dalmadi, A., Dulai, S., and Koncz, C.** (2004). A mutation in the cap binding protein 20 gene confers drought tolerance to *Arabidopsis*. *Plant Mol. Biol.* **55**, 679-686.
- Park, E.H., Walker, S.E., Lee, J.M., Rothenburg, S., Lorsch, J.R., and Hinnebusch, A.G.** (2011). Multiple elements in the eIF4G1 N-terminus promote assembly of eIF4G1•PABP mRNPs *in vivo*. *EMBO J.* **30**, 302-316.
- Park, E.H., Walker, S.E., Zhou, F., Lee, J.M., Rajagopal, V., Lorsch, J.R., and Hinnebusch, A.G.** (2012). Yeast eukaryotic initiation factor (eIF) 4B enhances complex assembly between eIF4A and eIF4G *in vivo*. *J. Biol. Chem.* **288**, 2340-2354.
- Park, H.S., Browning, K.S., Hohn, T., and Ryabova, L.A.** (2004). Eukaryotic initiation factor 4B controls eIF3-mediated ribosomal entry of viral reinitiation factor. *EMBO J.* **23**, 1381-1391.
- Park, H.S., Himmelbach, A., Browning, K.S., Hohn, T., and Ryabova, L.A.** (2001). A plant viral "reinitiation" factor interacts with the host translational machinery. *Cell* **106**, 723-733.
- Parsyan, A., Svitkin, Y., Shahbazian, D., Gkogkas, C., Lasko, P., Merriker, W.C., and Sonenberg, N.** (2011). mRNA helicases: the tacticians of translational control. *Nat. Rev. Mol. Cell Biol.* **12**, 235-245.
- Patrick, R.M., and Browning, K.S.** (2012). The eIF4F and eIFiso4F complexes of plants: An evolutionary perspective. *Comp Funct. Genomics* **2012**, 287814.
- Patrick, R.M., Mayberry, L.K., Choy, G., Woodard, L.E., Liu, J.S., White, A., Mullen, R.A., Tanavin, T.M., Latz, C.A., and Browning, K.S.** (2014). Two *Arabidopsis* loci encode novel eukaryotic initiation factor 4E isoforms that are functionally distinct from the conserved plant eukaryotic initiation factor 4E. *Plant Physiol.* **164**, 1820-1830.
- Paz-Avram, T., Yahalom, A., and Chamovitz, D.A.** (2008). *Arabidopsis* eIF3e interacts with subunits of the ribosome, Cop9 signalosome and proteasome. *Plant Signal. Behav.* **3**, 409-411.
- Petsch, K.A., Mylne, J., and Botella, J.R.** (2005). Cosuppression of eukaryotic release factor 1-1 in *Arabidopsis* affects cell elongation and radial cell division. *Plant Physiol.* **139**, 115-126.
- Pick, E., Hofmann, K., and Glickman, M.H.** (2009). PCI complexes: Beyond the proteasome, CSN, and eIF3 trioka. *Mol. Cell* **35**, 260-264.
- Piques, M., Schulze, W.X., Höhne, M., Usadel, B., Gibon, Y., Rohwer, J., and Stitt, M.** (2009). Ribosome and transcript copy numbers, polysome occupancy and enzyme dynamics in *Arabidopsis*. *Mol. Syst. Biol.* **5**, 314.
- Pisarev, A.V., Hellen, C.U., and Pestova, T.V.** (2007). Recycling of eukaryotic posttermination ribosomal complexes. *Cell* **131**, 286-299.
- Pisarev, A.V., Skabkin, M.A., Pisareva, V.P., Skabkina, O.V., Rakotondrifara, A.M., Hentze, M.W., Hellen, C.U., and Pestova, T.V.** (2010). The role of ABCE1 in eukaryotic posttermination ribosomal recycling. *Mol. Cell* **37**, 196-210.
- Pisareva, V.P., and Pisarev, A.V.** (2014). eIF5 and eIF5B together stimulate 48S initiation complex formation during ribosomal scanning. *Nucl. Acids Res.* **42**, 12052-12069.
- Preis, A., Heuer, A., Barrio-Garcia, C., Hauser, A., Eyler, D.E., Berninghausen, O., Green, R., Becker, T., and Beckmann, R.** (2014). Cryoelectron microscopic structures of eukaryotic translation termination complexes containing eRF1•eRF3 or eRF1•ABCE1. *Cell Rep.* **8**, 59-65.
- Putnam, A.A., and Jankowsky, E.** (2013). DEAD-box helicases as integrators of RNA, nucleotide and protein binding. *Biochim. Biophys. Acta.*

- 1829, 884-893.
- Pyl, E.T., Piques, M., Ivakov, A., Schulze, W., Ishihara, H., Stitt, M., and Sulpice, R.** (2012). Metabolism and growth in *Arabidopsis* depend on the daytime temperature but are temperature-compensated against cool nights. *Plant Cell* **24**, 2443-2469.
- Querol-Audi, J., Sun, C., Vogan, J.M., Smith, M.D., Gu, Y., Cate, J.H., and Nogales, E.** (2013). Architecture of human translation initiation factor 3. *Structure* **21**, 920-928.
- Raczynska, K.D., Stepien, A., Kierzkowski, D., Kalak, M., Bajczyk, M., McNicol, J., Simpson, C.G., Szweykowska-Kulinska, Z., Brown, J.W., and Jarmolowski, A.** (2014). The *SERRATE* protein is involved in alternative splicing in *Arabidopsis thaliana*. *Nucleic Acids Res* **42**, 1224-1244.
- Rahmani, F., Hummel, M., Schuurmans, J., Wiese-Klinkenberg, A., Smeekens, S., and Hanson, J.** (2009). Sucrose control of translation mediated by an upstream open reading frame-encoded peptide. *Plant Physiol.* **150**, 1356-1367.
- Rasheedi, S., Suragani, M., Haq, S.K., Sachchidanand, Bhardwaj, R., Hasnain, S.E., and Ehtesham, N.Z.** (2010). Expression, purification and ligand binding properties of the recombinant translation initiation factor (PeIF5B) from *Pisum sativum*. *Mol. Cell Biochem.* **344**, 33-41.
- Rausell, A., Kanhonou, R., Yenush, L., Serrano, R., and Ros, R.** (2003). The translation initiation factor eIF1A is an important determinant in the tolerance to NaCl stress in yeast and plants. *Plant J.* **34**, 257-267.
- Raychaudhuri, P., Stringer, E.A., Valenzuela, D.M., and Maitra, U.** (1984). Ribosomal subunit antiassociation activity in rabbit reticulocyte lysates. Evidence for a low molecular weight ribosomal subunit antiassociation protein factor (Mr = 25,000). *J. Biol. Chem.* **259**, 11930-11935.
- Reddy, A.S.** (2007). Alternative splicing of pre-messenger RNAs in plants in the genomic era. *Annu. Rev. Plant Biol.* **58**, 267-294.
- Reddy, A.S., Marquez, Y., Kalyna, M., and Barta, A.** (2013). Complexity of the alternative splicing landscape in plants. *Plant Cell* **25**, 3657-3683.
- Ren, B., Chen, Q., Hong, S., Zhao, W., Feng, J., Feng, H., and Zuo, J.** (2013). The *Arabidopsis* eukaryotic translation initiation factor eIF5A-2 regulates root protoxylem development by modulating cytokinin signaling. *Plant Cell* **25**, 3841-3857.
- Ren, M., Qiu, S., Venglat, P., Xiang, D., Feng, L., Selvaraj, G., and Datla, R.** (2011). Target of rapamycin regulates development and ribosomal RNA expression through kinase domain in *Arabidopsis*. *Plant Physiol.* **155**, 1367-1382.
- Ren, M., Venglat, P., Qiu, S., Feng, L., Cao, Y., Wang, E., Xiang, D., Wang, J., Alexander, D., Chalivendra, S., Logan, D., Mattoo, A., Selvaraj, G., and Datla, R.** (2012). Target of rapamycin signaling regulates metabolism, growth, and life span in *Arabidopsis*. *Plant Cell* **24**, 4850-4874.
- Rhoads, R.E.** (2009). eIF4E: new family members, new binding partners, new roles. *J. Biol. Chem.* **284**, 16711-16715.
- Richter, J.D., and Sonenberg, N.** (2005). Regulation of cap-dependent translation by eIF4E inhibitory proteins. *Nature* **433**, 477-480.
- Riechmann, J.L., Ito, T., and Meyerowitz, E.M.** (1999). Non-AUG initiation of *AGAMOUS* mRNA translation in *Arabidopsis thaliana*. *Mol. Cell Biol.* **19**, 8505-8512.
- Robaglia, C., and Caranta, C.** (2006). Translation initiation factors: a weak link in plant RNA virus infection. *Trends Plant Sci.* **11**, 40-45.
- Robaglia, C., Thomas, M., and Meyer, C.** (2012). Sensing nutrient and energy status by SnRK1 and TOR kinases. *Curr. Opin. Plant Biol.* **15**, 301-307.
- Rogers, G.W., Jr., Komar, A.A., and Merrick, W.C.** (2002). eIF4A: The godfather of the DEAD box helicases. *Prog. Nucl. Acid Res. Mol. Biol.* **72**, 307-331.
- Rogers, K., and Chen, X.** (2013). Biogenesis, turnover, and mode of activation of plant microRNAs. *Plant Cell* **25**, 2383-2399.
- Rom, E., Kim, H.C., Gingras, A.C., Marcotrigiano, J., Favre, D., Olsen, H., Burley, S.K., and Sonenberg, N.** (1998). Cloning and characterization of 4EHP, a novel mammalian eIF4E-related cap-binding protein. *J. Biol. Chem.* **273**, 13104-13109.
- Rosado, A., and Raikhel, N.V.** (2010). Application of the gene dosage balance hypothesis to auxin-related ribosomal mutants in *Arabidopsis*. *Plant Signal. Behav.* **5**, 450-452.
- Rosado, A., Li, R., van de Ven, W., Hsu, E., and Raikhel, N.V.** (2012). *Arabidopsis* ribosomal proteins control developmental programs through translational regulation of auxin response factors. *Proc. Natl. Acad. Sci. USA* **109**, 19537-19544.
- Roy, B., and von Arnim, A.G.** (2013). Translational regulation of cytoplasmic mRNAs. *The Arabidopsis Book* **11**, e0165.
- Roy, B., Vaughn, J.N., Kim, B.H., Zhou, F., Gilchrist, M.A., and Von Arnim, A.G.** (2010). The h subunit of eIF3 promotes reinitiation competence during translation of mRNAs harboring upstream open reading frames. *RNA* **16**, 748-761.
- Russell, D.W., and Spremulli, L.L.** (1978). Identification of a wheat germ ribosome dissociation factor distinct from initiation factor eIF-3. *J. Biol. Chem.* **253**, 6647-6649.
- Russell, D.W., and Spremulli, L.L.** (1979). Purification and characterization of a ribosome dissociation factor (eukaryotic initiation factor 6) from wheat germ. *J. Biol. Chem.* **254**, 8796-8800.
- Russell, D.W., and Spremulli, L.L.** (1980). Mechanism of action of the wheat germ ribosome dissociation factor: interaction with the 60 S subunit. *Arch. Biochem. Biophys.* **201**, 518-526.
- Ruud, K.A., Kuhlow, C., Goss, D.J., and Browning, K.S.** (1998). Identification and characterization of a novel cap-binding protein from *Arabidopsis thaliana*. *J. Biol. Chem.* **273**, 10325-10330.
- Safer, B.** (1989). Nomenclature of initiation, elongation and termination factors for translation in eukaryotes. *Eur. J. Biochem.* **186**, 1-3.
- Sahoo, R.K., Gill, S.S., and Tuteja, N.** (2012). Pea DNA helicase 45 promotes salinity stress tolerance in IR64 rice with improved yield. *Plant Signal. Behav.* **7**, 1042-1046.
- Saini, A.K., Nanda, J.S., Martin-Marcos, P., Dong, J., Zhang, F., Bhardwaj, M., Lorsch, J.R., and Hinnebusch, A.G.** (2014). Eukaryotic translation initiation factor eIF5 promotes the accuracy of start codon recognition by regulating Pi release and conformational transitions of the preinitiation complex. *Nucl. Acids Res.* **42**, 9623-9640.
- Sasikumar, A.N., Perez, W.B., and Kinzy, T.G.** (2012). The many roles of the eukaryotic elongation factor 1 complex. *Wiley Interdiscip. Rev. RNA* **3**, 543-555.
- Sasvari, Z., Izotova, L., Kinzy, T.G., and Nagy, P.D.** (2011). Synergistic roles of eukaryotic translation elongation factors 1By and 1A in stimulation of tombusvirus minus-strand synthesis. *PLoS Pathog.* **7**, e1002438.
- Schepetilnikov, M., Dimitrova, M., Mancera-Martinez, E., Geldreich, A., Keller, M., and Ryabova, L.A.** (2013). TOR and S6K1 promote translation reinitiation of uORF-containing mRNAs via phosphorylation of eIF3h. *EMBO J.* **32**, 1087-1102.
- Schepetilnikov, M., Kobayashi, K., Geldreich, A., Caranta, C., Robaglia, C., Keller, M., and Ryabova, L.A.** (2011). Viral factor TAV recruits TOR/S6K1 signalling to activate reinitiation after long ORF translation. *EMBO J.* **30**, 1343-1356.
- Schmitt, E., Naveau, M., and Mechulam, Y.** (2010). Eukaryotic and archaeal translation initiation factor 2: a heterotrimeric tRNA carrier. *FEBS Lett.* **584**, 405-412.
- Schmollinger, S., Mülhhaus, T., Boyle, N.R., Blaby, I.K., Casero, D., Mettler, T., Moseley, J.L., Kropat, J., Sommer, F., Strenkert, D., Hemme, D., Pellegrini, M., Grossman, A.R., Stitt, M., Schroda, M.,**

- and Merchant, S.S. (2014). Nitrogen-sparing mechanisms in *Chlamydomonas* affect the transcriptome, the proteome, and photosynthetic metabolism. *Plant Cell* **26**, 1410-1435.
- Schütz, P., Bumann, M., Oberholzer, A.E., Bieniossek, C., Trachsel, H., Altmann, M., and Baumann, U. (2008). Crystal structure of the yeast eIF4A-eIF4G complex: an RNA-helicase controlled by protein-protein interactions. *Proc. Natl. Acad. Sci. USA* **105**, 9564-9569.
- Seal, S.N., Schmidt, A., and Marcus, A. (1983). Wheat Germ eIF-2 and Co-eIF-2: Resolution and functional characterization in *in vitro* protein synthesis. *J. Biol. Chem.* **258**, 10573-10576.
- Shaikhin, S.M., Smailov, S.K., Lee, A.V., Kozhanov, E.V., and Iskakov, B.K. (1992). Interaction of wheat germ translation initiation factor 2 with GDP and GTP. *Biochimie* **74**, 447-454.
- Shin, Y.J., Kim, S., Du, H., Choi, S., Verma, D.P., and Cheon, C.I. (2012). Possible dual regulatory circuits involving AtS6K1 in the regulation of plant cell cycle and growth. *Mol. Cells* **33**, 487-496.
- Si, K., Chaudhuri, J., Chevesich, J., and Maitra, U. (1997). Molecular cloning and functional expression of a human cDNA encoding translation initiation factor 6. *Proc. Natl. Acad. Sci.* **94**, 14285-14290.
- Siddiqui, N., Osborne, M.J., Gallie, D.R., and Gehring, K. (2007). Solution structure of the PABC domain from wheat poly (A)-binding protein: An insight into RNA metabolic and translational control in plants. *Biochemistry* **46**, 4221-4231.
- Simon, A.E., and Miller, W.A. (2013). 3' cap-independent translation enhancers of plant viruses. *Annu. Rev. Microbiol.* **67**, 21-42.
- Simpson, G.G., Laurie, R.E., Dijkwel, P.P., Quesada, V., Stockwell, P.A., Dean, C., and Macknight, R.C. (2010). Noncanonical translation initiation of the *Arabidopsis* flowering time and alternative polyadenylation regulator *FCA*. *Plant Cell* **22**, 3764-3777.
- Singh, B., Chauhan, H., Khurana, J.P., Khurana, P., and Singh, P. (2013). Evidence for the role of wheat eukaryotic translation initiation factor 3 subunit g (*TaeIF3g*) in abiotic stress tolerance. *Gene* **532**, 177-185.
- Singh, C.R., Watanabe, R., Chowdhury, W., Hiraishi, H., Murai, M.J., Yamamoto, Y., Miles, D., Ikeda, Y., Asano, M., and Asano, K. (2012). Sequential eukaryotic translation initiation factor 5 (eIF5) binding to the charged disordered segments of eIF4G and eIF2 β stabilizes the 48S preinitiation complex and promotes its shift to the initiation mode. *Mol. Cell Biol.* **32**, 3978-3989.
- Singh, G., M., J., Kulshreshtha, R., Khurana, J.P., Kumar, S., and Singh, P. (2007). Expression analysis of genes encoding translation initiation factor 3 subunit g (*TaeIF3g*) and vesicle-associated membrane protein-associated protein (*TaVAP*) in drought tolerant and susceptible cultivars of wheat. *Plant Sci.* **173**, 660-669.
- Siridechadilok, B., Fraser, C.S., Hall, R.J., Doudna, J.A., and Nogales, E. (2005). Structural roles for human translation factor eIF3 in initiation of protein synthesis. *Science* **310**, 1513-1515.
- Skabkin, M.A., Skabkina, O.V., Hellen, C.U., and Pestova, T.V. (2013). Reinitiation and other unconventional posttermination events during eukaryotic translation. *Mol. Cell* **51**, 249-264.
- Smailov, S.K., Lee, A.V., and Iskakov, B.K. (1993). Study of phosphorylation of translation elongation factor 2 (EF-2) from wheat germ. *FEBS Lett.* **321**, 219-223.
- Smith, R.W., and Gray, N.K. (2010). Poly(A)-binding protein (PABP): a common viral target. *Biochem J.* **426**, 1-12.
- Sokabe, M., Fraser, C.S., and Hershey, J.W. (2012). The human translation initiation multi-factor complex promotes methionyl-tRNA_i binding to the 40S ribosomal subunit. *Nucl. Acids Res.* **40**, 905-913.
- Sonenberg, N., and Hinnebusch, A.G. (2009). Regulation of translation initiation in eukaryotes: mechanisms and biological targets. *Cell* **136**, 731-745.
- Sorenson, R., and Bailey-Serres, J. (2014). Selective mRNA sequestration by *OLIGOURIDYLATE-BINDING PROTEIN 1* contributes to translational control during hypoxia in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* **111**, 2373-2378.
- Sormani, R., Masclaux-Daubresse, C., Daniel-Vedele, F., Daniele-Vedele, F., and Chardon, F. (2011a). Transcriptional regulation of ribosome components are determined by stress according to cellular compartments in *Arabidopsis thaliana*. *PLoS One* **6**, e28070.
- Sormani, R., Delannoy, E., Lageix, S., Bitton, F., Lanet, E., Saez-Vasquez, J., Deragon, J.M., Renou, J.P., and Robaglia, C. (2011b). Sublethal cadmium intoxication In *Arabidopsis thaliana* impacts translation at multiple levels. *Plant Cell Physiol.* **52**, 436-447.
- Speth, C., Willing, E.M., Rausch, S., Schneeberger, K., and Laubinger, S. (2013). RACK1 scaffold proteins influence miRNA abundance in *Arabidopsis*. *Plant J.* **76**, 433-445.
- Staiger, D., and Brown, J.W. (2013). Alternative splicing at the intersection of biological timing, development, and stress responses. *Plant Cell* **25**, 3640-3656.
- Sulpice, R., Flis, A., Ivakov, A.A., Apelt, F., Krohn, N., Encke, B., Abel, C., Feil, R., Lunn, J.E., and Stitt, M. (2014). *Arabidopsis* coordinates the diurnal regulation of carbon allocation and growth across a wide range of photoperiods. *Mol. Plant* **7**, 137-155.
- Sun, Y.L., and Hong, S.K. (2013). Sensitivity of translation initiation factor eIF1 as a molecular target of salt toxicity to sodic-alkaline stress in the halophytic grass *Leymus chinensis*. *Biochem. Genet.* **51**, 101-118.
- Suragani, M., Rasheedi, S., Hasnain, S.E., and Ehtesham, N.Z. (2011). The translation initiation factor, PeIF5B, from *Pisum sativum* displays chaperone activity. *Biochem. Biophys. Res. Commun.* **414**, 390-396.
- Syed, N.H., Kalyna, M., Marquez, Y., Barta, A., and Brown, J.W. (2012). Alternative splicing in plants--coming of age. *Trends Plant Sci.* **17**, 616-623.
- Szamecz, B., Rutkai, E., Cuchalová, L., Munzarová, V., Herrmannová, A., Nielsen, K.H., Burela, L., Hinnebusch, A.G., and Valásek, L. (2008). eIF3a cooperates with sequences 5' of uORF1 to promote resumption of scanning by post-termination ribosomes for reinitiation on GCN4 mRNA. *Genes Dev.* **22**, 2414-2425.
- Szick, K., Springer, M., and Bailey-Serres, J. (1998). Evolutionary analyses of the 12-kDa acidic ribosomal P-proteins reveal a distinct protein of higher plant ribosomes. *Proc. Natl. Acad. Sci. USA* **95**, 2378-2383.
- Szick-Miranda, K., and Bailey-Serres, J. (2001). Regulated heterogeneity in 12-kDa P-protein phosphorylation and composition of ribosomes in maize (*Zea mays* L.). *J. Biol. Chem.* **276**, 10921-10928.
- Tajrish, M.M., Vaid, N., Tuteja, R., and Tuteja, N. (2011). Overexpression of a pea DNA helicase 45 in bacteria confers salinity stress tolerance. *Plant Signal. Behav.* **6**, 1271-1275.
- Tcherkezian, J., Cargnello, M., Romeo, Y., Huttlin, E.L., Lavoie, G., Gygi, S.P., and Roux, P.P. (2014). Proteomic analysis of cap-dependent translation identifies LARP1 as a key regulator of 5'TOP mRNA translation. *Genes Dev.* **28**, 357-371.
- Thiebauld, O., Schepetilnikov, M., Park, H.S., Geldreich, A., Kobayashi, K., Keller, M., Hohn, T., and Ryabova, L.A. (2009). A new plant protein interacts with eIF3 and 60S to enhance virus-activated translation re-initiation. *EMBO J.* **28**, 3171-3184.
- Thoreen, C.C., Chantranupong, L., Keys, H.R., Wang, T., Gray, N.S., and Sabatini, D.M. (2012). A unifying model for mTORC1-mediated regulation of mRNA translation. *Nature* **485**, 109-113.
- Tiller, N., and Bock, R. (2014). The translational apparatus of plastids and its role in plant development. *Mol. Plant* **7**, 1105-1120.
- Timmer, R.T., Lax, S.R., Hughes, D.L., Merrick, W.C., Ravel, J.M., and Browning, K.S. (1993). Characterization of wheat germ protein synthesis initiation factor eIF-4C and comparison of eIF-4C from wheat germ

- and rabbit reticulocytes. *J. Biol. Chem.* **268**, 24863-24867.
- Tiruneh, B.S., Kim, B.H., Gallie, D.R., Roy, B., and von Arnim, A.G.** (2013). The global translation profile in a ribosomal protein mutant resembles that of an eIF3 mutant. *BMC Biol.* **11**, 123.
- Topisirovic, I., Svitkin, Y.V., Sonenberg, N., and Shatkin, A.J.** (2011). Cap and cap-binding proteins in the control of gene expression. *Wiley Interdiscip. Rev. RNA* **2**, 277-298.
- Turck, F., Zilbermann, F., Kozma, S.C., Thomas, G., and Nagy, F.** (2004). Phytohormones participate in an S6 kinase signal transduction pathway in *Arabidopsis*. *Plant Physiol.* **134**, 1527-1535.
- Turkina, M.V., Klang Årstrand, H., and Vener, A.V.** (2011). Differential phosphorylation of ribosomal proteins in *Arabidopsis thaliana* plants during day and night. *PLoS One* **6**, e29307.
- Ude, S., Lassak, J., Starosta, A.L., Kraxenberger, T., Wilson, D.N., and Jung, K.** (2013). Translation elongation factor EF-P alleviates ribosome stalling at polyproline stretches. *Science* **339**, 82-85.
- Vain, P., Thole, V., Worland, B., Opanowicz, M., Bush, M.S., and Doonan, J.H.** (2011). A T-DNA mutation in the RNA helicase eIF4A confers a dose-dependent dwarfing phenotype in *Brachypodium distachyon*. *Plant J.* **66**, 929-940.
- Valasek, L.S.** (2012). 'Ribozoomin'--translation initiation from the perspective of the ribosome-bound eukaryotic initiation factors (eIFs). *Curr. Protein Pept. Sci.* **13**, 305-330.
- Valenzuela, D.M., Chaudhuri, A., and Maitra, U.** (1982). Eukaryotic ribosomal subunit anti-association activity of calf liver is contained in a single polypeptide chain protein of Mr = 25,500 (eukaryotic initiation factor 6). *J. Biol. Chem.* **257**, 7712-7719.
- Valouev, I.A., Kushnirov, V.V., and Ter-Avanesyan, M.D.** (2002). Yeast polypeptide chain release factors eRF1 and eRF3 are involved in cytoskeleton organization and cell cycle regulation. *Cell Motil. Cytoskeleton* **52**, 161-173.
- van Heerden, A., and Browning, K.S.** (1994). Expression in *Escherichia coli* of the two subunits of the isozyme form of wheat germ protein synthesis initiation factor 4F. Purification of the subunits and formation of an enzymatically active complex. *J. Biol. Chem.* **269**, 17454-17457.
- Verma, D.P.S., and Chatterjee, J.** (2009). TORing with cell cycle, nutrients, stress and growth. In *Signal Crosstalk in Plant Stress Responses*, K. Yoshioka and K. Shinozaki, eds (New York, NY: Wiley), pp. 161-180.
- Verschoor, A., Srivastava, S., Grassucci, R., and Frank, J.** (1996). Native 3D structure of eukaryotic 80s ribosome: morphological homology with *E. coli* 70S ribosome. *J. Cell Biol.* **133**, 495-505.
- Voigts-Hoffmann, F., Klinge, S., and Ban, N.** (2012). Structural insights into eukaryotic ribosomes and the initiation of translation. *Curr. Opin. Struct. Biol.* **22**, 768-777.
- von Arnim, A.G., and Chamovitz, D.A.** (2003). Protein homeostasis: a degrading role for Int6/eIF3e. *Curr. Biol.* **13**, R323-325.
- von Arnim, A.G., Jia, Q., and Vaughn, J.N.** (2014). Regulation of plant translation by upstream open reading frames. *Plant Sci.* **214**, 1-12.
- Walker, S.E., Zhou, F., Mitchell, S.F., Larson, V.S., Valasek, L., Hinnebusch, A.G., and Lorsch, J.R.** (2012). Yeast eIF4B binds to the head of the 40S ribosomal subunit and promotes mRNA recruitment through its N-terminal and internal repeat domains. *RNA* **9**, 191-207.
- Wamboldt, Y., Mohammed, S., Elowsky, C., Wittgren, C., de Paula, W.B., and Mackenzie, S.A.** (2009). Participation of leaky ribosome scanning in protein dual targeting by alternative translation initiation in higher plants. *Plant Cell* **21**, 157-167.
- Wang, A., and Krishnaswamy, S.** (2012). Eukaryotic translation initiation factor 4E-mediated recessive resistance to plant viruses and its utility in crop improvement. *Mol. Plant Pathol.* **13**, 795-803.
- Wang, G., Zhang, J., Fan, X., Sun, X., Qin, H., Xu, N., Zhong, M., Qiao, Z., Tang, Y., and Song, R.** (2014). *Proline responding1* plays a critical role in regulating general protein synthesis and the cell cycle in maize. *Plant Cell* **26**, 2582-2600.
- Wang, J., Lan, P., Gao, H., Zheng, L., Li, W., and Schmidt, W.** (2013). Expression changes of ribosomal proteins in phosphate- and iron-deficient *Arabidopsis* roots predict stress-specific alterations in ribosome composition. *BMC Genomics* **14**, 783.
- Wang, L., Xu, C., Wang, C., and Wang, Y.** (2012). Characterization of a eukaryotic translation initiation factor 5A homolog from *Tamarix androssowii* involved in plant abiotic stress tolerance. *BMC Plant Biol.* **12**, 118.
- Wang, T.W., Lu, L., Zhang, C.G., Taylor, C., and Thompson, J.E.** (2003). Pleiotropic effects of suppressing deoxyhypusine synthase expression in *Arabidopsis thaliana*. *Plant Mol. Biol.* **52**, 1223-1235.
- Wang, X., and Grumet, R.** (2004). Identification and characterization of proteins that interact with the carboxy terminus of poly(A)-binding protein and inhibit translation *in vitro*. *Plant Mol. Biol.* **54**, 85-98.
- Warner, J.R., and McIntosh, K.B.** (2009). How common are extraribosomal functions of ribosomal proteins? *Mol. Cell* **34**, 3-11.
- Webster, C., Gaut, R.L., Browning, K.S., Ravel, J.M., and Roberts, J.K.M.** (1991). Hypoxia enhances phosphorylation of eukaryotic initiation factor 4A in maize root tips. *J. Biol. Chem.* **266**, 23341-23346.
- Wei, C.C., Balasta, M.L., Ren, J.H., and Goss, D.J.** (1998). Wheat germ poly(A) binding protein enhances the binding affinity of eukaryotic initiation factor 4F and (iso)4F for cap analogues. *Biochemistry* **37**, 1910-1916.
- Wei, Z., Xue, Y., Xu, H., and Gong, W.** (2006). Crystal structure of the C-terminal domain of *S. cerevisiae* eIF5. *J. Mol. Biol.* **359**, 1-9.
- Williams, A.J., Werner-Fraczek, J., Chang, I.F., and Bailey-Serres, J.** (2003). Regulated phosphorylation of 40S ribosomal protein S6 in root tips of maize. *Plant Physiol.* **132**, 2086-2097.
- Wilson, D.N., and Doudna Cate, J.H.** (2012). The structure and function of the eukaryotic ribosome. *Cold Spring Harb. Perspect. Biol.* **4**, a011536.
- Wu, X., Liu, M., Downie, B., Liang, C., Ji, G., Li, Q.Q., and Hunt, A.G.** (2011). Genome-wide landscape of polyadenylation in *Arabidopsis* provides evidence for extensive alternative polyadenylation. *Proc. Natl. Acad. Sci. USA* **108**, 12533-12538.
- Xia, C., Wang, Y.J., Li, W.Q., Chen, Y.R., Deng, Y., Zhang, X.Q., Chen, L.Q., and Ye, D.** (2010). The *Arabidopsis* eukaryotic translation initiation factor 3, subunit f (AtelF3f), is required for pollen germination and embryogenesis. *Plant J.* **63**, 189-202.
- Xiong, Y., and Sheen, J.** (2012). Rapamycin and glucose-target of rapamycin (TOR) protein signaling in plants. *J. Biol. Chem.* **287**, 2836-2842.
- Xiong, Y., and Sheen, J.** (2013). Moving beyond translation: glucose-TOR signaling in the transcriptional control of cell cycle. *Cell Cycle* **12**, 1989-1990.
- Xiong, Y., and Sheen, J.** (2014). The role of target of rapamycin signaling networks in plant growth and metabolism. *Plant Physiol.* **164**, 499-512.
- Xiong, Y., McCormack, M., Li, L., Hall, Q., Xiang, C., and Sheen, J.** (2013). Glucose-TOR signalling reprograms the transcriptome and activates meristems. *Nature* **496**, 181-186.
- Xue, S., and Barna, M.** (2012). Specialized ribosomes: a new frontier in gene regulation and organismal biology. *Nat. Rev. Mol. Cell Biol.* **13**, 355-369.
- Yahalom, A., Kim, T.H., Winter, E., Karniol, B., Von Arnim, A.G., and Chamovitz, D.A.** (2001). *Arabidopsis* eIF3e (INT-6) associates with both eIF3c and the COP9 signalosome subunit CSN7. *J. Biol. Chem.* **276**, 334-340.
- Yahalom, A., Kim, T.H., Roy, B., Singer, R., Von Arnim, A.G., and Chamovitz, D.A.** (2008). *Arabidopsis* eIF3e is regulated by the COP9 signalosome and has an impact on development and protein transla-

- tion. *Plant J.* **53**, 300-311.
- Yamamoto, Y.Y., Yoshitsugu, T., Sakurai, T., Seki, M., Shinozaki, K., and Obokata, J.** (2009). Heterogeneity of *Arabidopsis* core promoters revealed by high-density TSS analysis. *Plant J.* **60**, 350-362.
- Yumak, H., Khan, M.A., and Goss, D.J.** (2010). Poly(A) tail affects equilibrium and thermodynamic behavior of tobacco etch virus mRNA with translation initiation factors eIF4F, eIF4B and PABP. *Biochim. Biophys. Acta.* **1799**, 653-658.
- Yusupova, G., and Yusupov, M.** (2014). High-resolution structure of the eukaryotic 80S ribosome. *Annu. Rev. Biochem.* **83**, 467-486.
- Zanetti, M.E., Chang, I.F., Gong, F., Galbraith, D.W., and Bailey-Serres, J.** (2005). Immunopurification of polyribosomal complexes of *Arabidopsis* for global analysis of gene expression. *Plant Physiol.* **138**, 624-635.
- Zhang, J., Mao, Z., and Chong, K.** (2013). A global profiling of uncapped mRNAs under cold stress reveals specific decay patterns and endonucleolytic cleavages in *Brachypodium distachyon*. *Genome Biol.* **14**, R92.
- Zhang, Y., Liu, S., Lajoie, G., and Merrill, A.R.** (2008a). The role of the diphthamide-containing loop within eukaryotic elongation factor 2 in ADP-ribosylation by *Pseudomonas aeruginosa* exotoxin A. *Biochem. J.* **413**, 163-174.
- Zhang, Y., Wang, Y., Kanyuka, K., Parry, M.A., Powers, S.J., and Halford, N.G.** (2008b). GCN2-dependent phosphorylation of eukaryotic translation initiation factor-2 α in *Arabidopsis*. *J. Exp. Bot.* **59**, 3131-3141.
- Zhang, Y.H., Dickinson, J.R., Paul, M.J., and Halford, N.G.** (2003). Molecular cloning of an *Arabidopsis* homologue of GCN2, a protein kinase involved in co-ordinated response to amino acid starvation. *Planta* **217**, 668-675.
- Zhou, C., Arslan, F., Wee, S., Krishnan, S., Ivanov, A.R., Oliva, A., Leatherwood, J., and Wolf, D.A.** (2005). PCI proteins eIF3e and eIF3m define distinct translation initiation factor 3 complexes. *BMC Biol.* **3**, 14.
- Zhou, F., Roy, B., and Von Arnim, A.G.** (2010a). Translation reinitiation and development are compromised in similar ways by mutations in translation initiation factor eIF3h and the ribosomal protein RPL24. *BMC Plant Biol.* **10**, 193.
- Zhou, F., Roy, B., Dunlap, J.R., Enganti, R., and von Arnim, A.G.** (2014a). Translational control of *Arabidopsis* meristem stability and organogenesis by the eukaryotic translation factor eIF3h. *PLoS One* **9**, e95396.
- Zhou, F., Walker, S.E., Mitchell, S.F., Lorsch, J.R., and Hinnebusch, A.G.** (2014b). Identification and characterization of functionally critical, conserved motifs in the internal repeats and N-terminal domain of yeast translation initiation factor 4B (yeIF4B). *J. Biol. Chem.* **289**, 1704-1722.
- Zhou, X., Cooke, P., and Li, L.** (2010b). Eukaryotic release factor 1-2 affects *Arabidopsis* responses to glucose and phytohormones during germination and early seedling development. *J. Exp. Bot.* **61**, 357-367.
- Zhou, X., Sun, T.H., Wang, N., Ling, H.Q., Lu, S., and Li, L.** (2010c). The cauliflower *Orange* gene enhances petiole elongation by suppressing expression of eukaryotic release factor 1. *New Phytol.* **190**, 89-100.
- Zoncu, R., Efeyan, A., and Sabatini, D.M.** (2011). mTOR: from growth signal integration to cancer, diabetes and ageing. *Nat. Rev. Mol. Cell Biol.* **12**, 21-35.