

A Key to the Pacific Genera of Eumeninae (Hymenoptera: Vespidae)

Author: Carpenter, James M.

Source: American Museum Novitates, 2023(3995) : 1-16

Published By: American Museum of Natural History

URL: <https://doi.org/10.1206/3995.1>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

A Key to the Pacific Genera of Eumeninae (Hymenoptera: Vespidae)

JAMES M. CARPENTER¹

ABSTRACT

A key to the genera of the Eumeninae occurring in Oceania is presented. Hitherto there has been no published key to these genera, making routine identification difficult. New combinations are *Pararrhynchium atrum emifasciatum* (Giordani Soika), *Pararrhynchium simillimum* (Giordani Soika), *Parodynerus mariannensis* (Bequaert and Yasumatsu), and *Phimenes solomonis malaitensis* (van der Vecht).

INTRODUCTION

The potter wasp fauna of Oceania has been very incompletely studied. These wasps have been included in historical lists of some prominent islands, e.g., Fiji (Turner, 1919; Williams, 1947; Fullaway, 1957) and Samoa (Perkins and Cheesman, 1928), and are listed in contemporary treatments of, e.g., the Cook Islands (Kuhlmann, 2006) and French Polynesia (Ramage, 2017). There have been descriptive papers that included several taxa from a subregion, e.g., Micronesia (Bequaert and Yasumatsu, 1939; Yasumatsu, 1945). Giordani Soika (1958) provided a systematic treatment of the fauna of southern Polynesia. Carpenter (2008) revised the fauna of Hawaii, which is by far the most species-rich part of Polynesia. The aim of the present paper is to extend that treatment to other parts of Oceania by providing a key to genera and taxonomic notes with new combinations.

For the purposes of the present paper Oceania is construed to comprise Melanesia (less New Guinea), Micronesia and Polynesia. Australasia is not included because Australia has been recently treated (Carpenter and Brown, 2021a, 2021b) and New Zealand has no native vespidae fauna. New Guinea is excluded because the Indonesian part has been treated by Nugroho et al. (2012).

¹ American Museum of Natural History, Division of Invertebrate Zoology, New York.

KEY TO GENERA OF PACIFIC EUMENINAE

1. Metasoma petiolate: segment I in dorsal view with width half or less that of II, and at least twice as long as wide, usually longer (figs. 1, 3-4).....Tribe Eumenini, 2
- Metasoma not petiolate: segment I with width more than half that of II, much less than twice as long as wide (figs. 2, 14-16, 19). Tribe Odynerini, 4
2. Pronotum with pretegular carina absent (fig. 24); Tergum II with translucent apical lamella (fig. 3). *Eumenes* Latreille
- Pronotum with pretegular carina present (fig. 23); Tergum II without apical lamella (fig. 4)... . . 3
3. Tergum I with section after spiracles shorter than section before spiracles. . . . *Delta* de Saussure
- TI with section after spiracles longer than section before spiracles. . . . *Phimenes* Giordani Soika
4. Forewing with second submarginal cell petiolate (fig. 5)... *Paralastor* de Saussure
- Forewing with second submarginal cell not petiolate (figs. 6, 7). 5
5. Tegula campanulate: laterally semicircular in shape (fig. 8). 6
- Tegula not evenly curved laterally, with length exceeding width (fig. 10, 17, 18, 22)... 8
6. Submarginal carina and propodeal vavlula fused, with notch between them and not projecting (fig. 9); female cephalic foveae absent; anterior face of pronotum usually with weak impressions. *Nesodynerus* Perkins
- Submarginal carina and propodeal vavlula not fused, projecting, elongate (fig. 12); female with cephalic foveae (fig. 11); anterior face of pronotum with two close-set, deeply impressed pits (fig. 13)... 7
7. Tergum I with two transverse carinae; Tergum II without acarinarium, ridged basally (fig. 14)... *Subancistrocerus* de Saussure
- Tergum I with one rudimentary carina (figs. 15, 16); Tergum II smooth basally (fig. 15), forming acarinarium beneath apex of first tergum that is often full of mites (often concealed, tergum should be bent backward to expose acarinarium). *Parancistrocerus* Bequaert
8. Tergum I transversely carinate (figs. 15, 16). *Pararrhynchium* de Saussure
- TI not carinate (figs. 2, 14, 19)... 9
9. Tegula shorter than parategula posteriorly (figs. 10, 17); axillary fossa in dorsal view much narrower than long, slitlike (fig. 17). 10
- Tegula usually exceeding parategula posteriorly (figs. 18, 22), or at least equaling it; axillary fossa in dorsal view not slitlike, at least as wide as long, oval (fig. 18)... 12
10. Metasomal segment I subcylindrical, petiolate basally before expanding abruptly to a width slightly more than half that of segment II (fig. 19); color bright metallic... *Eudiscoelius* Friese
- Metasomal segment I sessile, about as wide as II; color not bright metallic (figs. 2, 14-16)... . . 11
11. Scutum posteriorly and scutellum impunctate; metanotum depressed medially; male midfemur basally emarginate (fig. 20)... *Rhynchium* Spinola
- Scutum and scutellum punctate; metanotum not depressed; male midfemur not basally emarginate. *Anterrhynchium* de Saussure
12. Tegula long, with posterior lobe covering parategula and pointed medially (fig. 21)... *Knemodynerus* Blüthgen
- Tegula shorter, with posterior lobe not covering parategula, never pointing medially (fig. 18). . . . 13
13. Propodeum without superior carinae; second submarginal cell with second recurrent vein nearly or completely interstitial with third submarginal cell (fig. 7). *Epsilon* de Saussure
- Propodeum with superior carinae (fig. 22; sometimes faint); second submarginal cell with second recurrent vein far from third submarginal cell (figs. 5-6). 14
14. Pronotum with oblique humeral carina (fig. 23)... *Pachodynerus* de Saussure

- Pronotum without humeral carina (fig. 24).....15
- 15. Tegula exceeding parategula (fig. 18); Tergum I about as wide as Tergum II (fig. 16); punctuation strong. *Euodynerus* Dalla Torre
- Tegula not exceeding parategula; Tergum I narrower than II (fig. 2); punctuation reduced.
..... *Parodynerus* de Saussure

TAXONOMIC NOTES

Tribe Eumenini Leach

Genus *Delta* de Saussure

Delta de Saussure, 1855: 130, 132, 143, name for divisions II and III of genus *Eumenes* Latreille in de Saussure, 1852: 44, 60 (26 + 3 species).

Type species: *Vespa maxillosa* DeGeer, 1775 [= *Vespa emarginata* Linnaeus, 1758], by subsequent designation of Bequaert, 1925a: 137 [erroneously as *Sphex maxillosus* DeGeer, correctly in 1926: 487].

Erinys Zirngiebl, 1953: 173, subgenus of *Eumenes* Latreille (junior homonym of *Erinys* Rye, 1876).

Type species: *Vespa unguiculata* Villers, 1789, by monotypy.

Alfieria Giordani Soika, 1934: 436, genus.

Type species: *Eumenes anomalus* Zavattari, 1909, by original designation and monotypy.

This is an Old World genus, consisting of nearly 50 species, with most of them being African or Asian. All of the species in Hawaii are introduced, as is one in New Caledonia, but there are a half dozen indigenous species distributed from the Solomon Islands to Fiji: *Delta campaniforme* (Fabricius) and *D. pyriforme philippinense* (Bequaert) are adventive in Hawaii (Carpenter, 2008), *D. esuriens esuriens* (Fabricius) is adventive in New Caledonia (Gusenleitner and Madl, 2011), and *D. esuriens okinawae* Giordani Soika adventive in the Cook Islands (Kuhlmann, 2006) (and probably Palau). *Delta latreillei petiolare* (Schulz) is adventive in Hawaii as well (and Fiji; Evenhuis, 2007), but is presumably indigenous to the Admiralty Islands, Trobriand Islands, and Solomon Islands (it is also found in New Guinea); *D. latreillei silaceum* (van der Vecht) is endemic to the Solomon Islands, New Britain and New Ireland (van der Vecht, 1960). *Delta insulare* (Smith) is endemic to Fiji; *D. pagdeni* (van der Vecht) and *D. versicolor* van der Vecht are both endemic to the Solomon Islands (van der Vecht, 1981); and six subspecies of *D. xanthura* (de Saussure) are found in Vanuatu and New Caledonia (Giordani Soika, 1958).

Genus *Eumenes* Latreille

Eumenes Latreille, 1802: 360, genus.

Type species: "*Eumenes coarctata* Fab." [= *Vespa coarctata* Linnaeus, 1758], by subsequent designation of Latreille, 1810: 438.




FIG. 1. First metasomal segment in dorsal view, *Delta campaniforme*. Fig. 2. Metasoma in dorsal view, *Nesodynerus rudolphi*. Figs. 3, 4. Metasoma in lateral view: 3. *Pachymenes novarae*; 4. *Pachymenes ater*. Figs. 5, 6. Forewing: 5. *Paralastor* sp.; 6. *Diemodynerus pseudacarodynerus*. Scale bars = 1 mm.




FIG. 7. Forewing, *Pararhaphidoglossa invenusta*. Fig. 8. Tegula, *Parancistrocerus pedestris*. Fig. 9. Mesosoma in lateral view, *Epsilon laboriosum*. Fig. 10. Tegula, *Rhynchium superbum*. Fig. 11. Head in dorsal view, *Ectopioglossa polita australensis*. Fig. 12. Mesosoma in oblique lateral view, *Leptochilus rufinodius*. Scale bars = 1 mm.




FIG. 13. head and pronotum in anterodorsal view, *Subancistrocerus monsticornis*. Figs. 14, 15. Metasoma segments in dorsolateral view: 14. *Stenodynerus papagorum*; 15. *Parancistrocerus* sp. Fig. 16. Metasoma in dorsal view, *Ancistroceroides levis*. Figs. 17, 18. Mesosoma in dorsal view: 17. *Stenonartonia apicipennis*; 18. *Ancistrocerus flavomarginatus*. Scale bars = 1 mm.




FIG. 19. Metasoma in dorsal view, *Eudiscoelius jacinthae*. Fig. 20. Midleg, *Rhynchium superbum*. Fig. 21. Mesosoma in dorsal view, *Knemodynerus excellens*. Fig. 22. Propodeum in dorsoposterior view, *Pseudodynerus griseolus*. Figs. 23, 24. Pronotum in anterolateral view; 23. *Pachodynerus nasidens*. 24. *Euodynerus localis*. Scale bars = 1 mm.

Alpha de Saussure, 1855: 128, 137, name for division I of genus *Eumenes* Latreille in de Saussure, 1852: 28 (junior homonym of *Alpha* de Saussure, 1854).

Type species: *Vespa coarctata* Linnaeus, 1758, by subsequent designation of Bequaert, 1926: 485.

Eumenis Kriechbaumer, 1879: 57 (unjustified emendation of *Eumenes* Latreille).

Eumenidion von Schulthess, 1913: 2, subgenus of *Eumenes* Latreille.

Type species: "*Eumenes coarctatus* L." [= *Vespa coarctata* Linnaeus, 1758], by original designation.

The genus is cosmopolitan, with more than 100 described species. A single species, *Eumenes mediterraneus* Kriechbaumer, has been introduced into Tahiti. This is a Palearctic species, first reported by Cheesman (1928) as *E. pomiformis*. Gusenleitner (2011) identified the species from Tahiti.

Genus *Phimenes* Giordani Soika

Phimenes Giordani Soika, 1992: 41, 66, genus, replacement name for *Phi* de Saussure, 1855, non de Saussure, 1854.

Type species: *Vespa arcuata* Fabricius, 1775, by subsequent designation of Bequaert, 1926: 487, as type species of *Phi* de Saussure, 1855.

Phi de Saussure, 1855: 132, name for division IV of genus *Eumenes* Latreille in de Saussure, 1852: 63, 145 (junior homonym of *Phi* de Saussure, 1854).

Type species: *Vespa arcuata* Fabricius, 1775, by subsequent designation of Bequaert, 1926: 487.

This is a primarily Asian genus consisting of 15 species, with one species, *Phimenes solomonis* (van der Vecht), in the Solomon Islands (van der Vecht, 1959), with two subspecies, one of which is a new combination in *Phimenes*: *P. solomonis malaitensis* (van der Vecht), NEW COMBINATION, and *P. solomonis solomonis* (van der Vecht). *Phimenes curvatus* (de Saussure) was introduced into Hawaii, and I earlier (Carpenter, 2008) suggested that it may no longer occur there, but later I collected it on Kauai.

Tribe Odynerini Lepeletier

Genus *Anterhynchium* de Saussure

Anterhynchium de Saussure, 1863: 205, name for division I of genus *Rhynchium* Spinola in de Saussure, 1852: 103, and 1855: 175.

Type species: *Rygius synagroides* de Saussure, 1852, by subsequent designation of van der Vecht, 1963: 73.

This is an Old World genus consisting of nearly 50 species. There are four species in the subgenus *Epiodynerus* distributed from Indonesia, Australia, and the Solomon Islands (*Anterhynchium nigrocinctum* (de Saussure); *Anterhynchium woodfordi* (Meade-Waldo) is endemic to

the Solomon Islands) to Vanuatu (*A. alecto lalepi* (Giordani Soika) and *A. alecto parallelum* (Giordani Soika)) and New Caledonia (*A. alecto alecto* (Lepelletier) and *A. chozali* (Cheesman)), and further to Fiji and French Polynesia (*A. rufipes* (Fabricius)) (see Giordani Soika, 1958; van der Vecht, 1963; Gusenleitner, 2011; Gusenleitner and Madl, 2011; Carpenter and Brown, 2021a).

Genus *Epsilon* de Saussure

Epsilon de Saussure, 1855: 229, 252, name for division III of subgenus *Leionotus* de Saussure of genus *Odynerus* Latreille in de Saussure, 1852: 169; declared available from date of publication by Opinion 893 (ICZN, 1970).

Type species: *Odynerus dyscherus* de Saussure, 1852, by subsequent designation of van der Vecht, 1967: 31; confirmed by Opinion 893 (ICZN, 1970).

This is an Oriental and Australian genus consisting of nearly 20 species. There is one species in the Solomon Islands (Giordani Soika, 1994), *Epsilon achterbergi* Giordani Soika.

Genus *Eudiscoelius* Friese

Eudiscoelius Friese, 1904: 16, genus.

Type species: *Eudiscoelius metallicus* Friese, 1904, by monotypy.

Euchalcomenes Turner, 1908: 90, genus.

Type species: *Euchalcomenes gilberti* Turner, 1908, by original designation.

This is an Oriental and Australian genus consisting of 12 species. There are four species in the Bismarck and Solomon Islands (Giordani Soika, 1994): *Eudiscoelius bismarcki* Giordani Soika is from New Britain and the Solomon Islands; *E. ferrugineipes* Giordani Soika, *E. lucens* Giordani Soika and *E. solomon* Giordani Soika are all endemic to the Solomon Islands.

Genus *Euodynerus* Dalla Torre

Euodynerus Dalla Torre, 1904: 38, name for section II of division III of subgenus *Leionotus* of genus *Odynerus* Latreille in de Saussure, 1853: 177; declared available from date of publication by Opinion 893 (ICZN, 1970).

Type species: *Vespa dantici* Rossi, 1790, by subsequent designation of Blüthgen, 1938: 277; confirmed by Opinion 893 (ICZN, 1970).

This is a cosmopolitan genus, of more than 100 species, with four species of the nominate subgenus endemic to Hawaii (Carpenter, 2008): *Euodynerus epipseustes* (Perkins), *E. localis* (Smith), *E. nigripennis* (Holmgren) and *E. radula* (Fabricius).

Genus *Knemodynerus* Blüthgen

Knemodynerus Blüthgen, 1940: 43, subgenus of “*Euodynerus* Blüthgen” [= *Euodynerus* Dalla Torre].

Type species: *Odynerus excellens* Pérez, 1907, by original designation.

This is an Old World genus consisting of more than 40 species. There are three species in Oceania: *Knemodynerus circumspectus derufatus* Giordani Soika was described from the Solomon Islands (Giordani Soika, 1994), and *K. paganensis* (Yasumatsu) and *K. saipanensis* (Yasumatsu) were described from the Marianas (Yasumatsu, 1945).

Genus *Nesodynerus* Perkins

Nesodynerus Perkins, 1901: 267, genus, in key.

Type species: *Odynerus rudolphi* Dalla Torre, 1889 (a replacement name for *Odynerus cardinalis* Blackburn, 1886, *non* Morawitz, 1885), by subsequent designation of Carpenter, 1986: 76.

Pseudopterocheilus Perkins, 1901: 266, genus.

Type species: *Odynerus pterocheiloides* Perkins, 1899, by original designation.

Chelodynerus Perkins, 1902: 136, genus.

Type species: *Odynerus chelififer* Perkins, 1899, by monotypy.

This genus is endemic to Hawaii (Carpenter, 2008). With more than 100 described species, it is much the largest eumenine genus in Oceania.

Genus *Pachodynerus* de Saussure

Pachodynerus de Saussure, 1870: 56, division of subgenus *Odynerus* of genus *Odynerus* Latreille; declared available from date of publication by Opinion 893 (ICZN, 1970).

Type species: *Odynerus californicus* de Saussure, 1870, by subsequent designation of Bohart, 1951: 892; confirmed by Opinion 893 (ICZN, 1970).

Monobiella Ashmead, 1900: 312, genus.

Type species: *Vespa atrata*, 1798, by monotypy.

This is an American genus consisting of nearly 50 species; the so-called keyhole wasp, *Pachodynerus nasidens* (Latreille), has been introduced on islands all over the Pacific (e.g., Krombein, 1949; Yamane et al., 1996; Gusenleitner, 2011), thanks to its indiscriminate choice of preexisting cavities as nesting sites. At Brisbane airport these nesting habits now present an aviation hazard (House et al., 2020a, 2020b)!

Genus *Paralastor* de Saussure

Paralastor de Saussure, 1856: 328, division of subgenus *Alastor* of genus *Alastor* Lepeletier; declared available from date of publication by Opinion 893 (ICZN, 1970).

Type species: *Alastor tuberculatus* de Saussure, 1853, by subsequent designation of van der Vecht, 1967: 31; confirmed by Opinion 893 (ICZN, 1970).

Alastoroides de Saussure, 1856: 327, subgenus of *Alastor* Lepeletier.

Type species: *Alastor clotho* Lepeletier, 1841, by subsequent designation of Ashmead, 1902: 210.

Paralastoroides de Saussure, 1856: 328, division of subgenus *Alastoroides* de Saussure of genus *Alastor* Lepeletier; declared available from date of publication by Opinion 893 (ICZN, 1970).

Type species: *Alastor clotho* Lepeletier, 1841, by monotypy.

This is a primarily Australian genus, consisting of well over 100 species (Carpenter and Brown, 2021a), with a single species described from Fiji, *Paralastor graeffei* (de Saussure) (see Perkins, 1914).

Genus *Parancistrocerus* Bequaert

Parancistrocerus Bequaert, 1925b: 64, subgenus of *Ancistrocerus* Wesmael.

Type species: *Odynerus fulvipes* de Saussure, 1855 [= *O. "flavipes* Fabricius" sensu de Saussure, 1852, non *Vespa flavipes* Fabricius, 1775], by original designation.

This genus consists of more than 100 species from the Americas and the Oriental Region. There are more than 50 species recorded from the Oriental Region (Kumar et al., 2016; Li and Carpenter, 2019), but the only Pacific species is the American species *Parancistrocerus fulvipes* (de Saussure), which has been introduced into Midway (Nishida and Beardsley, 2002).

Genus *Pararrhynchium* de Saussure

Pararrhynchium de Saussure, 1855: 173, division of genus *Rhynchium* Spinola.

Type species: *Rhynchium ornatum* Smith, 1852, by monotypy.

Prorrhynchium de Saussure, 1855: 174, division of genus *Rhynchium* Spinola. Rejected by van der Vecht, 1963: 94, acting as first reviser.

Type species: *Rhynchium smithii* de Saussure, 1855, by monotypy.

Lissodynerus Giordani Soika, 1973: 119, used as generic name for *Odynerus septemfasciatus* var. *feanus* Giordani Soika, 1941. Unavailable under Article 13.1.1 of the Code.

Trichodynerus Giordani Soika and Kojima, 1988: 178, used as a generic name in the combination *Trichodynerus agilis cursor* Giordani Soika and Kojima, 1988. Unavailable under Article 13.1.1 of the Code.

Lissodynerus Giordani Soika, 1993: 135, genus; reference to *Lissodynerus* Giordani Soika, 1973, and *Trichodynerus* Giordani Soika and Kojima, 1988.

Type species: *Odynerus septemfasciatus* Smith, 1857, by original designation.

Note: Redescribed as a "new" genus by Giordani Soika, 1994: 301.

This is a primarily Oriental genus, consisting of more than 30 species. Three species have been described, in the genus *Lissodynerus*, from the Solomon Islands (Giordani Soika, 1994): *Pararrhynchium atrum atrum* (Giordani Soika), *P. atrum emifasciatum* (Giordani Soika) NEW COMBINATION, *P. simillimum* (Giordani Soika) NEW COMBINATION and *P. solomon* (Giordani Soika).

Genus *Parodynerus* de Saussure

Parodynerus de Saussure, 1855: 245, name for division I of subgenus *Odynerus* of genus *Odynerus* Latreille, in de Saussure, 1852: 155; declared available from date of publication by Opinion 893 (ICZN, 1970).

Type species: “*Odynerus bicinctus* (F.)” [= *Vespa bicincta* Fabricius, 1781], by subsequent designation of Giordani Soika, 1958: 214, who in the same paper listed the originally included *Odynerus bizonatus* Boisduval, 1833, as a synonym of *Odynerus bicinctus* (F.); confirmed by Opinion 893 (ICZN, 1970).

This genus is endemic to Oceania, with a total of five species: four in Fiji (*P. bicinctus* (Fabricius), *P. mediocinctus mediocinctus* (Turner), *P. mediocinctus taveunensis* Giordani Soika, and *P. nigropetiolatus nigropetiolatus* Giordani Soika); two in Samoa (*P. bicinctus* (Fabricius) and *P. quodi quodi* (Vachal)); two in Tonga (*P. bicinctus* (Fabricius) and *P. quodi quodi* (Vachal)); one in Tuvalu (*P. bicinctus* (Fabricius)); five in Vanuatu (*P. cheesmani* Giordani Soika, *P. nigropetiolatus nigerrimus* Giordani Soika, *P. quodi hebridensis* Giordani Soika, *P. quodi iasafui* Giordani Soika, and *P. quodi malekulensis* Giordani Soika); two in New Caledonia (*P. quodi lifuensis* Giordani Soika and *P. quodi quodi* (Vachal)); one in the Cook Islands (*P. bicinctus* (Fabricius)); and two in French Polynesia (*P. bicinctus* (Fabricius) and *P. quodi quodi* (Vachal)) (see Giordani Soika, 1958, 1971). A sixth species is a new combination in the genus, from the Marianas, *Parodynerus mariannensis* (Bequaert and Yasumatsu), NEW COMBINATION (Bequaert and Yasumatsu, 1939).

Genus *Rhynchium* Spinola

Rhynchium Spinola, 1806, Ins. Ligur. 1: 84, genus, emendation of *Ryghium* validated by Opinion 747 (ICZN, 1965) (no. 1688 of Official List of Generic Names in Zoology).

Type species: *Ryghium europaeum* Spinola, 1806 [= *Vespa oculata* Fabricius, 1781], by monotypy (no. 2095 of Official List of Specific Names in Zoology).

Ryghium Spinola, 1806: 84, genus. Incorrect original spelling of *Rhynchium* Spinola, placed on Official Index of Rejected and Invalid Generic Names in Zoology (as no. 1768) by Opinion 747 (ICZN, 1965).

Rhynchium Billberg, 1820: 109. Emendation of *Ryghium* [!] Spinola; validated and placed on Official List of Generic Names in Zoology (as no. 1688) by Opinion 747 (ICZN, 1965).

Ryghium [!] Billberg, 1820: 109. Placed on Official Index of Rejected and Invalid Generic Names in Zoology (as no. 1769) by Opinion 747 (ICZN, 1965).

Rynchium Sturm, 1829: 12. Unjustified emendation of *Rhynchium* Spinola, placed on Official Index of Rejected and Invalid Generic Names in Zoology (as no. 1770) by Opinion 747 (ICZN, 1965).

Rhyghium [!] de Saussure, 1853: xxxi, 276. Placed on Official Index of Rejected and Invalid Generic Names in Zoology (as no. 1771) by Opinion 747 (ICZN, 1965).

Eurrhynchium Dalla Torre, 1904: 33, name for division II of *Rhynchium* Spinola in de Saussure, 1852: 105.

Type species: *Vespa oculata* Fabricius, 1781, by subsequent designation of van der Vecht and Carpenter, 1990: 23.

This Old World genus, consisting of more than 20 species, has one widespread species distributed from the Seychelles to the Marianas and possibly Palau (*Rhynchium brunneum brunneum* (Fabricius)), with two other species in the Solomon Islands (*R. salomonis* Meade-Waldo and *R. vittatum* du Buysson), and one in New Caledonia and French Polynesia (*R. quinquecinctum tahitense* de Saussure) (see Bequaert and Yasumatsu, 1939; van der Vecht, 1960; Madl et al., 1996; Gusenleitner, 2011).

Genus *Subancistrocerus* de Saussure

Subancistrocerus de Saussure, 1855: 206, name for division I of subgenus *Ancistrocerus* Wesmæl of genus *Odynerus* Latreille in de Saussure, 1852: 126, declared available from date of publication by Opinion 893 (ICZN, 1970).

Type species: *Odynerus sichelii* de Saussure, 1855, by subsequent designation of Bequaert, 1925b: 61; confirmed by Opinion 893 (ICZN, 1970).

Epancistrocerus de Saussure, 1856: 352, in Errata, substitute name for *Subancistrocerus* de Saussure.

Type species: *Odynerus sichelii* de Saussure, 1855, by subsequent designation of Bequaert, 1925b: 61.

This Old World genus, consisting of more than 30 species, has two species on Palau (*Subancistrocerus esakii* (Bequaert and Yasumatsu) and *S. palauensis* (Bequaert and Yasumatsu)), one in the Solomon Islands (*S. solomonis gizensis* Giordani Soika and *S. solomonis solomonis* Giordani Soika), and one on Yap (*S. yapensis* (Yasumatsu)) (see Giordani Soika, 1994).

ACKNOWLEDGMENTS

I thank Valerie Giles, Molly Rightmyer, and Ann Sanderson for assistance with illustrations and Lien Nguyen for reviewing the manuscript.

REFERENCES

- Ashmead, W.H. 1900. Report upon the Aculeate Hymenoptera of the island of St. Vincent and Grenada, with additions to the parasitic Hymenoptera and a list of the described Hymenoptera of the West Indies. Transactions of the Entomological Society of London 1900: 207–367.
- Ashmead, W.H. 1902. Classification of the fossorial, predacious and parasitic wasps, or the superfamily Vespoidea. (Paper no. 7). Family XXIX. Eumenidae. Canadian Entomologist 34: 203–210.
- Bequaert, J. 1925a. *Eumenes dyscherus* H. de Saussure, a Neotropical, not an African wasp, and other notes on synonymy. Bulletin of the Brooklyn Entomological Society 20: 134–140.

- Bequaert, J. 1925b. The genus *Ancistrocerus* (Hymenoptera, Vespidae) in North America, with a partial key to the species. Transactions of the American Entomological Society 57: 57–117.
- Bequaert, J. 1926. The genus *Eumenes* Latreille. South Africa, with a revision of the Ethiopian species (Hymenoptera). Annals of the South African Museum 23: 483–577.
- Bequaert, J., and K. Yasumatsu. 1939. Vespoidea of Micronesia. Tenthredo 2: 314–328.
- Billerg, G.J. 1820. Enumeratio insectorum in Museo. Stockholm: Typis Gadelianis.
- Blüthgen, P. 1938 (“1937”). Systematisches Verzeichnis der Faltenvespen Mitteleuropas, Skandinaviens und Englands. Konowia 16: 270–295.
- Blüthgen, P. 1940. Über *Odynerus meyeri* CAM., *Od. meyeri* CAM. var. *albolimbatus* SCHULTHESS und *Od. pseudolateralis* M.W. (Hym. Vespidae). Entomologisk Tidskrift 61 (1–2): 41–45.
- Bohart, R.M. 1951. Vespidae. In C.F. Muesebeck, K.V. Krombein and H.K. Townes (eds.), Hymenoptera of America North of Mexico synoptic catalog. United States Department of Agriculture, Agriculture Monograph 2: 875–907.
- Carpenter, J.M. 1986. A synonymic generic checklist of the Eumeninae (Hymenoptera: Vespidae). Psyche 93: 61–90.
- Carpenter, J.M. 2008. Review of the Hawaiian Vespidae (Hymenoptera). Bishop Museum Occasional Papers 99: 1–18.
- Carpenter, J.M., and G.R. Brown. 2021a. Catalogue of the Australian Eumeninae (Hymenoptera: Vespidae). Zootaxa 4919 (1): 1–68.
- Carpenter, J.M., and G.R. Brown. 2021b. A key to the Australian genera of Eumeninae (Hymenoptera: Vespidae). Records of the Australian Museum 73 (3): 87–101.
- Cheesman, L.E. 1928. A contribution towards the insect fauna of French Oceania, II. Annals and Magazine of Natural History (10) 1: 169–194.
- Dalla Torre, K.W. 1904. Hymenoptera Fam. Vespidae. Genera Insectorum 19: 1–108.
- Evenhuis, N.L. 2007. Checklist of Fiji Hymenoptera. Bishop Museum Technical Report 38 (11): 1–29.
- Friese, H. 1904. Eine metallisch gefarbte Vespidae. Zeitschrift für Systematische Hymenopterologie und Dipterologie 4: 16.
- Fullaway, D.T. 1957. Checklist of the Hymenoptera of Fiji. Proceedings of the Hawaiian Entomological Society 16: 269–280.
- Giordani Soika, A. 1934. Un nuovo genere egiziano di eumenini (Hymenoptera: Vespidae). Bulletin de la Societe Royale Entomologique d’Égypte 4: 436–439.
- Giordani Soika, A. 1941. Studi sui vespidi solitari. Bollettino della Società Veneziana di Storia Naturale 2 (3): 130–280.
- Giordani Soika, A. 1958 (“1957”). Biogeografia, evoluzione e sistematica dei vespidi solitari della Polinesia meridionale. Bollettino del Museo Civico di Storia Naturale di Venezia 10: 183–221.
- Giordani Soika, A. 1971. Notulae vespilogicae XXX. Nuovo contributo alla conoscenza degli eumenidi della Polinesia (Hymenoptera). Bollettino della Società Entomologica Italiana 103 (3–4): 78–80.
- Giordani Soika, A. 1973. Notulae vespilogicae XXXV. Descrizione di nuovi eumenidi. Bollettino del Museo Civico di Storia Naturale di Venezia 24: 97–131.
- Giordani Soika, A. 1992. Di alcuni eumenidi nuovi o poco noti (Hymenoptera Vespoidea). Lavori della Società Veneziana di Scienze Naturali 17: 41–68.
- Giordani Soika, A. 1993. Contributo alla conoscenza degli eumenidi dell’Australia e della Nuova Guinea (Hym. Vespoidea). Bollettino del Museo Civico di Storia Naturale di Venezia 42: 125–149.

- Giordani Soika, A. 1994. Ricerche sistematiche su alcuni generi di eumenidi della regione orientale e della Papuasias (Hymenoptera, Vespoidea). *Annali del Museo Civico di Storia Naturale "Giacomo Doria"* 90: 1–348.
- Giordani Soika, A., and J. Kojima. 1988. Eumeninae wasps collected in Papua New Guinea by J. Kojima (Hymenoptera Vespidae). *Bolletino del Museo Civico di Storia Naturale di Venezia* 38: 175–182.
- Gusenleitner, J. 2011. Über Vespidae auf pazifischen Inseln (Hymenoptera: Vespidae: Polistinae, Eumeninae). *Linzer Biologische Beiträge* 43 (2): 1291–1293.
- Gusenleitner, J., and M. Madl. 2011. Annotated catalogue of the Vespidae (Hymenoptera: Vespoidea) of New Caledonia. *Zeitschrift der Arbeitsgemeinschaft Österreichischer Entomologen* 63: 133–140.
- House, A.P.N., J.G. Ring, M.J. Hill and P.P. Shaw. 2020a. Insects and aviation safety: the case of the keyhole wasp *Pachodynerus nasidens* (Hymenoptera: Vespidae) in Australia. *Transportation Research Interdisciplinary Perspectives* 4: 100096.
- House, A.P.N., J.G. Ring, and P.P. Shaw. 2020b. Inventive nesting behaviour in the keyhole wasp *Pachodynerus nasidens* Latreille (Hymenoptera: Vespidae) in Australia, and the risk to aviation safety. *PLoS ONE* 15 (11): e0242063.
- ICZN. 1965. Opinion 747. *Rygius* Spinola 1806 (Insecta, Hymenoptera): Validation of emendation to *Rhynchium*. *Bulletin of Zoological Nomenclature* 22: 186–187.
- ICZN. 1970. Opinion 893. Eumenidae names of Saussure (Hymenoptera): Grant of availability to certain names proposed for secondary divisions of genera. *Bulletin of Zoological Nomenclature* 26: 187–191.
- Kriechbaumer, J. 1879. Eumeniden–Studien. *Entomologische Nachrichten* 5: 57–59 + 85–89 + 201–204 + 309–312.
- Krombein, K.V. 1949. The aculeate Hymenoptera of Micronesia. I. Scoliidae, Mutillidae, Pompilidae and Sphecidae. *Proceedings of the Hawaiian Entomological Society* 13: 367–410.
- Kuhlmann, M. 2006. Fauna and biogeography of the bees and wasps of the Cook Islands (Hymenoptera Aculeata). *Journal of Hymenoptera Research* 15 (1): 26–37.
- Kumar, P.G., J.M. Carpenter, and P.M. Sureshan. 2016. A taxonomic review of the genus *Parancistrocerus* Bequaert (Hymenoptera: Vespidae: Eumeninae) from the Indian subcontinent with the description of three new species. *Halteres* 7: 136–156.
- Latreille, P.A. 1802. *Histoire naturelle, générale et particulière des crustacés et des insectes*, 3. Paris: F. Dufart.
- Latreille, P.A. 1810. *Considérations générales sur l'ordre naturel des animaux composant les classes des crustacés, des arachnides, et des insectes*. Paris: F. Schoell.
- Li, T.-J., and J.M. Carpenter. 2019. Descriptions of eight new species of the genus *Parancistrocerus* Bequaert (Hymenoptera: Vespidae: Eumeninae), with a key to the Oriental species. *Zootaxa* 4551 (3): 251–274.
- Madl, M., P. Matyot, and S. Schödl. 1996. Vespidae and Sphecidae from the Seychelles (Insecta, Hymenoptera). *Linzer Biologische Beiträge* 28 (2): 829–834.
- Nishida, G.M., and J.W. Beardsley. 2002. A review of the insects and related arthropods of Midway Atoll. *Records of the Hawaii Biological Survey for 2000. Bishop Museum Occasional Papers* 68: 25–69.
- Nugroho, H., J. Kojima, and J.M. Carpenter. 2012 ("2011"). Checklist of vespid species (Insecta: Hymenoptera: Vespidae) occurring in Indonesian Archipelago. *Treubia* 38: 71–186.
- Perkins, R.C.L. 1901. Notes on Hawaiian aculeate Hymenoptera. *Entomologists Monthly Magazine* (2) 12: 264–268.

- Perkins, R.C.L. 1902. Notes on Hawaiian wasps, with descriptions of new species. *Transactions of the Entomological Society of London* 1902: 131–140.
- Perkins, R.C.L. 1914. On the species of *Alastor* (*Paralastor*) Sauss. and some other Hymenoptera of the family Eumenidae. *Proceedings of the Zoological Society of London* 1914: 563–624.
- Perkins, R.C.L., and L.E. Cheesman. 1928. Apoidea, Sphecoidea, and Vespoidea. *Insects of Samoa V. Hymenoptera*, Fasc. 1: 1–32.
- Ramage, T. 2017. Checklist of the terrestrial and freshwater arthropods of French Polynesia (Chelicerata; Myriapoda; Crustacea; Hexapoda). *Zoosystema* 39 (2): 213–225.
- Saussure, H.F. de. 1852–1858. Études sur la famille des vespides, vols. 1–3. V. Paris: Masson & J. Cherbuliez. [for publication dates of specific pages see F.J. Griffin, 1939, On the dates of publication of de Saussure (H. de): Etudes sur la famille des vespides 1–3. 1852–1858, in *Journal of the Society for the Bibliography of Natural History* 1: 211–212]
- Saussure, H. de. 1863. Mélanges hyménoptérologiques II. *Mémoires de la Société de Physique et d'histoire Naturelle de Genève* 17 (1): 171–244.
- Saussure, H. de. 1870. Vespidae americanae novae nonnullae. *Revue et Magasin de Zoologie* (2) 22: 55–62.
- Schulthess, A. von. 1913. Neue aethiopsiche Eumeniden. *Societas Entomologica* 28: 1–3, 6–78.
- Spinola, M. 1806. *Insectorum Liguriae*, 1. Genuae: Y. Gravier.
- Sturm, J. 1829. Verzeichniss einiger zum Umstausch oder Verkauf vorrathigen Insecten. Nurmberg: [self-published].
- Turner, R.E. 1908. Two new diplopterous Hymenoptera from Queensland. *Transactions of the Royal Entomological Society of London* 1908: 89–91.
- Turner, R.E. 1919. The Hymenoptera of Fiji. *Transactions of the Royal Entomological Society of London* (1918) 1919: 334–346.
- Vecht, J. van der, 1959. On *Eumenes arcuatus* (Fabricius) and some allied Indo-Australian wasps (Hymenoptera, Vespidae). *Zoologische Verhandelingen* 41: 1–71.
- Vecht, J. van der, 1960. On *Abispa* and some other Eumenidae from the Australian region (Hymenoptera, Vespidae). *Nova Guinea, Zoology* 6: 91–115.
- Vecht, J. van der. 1963. Studies on Indo-Australian and East Asiatic Eumenidae (Hymenoptera Vespidae). *Zoologische Verhandelingen* 60: 1–116.
- Vecht, J. van der. 1967. The status of certain genus-group names in the Eumenidae (Hymenoptera, Vespidae) Z.N. (S.) 1689. *Bulletin of Zoological Nomenclature* 24: 27–33.
- Vecht, J. van der. 1981. Studies of Indo-Australian solitary wasps. *Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen (Series C)* 84: 443–463.
- Vecht, J. van der, and J.M. Carpenter. 1990. A catalog of the genera of the Vespidae (Hymenoptera). *Zoologische Verhandelingen* 260: 3–62.
- Williams, F.X. 1947. Aculeate wasps of Fiji. *Occasional Papers of the Bernice P. Bishop Museum* 18 (21): 317–336.
- Yamane, S., J. Gusenleitner and A.S. Menke. 1996. *Pachodynerus nasidens* (Latreille) (Hymenoptera, Vespoidea), an adventive potter wasp new to Japan. *Species Diversity* 1 (2): 93–97.
- Yasumatsu, K. 1945. Vespoidea of Micronesia 2 (Hymenoptera). *Mushi* 16: 35–45.
- Zavattari, E. 1912. Materialien für eine Monographie der Neotropischen Eumeniden. *Archiv für Naturgeschichte* 78A (4): 1–272.
- Zirngiebl, L. 1953. Zur Wespen-Fauna der Pfalz. *Mitteilungen Pollichia* (3) 1: 160–179.